



Title	A characterization of some partial geometric spaces
Author(s)	Kimura, Hiroshi; Ohmori, Hiroyuki
Citation	Osaka Journal of Mathematics. 1985, 22(3), p. 401-410
Version Type	VoR
URL	<a href="https://doi.org/10.18910/7131">https://doi.org/10.18910/7131</a>
rights	
Note	

*The University of Osaka Institutional Knowledge Archive : OUKA*

<https://ir.library.osaka-u.ac.jp/>

The University of Osaka

Kimura, H. and Ohmori, H.  
Osaka J. Math.  
22 (1985), 401-410

## A CHARACTERIZATION OF SOME PARTIAL GEOMETRIC SPACES

Dedicated to Professor Hirosi Nagao on his 60th birthday

HIROSHI KIMURA AND HIROYUKI OHMORI

(Received April 17, 1984)

### 1. Introduction

A partial geometric space  $S$  of dimension  $m \geq 2$  defined in [2, 6] consists of the sets  $\{A_i\}_{i=-1}^m$  and the set  $T$  such that the following eight axioms are satisfied:

- (1)  $A_i \cap A_j = \emptyset$  whenever  $i \neq j$  and  $-1 \leq i, j \leq m$ .
- (2)  $|A_{-1}| = |A_m| = 1$ .
- (3)  $T \subset \prod_{i=-1}^m A_i$ .

The elements of  $A_i$ ,  $-1 \leq i \leq m$ , are called  $i$  elements of  $S$ . The elements of  $T$  are called flags of  $S$ . There is a property called incidence which is a relation between the elements of  $S$  based on the flags.

(4) For each  $i$  element  $x_i$  there is a flag  $(t_{-1}, \dots, t_m) \in T$  such that  $x_i = t_i$ , where  $-1 \leq i \leq m$ .

(5) Whenever  $(y_{-1}, \dots, y_m) \in T$  and  $(z_{-1}, \dots, z_m) \in T$  and  $y_k = z_k$  for some  $k$ ,  $-1 \leq k \leq m$ , then there exists a flag  $(t_{-1}, \dots, t_m) \in T$ , where  $t_i = y_i$  for  $-1 \leq i \leq k$ , and  $t_j = z_j$  for  $k \leq j \leq m$ .

(6) If  $x_i \in A_i$  and  $x_j \in A_j$ , then  $x_i$  and  $x_j$  have an  $l$  intersection  $x_l \in A_l$  and an  $s$  join  $x_s \in A_s$ . Here  $x_i$  and  $x_j$  are said to have an  $l$  intersection  $x_l$  ( $s$  join  $x_s$ ), where  $-1 \leq l \leq \min\{i, j\}$  ( $\max\{i, j\} \leq s \leq m$ ) if and only if  $x_l$  ( $x_s$ ) is incident with  $x_i$  and  $x_j$  such that whenever  $x_n$  is an  $n$  element of  $S$  for  $-1 \leq n \leq \min\{i, j\}$  ( $\max\{i, j\} \leq n \leq m$ ) which is incident with  $x_i$  and  $x_j$ , then  $x_n$  is incident with  $x_l$  ( $x_s$ ) and  $-1 \leq n \leq l$  ( $s \leq n \leq m$ ). By the definition,  $x_i$  and  $x_j$  have unique intersection and unique join.

(7) If  $x_{i-1} \in A_{i-1}$  and  $x_{i+1} \in A_{i+1}$  are incident, then there are  $k(i)$   $i$  elements which are incident with  $x_{i-1}$  and  $x_{i+1}$ , where  $2 \leq k(i) < \infty$ , for  $0 \leq i \leq m-1$ . The number  $k(i)$  is independent of the choice of  $x_{i-1}$  and  $x_{i+1}$ , and depends only on  $i$ .  $k(0), k(1), \dots, k(m-1)$  are called the configuration parameters of  $S$ .

(8) Let  $m \geq 2$ . If  $x_i \in A_i$  and  $x_{i+1} \in A_{i+1}$  have an  $(i-1)$  intersection  $x_{i-1}$  and an  $s$  join  $x_s$ , where  $0 \leq i \leq m-2$  and  $i+2 \leq s \leq m$ , then there are  $t(i, s, k)$   $i$

elements  $y_i$ , which are incident with  $x_{i-1}$  and  $x_{i+1}$  such that  $y_i$  and  $x_i$  have an  $(i+k)$  join for  $1 \leq k \leq s-i-1$ . Also  $\sum_{k=1}^{s-i-1} t(i, s, k) \geq 1$  for  $0 \leq i \leq m-2$  and  $i+2 \leq s \leq m$ . The numbers  $t(i, s, k)$  are called the geometric parameters of  $S$ .

The concept of a partial geometric space of dimension  $m$  is an extension of the concept of a partial geometry introduced by R.C. Bose [1]. A partial geometry of dimension three introduced by R. Lasker and J. Dunbar [5] is called an L.D. partial geometric space of dimension three in [6].

We have two examples of partial geometric spaces of dimension  $m$ .

**EXAMPLE 1** [6]. Let  $A$  be a set consisting  $(m+1)$  distinct symbols, where  $m \geq 2$ . Let  $A_{-1} = \{\phi\}$ . For  $0 \leq j \leq m$ ,  $A_j = \{B \subset A \mid |B| = j+1\}$ . Note that  $A_m = \{A\}$ . Let  $T = \{(t_{-1}, \dots, t_m) \in \prod_{i=-1}^m A_i \mid t_i \subset t_{i+1} \text{ for } -1 \leq i \leq m-1\}$ . Then  $S_1 = (\{A_i\}_{i=-1}^m, T)$  is a partial geometric space of dimension  $m$ . The configuration parameters are  $k(i) = 2$  for  $0 \leq i \leq m-1$ . The geometric parameters are  $t(i, i+2, 1) = 2$  for  $0 \leq i \leq m-2$  and the rest geometric parameters need not be defined.

**EXAMPLE 2.** Let  $PG(m, q)$  be the finite projective geometry of dimension  $m$  and of order  $q$ , where  $m \geq 2$  and  $q$  is a prime power. Let  $A_{-1} = \{\phi\}$ . For  $0 \leq j \leq m$ ,  $A_j = \{B \mid B \text{ is a } j \text{ dimensional subspace of } PG(m, q)\}$ . Let  $T = \{(t_{-1}, \dots, t_m) \in \prod_{i=-1}^m A_i \mid t_i \subset t_{i+1} \text{ for } 0 \leq i \leq m-1\}$ . Then  $S_2 = (\{A_i\}_{i=-1}^m, T)$  is a partial geometric space of dimension  $m$ . The configuration parameters are  $k(i) = q+1$  for  $0 \leq i \leq m-1$ . The geometric parameters are  $t(i, i+2, 1) = q+1$  for  $0 \leq i \leq m-2$  and the rest geometric parameters need not be defined.

Two partial geometric spaces  $S_1$  and  $S_2$  of dimension  $m$  have common property:

$$(\#) \quad \begin{cases} \text{(i) } k(i) \text{ is constant for } 0 \leq i \leq m-1 \\ \text{(ii) } t(i, s, k) = k(i) \text{ for } 0 \leq i \leq m-2, \text{ where } s = i+2 \text{ and } k = 1, \\ \text{and the rest geometric parameters need not be defined.} \end{cases}$$

From (ii) of the property, we note that for any  $i$  element and  $i+1$  element which have an  $(i-1)$  intersection and are not incident, they have an  $(i+2)$  join.

In section 2, we shall prove the following theorem.

**Theorem.** Let  $S = (\{A_i\}_{i=-1}^m, T)$  be a partial geometric space of dimension  $m \geq 2$  satisfying property  $(\#)$ . Then  $S = S_1$  if  $k(i) = 2$ , and  $S = S_2$  if  $k(i) = \alpha + 1 > 2$  and  $m \geq 3$ .

In section 3, we shall give an another example of partial geometric space of dimension  $m \geq 3$ .

## 2. Proof of Theorem

Let  $S = (\{A_i\}_{i=-1}^m, T)$  be a partial geometric space of dimension  $m \geq 2$ . Let  $x_i \in A_i$  and  $x_j \in A_j$ , where  $-1 \leq i, j \leq m$ .  $x_i$  is said to be incident with  $x_j$  if and only if there exists a flag  $(t_{-1}, \dots, t_m) \in T$  such that  $x_i = t_i$  and  $x_j = t_j$ . Let  $x_j \in A_j$  and  $x_k \in A_k$  such that  $x_j$  and  $x_k$  are incident, where  $-1 \leq j < k \leq m$ .  $\phi(i, x_j, x_k)$  is the number of  $i$  elements of  $S$  which are incident with  $x_j$  and  $x_k$ , where  $-1 \leq i \leq m$ . The number  $\phi(i, x_j, x_k)$  is a finite positive integer which is independent of the choice of the  $j$  element  $x_j$  and the  $k$  element  $x_k$  [2]. Therefore put  $\phi(i, j, k) = \phi(i, x_j, x_k)$ .

From now on in this section, we assume that  $S$  satisfies the property (#).

**Lemma 1.** *Let  $x_i$  and  $y_i$  be two distinct  $i$  elements such that they have an  $(i-1)$  intersection  $x_{i-1}$  for  $0 \leq i \leq m-1$ . Then  $x_i$  and  $y_i$  have an  $(i+1)$  join.*

Proof. Let  $x_{i+1}$  be a join of  $x_i$  and  $y_i$ , where  $i > 1$ . Then there exists an  $(i+1)$  element  $y_{i+1}$  which is incident with  $x_{i-1}$  and  $x_{i+1}$  and is not incident with  $x_i$ . From the property (#), we have  $i=2$  and there are  $k(i)$   $i$  elements  $z_i$ , which are incident with  $x_{i-1}$  and  $y_{i+1}$ , such that  $z_i$  and  $x_i$  have an  $(i+1)$  join. Those  $k(i)$   $i$  elements are distinct from  $y_i$ . Consequently, there are  $(k(i)+1)$   $i$  elements which are incident with  $x_{i-1}$  and  $y_{i+1}$ . This is a contradiction. Therefore  $i=1$ , i.e.  $x_i$  and  $y_i$  have an  $(i+1)$  join.

**Lemma 2.**  $\phi(i, x_{i-1}, x_k) = \phi(i, i-1, k) = k(i)(k(i)-1)^{k-i-1} + (k(i)-1)^{k-i-2} + \dots + (k(i)-1)+1$ , where  $0 \leq i < k \leq m$ , and  $x_{i-1} \in A_{i-1}$  and  $x_k \in A_k$ .

Proof. It shall be proved by induction on  $k-i+1$ , say  $t$ . When  $t=2$ , from the definition,  $\phi(i, x_{i-1}, x_{i+1}) = \phi(i, i-1, i+1) = k(i)$ . Therefore the lemma holds when  $t=2$ . Suppose that  $t > 2$  and assume that the lemma holds whenever  $k-i+1 < t$ , where  $0 \leq i < k \leq m$ , and  $2 < t \leq m+1$ . Let  $x_{i-1}$  be an  $(i-1)$  element and  $x_k$  be a  $k$  element such that  $x_{i-1}$  and  $x_k$  are incident in  $S$ , where  $0 \leq i \leq m-2$ ,  $i+2 \leq k \leq m$  and  $k-i+1=t$ . Count triples  $(x_i, x_{i+1}, x_{i+2})$ , where  $x_l$  ( $i \leq l \leq i+2$ ) is an  $l$  element such that  $x_l$  and  $x_{l+1}$  ( $i-1 \leq l \leq i+1$ ), and  $x_{i+2}$  and  $x_k$  are incident in  $S$ .

Given a fixed  $i$  element  $x_i$  which is incident with  $x_{i-1}$  and  $x_k$ , there are  $(k(i)(k(i)-1)^{k-i-2} + (k(i)-1)^{k-i-3} + \dots + (k(i)-1)+1)$   $(i+1)$  elements  $x_{i+1}$ , which are incident with  $x_i$  and  $x_k$ , by the induction hypothesis.

Similarly, given a fixed pair  $(x_i, x_{i+1})$ , where  $x_{i-1}$  and  $x_i$ ,  $x_i$  and  $x_{i+1}$ , and  $x_{i+1}$  and  $x_k$  are incident in  $S$ , there are  $(k(i)(k(i)-1)^{k-i-3} + (k(i)-1)^{k-i-4} + \dots + (k(i)-1)+1)$   $(i+2)$  elements, which are incident with both  $x_{i+1}$  and  $x_k$ .

Therefore the number of triples is

$$(k(i)(k(i)-1)^{k-i-2} + (k(i)-1)^{k-i-3} + \dots + (k(i)-1)+1) \times \\ (k(i)(k(i)-1)^{k-i-3} + (k(i)-1)^{k-i-4} + \dots + (k(i)-1)+1) \phi(i, x_{i-1}, x_k).$$

On the other hand, count pairs  $(x_i, x_{i+1})$ , where  $x_l$  ( $i \leq l \leq i+1$ ) is an  $l$  element such that  $x_{i'}$  and  $x_{i'+1}$  ( $i-1 \leq l' \leq i$ ) are incident in  $S$ . Let  $x_i$  and  $y_i$  be distinct two  $i$  elements which are incident with  $x_{i-1}$  and  $x_k$ , then  $x_i$  and  $y_i$  have an  $(i+1)$  join, say  $y_{i+1}$ . For  $y_{i+1}$ , there are  $\binom{k(i)}{2}$  pairs  $(x'_i, y'_i)$  such that an  $i$  elements  $x'_i$  and  $y'_i$  have an  $(i+1)$  join  $y_{i+1}$  and an intersection  $x_{i-1}$ , by the definition of  $k(i)$ . Consequently there are  $\binom{\phi(i, x_{i-1}, x_k)}{2} k(i)/\binom{k(i)}{2}$  pairs  $(x_i, x_{i+1})$  such that  $x_{i-1}$  and  $x_i$ ,  $x_i$  and  $x_{i+1}$ , and  $x_{i+1}$  and  $x_k$  are incident in  $S$ . The contribution to triples of such a pair  $(x_i, x_{i+1})$  is  $(k(i)(k(i)-1)^{k-i-3} + k(i)-1)^{k-i-4} + \dots + (k(i)-1)+1$  by the induction hypothesis. Therefore we get

$$\begin{aligned} & (k(i)(k(i)-1)^{k-i-2} + (k(i)-1)^{k-i-3} + \dots + (k(i)-1)+1) \times \\ & (k(i)(k(i)-1)^{k-i-3} + (k(i)-1)^{k-i-4} + \dots + (k(i)-1)+1) \phi(i, x_{i-1}, x_k) \\ & = (k(i)(k(i)-1)^{k-i-3} + (k(i)-1)^{k-i-4} + \dots + (k(i)-1)+1) \times \\ & \left( \phi(i, x_{i-1}, x_k) \right) k(i)/\binom{k(i)}{2}. \end{aligned}$$

Consequently we have the lemma.

REMARK. This lemma can be obtained from Theorem 7.1 in [6].

**Lemma 3.** If  $k(i)=2$ ,  $\phi(i, i-1, k)=k-i+1$ , and if  $k(i)=\alpha+1>2$ ,  $\phi(i, i-1, k)=(\alpha^{k-i+1}-1)/(\alpha-1)$ , for  $0 \leq i < k \leq m$ .

Proof. It is obvious from Lemma 2.

**Lemma 4.** If  $k(i)=2$ ,  $\phi(i, j, k)=\binom{k-j}{i-j}$  for  $-1 \leq j < i < k \leq m$ .

Proof. Let  $x_j$  be a  $j$  element and  $x_k$  be a  $k$  element such that  $x_j$  and  $x_k$  be incident in  $S$ . Count  $(k-j+1)$ -tuples  $(x_j, \dots, x_i, \dots, x_k)$ , where  $x_l$  ( $j \leq l \leq k-1$ ) is an  $l$  element such that  $x_l$  and  $x_{l+1}$  are incident. By Lemma 2, there are  $(k-j)$   $(j+1)$  elements  $x_{j+1}$  which are incident with  $x_j$  and  $x_k$ . For such  $x_{j+1}$ , there are  $(k-j-1)$   $(j+2)$  elements which are incident with  $x_{j+1}$  and  $x_k$ , and so on. Consequently there are  $(k-j)!(k-j+1)$ -tuples. On the other hand, given a fixed  $i$  element  $x_i$  which is incident with  $x_j$  and  $x_k$ , there are  $(i-j)!(i-j+1)$ -tuples  $(x_j, \dots, x_i)$  where an  $l$  element  $x_l$  and an  $(l+1)$  element  $x_{l+1}$  are incident ( $j \leq l \leq i-1$ ), and there are  $(k-i)!(k-i+1)$ -tuples  $(x_i, \dots, x_k)$  where an  $l'$  element  $x_{l'}$  and an  $(l'+1)$  element  $x_{l'+1}$  are incident in  $S$ , for  $i \leq l' \leq k-1$ . Therefore we get  $\phi(i, x_j, x_k)(k-i)!(i-j)! = (k-j)!$ . Thus the proof is complete.

**Lemma 5.** If  $k(i)=\alpha+1>2$ ,  $\phi(i, j, k)=\prod_{l=1}^{i-j} (\alpha^{k-j+l}-1)/(\alpha^l-1)$  for  $-1 \leq j < i < k \leq m$ .

Proof. It is similar to the proof of Lemma 4. So, we shall omit a proof.

We note that  $\phi(i, j, k) = \phi(i, -1, j)$  when  $i < j$ , and  $\phi(i, j, k) = \phi(i, k, m)$  when  $k < i$ . So,  $\phi(i, j, k)$  is defined for  $i, j$  and  $k$  such that  $-1 \leq j < k \leq m$ ,  $-1 \leq i \leq m$  and  $i \neq j, k$ .

By the incidence structure in  $S$ , an  $i$  element  $x_i$  can be corresponded to a subset  $b(x_i)$  of  $A_0$  consisting of 0 elements which are incident with  $x_i$ , where  $0 \leq i \leq m$ .

**Lemma 6.** *The above correspondence of  $A_i$  to a family consisting of subsets of  $A_0$  is injective.*

Proof. Assume that  $b(x_i) = b(y_i)$  for an  $i$  element  $y_i$  ( $\neq x_i$ ). Let  $x_l$  be an  $l$  intersection of  $x_i$  and  $y_i$ . Then  $l < i$  and  $b(x_l) \supseteq b(x_i)$ . On the other hand,  $|b(z_j)| = \phi(0, -1, j)$  for every  $j$  element  $z_j$ . This contradicts  $\phi(0, -1, l) < \phi(0, -1, i)$ .

REMARK. Similarly we can prove that  $b(x_i) \neq b(x_j)$  for  $x_i \in A_i$  and  $x_j \in A_j$ , where  $i \neq j$ .

**Lemma 7.** *If  $k(i) = 2$ , then  $S = S_1$ .*

Proof.  $|A_0| = \phi(0, -1, m) = m + 1$ . Since  $\phi(0, -1, i) = i + 1$ , every element of  $A_i$  is a subset of  $A_0$  consisting of  $i + 1$  elements. By Lemma 4 and Lemma 6,  $A_i$  is a family of all subsets of  $A_0$  containing  $i + 1$  elements. By the definition, for  $i < j$ ,  $x_i \in A_i$  and  $x_j \in A_j$  are incident if and only if  $b(x_i) \subset b(x_j)$ . Thus the proof is complete.

Next we assume that  $k(i) = \alpha + 1 \geq 3$  for  $0 \leq i \leq m - 1$ . By Lemma 6, an  $i$  element  $x_i$  is identified with a subset of  $A_0$ .

**Lemma 8.** *A incidence structure  $D = (A_0, A_{m-1})$  is a symmetric 2- $(v, k, \lambda)$  design, where  $v = (\alpha^{m+1} - 1)/(\alpha - 1)$ ,  $k = (\alpha^m - 1)/(\alpha - 1)$  and  $\lambda = (\alpha^{m-1} - 1)/(\alpha - 1)$ .*

Proof. By the definition,  $v = \phi(0, -1, m)$  and  $k = \phi(0, -1, m - 1)$ . Let  $x_0$  and  $y_0$  be two elements of  $A_0$ . Then there exists a 1 element  $x_1$  by Lemma 1 which is a join of  $x_0$  and  $y_0$ . But every element of  $A_{m-1}$  containing  $x_0$  and  $y_0$  has to contain  $x_1$ . Thus we have  $\lambda = \phi(m - 1, 1, m)$ . By Lemma 5, we have the lemma.

Elements of  $A_0$  and elements of  $A_{m-1}$  are called points and blocks in  $D$ , respectively. For  $x_i \in A_i$  and  $y_j \in A_j$ , where  $0 \leq i \leq j \leq m - 1$ , we define  $\langle x_i, y_j \rangle$

be an intersection of all blocks of  $D$  containing  $x_i$  and  $y_j$ . Especially  $\langle x_0, y_0 \rangle$  is called a line spanned by  $x_0$  and  $y_0$ , where  $x_0 \in A_0$  and  $y_0 \in A_0$ .

**Lemma 9.** *Let  $x_1$  and  $y_1$  be two elements of  $A_1$ . Then there is an element of  $A_{m-1}$  which is incident with  $x_1$  and not incident with  $y_1$ .*

Proof. Let  $x_l$  be an  $l$  join of  $x_1$  and  $y_1$ . Then  $l > 1$ . By the property of  $x_l$ , the number of elements of  $A_{m-1}$  which are incident with  $x_1$  and  $y_1$  equals to the number of elements of  $A_{m-1}$  which are incident with  $x_l$ . This number is  $\phi(m-1, l, m)$  which is smaller than  $\phi(m-1, 1, m)$  by Lemma 5. This proves the lemma.

**Lemma 10.**  *$D$  is a design such that its points and blocks are points and hyperplanes of a finite projective geometry  $P$  of dimension  $m$ , respectively.*

Proof. Let  $x_1$  be a 1 join of  $x_0$  and  $y_0$ , where  $x_0, y_0 \in A_0$ . By Lemma 1,  $x_1$  is contained in every block of  $D$  which is incident with  $x_0$  and  $y_0$ . Therefore  $\langle x_0, y_0 \rangle \supseteq x_1$ . If  $\langle x_0, y_0 \rangle \neq x_1$ , then there is an element  $z_0$  of  $\langle x_0, y_0 \rangle$  which is not incident with  $x_1$ . Let  $x_l$  be an  $l$  join of  $z_0$  and  $x_1$ , where  $l > 1$ . Let  $z_1$  be an element of  $A_1$  which is incident with  $x_l$  and  $z_0$ . Then  $z_1 \neq x_1$  and  $z_1$  is contained in all blocks which contain  $x_0$  and  $y_0$ . But by Lemma 9, there exists a block of  $D$  which is incident with  $x_1$  and not incident with  $z_1$ , and hence  $z_1$  is not contained in  $\langle x_0, y_0 \rangle$ . Hence  $\langle x_0, y_0 \rangle = x_1$ . Therefore  $(v-\lambda)/(k-\lambda) = \alpha + 1 = |x_1|$ . By using a result in [4], we have the lemma.

**Lemma 11.** *An  $i$  element  $x_i$  is a subspace of  $P$  of dimension  $i$  for  $1 \leq i \leq m$ .*

Proof. We shall prove the lemma by the induction on  $i$ . By Lemma 10, the case of  $i=1$  is true. Let  $i \geq 2$ . Then there exist elements  $x_{i-1}$  and  $y_{i-1}$  of  $A_{i-1}$ , and an element  $x_{i-2}$  of  $A_{i-2}$  such that they are incident with  $x_i$ , and that  $x_{i-2}$  is incident with  $x_{i-1}$  and  $y_{i-1}$ . By Lemma 6, there exists an element  $y_0$  of  $y_{i-1}$  which is not contained in  $x_{i-1}$ . By the induction hypothesis,  $y_{i-2} = \langle x_{i-2}, y_0 \rangle$  which is a subspace of  $P$  spanned by  $y_0$  and all elements of  $x_{i-2}$ . Therefore we have  $\langle x_{i-1}, y_0 \rangle = \langle x_{i-1}, y_{i-1} \rangle$ . Since  $A_m$  is a projective space and  $x_{i-1}$  is an  $i-1$  dimensional subspace,  $\langle x_{i-1}, y_0 \rangle$  is an  $i$  dimensional subspace, and hence  $|\langle x_{i-1}, y_0 \rangle| = (\alpha^{i+1} - 1)/(i-1)$ . On the other hand, we have  $\langle x_{i-1}, y_{i-1} \rangle \supseteq x_i$ , because  $x_i$  is contained in every elements of  $A_{m-1}$  containing  $x_{i-1}$  and  $y_{i-1}$ . By Lemma 3,  $|x_i| = |\langle x_{i-1}, y_0 \rangle|$ . Therefore we have  $x_i = \langle x_{i-1}, y_0 \rangle$ . Thus the proof is complete.

By Lemma 7 and Lemma 11, a proof of Theorem completes.

### 3. Another example

EXAMPLE 3. Let  $V$  be an  $m$  dimensional vector space over  $GF(2)$  ( $m \geq 3$ ),

and  $H$  the set consisting of all  $m-1$  dimensional subspaces of  $V$ . Put  $A_{-1} = \{\phi\}$ ,  $A_m = \{V - \{0\}\}$  and  $A_i = \{M_i^c \cap \dots \cap M_{m-i}^c \mid M_1 \supseteq M_2 \supseteq \dots \supseteq \bigcap_{u=1}^{m-i} M_u, M_u \in H\}$  for  $0 \leq i \leq m-1$ , where  $M_u^c = V - M_u$ . We say that  $x_i \in A_i$  is incident with  $x_j \in A_j$  if and only if  $x_i \subset x_j$  ( $i \leq j$ ). We shall show that  $S_3 = (\{A_i\}_{i=-1}^m, T)$  is a partial geometric space of dimension  $m$ , where

$$T = \{(x_{-1}, \dots, x_i, \dots, x_j, \dots, x_m) \in \prod_{i=-1}^m A_i \mid x_i \text{ is incident with } x_j \ (-1 \leq i < j \leq m)\}.$$

**Lemma 12.** *For  $x_i \in A_i$ ,  $|x_i| = 2^i$  ( $i \geq 0$ ).*

Proof. Let  $x_i = \bigcap_{u=1}^{m-i} M_u^c$ , then  $\bigcap_{u=1}^{m-i} M_u$  is a subspace of dimension  $i$ . Therefore we have that by the principle of inclusion and exclusion  $|x_i| = 2^m + \sum_{u=1}^{m-i} \binom{m-i}{u} (-1)^u 2^{m-u} = 2^i (2-1)^{m-i} = 2^i$ .

**REMARK A.** Let  $x_0 \in A_0$  and  $M \in H$  ( $x_0 \notin M$ ). Since  $V - \{0\}$  is a projective space,  $M^c$  is an affine space. Thus  $M^c - \{x_0\}$  is a projective space over  $GF(2)$ .

At first, we define the intersection and the join. For  $z_l \in A_l$  ( $0 \leq l \leq m-1$ ), put  $K(z_l) = \{M \in H \mid M^c \supset z_l\}$ , and  $K(z_{-1}) = H$  and  $K(z_m) = \phi$ , where  $z_{-1} \in A_{-1}$  and  $z_m \in A_m$ . Let  $x_i$  and  $y_j$  ( $-1 \leq i, j \leq m$ ) be elements of  $A_i$  and  $A_j$ , respectively. Then a set  $\bigcap_{u=1}^{m-l} L_u^c$  is defined to be an  $l$  intersection of  $x_i$  and  $y_j$  where elements  $L_u$  ( $1 \leq u \leq m-l$ ) of  $K(x_i) \cup K(y_j)$  satisfy  $L_1 \supseteq L_2 \supseteq \dots \supseteq \bigcap_{u=1}^{m-l} L_u$  and  $\bigcap_{u=1}^{m-l} L_u \subset L$  for any element  $L$  of  $K(x_i) \cup K(y_j)$ . We denote  $\bigcap_{u=1}^{m-l} L_u^c$  by  $x_i \wedge y_j$ . We note that if there exists an element  $L_{m-l+1}$  of  $K(x_i) \cup K(y_j)$  such that  $\bigcap_{u=1}^{m-l} L_u = \bigcap_{u=1}^{m-l+1} L_u$  and  $\bigcap_{u=1}^{m-l} L_u^c = \bigcap_{u=1}^{m-l+1} L_u^c$ , then  $x_i$  and  $y_j$  have a  $-1$  intersection. Because let  $\bar{V} = V / L_1 \cap \dots \cap L_{m-l}$  and  $\bar{L}_u = L_u / L_1 \cap \dots \cap L_{m-l}$  ( $1 \leq u \leq m-l$ ). By Lemma 12,  $|\bar{L}_1^c \cap \dots \cap \bar{L}_{m-l}^c| = 1$ , and hence  $\bar{L}_1^c \cap \dots \cap \bar{L}_{m-l+1}^c = \phi$ . This implies  $x_i \wedge y_j = \phi$ .

Next, a set  $\bigcap_{w=1}^{m-s} J_w^c$  is defined to be an  $s$  join of  $x_i$  and  $y_j$  where element  $J_w$  ( $1 \leq w \leq m-s$ ) of  $K(x_i) \cap K(y_j)$  ( $\neq \phi$ ) are satisfy  $J_1 \supseteq J_2 \supseteq \dots \supseteq \bigcap_{w=1}^{m-s} J_w$  and  $\bigcap_{w=1}^{m-s} J_w \subset J$  for any element  $J$  of  $K(x_i) \cap K(y_j)$ . We denote  $\bigcap_{w=1}^{m-s} J_w^c$  or  $V - \{0\}$  by  $x_i \vee y_j$  according to  $K(x_i) \cap K(y_j) \neq \phi$  or  $= \phi$ . It is obvious that the intersection and the join of  $x_i$  and  $y_j$  is well-defined.

By the above paragraph, we have the following lemma.

**Lemma 13.** *Let  $K$  be a subset of  $H$ . Then  $\bigcap_{N \in K} N^c$  is an element of  $A_l$  for some  $l$ .*

**Lemma 14.** Let  $x_i = \bigcap_{w=1}^{m-i} M_w^c$  and  $x_j = \bigcap_{u=1}^{m-j} N_u^c$  be elements of  $A_i$  and  $A_j$ , respectively. If  $x_i \subset x_j$ , then  $\bigcap_{w=1}^{m-i} M_w \subset \bigcap_{u=1}^{m-j} N_u$ .

Proof. Suppose that there exists  $N_z$  ( $1 \leq z \leq m-j$ ) such that  $\bigcap_{w=1}^{m-i} M_w \not\subset N_z$ . Then  $\bigcap_{w=1}^{m-i} M_w \not\supseteq \bigcap_{w=1}^{m-i} M_w \cap N_z$ , so  $x_{i-1} = \bigcap_{w=1}^{m-i} M_w^c \cap N_z^c$  is an element of  $A_{i-1}$ . Hence  $\bigcup_{w=1}^{m-i} M_w \cup N_z \not\supseteq \bigcup_{w=1}^{m-i} M_w$  by Lemma 12. On the other hand, by the hypothesis  $x_i \subset x_j$ ,  $\bigcup_{w=1}^{m-i} M_w \supset \bigcup_{u=1}^{m-j} N_u$ . Hence  $\bigcup_{w=1}^{m-i} M_w \supset N_z$ . This is a contradiction.

**Lemma 15.** Let  $W$  be an  $i$  dimensional subspace of  $V$ . Then  $|\{x_i \in A_i \mid x_i = \bigcap_{w=1}^{m-i} M_w^c, \text{ where } \bigcap_{w=1}^{m-i} M_w = W\}| = 2^{m-i} - 1$ .

Proof. Put  $\bar{V} = V/W$ . By Lemma 12,  $|\bar{M}_1^c \cap \dots \cap \bar{M}_{m-i}^c| = 1$ . Since  $GL(m-i, 2)$  acts transitively on  $\bar{V} - \{\bar{0}\}$ , we have the lemma.

By Lemmas 14 and 15, we have the following:

**Lemma 16.**  $|A_0| = 2^m - 1$  and  $|A_i| = \left( \prod_{u=1}^i \frac{2^{m+1-u} - 1}{2^u - 1} \right) (2^{m-i} - 1)$  for  $m > i > 0$ .

**Lemma 17.** Let  $x_i = \bigcap_{u=1}^{m-i} M_u^c$  be an element of  $A_i$  ( $0 \leq i \leq m-1$ ), then  $|K(x_i)| = 2^{m-i-1}$ .

Proof. Without loss of generality, we may assume  $i=0$ . By Lemma 12, put  $M_1^c \cap \dots \cap M_m^c = \{a\}$ , that is every elements of  $H$  contained in  $\bigcup_{u=1}^m M_u$  does not contain  $\{a\}$ . Since the number of hyperplanes of  $V/\langle a \rangle$  equals  $2^{m-1} - 1$ , the number in the lemma equals  $(2^m - 1) - (2^{m-1} - 1) = 2^{m-1}$ .

**Lemma 18.**  $k(0) = k(m-1) = 2$  and  $k(i) = 3$  for  $0 < i < m-1$ .

Proof. Let  $x_i$  be an element of  $A_i$ . Since  $|A_0| = 2^m - 1$  by Lemma 16 and  $|x_1| = 2$  by Lemma 12, we have  $k(0) = 2$ . For  $k(m-1)$ , consider a factor space. Then we have similarly that  $k(m-1) = 2$ . For  $0 < i < m-1$ , the lemma follows from Remark A and Example 2.

**Lemma 19.** Let  $a$  and  $x_i$  be elements of  $V - \{0\}$  and  $A_i$ , respectively. Assume that there exist elements  $M$  and  $N$  of  $K(x_i)$  such that  $a \in M$  and  $a \notin N$ , where  $i \geq 0$ . Then  $|\{L \in K(x_i) \mid a \in L\}| = |\{L \in K(x_i) \mid a \notin L\}|$ .

Proof. Without loss of generality, we may assume  $\bigcap_{L \in K(x_i)} L = \{0\}$ , that is,  $i=0$ . Put  $X = \{L \in K(x_0) \mid a \in L\}$  and  $Y = K(x_0) - X$ . Let  $y_j = \bigcap_{L \in X} L^c$  and  $z_l = \bigcap_{L \in Y} L^c$ . Since  $\bigcap_{L \in X} L \ni a$ ,  $j > 0$ . Since  $z_l \ni a$ ,  $z_l \neq x_0$ , and hence  $l > 0$ . Since  $|K(x_0)| > |$

$|K(y_j)| = 2^{m-j-1} \geq |X|$  and  $|K(x_0)| > |K(z_l)| = 2^{m-l-1} \geq |Y|$ , we have that  $2^{m-1} = |X| + |Y| \leq 2^{m-j-1} + 2^{m-l-1}$ . Hence  $j=l=1$ . This proves the lemma.

**Lemma 20.** *The geometric parameters are the following:*

(1)  $t(i, i+2, 1) = 3$  for  $2 < i+2 < m$ ,

(2)  $t(i, m, m-1) = 1$  and  $t(i, m, 1) = 2$  for  $0 < i \leq m-2$ ,

(3)  $t(0, 2, 1) = 1$  if  $\langle x_0, x_1 \rangle$  is a subspace of dimension 3 and  $t(0, m, 1) = 2$  if  $x_0 \subset \langle x_1 \rangle$ , where  $x_u$  ( $u=0, 1$ ) are elements of  $A_u$  such that  $x_0$  is not incident with  $x_1$ . The rest geometric parameters need not be defined.

Proof. (1) follows from Example 2 and Remark A. Let  $x_i$  and  $x_{i+1}$  be elements of  $A_i$  and  $A_{i+1}$ , respectively, such that they have an  $(i-1)$  intersection  $x_{i-1}$  and an  $m$  join  $x_m$ . Considering a factor space, we may assume  $i=1$ . Put  $x_0 = \{a\}$ ,  $x_1 = \{a, b\}$  and  $x_2 = \{a, c, d, e\}$  by Lemma 12, where  $a, b, c, d$  and  $e$  are distinct elements of  $V - \{0\}$ . Since  $x_m = x_1 \vee x_2$ , there exist elements  $M$  and  $N$  of  $K(x_1)$  and  $K(x_2)$ , respectively, such that  $M$  does not contain  $a$  and  $b$ , and that  $N$  contains  $b$  and does not contain  $a, c, d$  and  $e$ . Let  $Y = K(x_0) - K(x_1)$ . Then  $|Y| = 2^{m-2}$  by Lemma 17 and  $N (\in H)$  is contained in  $Y$  if and only if  $N$  contains  $b$  and does not contain  $a$ . Put  $y_1 = \bigcap_{N \in Y} N$ , then  $x_0 \subset y_1 \subset x_2$  since  $\bigcap_{N \in Y} N \ni b$ . Thus  $y_1$  is an element of  $A_1$  and  $K(y_1) \cap K(x_1) = \emptyset$ , since  $Y = K(y_1)$ . Therefore  $y_1 \vee x_1$  is contained in  $A_m$  and  $t(1, m, m-1) \geq 1$ . Let  $z_1 = \{a, c\}$  and  $w_1 = \{a, d\}$ . Since  $K(y_1) \cap K(x_1) = \emptyset$  and  $|K(x_i)| = |K(y_1)| = |K(x_0)|/2$ ,  $K(z_1) \cap K(x_1) \neq \emptyset$  and  $K(w_1) \cap K(x_1) \neq \emptyset$ . This implies that there are elements  $M$  and  $N$  of  $K(x_1)$  such that  $c \in M$  and  $c \notin N$ . By Lemma 19,  $|K(x_1) \cap K(z_1)| = |K(x_1)|/2$ , and hence  $x_1 \vee z_1 \in A_2$ . Similarly  $x_1 \vee w_1 \in A_2$ . Therefore  $t(1, m, 1) \geq 2$ . By the definition,  $\sum_{u=1}^{m-1} t(1, m, u) = k(1) = 3$ . This implies (2). Next assume that  $i=0$ . Put  $x_0 = \{a\}$ ,  $x_1 = \{b, c\}$  and let  $x_s = x_0 \vee x_1$ . Since  $|A_1| = \binom{2^m-1}{2}$  by Lemma 16,  $\{a, b\}$  and  $\{a, c\}$  are contained in  $A_1$ . Thus  $t(0, s, 1) = 2$ . If  $\langle a, b \rangle \ni c$ , then  $|H| - 3|\{M \in H \mid a \in M\}| + 2|\{M \in H \mid M \supset \langle a, b \rangle\}| = (2^m-1) - 3(2^{m-1}-1) + 2(2^{m-2}-1) = 0$ . Therefore  $K(x_0) \cap K(x_1) = \emptyset$ , so  $s=m$ . If  $\langle a, b \rangle \ni c$ , then

$$\begin{aligned} & |H| - 3|\{M \in H \mid a \in M \text{ and } b, c \notin M\}| + 3|\{M \in H \mid a, b \in M \text{ and } c \notin M\}| - |\{M \in H \mid a, b, c \in M\}| \\ & = (2^{m-1}-1) - 3(2^{m-1}-1) + 3(2^{m-2}-1) - (2^{m-3}-1) = 2^{m-3}. \end{aligned}$$

Therefore  $|\{M \in H \mid a, b, c \notin M\}| = |K(x_0 \vee x_1)| = 2^{m-3}$  and hence  $x_0 \vee x_1$  is an element of  $A_2$ . This completes a proof of the lemma.

**References**

- [1] R.C. Bose: *Strongly regular graphs, partial geometries and partially balanced designs*, Pacific J. Math. **13** (1963), 389–419.
- [2] R.C. Bose and R. Miskimins: *Partial geometric spaces of  $m$  dimensions*, Algebraic methods in graph theory Vol. I, II (Szeged, 1978), 37–45, Colloq. Math. Soc. Janos Bolyai, 25, North-Holland, Amsterdam, 1981.
- [3] P. Dembowski: Finite geometries, Springer-Verlag, Berlin-New York, 1968.
- [4] P. Dembowski and A. Wagner: *Some characterization of finite projective spaces*, Arch. Math. **11** (1960), 465–469.
- [5] L. Lasker and J. Dunbar: *Partial geometry of dimension three*, J. Combin. Theory **24** (1978), 187–201.
- [6] R. Miskimins: *Partial geometries of dimension  $m$* , Ph. D. Thesis (Colorado State University: Summer, 1978).

Department of Mathematics  
Ehime University  
Matsuyama 790, Japan