



Title	On hypoelliptic operators with multiple characteristics of odd order
Author(s)	Morimoto, Yoshinori
Citation	Osaka Journal of Mathematics. 1983, 20(2), p. 237-255
Version Type	VoR
URL	https://doi.org/10.18910/7262
rights	
Note	

The University of Osaka Institutional Knowledge Archive : OUKA

<https://ir.library.osaka-u.ac.jp/>

The University of Osaka

ON HYPOELLIPTIC OPERATORS WITH MULTIPLE CHARACTERISTICS OF ODD ORDER

YOSHINORI MORIMOTO

(Received January 16, 1981)

Introduction. In the recent paper [4], Hörmander has clarified Egorov's work [3] on sub-elliptic operators, by improving several points. The purpose of the present paper is to show that the method in [4] is applicable to certain pseudodifferential operators with multiple characteristics of odd order. Rubinstein [13], Wenston [15], [16], Popivanov [12], Menikoff [6] and Popivanov-Popov [17] independently treated some class of differential (or pseudodifferential) operators with double (or multiple, see [17]) characteristics satisfying the conditions similar to those given by Nirenberg-Treves [10] for operators of principal type.

It should be noted that, roughly speaking, operators considered in those papers can be reduced micro-locally to $D_{x_1} + ix_1^k D_{x_2}$ (k integer), which was studied by Mizohata [8]. On the other hand, the operator considered in the present paper can not be reduced only to Mizohata type everywhere in the sense of micro-local. At some point it will be reduced even to Egorov type $D_{x_1} + i(x_1^s D_{x_2} + x_1^a x_2^b |D_x|)$, where $|D_x|$ denotes the square root of $D_{x_1}^2 + D_{x_2}^2 + D_{x_3}^2$ and s, a, b are integers.

The plan of this paper is as follows. In Section 1 we state the assumptions and result. In Section 2 we reduce the proof of main theorem to "sub-elliptic estimate" for a localized operator whose symbol has a parameter $0 < \lambda \leq 1$ (see (2.37) and (3.4)). To prove this estimate, in Section 3 we show that we can use the same method as in [4]. The most part of Section 3 is devoted to show that the symbol of the localized operator satisfies inequalities similar to those in [4, Section 4]. In final section we prove the non-hypoellipticity of some operator in order to show the importance of the notion of modified-null-bicharacteristic curve, which is introduced in Section 1.

1. Assumptions and result

We say that $p(x, \xi) \in \mathcal{B}^\infty(R_x^n \times R_\xi^n)$ belongs to \bar{S}^m when $p(x, \xi)$ is positively homogeneous of degree m in $|\xi| \geq 1/2$. (Clearly \bar{S}^m is the subset of $S^m = S_{1,0}^m$. We refer the definition of $S_{1,0}^m$ to Kumano-go [5, p. 50].) For a conic set $U \subset R_x^n \times R_\xi^n$ and $q(x, \xi) \in C^\infty(U)$ with positive homogeneous of degree m in $|\xi| \geq 1/2$

we write $q(x, \xi) \in \bar{S}^m(U)$.

Let $p_1(x, \xi)$ belong to \bar{S}^1 and be real principal type, that is, be real valued and satisfy

$$(1.1) \quad d_{x\xi} p_1(x, \xi) \neq 0 \quad \text{on } \Gamma = p_1^{-1}(0) \cap \{|\xi| \geq 1/2\}.$$

Let l be an odd integer ≥ 3 and let $a(x, \xi) \in \bar{S}^{l-1}$ be complex valued and satisfy

$$(1.2) \quad \operatorname{Re} a \neq 0 \quad \text{on } \Gamma$$

and moreover

$$(1.3) \quad H_{p_1} \operatorname{Re} a = 0 \quad \text{on } \Gamma,$$

where H_{p_1} denotes the Hamilton vector field of p_1 .

Then under certain conditions among p_1 , $\operatorname{Re} a$ and $\operatorname{Im} a$ we shall discuss the hypoellipticity for a pseudodifferential operator L of order l which has the form

$$(1.4) \quad \begin{aligned} L &= P(x, D_x) + A(x, D_x) \quad \text{in } R_x^*, \\ \sigma(P) &= (p_1(x, \xi))^l, \quad \sigma(A) = a(x, \xi). \end{aligned}$$

Here $\sigma(P)$ denotes the symbol of pseudodifferential operator $P(x, D_x)$.

First we assume that

$$(1.5) \quad \begin{aligned} &\text{for any } (x_0, \xi_0) \in \Gamma \text{ there exist a conic neighborhood} \\ &U \text{ of } (x_0, \xi_0) \text{ and } q_0(x, \xi) \in \bar{S}^0(U) \text{ such that} \end{aligned}$$

$$(1.6) \quad H_{p_1} q_0 = 1 \quad \text{in } U$$

$$\text{and for } j=1, \dots, l-2$$

$$(1.7) \quad H_{q_0}^j a = 0 \quad \text{on } \Gamma \cap U.$$

To state the second condition corresponding to (A) in Egorov [3], or (Ψ) in Nirenberg-Treves [10], we define the modified-null-bicharacteristic curve of p_1 through $(x_0, \xi_0) \in \Gamma$ by the curve

$$(1.8) \quad [-T, T] \ni t \mapsto (x(t), \xi(t)),$$

where $(x(t), \xi(t))$ is the solution to

$$(1.9) \quad \begin{aligned} dx/dt &= d_\xi(p_1 + {}^l\sqrt{\operatorname{Re} a}) \\ d\xi/dt &= -d_x(p_1 + {}^l\sqrt{\operatorname{Re} a}), \quad (x(0), \xi(0)) = (x_0, \xi_0), \end{aligned}$$

and ${}^l\sqrt{\operatorname{Re} a}$ denotes a unique real l power root of $\operatorname{Re} a$. It follows from (1.3) that if $(x_0, \xi_0) \in \Gamma$, then $(x(t), \xi(t)) \in \Gamma$. The right hand side of (1.9) are not homogeneous in ξ , so that the behavior of modified-null-bicharacteristic curve is not so. But we can define it on $[-T, T]$ for some $T > 0$ uniformly if $(x_0, \xi_0) \in \Gamma$ varies in a compact conic set ($|\xi_0| > 1$), because $d_\xi {}^l\sqrt{\operatorname{Re} a}$ and $|\xi|^{-1} d_x {}^l\sqrt{\operatorname{Re} a}$

are $\mathcal{O}(|\xi|^{-1/l})$. Second condition is that

$$(1.10) \quad \begin{cases} \text{for any } (x_0, \xi_0) \in \Gamma \text{ with } |\xi_0| \text{ sufficiently large,} \\ \text{if } \operatorname{Im} a(x(t_0), \xi(t_0)) > 0 \text{ for some } t_0 \in [-T, T], \\ \text{then } \operatorname{Im} a(x(t), \xi(t)) \geq 0 \text{ for all } t \in (t_0, T], \\ \text{where } (x(t), \xi(t)) \text{ is the modified-null-bicharacteristic} \\ \text{curve of } p_1 \text{ through } (x_0, \xi_0). \end{cases}$$

For a multi-index $I = (i_1, \dots, i_k)$ whose components i_j are 1 or 2, we use the following notations: $|I| = k$, $b(I) =$ "the number of j such that $i_j = 2$ ", $\mathcal{J} = \{I; b(I) \leq l-1\}$, $\mu(I) = (l-1)|I|/(l-b(I))$. For a $\mu > 0$ we put $\mathcal{J}_\mu = \{I \in \mathcal{J}; \mu(I) \leq \mu\}$. Set $\mathcal{Q}_0 = \{\mu(I); I \in \mathcal{J}\}$. Then we can write $\mathcal{Q}_0 = \{\mu_j; j=1, 2, \dots\}$ with an appropriate increasing sequence of rational numbers. For any $(x, \xi) \in \Gamma$, $\mu(x, \xi)$ denotes the rational $\mu_j \in \mathcal{Q}_0$ such that $p_I(x, \xi) \neq 0$ for some $I \in \mathcal{J}_{\mu_j}$ and $p_I(x, \xi) = 0$ for any $I \in \mathcal{J}_{\mu_{j-1}}$, where

$$(1.11) \quad p_I(x, \xi) = H_{i_1} H_{i_2} \cdots H_{i_{k-2}} p_{i_k}(x, \xi),$$

$H_1 = H_{p_1}$, $H_2 = H_{p_2}$ and $p_2(x, \xi) = \operatorname{Im} a(x, \xi)$. The third condition corresponding to (B) in Egorov [3] is that

$$(1.12) \quad \begin{cases} \text{for all } (x, \xi) \in \Gamma \text{ there exists some } \mu \in \mathcal{Q}_0 \text{ such} \\ \text{that } \mu(x, \xi) \leq \mu < \infty. \end{cases}$$

In what follows we denote the norm in Sobolev space H^s by $\|\cdot\|_s$. We write $\|\cdot\| = \|\cdot\|_0$.

Theorem 1.1. *Let (1.1)–(1.3) hold and let $L(x, D_x)$ in (1.4) satisfy (1.5)–(1.7), (1.10) and (1.12). Then for any compact set K of \mathbb{R}^n there exists a constant C_K such that*

$$(1.13) \quad \|u\|_{l-2+\sigma'} + \|P'u\|_\sigma \leq C_K(\|Lu\| + \|u\|), \quad u \in C_0^\infty(K),$$

where $\sigma(P') = (p_1(x, \xi))^{l-1}$, $\sigma = (l-1)/l\mu$, $\sigma' = \sigma + 1/l$.

REMARK 1. It follows from (1.13) that P is hypoelliptic. See Oleinik-Radkevich [11] and Morimoto [9, Theorem 2.2]. Theorem 2.2 of [9] is stated only for differential operators but its proof is also applicable to pseudo-differential operators.

REMARK 2. In differential operators we have the following example:

$$D_{x_1}^5 + D_{x_2}^4 + D_{x_3}^4 + i(x_1^2 x_2 (D_{x_2}^4 + D_{x_3}^4) + x_1^3 D_{x_2}^2 (D_{x_2}^2 + D_{x_3}^2)) \text{ in } \mathbb{R}_x^3.$$

All conditions of Theorem 1.1 are satisfied. Specially, the condition (1.10) is satisfied at $(0, \xi_0)$ with $\xi_0 = (0, 0, \xi_{03})$ as follows; the sign of $\operatorname{Im} a$ changes from —

to $+$ along the modified-null-bicharacteristic curve of p_1 through $(0, \xi_0)$. At the point $(0, \xi_0)$ this example must be reduced to Egorov type. Note that Egorov's operator of principal type is not the differential operator but the pseudodifferential operator. (See the introduction of [3].) For this example we have $\sigma=1/25$.

REMARK 3. The condition (1.10) is delicate and necessary in general for the hypoellipticity of P . Indeed, the operator with replaced $\operatorname{Re} a = \xi_2^4 + \xi_3^4$ in the above example by $(\xi_2 - \xi_3)^4 + \xi_3^4$ satisfies all conditions except (1.10), which is violated at $(0, \xi_0)$ with $\xi_0 = (0, 0, \xi_{03})$ and $\xi_{03} > 0$. Furthermore, we have

Proposition 1.2. *Differential operator \tilde{L}*

$$(1.14) \quad \tilde{L} = D_{x_1}^5 + (D_{x_2} - D_{x_4})^4 + D_{x_3}^4 \\ + i(x_1^2 x_2 (D_{x_2}^4 + D_{x_3}^4) + x_1^3 D_{x_2}^2 (D_{x_2}^2 + D_{x_3}^2)) \text{ in } R_x^3,$$

is not hypoelliptic at the origin.

2. Reduction to localized operator

Let $h(x) \in C_0^\infty(R_x^n)$ be 1 for $|x| < 1/2$ and vanish for $|x| > 1$. Set $h_\varepsilon(x) = h(x/\varepsilon)$ for a small $\varepsilon > 0$. For a $f(x, \xi) \in \bar{S}^m$ and $\gamma = (x_0, \bar{\xi}_0) \in R^n \times S^{n-1}$ we introduce a pseudodifferential operator $F_{\gamma, \varepsilon, \lambda}$ with a parameter $0 < \lambda \leq 1$ and a small $\varepsilon > 0$ as follows:

$$(2.1) \quad F_{\gamma, \varepsilon, \lambda}(y, D_y)v \\ = \lambda^{-2m} \int e^{iy\eta} h_\varepsilon(\lambda y) f(x_0 + \lambda y, \bar{\xi}_0 + \lambda \eta) h_\varepsilon(\lambda \eta) \hat{v}(\eta) d\eta, \\ v \in S_y, \quad d\eta = (2\pi)^{-n} d\eta,$$

where \hat{v} denotes the Fourier transform of v . Obviously, for a fixed $\varepsilon > 0$, $\{\lambda^{2m} \sigma(F_{\gamma, \varepsilon, \lambda})(y, \eta); 0 < \lambda \leq 1\}$ is a bounded set of $S_{0,0}^0$. Furthermore we obtain for a sufficiently small $\varepsilon > 0$

$$(2.2) \quad (F_{\gamma, \varepsilon, \lambda} v)(\lambda^{-1}(x - x_0)) \\ = e^{-i\lambda^{-2}x \cdot \bar{\xi}_0} h_\varepsilon(x - x_0) f(x, D_x) h_\varepsilon(\lambda^2 D_x - \bar{\xi}_0) u(x),$$

where $\hat{u}(\xi) = \lambda^n \hat{v}(\lambda(\xi - \lambda^{-2} \bar{\xi}_0)) \exp(i x_0(\lambda^{-2} \bar{\xi}_0 - \xi))$.

Lemma 2.1. *For any compact set $K \subset R^n$ there exists a constant C_K such that (1.13) holds if and only if for any $\gamma = (x_0, \bar{\xi}_0) \in R^n \times S^{n-1}$ one can find positives $\varepsilon_i = \varepsilon_{i, \gamma}$ ($i = 1, 2, 3$, $\varepsilon_1 < \varepsilon_2 < \varepsilon_3$) and a constant C_γ so that for any $0 < \lambda \leq 1$ the following estimate holds;*

$$(2.3) \quad \lambda^{-2(l-2+\sigma')} \|H_{\varepsilon_1, \lambda} v\| + \lambda^{-2\sigma} \|P'_{\gamma, \varepsilon_1, \lambda} v\| \\ \leq C_\gamma (\|P_{\gamma, \varepsilon_2, \lambda} + A_{\gamma, \varepsilon_2, \lambda} v\| + \|P'_{\gamma, \varepsilon_3, \lambda} v\| \\ + \lambda^{-2(l-2+\sigma)} \|v\|), \quad v \in S_y,$$

where $H_{\varepsilon,\lambda}, P_{\gamma,\varepsilon,\lambda}, P'_{\gamma,\varepsilon,\lambda}$ and $A_{\gamma,\varepsilon,\lambda}$ are defined by (2.1) with f replaced by $1, p_1', p_1'^{-1}$ and a , respectively.

Proof. In view of (2.2) it is not difficult to see the necessity of (2.3). We only show the sufficiency. The proof is the same way as in [4, p. 143], except the appearance of the second term in left hand side of (1.13) or (2.3). Note that for any $\xi_0 (|\xi_0|=1)$ and any small $\varepsilon>0$ and any real s

$$(2.4) \quad \begin{aligned} C^{-1} \|h_\varepsilon(\lambda D_y) v\| &\leq \|(\xi_0 + \lambda D_y)^s h_\varepsilon(\lambda D_y) v\| \\ &\leq C \|h_\varepsilon(\lambda D_y) v\|, \quad v \in S_y, \end{aligned}$$

holds for some $C=C_{s,\varepsilon}$ since $C^{-1} \leq |\xi_0 + \xi|^s \leq C$ on $\text{supp } h_\varepsilon(\xi)$. Substituting $\hat{v}(\eta) = h_{\varepsilon_0}(\lambda \eta) \hat{u}(\lambda^{-1} \eta + \lambda^{-2} \xi_0) \exp(i x_0 (\lambda^{-1} \eta + \lambda^{-2} \xi_0))$ for $u \in C_0^\infty(K)$ and some $\varepsilon_0 > \varepsilon_3$ into (2.3), we obtain by means of (2.2) and (2.4)

$$\begin{aligned} &\|h_1(x-x_0) h_1(\lambda^2 D_x - \xi_0) |D_x|^{t-2+\sigma'} u\| \\ &\quad + \|h_1(x-x_0) P'(x, D_x) |D_x|^\sigma h_1(\lambda^2 D_x - \xi_0) u\| \\ &\leq C (\|h_2(x-x_0) (P(x, D_x) + A(x, D_x)) h_2(\lambda^2 D_x - \xi_0) u\| \\ &\quad + \|h_3(x-x_0) P'(x, D_x) h_3(\lambda^2 D_x - \xi_0) u\| \\ &\quad + \|h_0(\lambda^2 D_x - \xi_0) |D_x|^{t-2+\sigma} u\|), \end{aligned}$$

where $h_j = h_{\varepsilon_j}$. Here we used the fact that, for a fixed $\varepsilon>0$, $\{\lambda^{-2} [h_\varepsilon(\lambda y), h_\varepsilon(\lambda D_y)]$; $0 < \lambda \leq 1\}$ is a bounded set of $S_{0,0}^0$. Since $[P, h_2(\lambda^2 D_x - \xi_0)]$ can be estimated by the second term of the left hand side, the proof is completed from the following proposition and usual finite covering argument over $K \times S^{n-1}$.

Proposition 2.2. Let $h(\xi) \in C_0^\infty(R^n)$ be 1 in a neighborhood of 0 and let ξ_0 belong to S^{n-1} . Then one can find some $\psi_j(\xi) \in \tilde{S}^0$ ($j=1,2$) such that

(2.5) $\psi_j(\xi_0) \neq 0$, $\text{supp } \psi_j \subset \text{some conic neighborhood of } \xi_0$ and for any N we have for some constant $C>0$

$$\begin{aligned} &C^{-1} \|\psi_1(D_x) u\|^2 \\ &\leq \int_0^1 \|h(\lambda^2 D_x - \xi_0) u\|^2 / \lambda \, d\lambda + \|u\|_{-N} \\ &\leq C (\|\psi_2(D_x) u\|^2 + \|u\|_{-N}), \quad u \in \mathcal{S}_x \end{aligned}$$

Proof. Put $r = |\xi|$, $\theta = \xi / |\xi|$. Then

$$\begin{aligned} &\int_0^1 \|h(\lambda^2 D_x - \xi_0) u\|^2 / \lambda \, d\lambda \\ &= \int d\theta \int_0^1 d\lambda \int_0^\infty h(\lambda^2 r \theta - \xi_0)^2 |\hat{u}(r\theta)|^2 / \lambda \, dr. \end{aligned}$$

It is easy to see that $\text{supp } h(\lambda^2 r \theta - \xi_0)$ is evaluated from above and below by

$$\{(\theta, r, \lambda); \theta \in \text{supp } \psi_j \cap S^{n-1} \text{ and } C_j^{-1} \leq r\lambda^2 \leq C_j\}$$

for some $\psi_j \in \bar{S}^0$ satisfying (2.5) and some $C_j > 0$ ($j=1, 2$). Therefore the integral is bounded by constant times

$$\int \psi_j^2(\theta) d\theta \int_{1/C}^{\infty} |\dot{u}(r\theta)|^2 dr \int_{(Cr)^{-1/2}}^{(Cr)^{1/2}} d\lambda,$$

where we used $(r/C)^{1/2} < 1/\lambda < (Cr)^{1/2}$ and $C = C_j$. This gives the desired estimate.

REMARK. The content of this proposition is briefly stated in [4, p. 143].

Since (2.3) is valid for $\gamma \notin \Gamma$, in view of Lemma 2.1 we now fix a $\gamma \in \Gamma$. Let a function $f_\lambda(y, \eta) \in \mathcal{B}^\infty(R_y^n \times R_\eta^n)$ with a parameter $0 < \lambda \leq 1$ satisfy

$$(2.6) \quad |\partial_y^\alpha \partial_\eta^\beta f_\lambda(y, \eta)| \leq C_{\alpha\beta} \lambda^{|\alpha+\beta|}$$

for any α, β , where $C_{\alpha\beta}$ is a constant independent of λ . Define a pseudodifferential operator $F_\lambda(y, D_y)$ by

$$(2.7) \quad F_\lambda v = \int e^{iy\eta} f_\lambda(y, \eta) \hat{v}(\eta) d\eta, \quad v \in \mathcal{S}.$$

If $f_\lambda(y, \eta)$ equals $f(\lambda y, \lambda \eta)$ for some $f(x, \xi) \in \mathcal{B}^\infty(R_x^n \times R_\xi^n)$, then we say that the operator $F_\lambda(y, D_y)$ has an original symbol $f(x, \xi)$. Under this notation, (2.3) for a fixed γ becomes

$$(2.8) \quad \begin{aligned} & \lambda^{-2(l-2+\sigma')} \|H_{\varepsilon_1, \lambda} v\| + \lambda^{-2(l-1+\sigma)} \|P'_{\varepsilon_1, \lambda} v\| \\ & \leq C (\|(\lambda^{-2l} P_{\varepsilon_2, \lambda} + \lambda^{-2(l-1)} A_{\varepsilon_2, \lambda}) v\| \\ & \quad + \lambda^{-2(l-1)} \|P'_{\varepsilon_3, \lambda} v\| + \lambda^{-2(l-2+\sigma)} \|v\|), \quad v \in \mathcal{S}, \end{aligned}$$

where the original symbols of $H_{\varepsilon, \lambda}$, $P_{\varepsilon, \lambda}$, $P'_{\varepsilon, \lambda}$ and $A_{\varepsilon, \lambda}$ are $h_\varepsilon(x, \xi) = h_\varepsilon(x) h_\varepsilon(\xi)$, $h_\varepsilon(p_{1, \gamma})^l$, $h_\varepsilon(p_{1, \gamma})^{l-1}$ and $h_\varepsilon a_\gamma$ respectively. Here $p_{1, \gamma}(x, \xi) = p_1(x + x_0, \xi + \xi_0)$, $a_\gamma(x, \xi) = a(x + x_0, \xi + \xi_0)$.

Lemma 2.3. *If (2.8) is valid and \mathcal{X} is a C^∞ canonical transformation keeping 0 fixed which is defined near 0, then (2.8) remains valid with some other ε_j and C if $P_{\varepsilon, \lambda}$, $A_{\varepsilon, \lambda}$ and $P'_{\varepsilon, \lambda}$ are replaced by $\tilde{P}_{\varepsilon, \lambda}$, $\tilde{A}_{\varepsilon, \lambda}$ and $\tilde{P}'_{\varepsilon, \lambda}$ respectively, whose original symbols are $h_\varepsilon(p_{1, \gamma} \circ \mathcal{X})^l$, $h_\varepsilon(a_\gamma \circ \mathcal{X})$ and $h_\varepsilon(p_{1, \gamma} \circ \mathcal{X})^{l-1}$, respectively.*

As pointed out in [4, the proof of Lemma 3.2] it suffices to prove the lemma when \mathcal{X} has a generating function $S(x, \xi)$, that is, $\mathcal{X}; (x, d_x S(x, \xi)) \mapsto (d_\xi S(x, \xi), \xi)$. The proof is based on several propositions on Fourier integral operators with phase function $S_\lambda(y, \eta) = \lambda^{-2} S(\lambda y, \lambda \eta)$.

DEFINITION 2.4. For any $f_\lambda(y, \eta) \in \mathcal{B}^\infty$ with (2.6) and for any $S(x, \xi) \in C^\infty(R_x^n \times R_\xi^n)$ satisfying

$$(2.9) \quad \det \partial_x \partial_{\xi} S(0, 0) \neq 0, \quad (d_x S(0, 0), d_{\xi} S(0, 0)) = (0, 0)$$

we define the Fourier integral operator $F_{s_{\lambda}}(y, D_y)$ with a parameter $0 < \lambda \leq 1$ by

$$(2.10) \quad F_{s_{\lambda}} v = \int e^{i s_{\lambda}(y, \eta)} k_{\lambda}(y, \eta) f_{\lambda}(y, \eta) \hat{v}(\eta) d\eta, \quad v \in \mathcal{S}$$

where $S_{\lambda}(y, \eta) = \lambda^{-2} S(\lambda y, \lambda \eta)$ and $k_{\lambda}(y, \eta) = k(\lambda y, \lambda \eta)$. Here we assume that $k(x, \xi) \in C_0^{\infty}$ is 1 in a neighborhood of 0 and $\det \partial_x \partial_{\xi} S(x, \xi) \neq 0$ on $\text{supp } k$. We define the conjugate Fourier integral operator $F_{s_{\lambda}}^*$ with a parameter $0 < \lambda \leq 1$ by

$$(2.11) \quad F_{s_{\lambda}}^*(y, D_y) v = \iint e^{i(y\eta - s_{\lambda}(\tilde{y}, \eta))} k_{\lambda}(\tilde{y}, \eta) f_{\lambda}(\tilde{y}, \eta) v(\tilde{y}) d\tilde{y} d\eta, \quad v \in \mathcal{S},$$

We call $f_{\lambda}(y, \eta)$ the symbol of $F_{s_{\lambda}}(y, D_y)$ ($F_{s_{\lambda}}^*(y, D_y)$), and moreover if $f_{\lambda}(y, \eta) = f(\lambda y, \lambda \eta)$ for some $f \in \mathcal{B}^{\infty}$, then we call $f(x, \xi)$ the original symbol of $F_{s_{\lambda}}(F_{s_{\lambda}}^*)$. We write $F_{s_{\lambda}} = I_{s_{\lambda}}$, $F_{s_{\lambda}}^* = I_{s_{\lambda}}^*$ if $f = 1$.

Put

$$(2.12) \quad \begin{aligned} \tilde{d}_{\xi} S(x, \tilde{\xi}, \xi) &= \int_0^1 d_{\xi} S(x, \tilde{\xi} + \theta(\xi - \tilde{\xi})) d\theta \\ \tilde{d}_x S(\tilde{x}, x, \xi) &= \int_0^1 d_x S(\tilde{x} + \theta(x - \tilde{x}), \xi) d\theta \end{aligned}$$

Put $\tilde{x} = \tilde{d}_{\xi} S(x, \tilde{\xi}, \xi)$ and $\tilde{\xi} = \tilde{d}_x S(\tilde{x}, x, \xi)$. Then the inverses

$$(2.13) \quad x = \phi(\tilde{x}; \tilde{\xi}, \xi) \quad \text{and} \quad \xi = \psi(\tilde{\xi}; \tilde{x}, x)$$

exist, respectively, in a neighborhood of 0 on account of (2.9) if $\text{supp } k$ is sufficiently small.

Proposition 2.5. *If $\text{supp } k$ in (2.10) and (2.11) is sufficiently small, then $I_{s_{\lambda}} I_{s_{\lambda}}^*$ and $I_{s_{\lambda}}^* I_{s_{\lambda}}$ are pseudodifferential operators whose symbols are*

$$(2.14) \quad \int e^{-i\tilde{y}\tilde{\eta}} r_{\lambda}(y, \eta + \tilde{\eta}, y + \tilde{y}) d\tilde{y} d\tilde{\eta}$$

and

$$(2.15) \quad \iint e^{-i\tilde{y}\tilde{\eta}} r_{\lambda}^*(\eta + \tilde{\eta}, y + \tilde{y}, \eta) d\tilde{y} d\tilde{\eta}$$

respectively, where $r_{\lambda}(y, \tilde{\eta}, \tilde{y})$ and $r_{\lambda}^*(\eta, \tilde{y}, \tilde{\eta})$ are given by

$$(2.16) \quad (k(x, \xi) k(\tilde{x}, \tilde{\xi}) | \det \int_0^1 \partial_x \partial_{\xi} S(\tilde{x} + \theta(x - \tilde{x}), \xi) d\theta |^{-1}) \quad \xi = \psi(\tilde{\xi}; \tilde{x}, x)$$

and

$$(2.17) \quad (k(x, \xi) k(x, \tilde{\xi}) | \det \int_0^1 \partial_x \partial_{\xi} S(x, \tilde{\xi} + \theta(\xi - \tilde{\xi})) d\theta |^{-1}) \quad x = \phi(\tilde{x}; \tilde{\xi}, \xi),$$

respectively. Here $(x, \tilde{x}, \xi, \tilde{\xi}) = (\lambda y, \tilde{y}, \eta, \tilde{\eta})$.

Proof is directly calculated by means of the change of variable; $\tilde{\xi} = \tilde{d}_x S(\tilde{x}, x, \xi)$ and $\tilde{x} = \tilde{d}_\xi S(x, \tilde{\xi}, \xi)$ respectively.

Corollary 2.6. *The operators F_{s_λ} and $F_{s_\lambda}^*$ for any f_λ with (2.6) are L_2 -bounded uniformly with respect to $0 < \lambda \leq 1$. If ε is small enough, then $I_{s_\lambda} I_{s_\lambda}^*$ and $I_{s_\lambda}^* I_{s_\lambda}$ are elliptic on $\text{supp } h_\varepsilon(\lambda y, \lambda \eta)$, that is, the estimates*

$$(2.18) \quad \|H_{\varepsilon, \lambda} v\| \leq C_1 (\|I_{s_\lambda} I_{s_\lambda}^* v\| + \lambda^2 \|v\|),$$

$$(2.19) \quad \|H_{\varepsilon, \lambda} v\| \leq C_2 (\|I_{s_\lambda}^* I_{s_\lambda} v\| + \lambda^2 \|v\|), \quad v \in \mathcal{S},$$

hold for some constants C_1 and C_2 .

Proof. The symbols of $F_{s_\lambda}^* F_{s_\lambda}$ and $F_{s_\lambda} F_{s_\lambda}^*$ are given by the versions of (2.14) and (2.15), respectively, which belong to a bounded set of $S_{0,0}^0$ uniformly on account of (2.6). The boundedness of $F_{s_\lambda}^* F_{s_\lambda}$ and $F_{s_\lambda} F_{s_\lambda}^*$ show the first statement. Note that for any $p(x, \tilde{\xi}, \tilde{x}, \xi) \in \mathcal{B}^\infty(R^{4n})$

$$(2.20) \quad O_s - \iint e^{-i\tilde{x}\tilde{\xi}} p(x, \tilde{\xi}, \tilde{x}, \xi) d\tilde{x} d\tilde{\xi} = \sum_{|\alpha| \leq N} p_\alpha(x, 0, 0, \xi) / \alpha! + N \sum_{|\beta| = N} \int_0^1 (1-\theta)^{N-1} \\ O_s - \iint e^{-i\tilde{x}\tilde{\xi}} p_\beta(x, \theta\tilde{\xi}, \tilde{x}, \xi) d\tilde{x} d\tilde{\xi} d\theta / \beta!$$

holds for any positive integer N , where $p_\alpha(x, \tilde{\xi}, \tilde{x}, \xi) = \partial_{\tilde{\xi}}^\alpha D_{\tilde{x}}^\alpha p(x, \tilde{\xi}, \tilde{x}, \xi)$. Here $O_s - \iint$ denotes the oscillatory integral (see [5, p. 42]). Applicatinos of (2.20) to (2.14) and (2.15) yield (2.18) and (2.19), respectively.

Proposition 2.7. *Let $F_\lambda(y, D_y)$ be the pseudodifferential operator with original symbol $f(x, \xi) \in \mathcal{B}^\infty$. Let G_{s_λ} and \tilde{G}_{s_λ} be Fourier integral operators whose original symbols are $f(x, d_x S(x, \xi))$ and $f(d_\xi S(x, \xi), \xi)$, respectively. Then $\lambda^{-2}(F_\lambda I_{s_\lambda} - G_{s_\lambda})$ and $\lambda^{-2}(I_{s_\lambda} F_\lambda - \tilde{G}_{s_\lambda})$ are L_2 -bounded operators uniformly with respect to $0 < \lambda \leq 1$.*

Proof. It is easy to check that the equations

$$F_\lambda I_{s_\lambda} v = \int e^{iS_\lambda(y, \eta)} r_\lambda(y, \eta) \hat{v}(\eta) d\eta$$

and

$$I_{s_\lambda} F_\lambda v = \int e^{iS_\lambda(y, \eta)} \tilde{r}_\lambda(y, \eta) \hat{v}(\eta) d\eta,$$

for

$$(2.21) \quad r_\lambda(y, \eta) = O_s - \iint e^{-i\tilde{y}\tilde{\eta}} f(x, \tilde{\xi} + \tilde{d}_x S(x, x + \tilde{x}, \xi)) k(x + \tilde{x}, \xi) d\tilde{y} d\tilde{\eta}$$

and

$$(2.22) \quad \tilde{r}_\lambda(y, \eta) = O_s - \iint e^{-i\tilde{y}\tilde{\eta}} f(\tilde{x} + \tilde{d}_\xi S(x, \xi + \tilde{\xi}, \xi), \xi) k(x, \xi + \tilde{\xi}) d\tilde{y} d\tilde{\eta},$$

respectively, where $(x, \tilde{x}, \xi, \tilde{\xi}) = \lambda(y, \tilde{y}, \eta, \tilde{\eta})$. Applications of (2.20) to (2.21) and (2.22) complete the proof.

Corollary 2.8. *For $f(x, \xi) \in C_0^\infty$ with support contained a sufficiently small neighborhood of 0, set $\tilde{f}(x, \xi) = (f \circ \mathcal{X})(x, \xi)$, where \mathcal{X} is defined by*

$$(x, d_x S(x, \xi)) \rightarrow (d_\xi S(x, \xi), \xi).$$

Then $\lambda^{-2}(F_\lambda I_{S_\lambda} - I_{S_\lambda} \tilde{F}_\lambda)$ is L_2 -bounded uniformly for $0 < \lambda \leq 1$. Furthermore, if $f(x, \xi) = (p_1(x, \xi))^{l'} h_\varepsilon(x, \xi)$ then for $\tilde{f} = (p_1^{l'} h_\varepsilon) \circ \mathcal{X}$, the estimate

$$(2.23) \quad \|(F_\lambda I_{S_\lambda} - I_{S_\lambda} \tilde{F}_\lambda)v\| \leq C(\lambda^2 \|F_\lambda' I_{S_\lambda} v\| + \lambda^2 \|I_{S_\lambda} \tilde{F}_\lambda' v\|), \quad v \in \mathcal{S},$$

holds for some constant C , where the original symbols of F_λ' and \tilde{F}_λ' are $(p_1)^{l'-1} h_\varepsilon$ and $((p_1)^{l'-1} h_\varepsilon) \circ \mathcal{X}$ for some $\varepsilon' > \varepsilon$.

Proof. The first part follows from Proposition 2.7. The second part is obtained by checking the second terms of the expansions of (2.21) and (2.22).

REMARK. It is clear that $\lambda^{-2}(I_{S_\lambda}^* F_\lambda - F_\lambda I_{S_\lambda}^*)$ is L_2 -bounded operator uniformly for $0 < \lambda \leq 1$. Indeed, this follows from $\|F_\lambda I_{S_\lambda} - I_{S_\lambda} \tilde{F}_\lambda\| = \|I_{S_\lambda}^* F_\lambda^* - \tilde{F}_\lambda^* I_{S_\lambda}^*\|$ and the fact that $\lambda^{-2}(F_\lambda - F_\lambda^*)$ and $\lambda^{-2}(\tilde{F}_\lambda - \tilde{F}_\lambda^*)$ are L_2 -bounded.

Proof of Lemma 2.3. Taking $I_{S_\lambda} v$ as v in (2.8) and noting Corollary 2.8, we obtain (2.8) for operators transformed, by using the fact that for any small $\varepsilon > 0$ the estimate

$$C^{-1} \|H_{\varepsilon', \lambda} v\| \leq \|I_{S_\lambda} H_{\varepsilon, \lambda} v\| + \lambda^2 \|v\| \leq C(\|H_{\varepsilon'', \lambda} v\| + \lambda^2 \|v\|), \quad v \in \mathcal{S},$$

holds for some $0 < \varepsilon' < \varepsilon < \varepsilon''$ and some constant C , which follows from (2.19).

Now we take a canonical transformation \mathcal{X} such that $p_{1, \gamma} \circ \mathcal{X} = \xi_1$ and $q_{0, \gamma} \circ \mathcal{X} = x_1$, where $q_{0, \gamma}$ is defined from q_0 given in (1.5) by the same way as in $p_{1, \gamma}$. Darboux theorem (see [7, Proposition 3.1]) shows that (1.6) guarantees the existence of such a \mathcal{X} . Application of Lemma 2.3 gives

Lemma 2.9. *The estimate (2.8) is valid if for some ε and ε' ($\varepsilon' > \varepsilon > 0$) the estimate*

$$(2.24) \quad \begin{aligned} & \lambda^{-2(l-2+\sigma')} \|h_\varepsilon(\lambda y', \lambda D_{y'}) v\| + \lambda^{-2\sigma} \|D_{y_1}^{l-1} h_\varepsilon(\lambda y', \lambda D_{y'}) v\| \\ & \leq C(\|(D_{y_1}^l + \lambda^{-2(l-1)} \tilde{A}_{\varepsilon', \lambda}(y, D_{y'}))v\| \\ & \quad + \|D_{y_1}^{l-1} v\| + \lambda^{-2(l-2+\sigma)} \|v\|), \\ & \text{if } v \in \mathcal{S} \text{ vanishes for } |y_1| > \varepsilon, \end{aligned}$$

holds for some C , where the symbol of $\tilde{A}_{\varepsilon, \lambda}(y, D_{y'})$ is $\tilde{a}(y_1, \lambda y', \lambda \eta') h_\varepsilon(\lambda y', \lambda \eta')$. Here $\tilde{a}(x, \xi') = (a_\gamma \circ \mathcal{X})(x, 0, \xi')$.

Proof. Condition (1.7) leads us to the

$$(2.25) \quad a_{\gamma} \circ \chi(x, \xi) = a_{\gamma} \circ \chi(x, 0, \xi') + b(x, \xi) \xi_1^{l-1}$$

for some b near origin since H_{q_0} is transformed to ∂_{ξ_1} . Therefore, substituting $h_{\varepsilon}(y_1, \lambda^2 D_{y_1}) h_{\varepsilon'}(\lambda y', \lambda D_{y'}) v$ ($\varepsilon < \varepsilon'' < \varepsilon'$) for v of (2.24) with replaced \tilde{a} by $a_{\gamma} \circ \chi(x, \xi)$ and changing variable (y_1, η_1) into $(\lambda y_1, \lambda^{-1} \eta_1)$ we get the estimate (2.8) canonically transformed.

Proposition 2.10. *Let $g_{\lambda}(y, \eta) \in C^{\infty}$ with a parameter $0 < \lambda \leq 1$ satisfy for any α, β*

$$(2.26) \quad |\partial_{\eta}^{\alpha} \partial_y^{\beta} g_{\lambda}| \leq C_{\alpha\beta} (|\eta_1|^2 + \lambda^{-4\delta})^{(m-|\alpha_1|)/2} \lambda^{|\alpha'| + \beta'}$$

where $0 < \delta \leq 1$ and m integer. Assume that, for some $h_{\lambda}(y', \eta') = h(\lambda y', \lambda \eta')$, g_{λ} satisfies for some $c_0 > 0$

$$(2.27) \quad |g_{\lambda}| \geq c_0 (|\eta_1|^2 + \lambda^{-4\delta})^{m/2} \text{ on } \{|y_1| < 1\} \times \text{supp } h_{\lambda}.$$

Then we get

$$(2.28) \quad \begin{aligned} & \|D_{y_1}^m h_{\lambda}(y', D_{y'}) v\| + \lambda^{-2\delta m} \|h_{\lambda}(y', D_{y'}) v\| \\ & \leq C (\|G_{\lambda}(y, D_y) v\| + \|v\|), \quad \text{if } v \in \mathcal{S} \text{ vanishes } |y_1| > 1. \end{aligned}$$

Proof is omitted. (for example, see [5, p. 77]).

Applying this proposition with $m=l$ and $\delta=(l-1)/l$ to $\eta_1^l + \lambda^{-2(l-1)} \tilde{a}(y_1, \lambda y', \lambda \eta') h_{\varepsilon}(\lambda y', \lambda \eta')$, we obtain (2.24) if $\text{Im } \tilde{a}(0) \neq 0$. From now on we assume $\text{Im } \tilde{a}(0) = 0$. Let $\omega(x, \xi')$ be a l power root of $(\tilde{a} h_{\varepsilon})(x, \xi')$ such that $\omega(0)$ is real (since $\text{Re } \tilde{a}(0) \neq 0$ by (1.2)).

Then we obtain the factorization

$$\xi_1^l + \tilde{a} h_{\varepsilon} = (\xi_1 + \omega) \sum_{j=1}^{l-1} (-\omega)^{j-1} \xi_1^{l-j-1}$$

Set $\omega_{\lambda}(y, \eta') = \omega(y_1, \lambda y', \lambda \eta')$ and set $l_{2,\lambda}(y, \eta) = \sum_{j=1}^{l-1} (-\lambda^{-2\delta} \omega_{\lambda})^{j-1} \eta_1^{l-j-1}$ ($\delta=(l-1)/l$).

Since $l_{2,\lambda}(y, \eta)$ satisfies (2.27) with $m=l-1$ and $\delta=(l-1)/l$, we get (2.24) if we show for some C

$$(2.29) \quad \begin{aligned} & \lambda^{-2\sigma} \|h_{\varepsilon,\lambda}(y', D_{y'}) v\| \leq C (\|(D_{y_1} + \lambda^{-2\delta} \omega_{\lambda}(y, D_{y'})) v\| + \|v\|), \\ & \text{if } v \in \mathcal{S} \text{ vanishes } |y_1| > \varepsilon. \end{aligned}$$

For brevity we denote $\tilde{a}(x, \xi') h_{\varepsilon'}(x', \xi')$ by $\tilde{a}(x, \xi')$ in what follows. Note that $\text{Re } \tilde{a}$ is independent of x_1 on account of (1.3) because H_{p_1} and Γ were transformed to ∂_{x_1} and $\xi_1=0$, respectively, by the χ . Using the expansion $(1+z)^{1/l} = 1 + z/l + O(z)^2$, we obtain

$$(2.30) \quad \begin{aligned} \omega(x, \xi') &= (\operatorname{Re} \tilde{a})^{1/l} (1 + \operatorname{Im} \tilde{a} / \operatorname{Re} \tilde{a})^{1/l} \\ &= r(x', \xi') + iq(x, \xi') + \mathcal{O}(|x| + |\xi'|) q(x, \xi') \end{aligned}$$

if we set $r = (\operatorname{Re} \tilde{a})^{1/l}$, $q = \operatorname{Im} \tilde{a} / l(\operatorname{Re} \tilde{a})^{(l-1)/l}$. Hence (2.29) follows if we show that for some C

$$(2.31) \quad \begin{aligned} &\lambda^{-2\sigma} \|h_{\varepsilon, \lambda}(y', D_{y'})v\| + \lambda^{-2\delta} \|q_{\lambda}(y, D_{y'})h_{\varepsilon, \lambda}(y', D_{y'})v\| \\ &\leq C(\|D_{y_1} + \lambda^{-2\delta}(r_{\lambda}(y', D_{y'}) + iq_{\lambda}(y, D_{y'}))v\| + \|v\|), \\ &\quad \text{if } v \in \mathcal{S} \text{ vanishes for } |y_1| > \varepsilon, \end{aligned}$$

where $q_{\lambda}(y, \eta') = q(y_1, \lambda y', \lambda \eta')$ and $r_{\lambda}(y', \eta') = r(\lambda y', \lambda \eta')$. Indeed, if we take $h_{\varepsilon'}(x', \xi')$ such that $h_{\varepsilon}(x', \xi') = 1$ on $\operatorname{supp} h_{\varepsilon'}$ and substitute $h_{\varepsilon', \lambda}v$ into (2.31), then we get (2.29) with replaced v by $h_{\varepsilon', \lambda}v$ since the part corresponding to third term of the left hand side of (2.30) can be estimated by the second term of the right hand side of (2.31) when ε is small enough. (Note that $\varepsilon'' < \varepsilon$).

Let $\Phi(x, \xi')$ be the solution to

$$(2.32) \quad \partial_{x_1} \Phi + r(x', d_{x'} \Phi) = 0, \quad \Phi(0, x', \xi') = x' \xi'$$

Without loss of generality we assume that the Φ exists on $\{|x_1| < \varepsilon\} \times \operatorname{supp} h_{\varepsilon}(x', \xi')$. Put $\Phi_{\lambda}(y, \eta') = \lambda^{-2} \Phi(\lambda^{2/l} y_1, y', \lambda \eta')$. Then Φ_{λ} of course exists on $\{|y_1| < \varepsilon\} \times \operatorname{supp} h_{\varepsilon, \lambda}(y', \eta')$. If we regard y_1 as a parameter, in the same way as in (2.10) and (2.11), we can define the Fourier integral operator and the conjugate Fourier integral operator with phase function $\Phi_{\lambda}(y, \eta')$ and the symbol $f_{\lambda}(y, \eta')$ satisfying

$$(2.6') \quad |\partial_y^{\alpha} \partial_{\eta'}^{\beta'} f_{\lambda}| \leq C_{\alpha\beta'} \lambda^{|\alpha' + \beta'|}$$

by

$$(2.33) \quad F_{\Phi_{\lambda}} v(y) = \int e^{i\Phi_{\lambda}(y, \eta')} k_{\lambda}(y', \eta') f_{\lambda}(y, \eta') \check{v}(y_1, \eta') d\eta'$$

and

$$(2.34) \quad \begin{aligned} F_{\Phi_{\lambda}}^* v(y) &= \iint e^{i(y' \eta' - \Phi_{\lambda}(y_1, \tilde{y}', \eta'))} k_{\lambda}(\tilde{y}', \eta') \\ &\quad f_{\lambda}(y_1, \tilde{y}', \eta') v(y_1, \tilde{y}') d\tilde{y}' d\eta', \end{aligned}$$

when $v \in \mathcal{S}$ vanishes for $|y_1| > \varepsilon$. Here $\check{v}(y_1, \eta')$ denotes the Fourier transform of $v(y)$ with respect to y' . Set

$$(2.35) \quad \Psi_{\lambda}(y, \eta) = y_1 \eta_1 + \Phi_{\lambda}(y, \eta').$$

Let \mathcal{X}_{λ} denote a canonical transformation with generating function Ψ_{λ} , that is, $\mathcal{X}_{\lambda}; (y, d_y \Psi_{\lambda}(y, \eta)) \mapsto (d_{\eta} \Psi_{\lambda}(y, \eta), \eta)$. Note that \mathcal{X}_{λ} and $\mathcal{X}_{\lambda}^{-1}$ are defined for $\{|y_1| < \varepsilon\} \times \operatorname{supp} h_{\varepsilon, \lambda}(y', \eta')$ if ε is small enough. Set $\tilde{q}_{\lambda}(y, \eta') = q_{\lambda} \circ \mathcal{X}_{\lambda}(y, \eta')$. It is easy to check that

$$(2.36) \quad \tilde{q}_{\lambda}(y, \eta') = q(y_1, \psi_{\lambda}(y, \eta'), d_{x'} \Phi(\lambda^{2/l} y_1, \psi_{\lambda}(y, \eta'), \lambda \eta'))$$

where $\psi_\lambda(y, \eta') = \psi(\lambda y'; \lambda^{2/l} y_1, \lambda \eta')$ and $\psi(x'; x_1, \xi')$ is defined as the inverse of $x' = d_{\xi'} \Phi(x_1, \cdot, \xi')$.

Lemma 2.11. *Assume that for some $\varepsilon_1 > 0$ and some constant C_1 the estimate*

$$(2.37) \quad \begin{aligned} & \lambda^{-2\sigma} \|h_{\varepsilon_1, \lambda}(y', D_{y'})v\| + \lambda^{-2\delta} \|\tilde{q}_\lambda(y, D_{y'})h_{\varepsilon_1, \lambda}(y', D_{y'})v\| \\ & \leq C_1 (\|(D_{y_1} + i\lambda^{-2\delta} \tilde{q}_\lambda(y, D_{y'}))v\| + \|v\|) \end{aligned}$$

holds if $v \in \mathcal{S}$ vanishes for $|y_1| > \varepsilon_1$. Then (2.3) holds for some $\varepsilon > 0$ and C .

Proof. Fix y_1 as a parameter and let $\Phi_\lambda(y_1, y', \eta')$ correspond to $S_\lambda(y, \eta)$ in Definition 2.4. By the remark after Corollary 2.8, we see that the $\lambda^{-2}(I_{\Phi_\lambda}^* q_\lambda - \tilde{q}_\lambda I_{\Phi_\lambda}^*)$ as an operator on $L_2(R_{y'}^{n-1})$ has a uniform bound with respect to $|y_1| < \varepsilon$ and $0 < \lambda \leq 1$, therefore it has a uniform bound as an operator on $L_2([-\varepsilon, \varepsilon] \times R_{y'}^{n-1})$ by integrating with respect to y_1 .

Note that

$$\begin{aligned} D_{y_1} I_{\Phi_\lambda}^* v &= I_{\Phi_\lambda}^* D_{y_1} v - (\partial_{y_1} \Phi_\lambda)^*_{\Phi_\lambda} v \\ &= I_{\Phi_\lambda}^* D_{y_1} v + \lambda^{-2\delta} T_{\Phi_\lambda}^* v, \\ &\text{if } v \in \mathcal{S} \text{ vanishes for } |y_1| > \varepsilon, \end{aligned}$$

where $t_\lambda(y, \eta') = r_\lambda(y', d_{y'} \Phi_\lambda(y, \eta'))$. In fact this follows from $\partial_{y_1} \Phi_\lambda(y, \eta') = -\lambda^{-2\delta} r_\lambda(y', d_{y'} \Phi_\lambda(y, \eta'))$. The adjoint form of Proposition 2.7 shows that the second term equals $I_{\Phi_\lambda}^* \lambda^{-2\delta} R_\lambda$ modulo L_2 -bounded operator. Hence, substitution $I_{\Phi_\lambda}^* v$ for v of (2.37) gives (2.31) because for $\tilde{h}_{\varepsilon, \lambda}(y_1, y', \eta')$ defined from $h_\varepsilon(x', \xi')$ in the same way as (2.36), there exists some ε' such that $\tilde{h}_{\varepsilon, \lambda} = 1$ on $\text{supp } h_\varepsilon(\lambda y', \lambda \eta') \times \{|y_1| < \varepsilon\}$, provided that $0 < \lambda \leq \lambda_0$ for some sufficiently small λ_0 .

In the rest of this section we investigate the properties of $\tilde{q}_\lambda(y, \eta')$ derived from the assumptions. It follows from (2.36) that for any α, β' and some $C_{\alpha\beta'}$ independent of λ

$$(2.38) \quad \begin{aligned} & |\partial_y^\alpha \partial_{\eta'}^{\beta'} \tilde{q}_\lambda(y, \eta')| \leq C_{\alpha\beta'} \lambda^{|\alpha'| + \beta'|} \\ & \text{on } \Omega_\varepsilon = \{|y_1| < \varepsilon, \quad |y'| + |\eta'| < \varepsilon \lambda^{-1}\}. \end{aligned}$$

The second property is that

$$(2.39) \quad \begin{aligned} & \tilde{q}_\lambda(y, \eta') \text{ does not change sign from } + \text{ to } - \text{ for} \\ & y_1 \text{ increasing if } \lambda \text{ is small enough.} \end{aligned}$$

This follows from (1.10). In fact, since it follows from (2.25) that

$$d_{x, \xi}(\text{Re } a_{\gamma \circ \mathcal{X}})^{1/l}(x, 0, \xi') = d_{x, \xi}(\text{Re } a_{\gamma \circ \mathcal{X}})^{1/l}(x, 0, \xi')$$

we obtain the property (2.39) because the modified-null-bicharacteristic curve is invariant under canonical transformations \mathcal{X} and \mathcal{X}_λ . The invariance of

Poisson brackets for canonical transformations and (2.25) give the following;

$$(2.40) \quad \begin{cases} \text{for any } (y, \eta') \in \Omega_\varepsilon = \{|y_1| < \varepsilon, |y'| + |\eta'| < \varepsilon \lambda^{-1}\} \\ \text{there exists a } I \in \mathcal{J}_\mu \text{ such that } \tilde{p}_{I,\lambda}(y, \eta') \neq 0 \text{ if} \\ \lambda \text{ is small enough.} \end{cases}$$

Here $\tilde{p}_{I,\lambda}(y, \eta')$ is defined in the same way as $p_I(x, \xi)$ of (1.11) with $p_1 = \eta_1$ and $p_2 = \lambda^{-2\delta} \tilde{q}_\lambda(y, \eta')$. Indeed, note that (1.12) is invariant under canonical transformations and in changing $p_i (i=1, 2)$ to $f_i p_i$ for non-vanishing functions $f_i (i=1, 2)$. By means of (2.25) and the definition of \mathcal{J} , we see that for any $(y, \eta') \in \Omega_\varepsilon$ there exist a $I \in \mathcal{J}_\mu$ and $c_I > 0$ such that

$$|p_{I,\lambda}^0(y, \eta')| \geq c_I \lambda^{-2+2b(I)/I},$$

where $p_{I,\lambda}^0$ is defined by setting $p_1 = \eta_1$ and $p_2 = \lambda^{-2\delta} q_\lambda(y, \eta')$. If $p_{I,\lambda}$ denotes $p_{I,\lambda}^0$ with replaced η_1 by $\eta_1 + \lambda^{-2\delta} r_\lambda(y', \eta')$, we obtain

$$|p_{I,\lambda} - p_{I,\lambda}^0| \leq C_I \lambda^{-2\delta+2b(I)/I}$$

for some constant C_I determined by the derivatives of r and q . Consequently, the invariance of Poisson brackets under \mathcal{X}_λ gives (2.40).

3. Proof of Theorem 1.1

In order to prove Theorem 1.1, as observed in the preceding section, it suffices to show (2.37). For the sake of simplicity, we denote $\lambda^{-2\delta} \tilde{q}_\lambda(y, \eta')$ by $q(x, \xi')$. Suppose that λ is small enough. By means of a constant scale change in the variables, we may assume from (2.38)–(2.40) that $q(x, \xi')$ satisfies the following conditions: For any α, β and some constant $C_{\alpha\beta}$ (independent of λ)

$$(3.1) \quad |D_\xi^\alpha D_x^\beta q(x, \xi')| \leq C_{\alpha\beta} \lambda^{|\alpha'| + \beta' - 2\delta} \quad \text{in } \Omega = \{|x_1| < 1, |(x', \xi')| < \lambda^{-1}\}$$

$$(3.2) \quad q(x, \xi') \text{ does not change sign from } + \text{ to } - \text{ for } x_1 \text{ increasing.}$$

$$(3.3) \quad \text{For any } (x, \xi') \in \Omega \text{ there exist some } \mu \in Q_0 \text{ and some } c_0 > 0 \text{ such that}$$

$$\sum_{I \in \mathcal{J}_\mu} \lambda^{2-2b(I)/I} |p_I(x, \xi')| \geq c_0 > 0$$

provided that $p_1 = \xi_1, p_2 = q(x, \xi')$. Then (2.37) is stated as follows; for some C

$$(3.4) \quad \begin{aligned} & \lambda^{-2\sigma} \|h(\lambda x', \lambda D')u\| + \|D_{x_1} h(\lambda x', \lambda D')u\| \\ & \leq C (\|(D_{x_1} + iq(x, D'))u\| + \|u\|), \\ & \quad \text{if } u \in \mathcal{S} \text{ vanishes for } |x_1| > 1/2, \end{aligned}$$

where h is C_0^∞ with support in a ball of radius $1/2$.

The proof of (3.4) for $q(x, \xi)$ with (3.1)–(3.3) is the same as in showing [4, (6.1) and (6.30)] except the difference of “weight”. (See [4, (4.1) and (4.3)].) To prove [4, (6.1) and (6.30)] it was important to obtain inequalities for q in [4, Section 4]. So we sketch the argument corresponding to [4, Section 4]. Put

$$(3.5) \quad M(x, \xi') = \max_{I \in \mathcal{J}_\mu} |p_I(x, \xi')/\rho|^{1/|I|}.$$

Here ρ is a large parameter, whose role is the same as in [4, Section 4] (see [4, p. 149]). By (3.1) and (3.3) we have

$$(3.6) \quad C_1 \lambda^{-2\delta/\mu} \rho^{-(l-1)/\mu} \leq M(x, \xi') \leq C_2 \lambda^{-2\delta} \rho.$$

Here and in what follows the constants are independent of λ and ρ .

The definition of $M=M(0)$ means in particular that

$$(3.7) \quad |D_{x_1}^j q(0)| < \rho M^{j+1}, \quad j \leq \mu-1,$$

and where (3.1) is valid we have by (3.6)

$$(3.8) \quad |D_{x_1}^j q| \leq \mathcal{O}(\lambda^c) \rho M^{[\mu]+1}$$

since $\rho^{l-2}/M^{[\mu+1]-\mu} \ll 1$ if λ is small enough, where c is some positive. If we set

$$F(t, y', \eta') = q(t/M, y'(\rho M)^{1/2\delta}, \eta'(\rho M)^{1/2\delta})/\rho M,$$

then the application of [4, Lemma 7.1] to $F(t, y', \eta')$ shows that

$$(3.9) \quad |D_\xi^\alpha D_x^\beta (\tilde{q}(x, \xi') - \xi_2 \partial \tilde{q}(x_1, 0)/\partial \xi_2)| \leq C_{\alpha\beta} \rho M^{\beta_1+1} (\rho M)^{-|\alpha'+\beta'|/2\delta} \\ \text{if } |x_1 M| < 1, \quad |(x', \xi')| < C(\rho M)^{1/2\delta},$$

where \tilde{q} is determined from q by a symplectic orthogonal transformation.

Let $\varepsilon = \lambda^\kappa$ with $0 < \kappa < 1/\mu$ (which is different from ε in the preceding section). Then

$$(3.10) \quad \varepsilon^2 (\rho M)^{1/2\delta} \gg 1.$$

As in [4], using this ε we consider the following two cases.

Case I. Assume that

$$(3.11) \quad |d_{x'\xi'} D_{x_1}^j q(0)| \leq \varepsilon \rho M^{j+1}, \quad j < \mu'.$$

where $\mu' = (l-2)(\mu-1)/2(l-1)$. In view of (3.1), (3.6) and (3.10) we get for some $c > 0$

$$(3.11)' \quad |d_{x'\xi'} D_{x_1}^j q(x, \xi')| \leq \mathcal{O}(\lambda^c) \varepsilon \rho M^{[\mu'] + 1}, \quad j > \mu', \quad (x, \xi') \in \Omega.$$

Then it is easy to check that the argument corresponding to Case I in [4, Section

4] follows with k and $k/2$ replaced by $[\mu-1]$ and μ' .

Case II. Assume now that (3.11) is not fulfilled. Choose $s < \mu'$ so that with $q^{(j)} = \partial^j q / \partial x_1^j$

$$(3.12) \quad d_{x' \xi'} q^{(s)}(0) = a$$

$$(3.13) \quad d_{x' \xi'} q^{(j)}(0) \leq a M^{j-s} \quad \text{for } j < \mu'$$

Then (3.1) and the fact that (3.11) is not valid give

$$(3.14) \quad \varepsilon \rho M^{s+1} < a \leq C \lambda^{1-2\delta}.$$

In view of (3.11)' we have then for every j

$$(3.13)' \quad |D_{x_1}^j \tilde{q}(x_1, 0) / \partial \xi_2| \leq C_j a M^{j-s} = C_j \rho M^{j+1} A_2, \quad |x_1 M| < 1,$$

where $A_2 = a / \rho M^{s+1}$. The equality of (3.13)' holds when $j=s$. From (3.9) we can therefore obtain an estimate of the form [4, (4.10)] with $B_2 = A_3 = B_3 = \dots = (\rho M)^{-1/2\delta}$ and $K = M$. However, $A_2 B_2 = a / M^{s+1} (\rho M)^{1/2\delta}$ so [4, Lemma 4.1] is not applicable if $a > M^{s+1} (\rho M)^{1/2\delta}$. In that case we shall replace the orthogonal symplectic transformation which led from q to \tilde{q} by a non-linear canonical transformation.

Thus assume for the moment that (3.12), (3.13) and

$$(3.14)' \quad M^{s+1} (\rho M)^{1/2\delta} < a \leq C \lambda^{1-2\delta}$$

are fulfilled. Let $b = a \lambda^{2(\delta-1)}$. Then the function

$$Q(x', \xi') = (q^{(s)}(0, bx', b\xi') - q^{(s)}(0, 0, 0)) / ab$$

is in a bounded subset of $C^\infty(U)$ ($U = \{|(x', \xi')| < C^{-1}\}$) since $|(bx', b\xi')| < \lambda^{-1}$. Hence there exists some canonical transformation \mathcal{X} belonging to a bounded set in C^∞ for $0 < \lambda \leq 1$ such that

$$Q \circ \mathcal{X}(x', \xi') = \xi_2$$

in a neighborhood of 0. If we put

$$\begin{aligned} \mathcal{X}_b(x', \xi') &= b\mathcal{X}(b^{-1}x', b^{-1}\xi') \\ \tilde{q}(x, \xi') &= q(x_1, \mathcal{X}_b(x', \xi')), \end{aligned}$$

then we obtain

$$(3.15) \quad \begin{aligned} \tilde{q}^{(s)}(x, \xi') &= a\xi_2 + \tilde{q}^{(s)}(0, 0) \\ &\text{when } x_1 = 0, |(x', \xi')| < cb. \end{aligned}$$

By the same way as in [4] we get

$$(3.16) \quad |a^j(\partial/\partial x_2)^j(\partial/\partial x_1)^i \tilde{q}(0)| \leq C_{ij} M^{j(s+1)} \rho M^{i+1}$$

for any i, j satisfying

$$j \leq l-2 \quad \text{and} \quad (i+1+j(s+1))(l-1)/(l-j-1) \leq \mu.$$

If we introduce

$$(3.17) \quad B_2 = M^{s+1}/a,$$

noting incidentally that $\rho A_2 B_2 = 1$ as required in [4, Lemma 4.1], this means that we have bounds for the derivatives of $\tilde{q}(x_1/M, x_2/B_2, 0)/\rho M$ at 0.

As stated in [4, p. 156], when we derive (3.16), we can replace the canonical transformation $\mathcal{X}_b(x', \xi')$ by another $\tilde{\mathcal{X}}_b(x', \xi')$ which is linear in all variables except x_2 , provided that the integral curve of the Hamilton field of $\tilde{q}^{(s)}(0, x', \xi')$ by new $\tilde{\mathcal{X}}_b$ is also the x_2 axis $x_2 = at, x_3 = \dots = \xi_n = 0$. We denote $\tilde{\mathcal{X}}_b$ by \mathcal{X}_b in what follows.

By the analogous calculation as in showing [4, (4.23) and (4.23)'] we obtain when $|x_1 M| < 1, |(x', \xi')| < cb$

$$(3.18) \quad |D_\xi^\alpha D_x^\beta \tilde{q}(x, \xi')| \leq C_{\alpha\beta} a M^{\beta_1} b^{1-|\alpha'+\beta'|} \quad \text{if } |\alpha' + \beta'| \neq 0$$

$$(3.18)' \quad |D_\xi^\alpha D_x^\beta \tilde{q}(x, \xi')| \leq C'_{\alpha\beta} \lambda^{-2\delta} b^{-|\alpha'+\beta'|} \quad \text{for any } \alpha', \beta'.$$

In particular (3.18)' is a much better estimate than (3.16) if $j \geq l-1$ or $(i+1+j(s+1))(l-1)/(l-j-1) > \mu$, and it is not only valid at 0. Hence (3.16) leads us to uniform bounds for $\tilde{q}(x_1/M, x_2/B_2, 0)/\rho M$ and all of its derivatives when $|x_1| < 1$ and $|x_2| < 1$.

Since $b \gg (\rho M)^{1/2\delta}$ we can apply [4, Lemma 7.1] to

$$F(t, x_2, y) = (M\rho)^{-1} \tilde{q}(t/M, x_2/B_2, x''(\rho M)^{1/2\delta}, \xi'(\rho M)^{1/2\delta})$$

where $x'' = (x_3, \dots, x_n)$ and $y = (x'', \xi')$. Therefore, by the same way as in getting [4, (4.25)] we obtain

$$(3.19) \quad |D_\xi^\alpha D_x^\beta (\tilde{q}(x, \xi') - \xi_2 \partial \tilde{q}(x_1, x_2, 0)/\partial \xi_2)| \\ \leq C M^{\beta_1+1} \rho B_2^{\beta_2} (\rho M)^{-|\alpha'+\beta'|/2\delta}$$

$$\text{when} \quad |x_1| M < 1, |(x'', \xi')| < (\rho M)^{1/2\delta}, |x_2| < 1/B_2.$$

As in [4], we obtain [4, (4.26)] and [4, (4.27)] with replaced the right hand side by $C b^{-\beta_2} \lambda^{2/l}$. Hence it follows from (3.12) that $(B_2/M) \partial \tilde{q}(x_1/M, 0)/\partial \xi_2$ is essentially a normalized polynomial in x_1 of degree $[\mu']$ and $\partial \tilde{q}(x_1/M, x_2, 0)/\partial \xi_2$ is almost independent of x_2 .

From (3.19) and these inequalities we obtain with $B_1 = M, A_j = B_j = (\rho M)^{-1/2\delta}$ when $j > 2$

$$(3.20) \quad |D_{\xi}^{\alpha} D_{\xi'}^{\beta} \tilde{q}(x, \xi')| \leq C_{\alpha\beta} \rho M A^{\alpha} B^{\beta}, \quad \text{if } |x_1 M| < 1, |x_2 B_2| < 1, \\ |\xi_2 A_2| < N, |(x'', \xi'')| < (\rho M)^{1/2\delta},$$

where N is a fixed but arbitrary constant. If we write $\tilde{M}(x, \xi') = M(x_1, \mathcal{X}_b(x', \xi'))$, it follows in view of [4, Lemma 4.1], where we take $A_1 = 1/M$ and $K = M$, that

$$(3.21) \quad \tilde{M}(x, \xi') \leq C_N M \quad \text{if } |x_1 M| < 1, |x_2 B_2| < 1, \\ |\xi_2 A_2| < N, |(x'', \xi'')| < (\rho M)^{1/2\delta}.$$

when (3.14) is fulfilled but not (3.14)' we get the same conclusion with B_2 replaced by $(\rho M)^{-1/2\delta}$ and \mathcal{X}_b equal to orthogonal symplectic transformation such that $\tilde{q}(x, \xi') = q(x_1, \mathcal{X}_b(x', \xi'))$.

The argument corresponding to the rest of [4, Section 4] can be done by the same way if we let (3.1)–(3.3), (3.16), (3.18), (3.18)', (3.19), (3.20), (3.21), $[\mu - 1]$, μ' , \mathcal{X}_b and $(\rho M)^{1/2\delta}$ correspond to [4, (4.1)–(4.3), (4.21), (4.23), (4.23)', (4.25), (4.28), (4.29), k , $k/2$, \mathcal{X}_a and $\sqrt{\rho M}$] respectively.

Because we have got the result corresponding to [4, Section 4] we can easily prove (3.4) by the same way as in [4, Section 6], if we take the above correspondence. The detail is left to the reader.

4. Proof of Proposition 1.2

The method of the proof is a version of [2] (, see also [15]). Suppose that \tilde{L} is hypoelliptic at the origin. Then, as well-known, there exist a positive integer s , a constant C and some neighborhood U of 0 such that

$$(4.1) \quad \|u\| \leq C \|\Lambda^s \tilde{L}u\|, \quad u \in C_0^\infty(U),$$

where the symbol of Λ is $(|\xi_1|^{10} + |\xi'|^8 + 1)$. Hence for any large N there exists a C_N such that

$$(4.1)' \quad \|u\| \leq C_N (\|\Lambda^s \tilde{L}u\| + \| |x|^{-N} u \| + \| |x|^N \Lambda^s \tilde{L}u \|), \quad u \in \mathcal{S}.$$

If we take the canonical change of variables (x, ξ) into

$$(4.2) \quad (\lambda^2 x_1, \lambda^9 x', \lambda^{-2\tau-2}(\xi_{01} + \lambda^{2\tau} \xi_1), \lambda^{-2\tau-9}(\xi'_0 + \lambda^{2\tau} \xi')), \\ (0 < \lambda \leq 1, \tau = 13),$$

with a fixed $\xi_0 = (\xi_{01}, 0, \xi_{03})$ satisfying

$$\xi_{01}^5 + 2\xi_{03}^4 = 0 \quad \text{and} \quad \xi_{03} > 0,$$

then it follows from (4.1)' that

$$(4.3) \quad \|u\| \leq C'_N (\lambda^{-M} \|L_\lambda u\| + \lambda^{2N-M} \|R_0(x, \lambda^{2\tau} D_x)u\|), \quad u \in \mathcal{S},$$

where

$$\begin{aligned}
L_\lambda &= L_1(\lambda^{2\tau} D_x) + i\lambda^\tau x_1^2 x_2 \xi_{03}^4 + \sum_{j=1}^2 \lambda^{\tau(2-j)} R_j(x, \lambda^{2\tau} D_x), \\
L_1(\xi) &= 5\xi_{01}^4 \xi_1 + 4\xi_{03}^3 (2\xi_3 - \xi_2), \\
R_j(x, \xi) &= \sum_{\substack{j \leq |\alpha| \\ |\alpha| + \beta \leq N_j}} a_{\alpha\beta j}(\lambda) x^\beta \xi^\alpha \quad (j = 0, 1, 2) \\
(a_{\alpha\beta j}(\lambda) &= \mathcal{O}(\lambda^c), c \geq 0).
\end{aligned}$$

Here M and $N_j (j=0, 1, 2)$ are some integers depending on s . Since $\xi_{03} > 0$ we can take $w(x)$ such that

$$\begin{aligned}
(4.4) \quad & L_1(\partial_x w) + i x_1^2 x_2 \xi_{03}^4 = 0 \\
& |\operatorname{Im} w| \geq c_0 |x|^4, \quad c_0 > 0.
\end{aligned}$$

Put $u_\lambda(x) = (\exp i\lambda^{-\tau} w(x)) \sum_{j=0}^N v_j(x, \lambda) \lambda^{\tau j}$, where v_j will be determined later such that they are polynomials of x whose coefficients have the same property as $a_{\alpha\beta j}$. Then we have

$$\begin{aligned}
L_\lambda u_\lambda &= \exp i\lambda^{-\tau} w \sum_{j=2}^N (L_1(D_x) + A(x, \partial_x w)) v_{j-2} + F_j(x) \lambda^{\tau j}, \\
(F_2 &= 0).
\end{aligned}$$

Here $A(x, \xi) = \sum_{|\alpha|=1} a_{\alpha\beta 1} x^\beta \xi^\alpha$ and $F_j(x)$ is a linear combination of the functions v_0, \dots, v_{j-3} and their derivatives. By means of Cauchy-Kowalewska theorem, solve the transport equations

$$(L_1(D_x) + A(x, \partial_x w)) v_j + F_{j+2}(x) = \mathcal{O}(|x|^{N-j})$$

under the condition $v_0(0, \lambda) = 1$ and $v_j(0, \lambda) = 0$ ($j \geq 1$), successively. Then the analytic solution obtained, which is defined in a certain neighborhood ω of the origin, is to be multiplied by a cut off function $\phi \in C_0^\infty(\omega)$ which equals 1 in another neighborhood of the origin. The multiplication does not affect (4.5) because $(1-\phi)v_j = \mathcal{O}(|x|^N)$. Substituting $u_\lambda(x) \in \mathcal{S}$ into (4.3) and changing variables x into $\lambda^{\tau/4} x$ give us the contradiction if $2N > M$ and if λ tends to 0.

References

- [1] Yu. V. Egorov: *Canonical transformations and pseudodifferential operators*, Trudy Moskov. Mat. Obšč. **24** (1971), 3–28. =Trans. Moscow Math. Soc. **24** (1971), 1–28.
- [2] Yu. V. Egorov: *On necessary conditions for the solvability of pseudodifferential equations of principal type*, Trudy Moskov. Mat. Obšč. **24** (1971), 29–41 =Trans. Moscow Math. Soc. **24** (1971), 29–42.
- [3] Yu. V. Egorov: *Subelliptic operators*, Uspehi. Mat. Nauk **30**; 2 (1975), 57–114, **30**; 3 (1975), 57–104 =Russian Math. Surveys **30**; 2 (1975), 59–118, **30**; 3

- (1975), 55–105.
- [4] L. Hörmander: *Subelliptic operators*, Seminar on Singularities of Solution of Linear Partial Differential Equations, Princeton University Press 1979, 3–49.
 - [5] H. Kumano-go: *Pseudo-differential operators*, Iwanami Publisher, Tokyo, 1974 (in Japanese).
 - [6] A. Menikoff: *On hypoelliptic operators with double characteristics*, Ann. Scuola Norm. Sup. Pisa Cl. Sci. (4) **4** (1977), 689–724.
 - [7] R.B. Melrose: *Equivalence of glancing surfaces*, Invent. Math. **37** (1976), 165–191.
 - [8] S. Mizohata: *Solutions nulles et solutions non-analytiques*, J. Math. Kyoto Univ. **1** (1961/2), 271–302.
 - [9] Y. Morimoto: *On the hypoellipticity for infinitely degenerate semi-elliptic operators*, J. Math. Soc. Japan **30** (1978), 327–358.
 - [10] L. Nirenberg and F. Trèves: *On local solvability of linear partial differential equations, Part I: Necessary conditions*, Comm. Pure Appl. Math. **23** (1970), 1–38, *Part II: Sufficient conditions*, Comm. Pure Appl. Math. **23** (1970), 459–509.
 - [11] O.A. Oleinik and E.V. Radkevich: *Second order equations with non-negative characteristic form*, Amer. Math. Soc., Providence, Rhode Island and Plenum Press, 1973.
 - [12] P.R. Popivanov: *On local solvability for a class of pseudodifferential equations with double characteristics*, Journal of I.G. Petrovskii Seminar **1** (1975), 237–278.
 - [13] R. Rubinstein: *Local solvability of the operators $u_{tt} + ia(t)u_x + b(t)u_t + c(t)u$* , J. Differential Equations **14** (1973), 185–194.
 - [14] F. Trèves: *A new method of proof of the subelliptic estimates*, Comm. Pure Appl. Math. **24** (1971), 71–115.
 - [15] P. Wenston: *A necessary conditions for the local solvability of the operator $P_m^2(x, D) + P_{2m-1}(x, D)$* , J. Differential Equations **25** (1977), 90–95.
 - [16] P. Wenston: *A sufficient condition for the local solvability of a linear partial differential operator with double characteristics*, J. Differential Equations **29** (1978), 374–387.
 - [17] P. Popivanov and G. Popov: *Microlocal properties of a class of pseudodifferential operators with multiple characteristics, Part I, Part II*, Serdica **6** (1980), 169–183, Serdica **7** (1981), 131–141. (in Russian).

Department of Engineering Mathematics
 Faculty of Engineering
 Nagoya University
 Furo-cho, Chikusa-ku,
 Nagoya 464, Japan

