

Title	Note on complex K-groups of compact Lie groups with fundamental group of prime order
Author(s)	Minami, Haruo
Citation	Osaka Journal of Mathematics. 1998, 35(3), p. 547-551
Version Type	VoR
URL	https://doi.org/10.18910/7263
rights	
Note	

Osaka University Knowledge Archive : OUKA

<https://ir.library.osaka-u.ac.jp/>

Osaka University

NOTE ON COMPLEX K -GROUPS OF COMPACT LIE GROUPS WITH FUNDAMENTAL GROUP OF PRIME ORDER

HARUO MINAMI

(Received April 9, 1997)

1. Let G be a compact connected simply-connected Lie group and Γ a central subgroup of prime order p . Held and Suter [1] and Hodgkin [3] present two kinds of methods which can be used to determine the structure of $K^*(G/\Gamma)$, where K denotes the complex K -functor.

The purpose of this note is to describe a simple method for the computation of $K^*(G/\Gamma)$, which uses equivariant complex K -theory. Special cases of this method have been applied for calculating, e.g., $K^*(PE_6)$ in [5] and $K^*(PSp(n))$ in [6]. We shall also use the structure theorem on $K^*(G)$ [2] and in addition that on $K^*(L^n(p))$ [4], where $L^n(p)$ denotes the lens space. As a result we can get the structure of $K^*(G/\Gamma)$ as an algebra. Using the notations explained later our result can be stated as follows.

Theorem ([1], [3]). *Let G , Γ and p be as above. Then*

$$K^*(G/\Gamma) = \Lambda_R(\beta(\zeta_2), \dots, \beta(\zeta_r), \beta(\rho_{r+1}), \dots, \beta(\rho_\ell), \beta(\kappa)) / ((V-1)\beta(\kappa))$$

where $\ell = \text{rank } G$ and $R = \mathbb{Z}[V]/(p^s(V-1), \dots, p^s(V-1)^{p-1}, V^p-1, (V-1)^{s(p-1)+1})$.

The rest of this note is devoted to our proof of the theorem together with the explanation of the symbols for the generators.

2. Let ρ_1, \dots, ρ_ℓ be the fundamental irreducible representations of G with $\ell = \text{rank } G$. Let V denote the canonical non-trivial complex one-dimensional representation of Γ . And let V^k denote the k -fold tensor product of V and qV^k the direct sum of q copies of V^k . Since G admits at least one faithful representation, all the ρ_i 's are not trivial on Γ . So by Schur's lemma we may assume that the restrictions of the first ρ_1, \dots, ρ_r to Γ can be written as

$$\rho_i|_\Gamma = p^{s_i} n_i V^{k_i}$$

with $1 \leq k_i \leq p-1$, $(p, n_i) = 1$ and $s_1 \leq s_i$ for all i and the rest are trivial on Γ .

Set $s = s_1$ and denote by V itself the line bundle $G \times_{\Gamma} V \rightarrow G/\Gamma$. Then the order of $V - 1$ in $\tilde{K}(G/\Gamma)$ is a power of p since V is induced from the canonical line bundle over a certain lens space $L^m(p)$. So using ρ_1 we see that $p^s(V - 1) = 0$ in $\tilde{K}(G/\Gamma)$, so that there exists a stable isomorphism

$$(2.1) \quad C : G \times p^s V \cong G \times \mathbf{C}^{p^s}$$

of Γ -vector bundles over G .

Let $B(qV)$ and $S(qV)$ be the unit ball and sphere in qV respectively and let $\Sigma^q V = B(qV)/S(qV)$ in which the pinched $S(qV)$ serves as base point. Let $L^m(p) = S((m+1)V)/\Gamma$ as usual and write again V for the line bundle $S((m+1)V) \times_{\Gamma} V \rightarrow L^m(p)$. Set $m = t(p-1) + r$ with $0 \leq r < p-1$. By [4] we then have

$$(2.2) \quad \tilde{K}(L^m(p)) = \mathbf{Z}_{p^{t+1}}\{V-1, \dots, (V-1)^r\} \oplus \mathbf{Z}_{p^t}\{(V-1)^{r+1}, \dots, (V-1)^{p-1}\}.$$

And its ring structure is given by the relations $V^p = 1$ and $(V-1)^{m+1} = 0$.

Let K_{Γ} denote the equivariant complex K -functor associated with Γ . Then for a free Γ -space X we have a canonical isomorphism $K_{\Gamma}(X) \cong K(X/\Gamma)$ which will be identified below. Let us put $n = s(p-1)$. Then by (2.2) we see that $p^s(V-1) = 0$ in $\tilde{K}(L^n(p))$ and hence we have a stable isomorphism

$$T : S((n+1)V) \times \mathbf{C}^{p^s} \cong S((n+1)V) \times p^s V$$

of Γ -vector bundles over $S((n+1)V)$. This gives rise to an element τ of $\tilde{K}_{\Gamma}(\Sigma^{(n+1)V})$ in a canonical manner such that its restriction to the origin of $B((n+1)V)$ is $p^s(1-V)$ in $R(\Gamma)$, the complex representation ring of Γ .

Consider the exact sequence for the pair $(B((n+1)V), S((n+1)V))$ in K_{Γ} -theory together with (2.2) when $m = n$. Then it is seen that τ equals the Thom element of $\tilde{K}_{\Gamma}(\Sigma^{(n+1)V})$ up to a multiple of unit of $R(\Gamma)$. The discussion proceeds by viewing this unit as 1 for brevity. Consider the exact sequence for the cofibration

$$S((n+1)V) \times G \xrightarrow{i} B((n+1)V) \times G \xrightarrow{j} \Sigma^{(n+1)V} \wedge G_+$$

where G_+ denotes the disjoint union of G and a single point $+$ which is taken to be the base point of G_+ . Then $j^* : \tilde{K}_{\Gamma}^*(\Sigma^{(n+1)V} \wedge G_+) \rightarrow K_{\Gamma}(B((n+1)V) \times G) = K^*(G/\Gamma)$ becomes a zero map because $j^*(\tau) = p^s(1-V) = 0$ by (2.1). Hence we have a short exact sequence

$$(2.3) \quad 0 \rightarrow K^*(G/\Gamma) \xrightarrow{I} K_{\Gamma}^*(S((n+1)V) \times G) \xrightarrow{\delta} K^*(G/\Gamma) \rightarrow 0$$

under the identification of the Thom isomorphism $K^*(G/\Gamma) \cong \tilde{K}_{\Gamma}^*(\Sigma^{(n+1)V} \wedge G_+)$. So what we next have to do is to determine the structure of the middle group of (2.3).

3. We now consider the generators of two groups of (2.3). Let X be a compact free Γ -space. Write V for the line bundle $X \times_{\Gamma} V \rightarrow X/\Gamma$ as in §2. Moreover let $f : X \rightarrow GL(n, \mathbb{C})$ be a Γ -map where $GL(n, \mathbb{C})$ is the general linear group with the trivial Γ -action. Then such a map defines a unique element of $K_{\Gamma}^{-1}(X)$, denoted by $\beta(f)$, as follows. The assignment $(x, v) \mapsto (x, f(x)v)$ with $x \in X, v \in \mathbb{C}^n$ yields an isomorphism $\theta : X \times \mathbb{C}^n \cong X \times \mathbb{C}^n$ of Γ -vector bundles over X . We get a Γ -vector bundle over the reduced suspension $S(X_+)$ of X_+ by clutching two n -dimensional product vector bundles over separate cones of X by θ . The reduced vector bundle of this is just $\beta(f)$.

Obviously ρ_i gives rise to $\beta(\rho_i) \in K^{-1}(G/\Gamma)$ for $r+1 \leq i \leq \ell$ and $\rho_i \circ \pi$ does $\beta(\rho_i \circ \pi) \in K_{\Gamma}^{-1}(S((n+1)V) \times G)$, denoted by the same symbol $\beta(\rho_i)$ for brevity, where π denotes the projection $S((n+1)V) \times G \rightarrow G$.

Define $\bar{\rho}_i : S((n+1)V) \times G \rightarrow GL(d_i, \mathbb{C})$ with $d_i = \text{degree } \rho_i$ for $1 \leq i \leq r$ by $\bar{\rho}_i(x, g)(v) = \pi((p^{s_i-s} n_i T)^{-k_i}(x, \rho_i(g)v))$ with $x \in S((n+1)V), g \in G, v \in \mathbb{C}^{d_i}$. Here qT denotes the direct sum of q copies of T and π the projection of $S((n+1)V) \times W$ to the second factor. Then $\bar{\rho}_i$ becomes a Γ -map, so that this gives rise to $\beta(\bar{\rho}_i) \in K_{\Gamma}^{-1}(S((n+1)V) \times G)$. Furthermore let us put $f(x, g)(v) = \pi(T^p(x, v))$ with $x \in S((n+1)V), g \in G, v \in \mathbb{C}^{p^s}$ and write ν for $\beta(f)$ which $f : S((n+1)V) \times G \rightarrow GL(p^s, \mathbb{C})$ defines. Then by definition and by making use of (2.1) we have the following.

$$(3.1) \quad \begin{aligned} I(\beta(\rho_i)) &= \beta(\rho_i) \quad (r+1 \leq i \leq \ell), \quad \delta(\nu) = 1 + V + \cdots + V^{p-1}, \\ \delta(\beta(\bar{\rho}_i)) &= -n_i(1 + V + \cdots + V^{k_i-1}) \quad (1 \leq i \leq r) \end{aligned}$$

From (3.1) and the fact that $(p, k_1) = 1, (p, n_1) = 1$ it follows that there exist two polynomials $a(X), b(X) \in \mathbb{Z}[X]$ such that if we put

$$\gamma = a(V)\beta(\bar{\rho}_1) + b(V)\nu \in K_{\Gamma}^{-1}(S((n+1)V) \times G)$$

then

$$(3.2) \quad \delta(\gamma) = 1.$$

By (3.1) and (3.2) we see that there exist more elements $\beta(\zeta_i)$ ($2 \leq i \leq r$) and $\beta(\kappa)$ of $K^{-1}(G/\Gamma)$ such that

$$(3.3) \quad \begin{aligned} I(\beta(\zeta_i)) &= \beta(\bar{\rho}_i) + n_i(1 + V + \cdots + V^{k_i-1})\gamma \quad (2 \leq i \leq r), \\ I(\beta(\kappa)) &= (1 + V + \cdots + V^{p-1})\beta(\bar{\rho}_1) + n_1 k_1 \nu. \end{aligned}$$

4. Let $1 \leq k \leq n+1$. Denote by the same symbol the images of $\beta(\bar{\rho}_i)$'s and $\beta(\rho_j)$'s of $K_{\Gamma}^{-1}(S((n+1)V) \times G)$ by $(i \times 1)^* : K_{\Gamma}^{-1}(S((n+1)V) \times G) \rightarrow K_{\Gamma}^{-1}(S(kV) \times$

G) where i denotes an inclusion $S(kV) \subset S((n+1)V)$. And write ν for the image of the Thom element of $K^{-1}(S^{2k-1}) \cong \mathbf{Z}$ by the transfer $K^{-1}(S^{2k-1}) \rightarrow K_F^{-1}(S(kV))$.

Let $k-1 = t(p-1) + r$ with $0 \leq r < p-1$ and set

$$R_k = \mathbf{Z}[V]/(p^{t+1}(V-1), \dots, p^{t+1}(V-1)^r, \\ p^t(V-1)^{r+1}, \dots, p^t(V-1)^{p-1}, V^p-1, (V-1)^k).$$

We will show that

$$(4.1) \quad K_F^*(S(kV) \times G) = \Lambda_{R_k}(\nu, \beta(\bar{\rho}_1), \dots, \beta(\bar{\rho}_r), \beta(\rho_{r+1}), \dots, \beta(\rho_\ell)) / ((V-1)\nu)$$

by induction on k .

Choose a circle subgroup S^1 of G which contains Γ and view this S^1 as $S(V)$. Then using the multiplication $S(V) \times G \rightarrow G$ yields a homeomorphism $S(V) \times_\Gamma G \approx S^1 \times G$ and so we have $K_F^*(S(V) \times G) \cong K^*(S^1 \times G)$. According to [2] $K^*(G) = \Lambda(\beta(\rho_1), \dots, \beta(\rho_\ell))$. From this and the definition of $\beta(\bar{\rho}_i)$'s and $\beta(\rho_j)$'s it can be easily checked that $K_F^*(S(V) \times G) = \Lambda(\nu, \beta(\bar{\rho}_1), \dots, \beta(\bar{\rho}_r), \beta(\rho_{r+1}), \dots, \beta(\rho_\ell))$.

Next consider the exact sequence for the pair $(S((k+1)V) \times G, S(V) \times G)$ in K_Γ -theory. Because of $S((k+1)V)/S(V) \approx \Sigma^V \wedge S(kV)_+$ we then obtain an exact sequence

$$\cdots \rightarrow K_F^*(S(V) \times G) \xrightarrow{\delta} K_F^*(S(kV) \times G) \\ \xrightarrow{J} K_F^*(S((k+1)V) \times G) \xrightarrow{I} K_F^*(S(V) \times G) \rightarrow \cdots$$

under the identification of the Thom isomorphism $K_F^*(S(kV) \times G) \cong \tilde{K}_F^*(\Sigma^V \wedge (S(kV) \times G)_+)$. And we see that there hold the equalities $J(1) = 1 - V$, $J(\nu) = \nu$ and $\delta(\nu) = 1 + V + \cdots + V^{p-1}$. By making use of these formulas we can proceed with the induction on k and we have (4.1) consequently.

Proof of Theorem. We have $\delta(\nu\beta(\bar{\rho}_1)) = \beta(\kappa)$ in addition to (3.1), (3.2) and (3.3). Using these formulas the theorem follows immediately from (4.1) when $k = n+1$ and the exactness of (2.3). \square

References

- [1] R.P. Held and U. Suter: *On the unitary K-theory of compact Lie groups with finite fundamental group*, Quart. J. Math. Oxford, (2) **24** (1973), 343–356.
- [2] L. Hodgkin: *On the K-theory of Lie groups*, Topology, **6** (1967), 1–36.
- [3] L. Hodgkin: *The equivariant Künneth theorem in K-theory*, Lecture Notes in Math. **496** (1975), 1–101.
- [4] T. Kambe: *The structure of K_Λ -rings of the lens space and their applications*, J. Math. Soc. Japan, **18** (1966), 135–146.

- [5] H. Minami: *On the K -theory of PE_6* , Osaka J. Math. **32** (1995), 1113–1130.
- [6] H. Minami: *On the K -theory of the projective symplectic groups*, Publ. RIMS, Kyoto Univ. **31** (1995), 1045–1063.

Department of Mathematics
Nara University of Education
Takabatake, Nara 630-8528, Japan

