Osaka University Knowledge Archive

Title	Hyperbolic lengths of some filling geodesics on Riemann surfaces with punctures
Author(s)	Zhang, Chaohui
Citation	Osaka Journal of Mathematics. 2008, 45(3), p. 773-787
Version Type	VoR
URL	https://doi.org/10.18910/7269
rights	
Note	

Osaka University Knowledge Archive : OUKA

https://ir.library.osaka-u.ac.jp/

Osaka University

Zhang, C.
Osaka J. Math.
45 (2008), 773–787

HYPERBOLIC LENGTHS OF SOME FILLING GEODESICS ON RIEMANN SURFACES WITH PUNCTURES

CHAOHUI ZHANG

(Received September 11, 2006, revised August 16, 2007)

Abstract

Let $\tilde{S}$ be a Riemann surface of type (p, n) with $3p - 3 + n > 0$ and $n \geq 1$. In this paper, we give a quantitative common lower bound for the hyperbolic lengths of all filling geodesics on $\tilde{S}$ generated by two parabolic elements in the fundamental group $\pi_1(\tilde{S}, a)$.

1. Introduction

Let $\tilde{c}$ be a non-trivial closed curve on a Riemann surface $\tilde{S}$ of type (p, n) with $3p - 3 + n > 0$. The length function

$$l_{\tilde{c}} \colon T(\tilde{S}) \to \mathbb{R}^+$$

on the Teichmüller space $T(\tilde{S})$ is defined by sending each hyperbolic structure $\sigma = \sigma(\tilde{S})$ of $T(\tilde{S})$ to the hyperbolic length $l_{\tilde{c}}(\sigma)$ of the closed geodesic homotopic to $\tilde{c}$ on $\sigma(\tilde{S})$.

It is well known ([11]) that the function $l_{\tilde{c}}$ achieves its positive minimum value when $\tilde{c}$ is a filling curve on $\tilde{S}$ in the sense that every component of $\tilde{S} \setminus \{\tilde{c}\}$ is either a disk or a once punctured disk. The extremal value, of course, depends only on the homotopy class of $\tilde{c}$. In this paper, we give a quantitative common lower bound through $T(\tilde{S})$ for the hyperbolic lengths of certain kind of filling curves on $\tilde{S}$.

Note that if $\tilde{S}$ contains punctures, then some elements $[\alpha]$ in the fundamental group $\pi_1(\tilde{S}, a)$, $a \in \tilde{S}$, are represented by loops α that pass through a and are boundaries of once punctured disks. Let $\mathcal{F}$ denote the set of those elements. $\mathcal{F} \subset \pi_1(\tilde{S}, a)$.

The main result of this paper is the following:

Theorem 1. *Let $\tilde{S}$ be a Riemann surface of type (p, n) with $3p - 3 + n > 0$ and $n \geq 1$. There are infinitely many homotopically independent filling curves $\tilde{c}$ on $\tilde{S}$ that can be expressed as products of two elements in $\mathcal{F}$. For each such $\tilde{c}$ and each hyperbolic structure σ on $\tilde{S}$, we have*:

$$l_{\tilde{c}}(\sigma) \geq 2\log(\kappa^2 - 5) - 4\log 2, \tag{1.1}$$

2000 Mathematics Subject Classification. Primary 32G15; Secondary 30F60.

where $\kappa = 16p + 8n - 21$ *if* $n \geq 3$; $16p + 3$ *if* $n = 2$; *and* $16p + 7$ *if* $n = 1$.

REMARK. From the definition of $\tilde{S}$ we know that $p \geq 0$ if $n \geq 4$; $p \geq 1$ if $n = 1, 2, 3$.

Let $\mathbb{H} = \{z \in \mathbb{C};\ \operatorname{Im} z > 0\}$ denote the upper half plane with the hyperbolic metric $\rho_{\mathbb{H}}$ given by

$$\rho_{\mathbb{H}}(z)|dz| = \frac{|dz|}{\operatorname{Im} z}. \tag{1.2}$$

Let $\varrho\colon \mathbb{H} \to \tilde{S}$ be the universal covering with a covering group G. G is a torsion free Fichsian group of the fist kind of type (p, n) so that $\mathbb{H}/G \cong \tilde{S}$. The set $\mathcal{F}$ is one-to-one correspondent with the set of parabolic elements of G.

Note that any hyperbolic element g is conjugate in $\mathrm{PSL}_2(\mathbb{R})$ to $z \mapsto \lambda_g\ z$, where $\lambda_g > 1$ is called the multiplier of g. Let A_g be the axis of g. Using (1.2) one calculates the hyperbolic length of $\varrho(A_g)$ is $\log \lambda_g$.

The hyperbolic element g is called essential if every component of $\tilde{S} \setminus \varrho(A_g)$ is either a disk or possibly a once punctured disk. Theorem 1 can be restated as follows:

Theorem 1′. *Let G be a finitely generated Fuchsian group of the first kind of type (p, n) with $3p-3+n > 0$ and $n \geq 1$. There are infinitely many essential hyperbolic elements g of G that are generated by two parabolic elements of G. Furthermore, for each such element g, the multiplier λ_g of g satisfies*:

$$\lambda_g \geq \left\{\frac{1}{4}(\kappa^2 - 5)\right\}^2, \tag{1.3}$$

where κ is given in Theorem 1.

2. Dilatations of pseudo-Anosov maps generated by two positive multi-twists

We first recall the definition and some basic properties of Teichmüller space $T(R)$ of a Riemann surface R of type (p, n), $3p - 3 + n > 0$. For more details see [3, 4, 8].

We define an equivalence class $[\sigma]$ of a conformal structure σ on $\tilde{S}$ as follows. Two conformal structures σ_1 and σ_2 on R are called strongly equivalent if there is an isometry h of $\sigma_1(R)$ onto $\sigma_2(R)$ such that $\sigma_2^{-1} \circ h \circ \sigma_1$, as a self-map of the underlying surface R, is isotopic to the identity. The collection of strong equivalence classes $[\sigma]$ of conformal structures σ form a Teichmüller space $T(R)$. $T(R)$ is naturally equipped with a complex structure.

Let $[\sigma_1]$ and $[\sigma_2]$ be two points in $T(R)$. Let $h\colon \sigma_1(R) \to \sigma_2(R)$ be a quasi-conformal mapping. Define the complex dilation $\nu(z) = \partial_{\bar{z}} h(z)/\partial_z h(z)$ and denote $\|\nu\| =$

ess.sup$\{|\nu(z)|;\ z \in \sigma_1(R)\}$. By definition, $\|\nu\| < 1$. The maximal dilatation $K(h)$ is defined by

$$K(h) = \frac{1 + \|\nu\|}{1 - \|\nu\|}.$$

The Teichmüller distance between $[\sigma_1]$ and $[\sigma_2]$ is defined as

$$\langle [\sigma_1], [\sigma_2] \rangle = \frac{1}{2} \inf \log K(h),$$

where $h\colon \sigma_1(R) \to \sigma_2(R)$ runs over all maps homotopic to $\sigma_2 \circ \sigma_1^{-1}$. $T(R)$ is a complete metric space with respect to the Teichmüller metric.

The mapping class group (or modular group) Mod_R is the group of isotopy classes f^* of self-maps f of R. f^* acts on $T(R)$ by sending each $[\sigma]$ to $[\sigma \circ f^{-1}]$. Mod_R is a group of isometries with respect to the Teichmüller metric defined above.

An element f^* of Mod_R is called hyperbolic if $\langle [\sigma], f^*[\sigma] \rangle$ assumes a positive minimum value at a point $[\sigma_0]$ in $T(R)$. An isometric image of the real line $\mathbb{R}$ into $T(R)$ is called a Teichmüller geodesic. Similarly, an isometric image of the unit disk $D = \{z \in \mathbb{C};\ |z| < 1\}$ into $T(R)$ is called a Teichmüller disk. According to Bers [4] and Kra [8], f^* is hyperbolic if and only if f^* keeps a Teichmüller geodesic l invariant, which is also equivalent to that f^* keeps a Teichmüller disk D invariant. In this case, for any point $[\sigma] \in l$, we let $[\sigma]$ be represented by R. Then f^* can be realized on R as an absolutely extremal Teichmüller mapping $f_0\colon R \to R$. Let $K(f_0)$ (> 1) denote the maximal dilatation of f_0.

Associated to l (or D), there is a integrable meromorphic quadratic differential ϕ that defines a singular Euclidean metric $|\phi|$ on R. Thus it determines a pair of singular measured foliation $(\mathcal{F}_h, \mathcal{F}_v)$, where $\mathcal{F}_h$ and $\mathcal{F}_v$ are horizontal and vertical foliations respectively.

The absolutely extremal map f_0 takes the two singular foliations into themselves. Away from all singularities, the map stretches the horizontal leaves by the stretching factor $\lambda(f_0) = K(f_0)^{1/2}$, and compress the vertical leaves by the factor $1/\lambda(f_0) = K(f_0)^{-1/2}$. By using the language of Thurston [15], such a map f_0 is also called a pseudo-Anosov diffeomorphism. We use the notations $l_\phi = l$ and $D_\phi = D$ to emphasis that those D and l are determined by ϕ.

In the homotopy class f^* of f, f_0 is a unique pseudo-Anosov diffeomorphism. So the action of f^* on $T(R)$ is analogous to the action of a hyperbolic Möbius transformation on $\mathbb{H}$. In particular, for each hyperbolic element f^*, there is a unique invariant Teichmüller geodesic and a unique invariant Teichmüller disk in $T(R)$.

Let $A = \{\alpha_1, \dots, \alpha_n\}$ and $B = \{\beta_1, \dots, \beta_m\}$, $n \geq 1$ and $m \geq 1$ be collections of disjoint simple closed geodesics on R. We assume that A and B intersect minimally, and $A \cup B$ fills R in the sense that every non-trivial loop on R intersects with either A or B or both. Let t_A and t_B denote the positive multi-twists along some elements of A and B,

respectively. $A \cup B$ is regarded as a graph on R. According to Proposition 6.4 of [9], when $A \cup B$ is dominant (see [9] for the definition), all elements in $\langle t_A, t_B \rangle$ except conjugates of powers of t_A^n, t_B^m, and possibly of $(t_A \circ t_B)^n$ and $(t_B \circ t_A)^n$, $n, m \in \mathbb{Z}$, are pseudo-Anosov maps. $(t_A \circ t_B)^n$ and $(t_B \circ t_A)^n$ are not pseudo-Anosov if the graph $A \cup B$ is critical (see also [9]).

Let $D_\phi \subset T(R)$ be a Teichmüller disk. We consider the stabilizer $\mathrm{Stab}(D_\phi)$ in Mod_R, which is called a Veech group on a surface in D_ϕ. Since the Teichmüller disk is isometrically the same as $\mathbb{H}$, $\mathrm{Stab}(D_\phi)$ actually determines a subgroup of $\mathrm{PSL}_2(\mathbb{R})$. Thus there defines a map

$$\mathcal{D}\colon\ \mathrm{Stab}(D_\phi) \to \mathrm{PSL}_2(\mathbb{R}).$$

By the main theorem of [10], there exist Teichmüller disks D_ϕ and D_ψ such that $t_A \in \mathrm{Stab}(D_\phi)$ and $t_B \in \mathrm{Stab}(D_\psi)$ and they determine elements $\mathcal{D}(t_A)$ and $\mathcal{D}(t_B)$ in $\mathrm{PSL}_2(\mathbb{R})$. Furthermore, t_A and t_B also act on a common Teichmüller disk D with respect to the flat structure constructed from the dual graph of $A \cup B$.

Define $N = N_{A,B}$ to be the $n \times m$ matrix whose (i, j)-entry is $i(\alpha_i, \beta_j)$, the intersection number of α_i and β_j, where $\alpha_i \in A$ and $\beta_j \in B$. By assumption, $A \cup B$ fills R. It follows that the graph defined by $A \cup B$ is connected. Hence NN^t is irreducible. Let $\mu(NN^t)$ be the maximum of moduli of the eigenvalues of NN^t (called the Perron-Frobenius eigenvalue in the literature), and set $\mu(A \cup B) = \sqrt{\mu(NN^t)}$. By [9, 16], we have the representations:

$$\mathcal{D}(t_A) = \begin{pmatrix} 1 & \mu(A \cup B) \\ 0 & 1 \end{pmatrix}$$

and

$$\mathcal{D}(t_B) = \begin{pmatrix} 1 & 0 \\ -\mu(A \cup B) & 1 \end{pmatrix}.$$

Let $\mu = \mu(A \cup B)$. The group $\langle \mathcal{D}(t_A), \mathcal{D}(t_B) \rangle$ generated by $\mathcal{D}(t_A)$ and $\mathcal{D}(t_B)$ is a discrete subgroup of $\mathrm{PSL}_2(\mathbb{R})$ if $\mu > 2$. In this case, $\langle \mathcal{D}(t_A), \mathcal{D}(t_B) \rangle$ is free of rank 2. Denote $\mathcal{M} = \mathbb{H}/\langle \mathcal{D}(t_A), \mathcal{D}(t_B) \rangle$. By Lemma 6.3 of [9], $\mathcal{M}$ has infinite area and its convex core is a twice punctured disk. These results will lead to the following Lemma 1.

Let ϵ_μ be the larger root of the quadratic equation

$$x^2 + (2 - \mu^2)x + 1 = 0.$$

That is,

$$\epsilon_\mu = \frac{1}{2}(\mu^2 - 2 + \mu\sqrt{\mu^2 - 4}).$$

It is easy to check that ϵ_μ is an increasing function with respect to μ. Notice that different metric scales were used in different papers, we need to review those arguments

presented in [9] and [8] for the sake of consistency.

Let f be a pseudo-Anosov element in $\langle t_A, t_B \rangle$. f determines a Teichmüller disk D isometric to $\mathbb{H}$ with respect to the Teichmüller metric on D and the hyperbolic metric on $\mathbb{H}$. Thus f induces a hyperbolic Möbius transformation $\mathcal{D}(f)$ on $\mathbb{H}$. Denote $F = \mathcal{D}(f)$. Let τ denote the translation length of F:

$$\tau = \inf_z \rho_{\mathbb{H}}(z, F(z)).$$

From Section 7.34 of Beardon [2], we know that

$$\frac{1}{2}|\mathrm{trace}(F)| = \cosh\frac{\tau}{2} = \frac{\exp(\tau/2) + \exp(-\tau/2)}{2}. \tag{2.1}$$

By isometry we obtain

$$\frac{1}{2}\log K(f) = \tau. \tag{2.2}$$

Since $\lambda(f) = K(f)^{1/2}$, from (2.2), we get

$$\log \lambda(f) = \tau. \tag{2.3}$$

Let $\xi = \exp(\tau/2)$. A simple calculation shows that ξ satisfies

$$\xi^2 - |\mathrm{trace}(F)|\xi + 1 = 0.$$

By Lemma 6.3 of [9], $\xi \geq \epsilon_\mu$, i.e., $\exp(\tau/2) \geq \epsilon_\mu$, or $\tau/2 \geq \log \epsilon_\mu$. It follows that

$$\tau \geq 2\log \epsilon_\mu.$$

Together with (2.3), we obtain

$$\log \lambda(f) \geq 2\log \epsilon_\mu.$$

Hence, we have $\lambda(f) \geq \epsilon_\mu^2$. We summarize the result in the following lemma.

Lemma 1 (Leininger [9]). *Assume that $\mu > 2$. For any pseudo-Anosov element f of $\langle t_A, t_B \rangle$, we have that*

$$\lambda(f) \geq \epsilon_\mu^2.$$

REMARK. Due to different metric scales the original result stated in [9] takes the inequality $\lambda(f) \geq \epsilon_\mu$.

Since $\lambda(f) = K(f)^{1/2}$, from Lemma 1, we obtain:

$$K(f) \geq \left\{\frac{1}{2}(\mu^2 - 2 + \mu\sqrt{\mu^2 - 4})\right\}^4. \tag{2.4}$$

3. Translation lengths of essential hyperbolic elements

Let $\tilde{S}$ be as in the introduction. Let $a \in \tilde{S}$ and let $S = \tilde{S} \setminus \{a\}$. S is of type $(p, n+1)$. Associated to $T(\tilde{S})$ there is a fiber space $F(\tilde{S})$ defined as follows. For each $[\nu] \in T(\tilde{S})$, by Ahlfors-Bers [1], there is normalized quasiconformal automorphism w^ν of the complex plane $\mathbb{C}$ such that the restriction $w^\nu|_{\mathbb{H}^*}$ to the lower half plane $\mathbb{H}^*$ is conformal, and its Beltrami coefficient $\partial_{\bar{z}} w^\nu / \partial_z w^\nu$ on $\mathbb{H}$ projects to a conformal structure that determines $[\nu]$. We form the Bers fiber space

$$F(\tilde{S}) = \{([\nu], z);\ [\nu] \in T(\tilde{S}),\ z \in w^\nu(\mathbb{H})\}.$$

Note that in this setting $\mathbb{H}$ is considered the central fiber and the group G acts on $F(\tilde{S})$ in a natural manner.

In [3], Bers established an isomorphism $\varphi\colon F(\tilde{S}) \to T(S)$ that is unique up to a modular transformation on $T(S)$.

The isomorphism φ determines an embedding φ^* of G into the mapping class group Mod_S such that each element in the image $\varphi^*(G)$ projects to the trivial mapping class on $\mathrm{Mod}_{\tilde{S}}$ defined by adding the puncture a back into S. Conversely, Bers [3] also showed that if a mapping class θ can be projected to the trivial one in $\mathrm{Mod}_{\tilde{S}}$, then θ lies in the image $\varphi^*(G)$.

It was shown in [8, 12] that $g \in G$ is parabolic if and only if $g^* = \varphi^*(g)$ is induced by a Dehn twist along a boundary curve $\partial\Delta$, where Δ is a twice punctured disk on S enclosing a and another puncture b, where b is regarded as a puncture of $\tilde{S}$ that is determined by the conjugacy class of g. In [8] Kra also proved that g is essential (that is, the complement of the projection of its axis A_g consists of disks and possibly once punctured disks) if and only if g^* is a pseudo-Anosov mapping class in Mod_S. By abuse of language, in the sequel we denote by $K(g^*)$ the dilatation of the corresponding absolutely extremal map on a surface S that realizes the mapping class g^*.

Let $g \in G$ be an essential hyperbolic element with axis A_g. For simplicity we denote $\tilde{c} = \varrho(A_g) \subset \tilde{S}$. Then $\tilde{c}$ is a filling geodesic on $\tilde{S}$ and by a theorem of [11], the length function $l_{\tilde{c}}\colon T(\tilde{S}) \to \mathbb{R}^+$ attains a minimum value at $[\mu_0] \in T(\tilde{S})$. We may assume that $[\mu_0] = [0]$. We need to see how the number $K(g^*)$ is dominated by $l_{\tilde{c}}([0])$. We review:

Lemma 2 (Kra [8]). *With the conditions above, we let λ_g denote the multiplier of g, i.e., g is conjugate to $z \mapsto \lambda_g z$. Then*

$$K(g^*) \leq \lambda_g^2. \tag{3.1}$$

Outline of proof. For any point $x \in A_g \subset \mathbb{H}$, the translation length $\rho_{\mathbb{H}}(x, g(x)) = l_{\tilde{c}}([0])$. By Royden's theorem [13] (see also Earle-Kra [6]), the Teichmüller metric on $T(S)$ coincides with the Kobayashi metric on $T(S)$. Since the restriction $\varphi|_{\mathbb{H}}\colon \mathbb{H} \to T(S)$ is holomorphic, it cannot increase the distance. Therefore,

$$\langle \varphi(x), g^* \circ \varphi(x)\rangle = \langle \varphi(x), \varphi(g(x))\rangle \leq \rho_{\mathbb{H}}(x, g(x)). \tag{3.2}$$

By definition, the translation length of g^* is no larger than the distance $\langle \varphi(x), g^* \circ \varphi(x)\rangle$. It follows from (3.2) that

$$\begin{aligned}\frac{1}{2}\log K(g^*) &\leq \langle \varphi(x), g^* \circ \varphi(x)\rangle \leq \rho_{\mathbb{H}}(x, g(x))\\ &= l_{\tilde{c}}([0]) = \int_1^{\lambda_g} \frac{1}{y}\, dy = \log \lambda_g.\end{aligned}$$

It follows that

$$K(g^*) \leq \lambda_g^2, \tag{3.3}$$

as asserted. □

Let $A = \{\alpha_1, \ldots, \alpha_n\}$ and $B = \{\beta_1, \ldots, \beta_m\}$, $n \geq 1$ and $m \geq 1$, be defined in Section 2. First we consider the case that A and B are restricted to contain only one element (the case of $n = m = 1$). In this case, it is well known (see [7]) that there are pseudo-Anosov maps f on S that are not represented by elements in the group $\langle t_A, t_B\rangle$ generated by t_A and t_B. In general case, it is not completely clear whether every essential element $g \in G$, g^* is in the group $\langle t_A, t_B\rangle$. However, there exist infinitely many pairs $\{\alpha_1, \beta_1\}$ so that the group $\langle t_{\alpha_1}, t_{\beta_1}\rangle$ contains pseudo-Anosov mapping classes of forms g^*, where $g \in G$ is essential. In particular, there are infinitely many pairs $\{\alpha_1, \beta_1\}$, where α_1 and β_1 are peripheral on $\tilde{S}$, so that $\langle t_{\alpha_1}, t_{\beta_1}\rangle$ contains pseudo-Anosov mapping classes of form g^* for $g \in G$ being essential hyperbolic.

By combining Lemma 1 and Lemma 2, we can readily obtain the following lemma:

Lemma 3. *Assume that a hyperbolic element $g \in G$ is essential and $g^* \in \langle t_A, t_B\rangle$ for certain $A = \{\alpha_1, \ldots, \alpha_n\}$ and $B = \{\beta_1, \ldots, \beta_n\}$, $n \geq 1$ and $m \geq 1$. Then $A \cup B$ fills S and*

$$\mu(A \cup B) \leq \max\left\{2, \frac{1}{2}\left(1 + \sqrt{5 + 4\sqrt{\lambda_g}}\right)\right\}.$$

Proof. Denote $\mu = \mu(A \cup B)$. By Lemma 1 and (3.3), we have

$$\lambda_g^2 \geq K(g^*) = \lambda(g^*)^2 \geq \left\{\frac{1}{2}(\mu^2 - 2 + \mu\sqrt{\mu^2 - 4})\right\}^4.$$

Note that $\lambda_g = \exp(l_{\tilde{c}}[0])$. We assume that $\mu > 2$. Then clearly $\mu - 2 \leq \sqrt{\mu^2 - 4}$. It follows that

$$\mu^2 - \mu - (1 + \sqrt{\lambda_g}) \leq 0.$$

The lemma then follows immediately. □

REMARK. The estimation in the above lemma can be sharpened by applying a theorem of [5] that states that if $\mu > 2$ then in fact $\mu > 2.0065936$. This implies that there is an integer N such that

$$\mu - \left(2 - \frac{1}{N}\right) \leq \sqrt{\mu^2 - 4}.$$

A simple calculation shows that $N \geq 7$.

4. Peripheral simple curves on punctured Riemann surfaces

In this section we prove that there are infinitely many essential elements $g \in G$ generated by two parabolic elements.

Let $\tilde{S}$ be of type (p, n) with $3p - 3 + n > 0$ and $n \geq 1$. Let $a \in \tilde{S}$ and $S = \tilde{S} \setminus \{a\}$. Then S is of type $(p, n + 1)$. Let $x_1 = a, x_2, \ldots, x_{n+1}$, $n \geq 1$, denote the punctures of S. Let $\mathcal{P}(S, a)$ denote the set of equivalence classes of paths α on S connecting a and another puncture, where two paths α_1 and α_2 are considered equivalent if they are homotopic to each other by a homotopy fixing the end punctures. Let $\mathcal{E}(S, a)$ denote the set of equivalence classes of twice punctured disks on S that enclose a and another puncture, where two such disks are equivalent if their boundary curves are homotopic to each other without interfering with any other punctures.

Given a path representative $\alpha \in \mathcal{P}(S, a)$, we can always fatten α, giving rise to an element in $\mathcal{E}(S, a)$. Conversely, for every element $\Delta \in \mathcal{E}(S, a)$, there is a path α connecting the two end punctures and lying entirely in Δ. α is unique up to a homotopy, i.e., any two such paths are homotopic within Δ and fix the end punctures. We thus obtain a bijection:

$$j : \mathcal{P}(S, a) \to \mathcal{E}(S, a). \tag{4.1}$$

two elements $\alpha, \beta \in \mathcal{P}(S, a)$ are called to fill S if every component of $S \setminus \{\alpha, \beta\}$ is either a disk or a once punctured disk. We need the following lemmas.

Lemma 4. *Let $\alpha, \beta \in \mathcal{P}(S, a)$ and assume that $\{\alpha, \beta\}$ fills S, then $\{\partial j(\alpha), \partial j(\beta)\}$ must also fill S in a regular sense.*

Proof. Denote $\Delta_\alpha = j(\alpha)$ and $\Delta_\beta = j(\beta)$. It is easy to see that $\Delta_\alpha \cap \Delta_\beta$ consists of quadrilateral (that are homeomorphic to disks) and two or one punctured disk components (depending on whether or not α and β share both end punctures). The rest of



Fig. 1.



Fig. 2.

components in $\Delta_\alpha \cup \Delta_\beta$ include components of $\Delta_\alpha \setminus \Delta_\beta$ and components of $\Delta_\beta \setminus \Delta_\alpha$, all of which are homeomorphic to disks. The remaining components of $S \setminus \{\Delta_\alpha \cup \Delta_\beta\}$ are essentially the same as the components in $\overline{S} \setminus \{\alpha \cup \beta\}$ which are either disks or punctured disks. This proves Lemma 4. $\square$

The following lemma comes from referee's comments:

Lemma 5. *Let S be of type $(p, n+1)$, $3p + n > 3$, $n \geq 1$. There are infinitely many pairs (α, β) of paths in $\mathcal{P}(S, a)$ so that $\{\alpha, \beta\}$ fills S.*

Proof. Observe that S can be thought of as a Riemann sphere with p handles and n punctures. Let H be a handle with $(\partial D_0, \partial D_0')$ the two boundary components. Let γ, δ be two curves on H that are not to be homotopic and $\{\gamma, \delta\}$ fills H. Note that γ can be winded around δ as many time as possible. The end points of γ are denoted by s, t, and the end points of δ are denoted by u, v. See Fig. 1.

We remove p pairs (D_i, D_i') of small disks and $n+1$ points $x_1 = a, x_2, \ldots x_{n+1}$, $n \geq 1$, from the Riemann sphere $\mathbb{S}^2$, obtaining S_0. S_0 is drawn in Fig. 2 in the case that p is even (if p is odd, the positions of D_p and D_p in Fig. 2 are switched).

For $i = 1, \ldots, p$, let (u_i, s_i) and (v_i, t_i) be pairs of marked points on ∂D_i and $\partial D_i'$ respectively. Paste p copies of H to S_0 in such a way that $(\partial D_0, \partial D_i)$ and $(\partial D_0', \partial D_i')$ are glued together with $u_i = u$, $v_i = v$, $s_i = s$, and $t_i = t$. Then we can define $\alpha \in \mathcal{P}(S, a)$ as follows. Connect $a = x_1$ and s_1, followed with γ, then connect t_1 and s_2, and then followed with γ again, and so forth. After p steps, we connect t_p and x_{n+1} by a path away from all punctures other than the end punctures. Similarly, we can define $\beta \in \mathcal{P}(S, a)$ to be a path that goes from $a = x_1$ to v_1, followed with the inverse



Fig. 3.

δ^{-1} of δ, then go to u_1, then connect u_1 and v_2, and so forth. After p steps, we draw a path connecting u_p to x_{n+1} in such a way that the component of $S\backslash\{\alpha, \beta\}$ that includes x_i, $i = 2, \ldots x_n$, is a once punctured disk (this could occur only when $n \geq 3$). Fig. 3 below shows the case that p is even and the two paths α and β are in $\mathcal{E}(S, a)$. One can easily check that any component of $S \setminus \{\alpha, \beta\}$ is either a disk or a once punctured disk, which says $\{\alpha, \beta\}$ fills S. □

From Lemma 4 and Lemma 5, one obtains:

Lemma 6. *There are infinitely many essential elements $g \in G$ that are generated by two parabolic elements.*

Proof. From the construction there are infinitely many pairs (α, β) that fills S. According to Lemma 4, there are infinitely many pairs $(j(\alpha), j(\beta))$ that fill S. By Lemma 3 of [17], there are parabolic elements T_i, $i = 1, 2$, so that $\varphi^*(T_1) = \partial j(\alpha)$ and $\varphi^*(T_2) = \partial j(\beta)$. Since $\partial j(\alpha)$ and $\partial j(\beta)$ are homotopic to trivial loops as a is filled in, we see that any finite product

$$\prod_i \left(t_{\partial j(\alpha)}^{n_i} \circ t_{\partial j(\beta)}^{m_i}\right), \quad n_i, m_i \in \mathbb{Z}, \tag{4.2}$$

projects to a trivial mapping class. It follows from Bers [4] that (4.2) is of form $\varphi^*(g)$ for an essential element $g \in G$. Clearly, g is generated by T_1 and T_2. This proves the lemma. □

5. Minimal intersections of two peripheral curves

In the previous section we constructed two curves α and β that are boundaries of twice punctured disks enclosing a. In this section we give an estimate of lower bound of intersections of α and β. We first prove:

Lemma 7. *Let $\alpha, \beta \in \mathcal{P}(S, a)$. Suppose that $\{\alpha, \beta\}$ fills S. Then in addition to a and another end puncture, α intersects with β at least $2p - 3 + n$ points. In particular, if $n = 1, 2$, then α intersects with β at least $2p$ points.*

Proof. Note that if punctures on S are considered distinguished points on the compactification $\overline{S}$ of S, $\alpha \cup \beta$ defines a graph on S with a number E of edges, a number F of vertices, and a number V of vertices. We know that the Euler characteristic $\chi(\overline{S}) = 2 - 2p$.

Assume that in addition to a and possible another end point, α intersects with β k times. Let n' be the number of once punctured disk components of $S \setminus \{\alpha, \beta\}$. If $k = 0$, then $\alpha \cup \beta$ is a binary tree or a circle. In former case, $E = 2$, $V = 3$. Since $F + V - E = 2 - 2p$, $F = 1 - 2p$, which implies that $F = 1$. In order for α, β to fill S, we must have $n' \leq 1$ and S is of type $(0, n + 1)$ for $n \leq 3$, contradicting to our hypothesis.

In later case, $E = 2$ and $V = 2$. Since $F + V - E = 2 - 2p$, $F = 2 - 2p$. This implies that $F = 2$. So we must have $n' \leq 2$, and S is of type $(0, n + 1)$ for $n \leq 3$. Again, this is a contradiction.

Now we assume that $k > 0$ and that all the intersections are distinct. There are two cases to consider.

CASE 1. α and β share only one endpoint a. In this case, we have $V = k + 3$, $E = 2(k+1)$. Since $\alpha \cup \beta$ fills S, $F \geq n'$, where $n' + 3 = n + 1$. Now from $\chi(\overline{S}) = 2 - 2p$ we obtain

$$2 - 2p = V + F - E \geq (k + 3) + (n - 2) - 2(k + 1).$$

It follows that $k \geq 2p + n - 3$.

CASE 2. α and β share both end punctures. In this case, $\alpha \cup \beta$ is closed when the two endpoints are added. We must have $V = k + 2$, $E = 2(k + 1)$ and $F \geq n'$, where $n' + 2 = n + 1$. Hence

$$2 - 2p = V + F - E \geq (k + 2) + (n - 1) - 2(k + 1).$$

It follows that $k \geq 2p + n - 3$.

In the case of $n = 1, 2$, the author was informed by the referee that $k \geq 2p$. In fact, we first assume that $n = 1$. Then α and β share the same end punctures. So $\alpha \cup \beta$ is a closed when the two endpoints are added. Since each component of $S \setminus \{\alpha, \beta\}$ is a disk, $S \setminus \{\alpha, \beta\}$ is not connected. Hence $F \geq 2$. Recall that $V = k + 2$, and $E = 2(k + 1)$. We have

$$2 - 2p = V + F - E \geq (k + 2) + 2 - 2(k + 1).$$

It follows that $k \geq 2p$. In the case of $n = 2$, when α, β share the same end punctures, by the same argument as above, we have $k \geq 2p$. Otherwise, we assume that α terminates x_2 and β terminates x_3 with $x_2 \neq x_3$. Then $V = k + 3$ and $E = 2(k + 1)$. Since $F \geq 1$, we have

$$2 - 2p = V + F - E \geq (k + 3) + 1 - 2(k + 1).$$

Thus $k \geq 2p$, as asserted. □

Let $\#\{c_1, c_2\}$ denote the set of the minimal intersection points of arbitrary two curves c_1, c_2 on S, and $i(c_1, c_2)$ the intersection number of c_1 and c_2. We have

Lemma 8. *Let α and β be defined as in* Lemma 4. *Then any point in $\#\{\alpha, \beta\}$ other than end punctures of α and β contributes at least* 4 *intersection points to $\#\{c_1, c_2\}$ for $c_1 = \partial j(\alpha)$, and $c_2 = \partial j(\beta)$.*

Proof. We only handle the case that α and β share both end punctures, as drawn in Fig. 3. Let y_i be such an intersection in $\#\{\alpha \cap \beta\}$. By hypothesis, $b = x_{n+1}$ is the other endpoint of α and β, respectively. Let $c_1' \sim c_1$, $c_2' \sim c_2$ be representatives of $\partial j(\alpha)$ and $\partial j(\beta)$, respectively. Assume that c_1' and c_2' are very close to α and β respectively. Observe that y_i contributes 4 intersections to $\#\{c_1', c_2'\}$. In fact, the intersection near y_i is a quadrilateral. Then the lemma follows from the fact that a homotopy does not decrease the intersection number. □

Together with Lemma 6, Lemma 7, and Lemma 8, we are able to prove the following:

Lemma 9. *Let $\tilde{S}$ be of type (p, n) with $3p-3+n > 0$ and $n \geq 1$. Then there are infinitely many pairs (c_1, c_2) of simple closed curves on S with the following properties*:
(1) *$c_1 = \partial\Delta_1$ and $c_2 = \partial\Delta_2$ for $\Delta_1, \Delta_2 \in \mathcal{E}(S, a)$,*
(2) *$\{c_1, c_2\}$ fills S in the regular sense, and*
(3) *the intersection number $i(c_1, c_2) \geq 8p+4n-10$ if $n \geq 3$; $i(c_1, c_2) \geq 8p+4$ if $n = 1$; and $i(c_1, c_2) \geq 8p+2$ if $n = 2$.*

Proof. First we consider the case of $n \geq 3$. If α, β share only one end puncture a, by Lemma 7, there are at least $2p-3+n$ distinct intersection points y_i in $\#\{\alpha, \beta\}$. By Lemma 8, each y_i, $1 \leq i \leq 2p-3+n$, contributes at least 4 intersections to $\#\{c_1, c_2\}$. The puncture a contributes at least 2 intersections in $\#\{c_1, c_2\}$. Therefore,

$$i(c_1, c_2) \geq 4(2p + n - 3) + 2 = 8p + 4n - 10.$$

If α and β share both end punctures (a and b), by Lemma 7 again, α and β cross at least $2p - 3 + n$ times. Let y_i, $1 \leq i \leq 2p - 3 + n$ denote these intersections. By Lemma 8, each y_i contributes at least 4 intersections to $\#\{c_1, c_2\}$, The punctures a and b each contributes at least 2 intersections to $\#\{c_1, c_2\}$. We conclude that

$$i(c_1, c_2) \geq 4(2p + n - 3) + 2 + 2 = 8p + 4n - 8.$$

It follows that $i(c_1, c_2) \geq 8p + 4n - 10$ if $k \geq 3$.

If $n = 1$, then S has only two punctures and $\alpha \cup \beta$ has to be closed as the two end punctures are filled in. By Lemma 7 and Lemma 8, we have $i(c_1, c_2) \geq 4(2p) + 4 = 8p + 4$. If $n = 2$, then S has three punctures. By Lemma 7 and Lemma 8 again, we have $i(c_1, c_2) \geq 4(2p) + 2$ if other than a α and β have different terminal punctures; and $i(c_1, c_2) \geq 4(2p) + 4$ if α and β have the same end punctures ($\alpha \cup \beta$ is closed as the end punctures are filled in). Overall we have $i(c_1, c_2) \geq 4(2p) + 2$ if $n = 2$. This proves the lemma. □

6. Proof of Theorem 1′

The fact that there are infinitely many essential elements g of G that are generated by two parabolic elements was proved in Section 4. Let $g \in G$ be an essential element generated by two parabolic elements T_1 and T_2. Let $t_1 = \varphi^*(T_1)$ and $t_2 = \varphi^*(T_2)$. By Theorem 2 of [8, 12], t_1 and t_2 are Dehn twists along c_1 and c_2 for $c_1 = \partial\Delta_1$ and $c_1 = \partial\Delta_2$, where $\Delta_1, \Delta_2 \in \mathcal{E}(S, a)$.

We remark that in our situation the fixed point z_i of T_i, $i = 1, 2$, cannot be vertices of a common fundamental region of G. For otherwise, let $\omega\colon G \to \pi_1(\tilde{S}, a)$ denote a canonical isomorphism. Then we have that $\omega(T_1^{\pm 1} \circ T_2^{\pm 1})$ is either a simple loop bounding 2 punctures of $\tilde{S}$, or a "figure 8" loop on $\tilde{S}$ that is not a filling loop unless $\tilde{S}$ is of type $(0, 3)$, which has been excluded by our assumption.

From Lemma 9, the intersection number $i(c_1, c_2) \geq \kappa_0$, where $\kappa_0 = 8p + 4n - 10$ if $n \geq 3$; $\kappa_0 = 8p + 4$ if $n = 1$; and $\kappa_0 = 8p + 2$ if $n = 2$. Since $A = \{c_1\}$ and $B = \{c_2\}$ consist single element, we have $n = m = 1$ in the discussion of Section 2. Hence by definition, $\mu(NN^t) = i(c_1, c_2)^2$. Thus

$$\mu(A \cup B) = \sqrt{\mu(NN^t)} = i(c_1, c_2) \geq \kappa_0. \tag{6.1}$$

In particular, since $p \geq 0$, and $p \geq 1$ if $n \leq 3$, we see that $\kappa_0 > 2$. It follows that $\mu(A \cup B) > 2$. Since $\lambda_g > 1$,

$$\frac{1}{2}\left(1 + \sqrt{5 + 4\sqrt{\lambda_g}}\right) > 2.$$

Now from Lemma 3 along with (6.1), we have that

$$\kappa_0 \leq \mu(A \cup B) \leq \frac{1}{2}\left(1 + \sqrt{5 + 4\sqrt{\lambda_g}}\right),$$

where $\lambda_g = \exp\{l_{\tilde{c}}([0])\}$. A simple calculation shows that

$$\lambda_g \geq \left\{\frac{1}{4}(\kappa^2 - 5)\right\}^2,$$

where $\kappa = 16p + 8n - 21$ if $n \geq 3$; $\kappa = 16p + 3$ if $n = 2$; and $\kappa = 16p + 7$ if $n = 1$. This proves Theorem 1′. Since $l_{\tilde{c}}([0]) = \log \lambda_g$, we obtain

$$l_{\tilde{c}}(\sigma) \geq 2\log(\kappa^2 - 5) - 4\log 2.$$

This proves Theorem 1. □

ACKNOWLEDGMENT. The initial version of the paper only deals with the case that $\tilde{S}$ is a punctured Riemann sphere. The author is greatly indebted to the referee for his/her generous contribution so that the extension from the case of Riemann spheres to the case of Riemann surfaces of general types becomes possible.

References

[1] L. Ahlfors and L. Bers: *Riemann's mapping theorem for variable metrics*, Ann. of Math. (2) **72** (1960), 385–404.
[2] A.F. Beardon: The Geometry of Discrete Groups, Springer, New York, 1983.
[3] L. Bers: *Fiber spaces over Teichmüller spaces*, Acta Math. **130** (1973), 89–126.
[4] L. Bers: *An extremal problem for quasiconformal mappings and a theorem by Thurston*, Acta Math. **141** (1978), 73–98.
[5] A.E. Brouwer and A. Neumaier: *The graphs with spectral radius between* 2 *and* $\sqrt{2+\sqrt{5}}$, Linear Algebra Appl. **114**/**115** (1989), 273–276.
[6] C.J. Earle and I. Kra: *On holomorphic mappings between Teichmüller spaces*; in Contributions to Analysis (a collection of papers dedicated to Lipman Bers), Academic Press, New York, 1974, 107–124.
[7] P. Hubert and E. Lanneau: *Veech groups without parabolic elements*, preprint (2005).
[8] I. Kra: *On the Nielsen-Thurston-Bers type of some self-maps of Riemann surfaces*, Acta Math. **146** (1981), 231–270.
[9] C.J. Leininger: *On groups generated by two positive multi-twists*: *Teichmüller curves and Lehmer's number*, Geom. Topol. **8** (2004), 1301–1359.
[10] A. Marden and H. Masur: *A foliation of Teichmüller space by twist invariant disks*, Math. Scand. **36** (1975), 211–228, and Addendum, **39** (1976), 232–238.
[11] B. Maskit and J.P. Matelski: *Minimum geodesics on surfaces*, manuscript.
[12] S. Nag: *Nongeodesic discs embedded in Teichmüller spaces*, Amer. J. Math. **104** (1982), 399–408.
[13] H.L. Royden: *Automorphisms and isometries of Teichmüller space*; in Advances in the Theory of Riemann Surfaces (Proc. Conf., Stony Brook, N.Y., 1969), Ann. of Math. Studies **66**, Princeton Univ. Press, Princeton, N.J., 1971, 369–383
[14] R.C. Penner: *Probing mapping class group using arcs*, manuscript, 2005.
[15] W.P. Thurston: *On the geometry and dynamics of diffeomorphisms of surfaces*, Bull. Amer. Math. Soc. (N.S.) **19** (1988), 417–431.
[16] W.A. Veech: *Teichmüller curves in moduli space, Eisenstein series and an application to triangular billiards*, Invent. Math. **97** (1989), 553–583.
[17] C. Zhang: *Hyperbolic mapping classes and their lifts on Bers fiber space*, to appear, 2005.

Department of Mathematics
Morehouse College
Atlanta, GA 30314
USA
e-mail: czhang@morehouse.edu