

Title	A GENERALIZATION OF FUNCTIONAL LIMIT THEOREMS ON THE RIEMANN ZETA PROCESS
Author(s)	Takanobu, Satoshi
Citation	Osaka Journal of Mathematics. 2019, 56(4), p. 843-882
Version Type	VoR
URL	<a href="https://doi.org/10.18910/73631">https://doi.org/10.18910/73631</a>
rights	
Note	

*Osaka University Knowledge Archive : OUKA*

<https://ir.library.osaka-u.ac.jp/>

Osaka University

## A GENERALIZATION OF FUNCTIONAL LIMIT THEOREMS ON THE RIEMANN ZETA PROCESS

SATOSHI TAKANOBU

(Received May 31, 2017, revised May 14, 2018)

### Abstract

$\zeta(\cdot)$  being the Riemann zeta function,  $\zeta_\sigma(t) := \frac{\zeta(\sigma+it)}{\zeta(\sigma)}$  is, for  $\sigma > 1$ , a characteristic function of some infinitely divisible distribution  $\mu_\sigma$ . A process with time parameter  $\sigma$  having  $\mu_\sigma$  as its marginal at time  $\sigma$  is called a Riemann zeta process. Ehm [2] has found a functional limit theorem on this process being a backwards Lévy process. In this paper, we replace  $\zeta(\cdot)$  with a Dirichlet series  $\eta(\cdot; a)$  generated by a nonnegative, completely multiplicative arithmetical function  $a(\cdot)$  satisfying (3), (4) and (5) below, and derive the same type of functional limit theorem as Ehm on the process corresponding to  $\eta(\cdot; a)$  and being a backwards Lévy process.

### Introduction

Let  $\zeta(\cdot)$  be the Riemann zeta function. Then  $\zeta_\sigma(t) := \frac{\zeta(\sigma+it)}{\zeta(\sigma)}$  is, for  $\sigma > 1$ , a characteristic function of some infinitely divisible distribution  $\mu_\sigma$ . This  $\mu_\sigma$  is called the Riemann zeta distribution indexed by parameter  $\sigma$ . We are interested in a (stochastic) process with time parameter  $\sigma$  whose marginal distribution at time  $\sigma$  is  $\mu_\sigma$ . Such a process is called a Riemann zeta (stochastic) process.

Ehm [2] has constructed this process so as to be a backwards Lévy process, and found a functional limit theorem on the process.

In this paper, we generalize the setting of Ehm. We replace  $\zeta(s)$  with a Dirichlet series  $\eta(s; a) = \sum_{n=1}^{\infty} \frac{a(n)}{n^s}$ , where  $a(\cdot)$  is a nonnegative, completely multiplicative arithmetical function satisfying (3), (4) and (5) below, and then derive the same type of functional limit theorem as Ehm on the process  $(-Z(\sigma; a))_{1 < \sigma < \infty}$  corresponding to  $\eta(\cdot; a)$  and being a backwards Lévy process, which is shortly called the  $\eta(\cdot; a)$ -process.

In Section 1, we review Ehm's result. In Section 2, we state our main result (cf. Theorem 1) and prove it, and in Section 3 give some examples of  $a(\cdot)$ .

In Section 4, we generalize  $a(\cdot)$  more, and then investigate limit theorems on  $Z(\sigma; a)$  as  $\sigma \searrow 1$  (cf. Theorems 2 ~ 4).

### 1. Review of Ehm's result

**1.1. Riemann zeta distribution.** The Riemann zeta function  $\zeta(\cdot)$  has two representations:

$$\zeta(s) = \begin{cases} \sum_{n=1}^{\infty} \frac{1}{n^s}, \\ \prod_{p:\text{prime}} \frac{1}{1 - \frac{1}{p^s}}. \end{cases}$$

Here  $s = \sigma + it, \sigma > 1, t \in \mathbb{R}$ . The former is a Dirichlet series representation and the latter is an Euler product representation. For fixed  $\sigma > 1, \zeta(\sigma + i \cdot)$  is positive definite as a function of  $\mathbb{R}$ . In other words,  $\zeta_{\sigma}(t) = \frac{\zeta(\sigma + it)}{\zeta(\sigma)}$  is a characteristic function of  $\mu_{\sigma} := \sum_{n=1}^{\infty} \frac{1}{\zeta(\sigma)n^{\sigma}} \delta_{\log \frac{1}{n}}$ . Indeed, by the former representation,

$$\zeta_{\sigma}(t) = \frac{1}{\zeta(\sigma)} \sum_{n=1}^{\infty} \frac{e^{i(\log \frac{1}{n})t}}{n^{\sigma}} = \int_{\mathbb{R}} e^{ixt} \sum_{n=1}^{\infty} \frac{1}{\zeta(\sigma)n^{\sigma}} \delta_{\log \frac{1}{n}}(dx) = \widehat{\mu_{\sigma}}(t).$$

$\mu_{\sigma}$  is called a Riemann zeta distribution with parameter  $\sigma$ . Furthermore, it is easy to see that  $\mu_{\sigma}$  is an infinitely divisible distribution: By the latter representation and

$$(1) \quad 1 + z = \exp\left\{ \int_0^1 \frac{z}{1 + zs} ds \right\}, \quad z \in \mathbb{C} \setminus (-\infty, -1],$$

it is checked that

$$\begin{aligned} \zeta_{\sigma}(t) &= \prod_{p:\text{prime}} \frac{1 - \frac{1}{p^{\sigma}}}{1 - \frac{1}{p^{\sigma+it}}} \\ &= \prod_{p:\text{prime}} \exp\left\{ \int_0^1 \frac{-\frac{1}{p^{\sigma}}}{1 - \frac{s}{p^{\sigma}}} ds - \int_0^1 \frac{-\frac{1}{p^{\sigma+it}}}{1 - \frac{s}{p^{\sigma+it}}} ds \right\} \\ &= \prod_{p:\text{prime}} \exp\left\{ \int_0^1 \sum_{n=1}^{\infty} \frac{s^{n-1}}{p^{n\sigma}} (e^{i(\log \frac{1}{p})nt} - 1) ds \right\} \\ &= \exp\left\{ \int_{\mathbb{R} \setminus \{0\}} (e^{ixt} - 1) \nu_{\sigma}(dx) \right\}, \end{aligned}$$

where

$$\nu_{\sigma}(dx) := \sum_{p:\text{prime}} \sum_{n=1}^{\infty} \frac{1}{np^{n\sigma}} \delta_{n \log \frac{1}{p}}(dx), \quad x \in \mathbb{R} \setminus \{0\}$$

is a Lévy measure.

$\zeta(\cdot)$  is extended meromorphically to the whole complex plane with only a simple pole at 1 with residue 1. Thus, asymptotically

$$(s - 1)\zeta(s) = 1 + O(|s - 1|) \quad \text{as } s \rightarrow 1.$$

By this, we easily have the following limit theorem for  $\mu_{\sigma}$  as  $\sigma \searrow 1$ :

**Claim 1.** As  $\sigma \searrow 1$ ,

$$\mu_{\sigma}\left(\frac{-1}{\sigma - 1} dx\right) \rightarrow \mathbf{1}_{(0,\infty)}(x)e^{-x} dx \quad (= \text{the exponential distribution with parameter } 1).$$

Proof. For  $\forall t \in \mathbb{R}$ ,

$$\begin{aligned} \int_{\mathbb{R}} e^{itx} \mu_{\sigma} \left( \frac{-1}{\sigma-1} dx \right) &= \int_{\mathbb{R}} e^{it(-(\sigma-1)y)} \mu_{\sigma}(dy) \\ &= \zeta_{\sigma}(-t(\sigma-1)) \\ &= \frac{\zeta(\sigma + i(-t(\sigma-1)))}{\zeta(\sigma)} \\ &\rightarrow \frac{1}{1-it} = \int_0^{\infty} e^{itx} e^{-x} dx \quad \text{as } \sigma \searrow 1. \end{aligned}$$

□

**1.2. Riemann zeta process.** A process with time parameter  $\sigma \in (1, \infty)$  having  $\mu_{\sigma}$  as its marginal at  $\sigma$  is called a Riemann zeta process. Following Ehm [2], we construct the process so as to be a backwards Lévy process.

DEFINITION 1. A process  $(Y(u))_{0 \leq u < 1}$  on some probability space  $(\Omega, \mathcal{F}, P)$  is called a geometric process if the following (a) ~ (d) hold:

- (a) For each  $u \in [0, 1)$ ,  $Y(u) \in \{0, 1, 2, \dots\}$ . Especially  $Y(0) = 0$ .
- (b)  $[0, 1) \ni u \mapsto Y(u) \in \mathbb{R}$  is right-continuous and non-decreasing.
- (c)  $(Y(u))_{0 \leq u < 1}$  is a Lévy process, i.e., for every  $0 < u_0 < u_1 < \dots < u_n < 1$ ,

$$Y(u_0), Y(u_1) - Y(u_0), \dots, Y(u_n) - Y(u_{n-1}) \text{ are independent,}$$

and, for each  $u \in (0, 1)$ ,  $Y(u) = Y(u-)$  a.s.

- (d) For each  $0 \leq u < v < 1$ ,

$$E[e^{it(Y(v)-Y(u))}] = \frac{1 - ue^{it}}{1 - u} \frac{1 - v}{1 - ve^{it}}.$$

In particular

$$E[e^{itY(u)}] = \frac{1 - u}{1 - ue^{it}}.$$

Thus  $P(Y(u) = n) = u^n(1 - u)$ ,  $n \in \{0, 1, 2, \dots\}$ , in other words,  $Y(u)$  is geometrically distributed with parameter  $1 - u$ .

DEFINITION 2. Let  $\{Y_p\}_{p:\text{prime}}$  be a sequence of independent geometric processes on some  $(\Omega, \mathcal{F}, P)$ . Then we define

$$Z(\sigma) := \sum_{p:\text{prime}} Y_p(p^{-\sigma}) \log p, \quad \sigma \in (1, \infty).$$

**Claim 2.**  $(-Z(\sigma))_{1 < \sigma < \infty}$  is a Riemann zeta process, and a backwards Lévy process. This means the following:

- (i)  $(1, \infty) \ni \sigma \mapsto Z(\sigma) \in [0, \infty)$  is left-continuous and non-increasing;
- (ii) For  $\sigma > 1$ ,  $Z(\sigma+) = Z(\sigma)$  a.s.,  $Z(1+) = \infty$  a.s. and  $Z(\infty) = 0$ ;
- (iii) For every  $\infty > \sigma_0 > \sigma_1 > \dots > \sigma_n > 1$ ,

$$Z(\sigma_0), Z(\sigma_1) - Z(\sigma_0), \dots, Z(\sigma_n) - Z(\sigma_{n-1}) \text{ are independent.}$$

Proof. Claim 2 is contained in Claim 7 below. So the proof of Claim 2 is omitted.  $\square$

**1.3. Ehm's functional limit theorem.** Claim 1 can be restated in terms of  $Z(\sigma)$ :

**Claim 3.** As  $\sigma \searrow 1$ ,

the distribution of  $(\sigma - 1)Z(\sigma) \rightarrow$  the exponential distribution with parameter 1.

Proof. As  $\sigma \searrow 1$ ,

$$P((\sigma - 1)Z(\sigma) \in dx) = \mu_\sigma\left(\frac{-1}{\sigma - 1}dx\right) \\ \rightarrow \text{the exponential distribution with parameter 1.}$$

$\square$

This limit theorem is generalized as a functional limit theorem:

**Fact 1** (cf. [2]). Let  $\varphi : (1, \infty) \rightarrow (0, \infty)$  be a  $C^1$ , strictly decreasing function such that

$$\varphi(1+) = \infty, \quad \varphi(\infty) = 0, \\ \varphi(\sigma) \sim \frac{1}{(\sigma - 1)^2} \quad \text{as } \sigma \searrow 1.$$

Let  $N(dsdu)$  be a Poisson random measure on  $(0, \infty) \times (0, \infty)$  with mean measure

$$n(dsdu) := \frac{1}{2}e^{-u/\sqrt{s}}s^{-3/2}dsdu, \quad s, u > 0.$$

Then the following holds:

$$\left(\frac{1}{\sqrt{T}}Z(\varphi^{-1}(Tt))\right)_{t \geq 0} \xrightarrow{D} \left(\int_0^{t+} \int_{(0, \infty)} uN(dsdu)\right)_{t \geq 0} \\ \text{in } D([0, \infty) \rightarrow \mathbb{R}) \quad \text{as } T \rightarrow \infty.$$

Here  $D([0, \infty) \rightarrow \mathbb{R})$  is the space of all real functions on  $[0, \infty)$  that are right-continuous and have left-hand limits. This space is endowed with the  $J_1$ -topology (cf. [8, 1]), so that it becomes a Polish space.  $(\frac{1}{\sqrt{T}}Z(\varphi^{-1}(Tt)))_{t \geq 0}$  and  $(\int_0^{t+} \int_{(0, \infty)} uN(dsdu))_{t \geq 0}$  are random elements of  $D([0, \infty) \rightarrow \mathbb{R})$ , that is, they are  $D([0, \infty) \rightarrow \mathbb{R})$ -valued random variables. In other words, almost all samples  $t \mapsto \frac{1}{\sqrt{T}}Z(\varphi^{-1}(Tt))$  and  $t \mapsto \int_0^{t+} \int_{(0, \infty)} uN(dsdu)$  belong to  $D([0, \infty) \rightarrow \mathbb{R})$ , and for each  $t \in [0, \infty)$ ,  $\frac{1}{\sqrt{T}}Z(\varphi^{-1}(Tt))$  and  $\int_0^{t+} \int_{(0, \infty)} uN(dsdu)$  are real random variables. The convergence above denotes the weak convergence of the distribution of  $(\frac{1}{\sqrt{T}}Z(\varphi^{-1}(Tt)))_{t \geq 0}$  to that of  $(\int_0^{t+} \int_{(0, \infty)} uN(dsdu))_{t \geq 0}$ .

The statement of this fact is different from that of Ehm [2]. Suited to our theorem stated in Section 2, the above fact has been presented.

**2. Our main result**

**2.1. Completely multiplicative arithmetical function  $a(\cdot)$ .** If an arithmetical function  $a : \mathbb{N} \rightarrow \mathbb{R}$  satisfies

$$(2) \quad a(mn) = a(m)a(n), \quad \forall m, \forall n \in \mathbb{N},$$

then  $a(\cdot)$  is said to be completely multiplicative.  $a(\cdot) = 0$  (i.e.,  $a(n) = 0 \ (\forall n \in \mathbb{N})$ ) and  $a(\cdot) = 1$  (i.e.,  $a(n) = 1 \ (\forall n \in \mathbb{N})$ ) are clearly such arithmetical functions. For a completely multiplicative  $a(\cdot)$ , it should be remarked that

$$a(\cdot) \neq 0 \text{ (i.e., } \exists n_0 \in \mathbb{N} \text{ s.t. } a(n_0) \neq 0) \Leftrightarrow_{\text{iff}} a(1) = 1,$$

in other words,  $a(\cdot) = 0 \Leftrightarrow_{\text{iff}} a(1) = 0$ . Since  $a(\cdot) = 0$  is too trivial, it is excluded from completely multiplicative companions. Thus, from now on  $a : \mathbb{N} \rightarrow \mathbb{R}$  is called completely multiplicative if  $a(1) = 1$  and (2) is satisfied. In this case, if  $n = \prod_p p^{\alpha_p(n)}$  is the prime factorization of  $n \in \mathbb{N}$ , where

$$\alpha_p(n) = \max\{m \in \{0, 1, 2, \dots\}; p^m \mid n\},$$

then

$$a(n) = \prod_p a(p)^{\alpha_p(n)}.$$

Here let  $x^0 = 1$  for  $x \in \mathbb{R}$ . Thus, the value of  $a(\cdot)$  is completely determined by that of  $(a(p))_{p:\text{prime}}$ .

In the following, let  $a : \mathbb{N} \rightarrow [0, \infty)$  be a completely multiplicative arithmetical function<sup>1</sup> such that

$$(3) \quad \sup_p a(p) < \infty, \quad \sup_p \frac{a(p)}{p} \leq 1,$$

$$(4) \quad \exists \tau \geq 0 \text{ s.t. } \sum_{p \leq x} \frac{a(p) \log p}{p} = (\tau + o(1)) \log x \quad \text{as } x \rightarrow \infty,$$

$$(5) \quad \tau + \#\{p; a(p) = p\} > 0.$$

Note that  $0 \leq \#\{p; a(p) = p\} < \infty$  since  $\sup_p a(p) < \infty$ . When  $\tau > 0$  in (4), (5) holds automatically. When  $\tau = 0$  in (4), (5) becomes  $\#\{p; a(p) = p\} > 0$ . By Mertens' first theorem:

$$(6) \quad \sum_{p \leq x} \frac{\log p}{p} = \log x + O(1) \quad \text{as } x \rightarrow \infty$$

(cf. [5, Theorem 425] or [10, Chapter I.1, Theorem 7]),  $a(\cdot) = 1$  is a typical example. In Section 3, we will give some other examples.

In what follows up to the end of Section 2, let us fix such an arithmetical function  $a(\cdot)$ .

**2.2. Presentation of Theorem 1.** To state our main result – Theorem 1, we need some definitions:

---

<sup>1</sup>For simplicity, we restrict completely multiplicative arithmetical functions appearing in this paper to be nonnegative.

DEFINITION 3. For  $s = \sigma + it$  ( $\sigma > 1, t \in \mathbb{R}$ ), we define

$$\eta(s; a) := \sum_{n=1}^{\infty} \frac{a(n)}{n^s} = \prod_p \frac{1}{1 - \frac{a(p)}{p^s}}.$$

By virtue of Claims 4 and 5 below, this is well-defined. When  $a(\cdot) = 1, \eta(\cdot; 1) = \zeta(\cdot)$ !

DEFINITION 4. For  $\sigma \in (1, \infty)$ , we define

$$\eta_{\sigma}(t; a) := \frac{\eta(\sigma + it; a)}{\eta(\sigma; a)}, \quad t \in \mathbb{R}.$$

If  $\mu_{\sigma}(dx; a)$  and  $\nu_{\sigma}(dx; a)$  are a 1-dimensional probability measure and a Lévy measure, respectively, defined by

$$\begin{aligned} \mu_{\sigma}(dx; a) &:= \sum_{n=1}^{\infty} \frac{a(n)}{\eta(\sigma; a)n^{\sigma}} \delta_{\log \frac{1}{n}}(dx), \\ \nu_{\sigma}(dx; a) &:= \sum_p \sum_{n=1}^{\infty} \frac{a(p)^n}{np^{n\sigma}} \delta_{n \log \frac{1}{p}}(dx), \end{aligned}$$

then

$$\eta_{\sigma}(t; a) = \widehat{\mu_{\sigma}(\cdot; a)}(t) = \exp \left\{ \int_{\mathbb{R} \setminus \{0\}} (e^{ixt} - 1) \nu_{\sigma}(dx; a) \right\}.$$

$\mu_{\sigma}(\cdot; a)$  is called the  $\eta(\cdot; a)$ -distribution with parameter  $\sigma$ .

DEFINITION 5. Let  $\{Y_p\}_p$  be a sequence of independent geometric processes on some  $(\Omega, \mathcal{F}, P)$ . Then we define

$$Z(\sigma; a) := \sum_p Y_p \left( \frac{a(p)}{p^{\sigma}} \right) \log p, \quad \sigma \in (1, \infty).$$

By Claim 7 below,  $(-Z(\sigma; a))_{1 < \sigma < \infty}$  is a backwards Lévy process whose marginal distribution at  $\sigma$  is  $\mu_{\sigma}(\cdot; a)$ . Thus, by imitating  $(-Z(\sigma))_{1 < \sigma < \infty}$ , this is called an  $\eta(\cdot; a)$ -process.

Our main result is the following:

**Theorem 1.** Let  $\varphi : (1, \infty) \rightarrow (0, \infty)$  be a  $C^1$ , strictly decreasing function such that

(7)  $\varphi(1+) = \infty, \varphi(\infty) = 0,$

(8)  $\varphi(\sigma) \sim \frac{1}{(\sigma - 1)^2}$  as  $\sigma \searrow 1.$

Let  $\rho := \tau + \#\{p; a(p) = p\} > 0$  (cf. Claim 6 below), and  $N^{(\rho)}(dsdu)$  be a Poisson random measure on  $(0, \infty) \times (0, \infty)$  with mean measure

(9)  $n^{(\rho)}(dsdu) := \frac{\rho}{2} e^{-\frac{u}{\sqrt{s}}} s^{-\frac{3}{2}} dsdu, \quad s, u > 0.$

Then the following holds:

$$\left( \frac{1}{\sqrt{T}} Z(\varphi^{-1}(Tt); a) \right)_{t \geq 0} \xrightarrow{D} \left( \int_0^{t+} \int_{(0, \infty)} u N^{(\rho)}(dsdu) \right)_{t \geq 0}$$

in  $D([0, \infty) \rightarrow \mathbb{R})$  as  $T \rightarrow \infty$ .

REMARK 1. In the definition of  $(Z(\sigma))_{1 < \sigma < \infty}$  (cf. Definition 2), we replaced  $p^{-\sigma}$  with  $\frac{a(p)}{p^\sigma}$  and obtained a functional limit theorem on the resultant process  $(Z(\sigma; a))_{1 < \sigma < \infty}$ . Since  $(Z(\sigma; a))_{1 < \sigma < \infty}$  is a process defined from  $\eta(\cdot; a)$ , we could even say that this functional limit theorem comes from a topic of the number theory. As a different generalization of [2], Ehm [3] replaced  $\log p$  with more general coefficient  $c_p$  and considered a functional limit theorem on the resultant process. In this case, though this process is of zeta type, its functional limit theorem is no longer related to the number theory.

**2.3. Some claims.** To make definitions given in the preceding subsection meaningful, we here present some claims:

**Claim 4.** (i)  $\sum_{p \leq x} \frac{a(p)}{p} = (\tau + o(1)) \log \log x$  as  $x \rightarrow \infty$ . When  $\tau > 0$ ,  $\sum_p \frac{a(p)}{p} = \infty$ .  
 (ii) For  $\sigma > 1$  and  $t \in \mathbb{R}$ ,  $\prod_p \frac{1 - \frac{a(p)}{p^{\sigma+it}}}{1 - \frac{a(p)}{p^{\sigma+it}}}$  is convergent. That is,  $\prod_{p \leq x} \frac{1 - \frac{a(p)}{p^{\sigma+it}}}{1 - \frac{a(p)}{p^{\sigma+it}}}$  is convergent as  $x \rightarrow \infty$ . As  $\sigma \searrow 1$ ,  $\prod_p \frac{1 - \frac{a(p)}{p^{\sigma+it}}}{1 - \frac{a(p)}{p^{\sigma+it}}} \nearrow \infty$ .

Proof. (i) For simplicity, let

$$(10) \quad C(x) := \sum_{p \leq x} \frac{a(p) \log p}{p}, \quad x \in \mathbb{R}.$$

$C(\cdot)$  is non-decreasing, right-continuous,  $C(x) = 0$  ( $\forall x < 2$ ), and

$$(11) \quad C(dt) = \sum_p \frac{a(p) \log p}{p} \delta_p(dt).$$

If, for  $x > 1$ , we set

$$(12) \quad \delta(x) := \frac{C(x)}{\log x} - \tau,$$

then  $\delta(\cdot)$  is of bounded variation on every bounded closed interval of  $(1, \infty)$ , and by (4),

$$(13) \quad \lim_{x \rightarrow \infty} \delta(x) = 0.$$

(11) and  $C(t) = \tau \log t + \delta(t) \log t$  tell us that for  $x \geq 3$ ,

$$\begin{aligned} \sum_{p \leq x} \frac{a(p)}{p} &= \sum_{p \leq x} \frac{1}{\log p} \frac{a(p) \log p}{p} \\ &= \int_{(2-\varepsilon, x]} \frac{C(dt)}{\log t} \quad [\text{where } 0 < \varepsilon < 1] \\ &= \tau \int_{2-\varepsilon}^x \frac{dt}{t \log t} + \int_{2-\varepsilon}^x \frac{\delta(t)}{t \log t} dt + \int_{(2-\varepsilon, x]} \delta(dt) \\ &= \tau \int_{2-\varepsilon}^x \frac{dt}{t \log t} + \int_{2-\varepsilon}^x \frac{\delta(t)}{t \log t} dt + \delta(x) + \tau. \end{aligned}$$

By letting  $\varepsilon \searrow 0$ ,

$$\sum_{p \leq x} \frac{a(p)}{p} = \tau \int_2^x \frac{dt}{t \log t} + \int_2^x \frac{\delta(t)}{t \log t} dt + \tau + \delta(x)$$



$$= \log \log x \left( \tau + \frac{\tau(-\log \log 2 + 1)}{\log \log x} + \frac{\delta(x)}{\log \log x} + \int_{\frac{\log \log 2}{\log \log x}}^1 \delta(e^{e^{(\log \log x)u}}) du \right)$$

[by the change of variables  $\frac{\log \log t}{\log \log x} = u$ ].

Since, by (13) and the bounded convergence theorem,

$$\lim_{x \rightarrow \infty} \int_{\frac{\log \log 2}{\log \log x}}^1 \delta(e^{e^{(\log \log x)u}}) du = 0,$$

we have

$$\sum_{p \leq x} \frac{a(p)}{p} = (\log \log x)(\tau + o(1)) \quad \text{as } x \rightarrow \infty.$$

(ii) First, (1) is rewritten as

$$(14) \quad 1 + z = e^z \exp\left\{-z^2 \int_0^1 \frac{s}{1 + zs} ds\right\}, \quad z \in \mathbb{C} \setminus (-\infty, -1].$$

Let  $\sigma > 1$  and  $t \in \mathbb{R}$ . Since, by (3),

$$(15) \quad \left| \frac{a(p)}{p^{\sigma+it}} \right| = \frac{1}{p^{\sigma-1}} \frac{a(p)}{p} \leq \frac{1}{p^{\sigma-1}} \leq \frac{1}{2^{\sigma-1}} < 1,$$

(14) implies that

$$(16) \quad \frac{1}{1 - \frac{a(p)}{p^{\sigma+it}}} = e^{\frac{a(p)}{p^{\sigma+it}}} \exp\left\{\frac{a(p)^2}{p^{2\sigma+i2t}} \int_0^1 \frac{s}{1 - \frac{a(p)}{p^{\sigma+it}} s} ds\right\}.$$

Multiplication in  $p \leq x$  yields that

$$\prod_{p \leq x} \frac{1}{1 - \frac{a(p)}{p^{\sigma+it}}} = \exp\left\{\sum_{p \leq x} \frac{a(p)}{p^{\sigma+it}}\right\} \exp\left\{\sum_{p \leq x} \frac{a(p)^2}{p^{2\sigma+i2t}} \int_0^1 \frac{s}{1 - \frac{a(p)}{p^{\sigma+it}} s} ds\right\}.$$

Here, by noting that

$$\begin{aligned} \sum_p \left| \frac{a(p)^2}{p^{2\sigma+i2t}} \int_0^1 \frac{s}{1 - \frac{a(p)}{p^{\sigma+it}} s} ds \right| &\leq \sum_p \frac{a(p)^2}{p^{2\sigma}} \int_0^1 \frac{s}{1 - \left| \frac{a(p)}{p^{\sigma+it}} \right| s} ds \\ &\leq \frac{(\sup_p a(p))^2}{2(1 - \frac{1}{2^{\sigma-1}})} \sum_p \frac{1}{p^{2\sigma}} \quad [\text{cf. (15)}] < \infty, \end{aligned}$$

the convergence of  $\prod_p \frac{1}{1 - \frac{a(p)}{p^{\sigma+it}}}$  is reduced to that of  $\sum_p \frac{a(p)}{p^{\sigma+it}}$ . Since

$$\sum_p \left| \frac{a(p)}{p^{\sigma+it}} \right| = \sum_p \frac{a(p)}{p^\sigma} \leq \sup_p a(p) \sum_p \frac{1}{p^\sigma} \leq \sup_p a(p) \sum_n \frac{1}{n^\sigma} < \infty,$$

$\prod_p \frac{1}{1 - \frac{a(p)}{p^{\sigma+it}}}$  is convergent.

Next we check the divergence of  $\prod_p \frac{1}{1 - \frac{a(p)}{p^\sigma}}$  as  $\sigma \searrow 1$ . First

$$\prod_p \frac{1}{1 - \frac{a(p)}{p^\sigma}} = \left( \prod_{p: a(p)=p} \frac{1}{1 - \frac{1}{p^{\sigma-1}}} \right) \left( \prod_{p: a(p)<p} \frac{1}{1 - \frac{a(p)}{p^\sigma}} \right)$$

$$= \left( \prod_{p:a(p)=p} \frac{1}{1 - \frac{1}{p^{\sigma-1}}} \right) \exp\left\{ \sum_p \frac{a(p)}{p^\sigma} \right\} \exp\left\{ - \sum_{p:a(p)=p} \frac{1}{p^{\sigma-1}} \right\} \\ \times \exp\left\{ \sum_{p:a(p)<p} \frac{a(p)^2}{p^{2\sigma}} \int_0^1 \frac{s}{1 - \frac{a(p)}{p^\sigma} s} ds \right\}.$$

Here note that

$$(17) \quad \sup_{q:a(q)<q} \frac{a(q)}{q} < 1.$$

This tells us that for  $p$  with  $a(p) < p$ ,

$$(18) \quad -\frac{a(p)}{p^\sigma} = -\frac{a(p)}{p} \frac{1}{p^{\sigma-1}} \geq -\left( \sup_{q:a(q)<q} \frac{a(q)}{q} \right) \frac{1}{p^{\sigma-1}} > -\left( \sup_{q:a(q)<q} \frac{a(q)}{q} \right) > -1,$$

so that

$$\sum_{p:a(p)<p} \frac{a(p)^2}{p^{2\sigma}} \int_0^1 \frac{s}{1 - \frac{a(p)}{p^\sigma} s} ds \leq \frac{(\sup_{q:a(q)<q} a(q))^2}{2(1 - \sup_{q:a(q)<q} \frac{a(q)}{q})} \sum_p \frac{1}{p^2} < \infty.$$

Thus

$$\lim_{\sigma \searrow 1} \exp\left\{ \sum_{p:a(p)<p} \frac{a(p)^2}{p^{2\sigma}} \int_0^1 \frac{s}{1 - \frac{a(p)}{p^\sigma} s} ds \right\} = \exp\left\{ \sum_{p:a(p)<p} \frac{a(p)^2}{p^2} \int_0^1 \frac{s}{1 - \frac{a(p)}{p} s} ds \right\} < \infty.$$

Clearly

$$\lim_{\sigma \searrow 1} \exp\left\{ - \sum_{p:a(p)=p} \frac{1}{p^{\sigma-1}} \right\} = e^{-\#\{p:a(p)=p\}} < \infty.$$

When  $\tau > 0$ ,

$$\lim_{\sigma \searrow 1} \exp\left\{ \sum_p \frac{a(p)}{p^\sigma} \right\} = \exp\left\{ \sum_p \frac{a(p)}{p} \right\} = \infty$$

by (i). When  $\tau = 0$ ,

$$\lim_{\sigma \searrow 1} \prod_{p:a(p)=p} \frac{1}{1 - \frac{1}{p^{\sigma-1}}} = \infty$$

since  $\{p; a(p) = p\} \neq \emptyset$ . Therefore, putting all together, we have

$$\lim_{\sigma \searrow 1} \prod_p \frac{1}{1 - \frac{a(p)}{p^\sigma}} = \infty.$$

□

**Claim 5.** For  $\sigma > 1$  and  $t \in \mathbb{R}$ ,  $\sum_{n=1}^\infty \frac{a(n)}{n^{\sigma+it}}$  is absolutely convergent, and coincides with  $\prod_p \frac{1}{1 - \frac{a(p)}{p^{\sigma+it}}}$ .

Proof. Fix  $\sigma > 1$  and  $t \in \mathbb{R}$ . Let  $p_j$  be the  $j$ th prime number. Note that

$$\mathbb{N}_L := \{p_1^{\alpha_1} \cdots p_L^{\alpha_L}; 0 \leq \alpha_1, \dots, \alpha_L \leq L\} \nearrow \mathbb{N} \quad \text{as } L \rightarrow \infty.$$

By the completely multiplicative property of  $a(\cdot)$ ,

$$\begin{aligned} \sum_{n \in \mathbb{N}_L} \frac{a(n)}{n^{\sigma+it}} &= \sum_{0 \leq \alpha_1, \dots, \alpha_L \leq L} \frac{a(p_1^{\alpha_1} \cdots p_L^{\alpha_L})}{(p_1^{\alpha_1} \cdots p_L^{\alpha_L})^{\sigma+it}} \\ &= \sum_{0 \leq \alpha_1, \dots, \alpha_L \leq L} \frac{a(p_1)^{\alpha_1}}{(p_1^{\sigma+it})^{\alpha_1}} \times \cdots \times \frac{a(p_L)^{\alpha_L}}{(p_L^{\sigma+it})^{\alpha_L}} \\ &= \frac{1 - (\frac{a(p_1)}{p_1^{\sigma+it}})^{L+1}}{1 - \frac{a(p_1)}{p_1^{\sigma+it}}} \times \cdots \times \frac{1 - (\frac{a(p_L)}{p_L^{\sigma+it}})^{L+1}}{1 - \frac{a(p_L)}{p_L^{\sigma+it}}} \quad [\text{cf. (15)}] \\ &= \left( \prod_{j=1}^L \frac{1}{1 - \frac{a(p_j)}{p_j^{\sigma+it}}} \right) \prod_{j=1}^L \left( 1 - \left( \frac{a(p_j)}{p_j^{\sigma+it}} \right)^{L+1} \right). \end{aligned}$$

By (1),

$$\prod_{j=1}^L \left( 1 - \left( \frac{a(p_j)}{p_j^{\sigma+it}} \right)^{L+1} \right) = \exp \left\{ \sum_{j=1}^L \left( - \left( \frac{a(p_j)}{p_j^{\sigma+it}} \right)^{L+1} \right) \int_0^1 \frac{1}{1 - \left( \frac{a(p_j)}{p_j^{\sigma+it}} \right)^{L+1} s} ds \right\}.$$

Since

$$\begin{aligned} &\left| \sum_{j=1}^L \left( - \left( \frac{a(p_j)}{p_j^{\sigma+it}} \right)^{L+1} \right) \int_0^1 \frac{1}{1 - \left( \frac{a(p_j)}{p_j^{\sigma+it}} \right)^{L+1} s} ds \right| \\ &\leq L \left( \frac{1}{2^{\sigma-1}} \right)^{L+1} \frac{1}{1 - \left( \frac{1}{2^{\sigma-1}} \right)^{L+1}} \quad [\text{cf. (15)}] \rightarrow 0 \quad \text{as } L \rightarrow \infty, \end{aligned}$$

we have

$$\lim_{L \rightarrow \infty} \prod_{j=1}^L \left( 1 - \left( \frac{a(p_j)}{p_j^{\sigma+it}} \right)^{L+1} \right) = 1,$$

which implies

$$\lim_{L \rightarrow \infty} \sum_{n \in \mathbb{N}_L} \frac{a(n)}{n^{\sigma+it}} = \prod_p \frac{1}{1 - \frac{a(p)}{p^{\sigma+it}}}.$$

When  $t = 0$ , the monotone convergence theorem tells us that

$$\sum_{n=1}^{\infty} \frac{a(n)}{n^{\sigma}} = \sum_{n \in \mathbb{N}} \frac{a(n)}{n^{\sigma}} = \lim_{L \rightarrow \infty} \sum_{n \in \mathbb{N}_L} \frac{a(n)}{n^{\sigma}} = \prod_p \frac{1}{1 - \frac{a(p)}{p^{\sigma}}} < \infty.$$

This shows the absolute convergence of  $\sum_{n=1}^{\infty} \frac{a(n)}{n^{\sigma+it}}$ , so that

$$\sum_{n=1}^{\infty} \frac{a(n)}{n^{\sigma+it}} = \sum_{n \in \mathbb{N}} \frac{a(n)}{n^{\sigma+it}} = \lim_{L \rightarrow \infty} \sum_{n \in \mathbb{N}_L} \frac{a(n)}{n^{\sigma+it}} = \prod_p \frac{1}{1 - \frac{a(p)}{p^{\sigma+it}}}$$

is obtained. □

**Claim 6.**  $-\frac{\eta'(\sigma; a)}{\eta(\sigma; a)} \sim \frac{\rho}{\sigma - 1}$  as  $\sigma \searrow 1$ , where  $\rho = \tau + \#\{p; a(p) = p\} > 0$  (cf. (5)).

Proof. It is divided into 5 steps.

1° Since, by Definition 3,

$$\log \eta(\sigma; a) = - \sum_p \log \left( 1 - \frac{a(p)}{p^\sigma} \right),$$

differentiating in  $\sigma$  yields that

$$\begin{aligned} - \frac{\eta'(\sigma; a)}{\eta(\sigma; a)} &= \sum_p \frac{-\frac{a(p)}{p^\sigma} \log \frac{1}{p}}{1 - \frac{a(p)}{p^\sigma}} \\ &= \sum_p \frac{a(p)}{p^\sigma} (\log p) \left( 1 + \frac{\frac{a(p)}{p^\sigma}}{1 - \frac{a(p)}{p^\sigma}} \right) \\ &= \sum_p \frac{a(p)}{p^\sigma} \log p \\ &\quad + \sum_{p; a(p)=p} \frac{\log p}{p^{2(\sigma-1)}} \frac{1}{1 - \frac{1}{p^{\sigma-1}}} + \sum_{p; a(p) < p} \frac{a(p)^2}{p^{2\sigma}} (\log p) \frac{1}{1 - \frac{a(p)}{p^\sigma}} \\ &=: \text{the first term} + \text{the second term} + \text{the third term.} \end{aligned}$$

2° Clearly

$$\begin{aligned} (\sigma - 1) \times \text{the second term} &= \sum_{p; a(p)=p} \frac{1}{p^{\sigma-1}} \frac{(\sigma - 1) \log p}{e^{(\sigma-1) \log p} - 1} \\ &\rightarrow \sum_{p; a(p)=p} 1 = \#\{p; a(p) = p\} \quad \text{as } \sigma \searrow 1. \end{aligned}$$

3° By (11),

$$\begin{aligned} \text{the first term} &= \sum_p \frac{1}{p^{\sigma-1}} \frac{a(p) \log p}{p} \\ &= \int_{(2-\varepsilon, \infty)} \frac{C(dx)}{x^{\sigma-1}} \quad [\text{where } 0 < \varepsilon < 1] \\ &= \tau \int_{2-\varepsilon}^\infty \frac{dx}{x^\sigma} + \int_{2-\varepsilon}^\infty \frac{\delta(x)}{x^\sigma} dx + \int_{(2-\varepsilon, \infty)} \frac{\log x}{x^{\sigma-1}} \delta(dx) \\ &= \tau \int_{2-\varepsilon}^\infty \frac{dx}{x^\sigma} + \int_{2-\varepsilon}^\infty \frac{\delta(x)}{x^\sigma} dx \\ &\quad + \int_{(2-\varepsilon, \infty)} \left( d \left( \frac{\log x}{x^{\sigma-1}} \delta(x) \right) - \delta(x) d(x^{-\sigma+1} \log x) \right) \\ &\quad [\text{by integration by parts}] \\ &= \tau \left( \int_{2-\varepsilon}^\infty \frac{dx}{x^\sigma} + \frac{\log(2-\varepsilon)}{(2-\varepsilon)^{\sigma-1}} \right) + (\sigma - 1) \int_{2-\varepsilon}^\infty \delta(x) \frac{\log x}{x^\sigma} dx. \end{aligned}$$

Letting  $\varepsilon \searrow 0$  yields that

$$\text{the first term} = \tau \left( \int_2^\infty \frac{dx}{x^\sigma} + \frac{\log 2}{2^{\sigma-1}} \right) + (\sigma - 1) \int_2^\infty \delta(x) \frac{\log x}{x^\sigma} dx$$

$$= \frac{1}{\sigma - 1} \left\{ \frac{\tau}{2^{\sigma-1}} (1 + (\sigma - 1) \log 2) + \int_{(\sigma-1) \log 2}^{\infty} \delta(e^{\frac{z}{\sigma-1}}) e^{-z} z dz \right\}$$

[by the change of variables  $(\sigma - 1) \log x = z$ ].

Since, by (13) and the Lebesgue convergence theorem,

$$\lim_{\sigma \searrow 1} \int_{(\sigma-1) \log 2}^{\infty} \delta(e^{\frac{z}{\sigma-1}}) e^{-z} z dz = 0,$$

we have

$$(\sigma - 1) \times \text{the first term} \rightarrow \tau \quad \text{as } \sigma \searrow 1.$$

4° Since, by (17),

$$\text{the third term} \leq \left( \sum_{p:a(p)<p} \frac{a(p)^2}{p^2} (\log p) \right) \frac{1}{1 - \sup_{q:a(q)<q} \frac{a(q)}{q}},$$

we have

$$(\sigma - 1) \times \text{the third term} \rightarrow 0 \quad \text{as } \sigma \searrow 1.$$

5° By putting 1° ~ 4° together,

$$(\sigma - 1) \times \left( -\frac{\eta'(\sigma; a)}{\eta(\sigma; a)} \right) \rightarrow \tau + \#\{p; a(p) = p\} = \rho \quad \text{as } \sigma \searrow 1.$$

□

**Claim 7.**  $(-Z(\sigma; a))_{1 < \sigma < \infty}$  is a backwards Lévy process whose marginal distribution at  $\sigma$  is  $\mu_\sigma(\cdot; a)$ .

Proof. It is divided into 3 steps.

1° Since, for  $\sigma > 1$ ,

$$\sum_p P\left(Y_p\left(\frac{a(p)}{p^\sigma}\right) > 0\right) = \sum_p \frac{a(p)}{p^\sigma} < \infty,$$

Borel-Cantelli's first lemma tells us that

$$P\left(\exists p_0: \text{prime s.t. } Y_p\left(\frac{a(p)}{p^\sigma}\right) = 0, \forall p > p_0\right) = 1.$$

This implies that

$$\sum_p Y_p\left(\frac{a(p)}{p^\sigma}\right) \log p \text{ is a finite sum a.s.}$$

Thus  $Z(\sigma; a)$  is well-defined.

2° Since, for each prime  $p$ ,  $(1, \infty) \ni \sigma \mapsto Y_p\left(\frac{a(p)}{p^\sigma}\right) \in \{0, 1, 2, \dots\}$  is left-continuous and non-increasing, so is  $(1, \infty) \ni \sigma \mapsto Z(\sigma; a) \in [0, \infty)$ .

Since  $\frac{a(p)}{p^{\sigma'}} \nearrow \frac{a(p)}{p^\sigma}$  as  $\sigma' \searrow \sigma (> 1)$  ( $\forall p; a(p) > 0$ ),

$$Y_p\left(\frac{a(p)}{p^{\sigma'}}\right) \nearrow Y_p\left(\frac{a(p)}{p^\sigma}\right) = Y_p\left(\frac{a(p)}{p^\sigma}\right) \text{ a.s. } (\forall p; a(p) > 0).$$

By the monotone convergence theorem,

$$Z(\sigma'; a) \nearrow \sum_p Y_p\left(\frac{a(p)}{p^{\sigma'}}\right) \log p \quad \text{a.s.}$$

Thus, in the case where  $\sigma > 1$ ,  $Z(\sigma+; a) = Z(\sigma; a)$  a.s.

For  $\sigma' > 1$ ,

$$\begin{aligned} Z(\sigma'; a) &= \sum_p Y_p\left(\frac{a(p)}{p^{\sigma'}}\right) \log p \\ &= \sum_{p:a(p)=p} Y_p\left(\frac{1}{p^{\sigma'-1}}\right) \log p + \sum_{p:a(p)<p} Y_p\left(\frac{a(p)}{p^{\sigma'}}\right) \log p \\ &=: \text{the first term} + \text{the second term.} \end{aligned}$$

In the same way as above,

$$\lim_{\sigma' \searrow 1} \text{the second term} = \sum_{p:a(p)<p} Y_p\left(\frac{a(p)}{p}\right) \log p \quad \text{a.s.}$$

When  $\tau > 0$ , Claim 4(i) tells us that

$$\sum_{p:a(p)<p} P\left(Y_p\left(\frac{a(p)}{p}\right) > 0\right) = \sum_{p:a(p)<p} \frac{a(p)}{p} = \infty.$$

By the independence of  $\{Y_p\}_p$  and Borel-Cantelli's second lemma,

$$\lim_{\sigma' \searrow 1} \text{the second term} = \infty \quad \text{a.s.}$$

When  $\tau = 0$ ,  $\{p; a(p) = p\} \neq \emptyset$ . By noting that  $Y_p(1-) = \infty$  a.s.,

$$\lim_{\sigma' \searrow 1} \text{the first term} = \infty \quad \text{a.s.}$$

Thus  $Z(1+; a) = \infty$  a.s.

As  $\sigma \nearrow \infty$ ,  $\frac{a(p)}{p^\sigma} \searrow 0$ , and hence  $Y_p\left(\frac{a(p)}{p^\sigma}\right) \searrow Y_p(0) = 0$ . By the Lebesgue convergence theorem,  $Z(\infty; a) = 0$ .

From Definition 1(c) and the independence of  $\{Y_p\}_p$ , it follows that for every  $\infty > \sigma_0 > \sigma_1 > \dots > \sigma_n > 1$ ,

$$Z(\sigma_0; a), Z(\sigma_1; a) - Z(\sigma_0; a), \dots, Z(\sigma_n; a) - Z(\sigma_{n-1}; a) \text{ are independent.}$$

3° For  $t \in \mathbb{R}$ ,

$$\begin{aligned} E[e^{it(-Z(\sigma;a))}] &= \prod_p E\left[e^{it(\log \frac{1}{p})Y_p\left(\frac{a(p)}{p^\sigma}\right)}\right] \\ &= \prod_p \frac{1 - \frac{a(p)}{p^\sigma}}{1 - \frac{a(p)}{p^{\sigma+it}}} \\ &= \frac{\eta(\sigma + it; a)}{\eta(\sigma; a)} = \eta_\sigma(t; a) = \int_{\mathbb{R}} e^{itx} \mu_\sigma(dx; a). \end{aligned}$$

□

**2.4. Proof and corollary of Theorem 1.** We are now in position to prove Theorem 1.

Proof of Theorem 1. Let us fix the  $\varphi(\cdot)$  in Theorem 1. The proof is divided into 2 steps.

1° Fix  $T > 0$ . By Claim 7,  $(X_T(t; a) := \frac{1}{\sqrt{T}}Z(\varphi^{-1}(Tt); a))_{t \geq 0}$  is, in the usual sense, a Lévy process with increasing paths (cf. Itô [6]). Here we set  $X_T(0; a) := 0$  by  $\varphi^{-1}(0) = \varphi^{-1}(0+) = \infty$  and  $Z(\infty; a) = 0$ .

Let  $N_T(dsdu)$  be a Poisson random measure on  $(0, \infty) \times (0, \infty)$  defined by  $X_T(\cdot; a)$ :

$$N_T(A) := \#\{t > 0; (t, X_T(t; a) - X_T(t-; a)) \in A\}, \quad A \in \mathcal{B}((0, \infty) \times (0, \infty)).$$

Then, the Lévy-Itô decomposition of  $X_T(\cdot; a)$  is given as

$$X_T(t; a) = \int_0^{t+} \int_{(0, \infty)} u N_T(dsdu), \quad t \geq 0.$$

And, a mean measure  $n_T(dsdu)$  of  $N_T(dsdu)$  is given as

$$(19) \quad n_T(dsdu) = \sum_p \sum_{n=1}^{\infty} (a(p)p^{-\varphi^{-1}(Ts)})^n (\log p) \frac{-T}{\varphi'(\varphi^{-1}(Ts))} ds \delta_{\frac{1}{\sqrt{T}}n \log p}(du).$$

Proof. Temporarily let  $n'_T(dsdu)$  be a right-hand side of (19). Clearly  $n'_T(dsdu)$  is a measure on  $(0, \infty) \times (0, \infty)$ , and

$$\begin{aligned} \iint_{\substack{0 < s \leq t \\ u > 0}} un'_T(dsdu) &= \sum_p \sum_{n=1}^{\infty} \int_0^t (a(p)p^{-\varphi^{-1}(Ts)})^n (\log p) \frac{-T}{\varphi'(\varphi^{-1}(Ts))} \frac{1}{\sqrt{T}} n(\log p) ds \\ &= \frac{1}{\sqrt{T}} \sum_p \sum_{n=1}^{\infty} \int_{\varphi^{-1}(Tt)}^{\infty} (a(p)p^{-r})^n n(\log p)^2 dr \\ &\quad [\text{by the change of variables } \varphi^{-1}(Ts) = r] \\ &= \frac{1}{\sqrt{T}} \sum_p \left( \sum_{n=1}^{\infty} (a(p)p^{-\varphi^{-1}(Tt)})^n \right) \log p \\ &= \frac{1}{\sqrt{T}} \sum_p \frac{\frac{a(p)}{p^{\varphi^{-1}(Tt)}} \log p}{1 - \frac{a(p)}{p^{\varphi^{-1}(Tt)}}} = \frac{1}{\sqrt{T}} \left( -\frac{\eta'(\sigma; a)}{\eta(\sigma; a)} \right) \Big|_{\sigma=\varphi^{-1}(Tt)} < \infty. \end{aligned}$$

By this and  $\iint_{\substack{0 < s \leq t \\ u > 0}} (u \wedge 1) n_T(dsdu) < \infty$ , it suffices to check that for  $t$  and  $\lambda > 0$ ,

$$(20) \quad E[e^{-\lambda X_T(t; a)}] = \exp\left\{- \iint_{\substack{0 < s \leq t \\ u > 0}} (1 - e^{-\lambda u}) n'_T(dsdu)\right\}.$$

For, let

$$X_T(t; a) = m(t) + \int_0^{t+} \int_{(0, \infty)} u N_T(dsdu)$$

be the Lévy-Itô decomposition of  $X_T(\cdot; a)$ , where  $m(\cdot)$  is a deterministic, continuous and non-decreasing process with  $m(0) = 0$ . Then

$$E[e^{-\lambda X_T(t;a)}] = \exp\left\{-\lambda m(t) - \iint_{\substack{0 < s \leq t \\ u > 0}} (1 - e^{-\lambda u}) n_T(dsdu)\right\}.$$

If, moreover, (20) holds, then

$$\begin{aligned} m(t) &= \iint_{\substack{0 < s \leq t \\ u > 0}} \frac{1 - e^{-\lambda u}}{\lambda} n'_T(dsdu) - \iint_{\substack{0 < s \leq t \\ u > 0}} \frac{1 - e^{-\lambda u}}{\lambda} n_T(dsdu) \\ &\rightarrow 0 \quad \text{as } \lambda \rightarrow \infty. \end{aligned}$$

This convergence immediately follows from the Lebesgue convergence theorem since  $0 < \frac{1 - e^{-\lambda u}}{\lambda} \leq \frac{1}{\lambda}((\lambda u) \wedge 1) = u \wedge (\frac{1}{\lambda}) \leq u \wedge 1$  ( $\forall \lambda \geq 1$ ),  $\iint_{\substack{0 < s \leq t \\ u > 0}} (u \wedge 1) n'_T(dsdu) < \infty$ ,  $\iint_{\substack{0 < s \leq t \\ u > 0}} (u \wedge 1) n_T(dsdu) < \infty$  and  $\lim_{\lambda \rightarrow \infty} \frac{1 - e^{-\lambda u}}{\lambda} = 0$  ( $\forall u > 0$ ). This implies that

$$X_T(t; a) = \int_0^{t+} \int_{(0, \infty)} u N_T(dsdu), \quad n_T(dsdu) = n'_T(dsdu).$$

(20) is shown in the following way: By Definition 5, Definition 1(d) and (1),

$$\begin{aligned} E[e^{-\lambda X_T(t;a)}] &= \prod_p E\left[e^{-\frac{\lambda}{\sqrt{T}}(\log p) Y_p(a(p)p^{-\varphi^{-1}(Tt)})}\right] \\ &= \prod_p \frac{1 - a(p)p^{-\varphi^{-1}(Tt)}}{1 - a(p)p^{-\varphi^{-1}(Tt)} p^{-\frac{\lambda}{\sqrt{T}}}} \\ &= \prod_p \exp\left\{\int_0^1 \frac{-a(p)p^{-\varphi^{-1}(Tt)}}{1 - a(p)p^{-\varphi^{-1}(Tt)} s} ds - \int_0^1 \frac{-a(p)p^{-\varphi^{-1}(Tt)} p^{-\frac{\lambda}{\sqrt{T}}}}{1 - a(p)p^{-\varphi^{-1}(Tt)} p^{-\frac{\lambda}{\sqrt{T}}}} ds\right\} \\ &= \prod_p \exp\left\{\int_0^t \frac{-a(p)p^{-\varphi^{-1}(Tr)}}{1 - a(p)p^{-\varphi^{-1}(Tr)}} p^{\varphi^{-1}(Tr)} p^{-\varphi^{-1}(Tr)} (-\log p) \frac{T dr}{\varphi'(\varphi^{-1}(Tr))}\right. \\ &\quad \left. + \int_0^t \frac{a(p)p^{-\varphi^{-1}(Tt)} p^{-\frac{\lambda}{\sqrt{T}}}}{1 - a(p)p^{-\varphi^{-1}(Tr)} p^{-\frac{\lambda}{\sqrt{T}}}} p^{\varphi^{-1}(Tr)} p^{-\varphi^{-1}(Tr)} (-\log p) \frac{T dr}{\varphi'(\varphi^{-1}(Tr))}\right\} \\ &\quad [\text{by the change of variables } r = \frac{1}{T}\varphi(\varphi^{-1}(Tt) + \frac{\log \frac{1}{s}}{\log p})] \\ &= \prod_p \exp\left\{\int_0^t \sum_{n=1}^{\infty} (a(p)p^{-\varphi^{-1}(Tr)})^n (-1 + e^{-\lambda \frac{n}{\sqrt{T}} \log p})(\log p) \frac{-T}{\varphi'(\varphi^{-1}(Tr))} dr\right\} \\ &= \exp\left\{-\iint_{\substack{0 < s \leq t \\ u > 0}} (1 - e^{-\lambda u}) n'_T(dsdu)\right\}. \quad \square \end{aligned}$$

2° By Claims 8 and 9 below,

$$\begin{aligned} n_T(dsdu) &\rightarrow n^{(\rho)}(dsdu) \quad \text{vaguely as } T \rightarrow \infty, \\ \lim_{k \rightarrow \infty} \limsup_{T \rightarrow \infty} \iint_{\substack{0 < s \leq t \\ u \in (0, \infty) \setminus [\frac{1}{k}, k]}} un_T(dsdu) &= 0, \quad \forall t > 0. \end{aligned}$$



Applying the general theory of Kasahara-Watanabe [7], we have

$$\left(\int_0^{t+} \int_{(0,\infty)} uN_T(dsdu)\right)_{t \geq 0} \xrightarrow{D} \left(\int_0^{t+} \int_{(0,\infty)} uN^{(\rho)}(dsdu)\right)_{t \geq 0}$$

in  $D([0, \infty) \rightarrow \mathbb{R})$  as  $T \rightarrow \infty$ ,

which is the assertion of the theorem. □

As for Claims 8 and 9, we begin with the following lemma:

**Lemma 1.** For  $\lambda \geq 0$  and  $t > 0$ ,

$$\lim_{T \rightarrow \infty} \iint_{\substack{0 < s \leq t \\ u > 0}} e^{-\lambda u} uN_T(dsdu) = \frac{\rho}{\lambda + \frac{1}{\sqrt{t}}} = \iint_{\substack{0 < s \leq t \\ u > 0}} e^{-\lambda u} uN^{(\rho)}(dsdu).$$

Proof. Fix  $\lambda \geq 0$  and  $t > 0$ . By (19),

$$\begin{aligned} & \iint_{\substack{0 < s \leq t \\ u > 0}} e^{-\lambda u} uN_T(dsdu) \\ &= \sum_p \sum_{n=1}^{\infty} \int_0^t e^{-\lambda \frac{1}{\sqrt{T}} n \log p} \frac{1}{\sqrt{T}} n(\log p)(a(p)p^{-\varphi^{-1}(Ts)})^n (\log p) \frac{-T}{\varphi'(\varphi^{-1}(Ts))} ds \\ &= \frac{1}{\sqrt{T}} \sum_p (\log p) \sum_{n=1}^{\infty} (a(p)p^{-\varphi^{-1}(Tt) - \frac{\lambda}{\sqrt{T}}})^n \\ &= \frac{1}{\sqrt{T}} \sum_p \frac{\frac{a(p)}{p^{\varphi^{-1}(Tt) + \frac{\lambda}{\sqrt{T}}}} \log p}{1 - \frac{a(p)}{p^{\varphi^{-1}(Tt) + \frac{\lambda}{\sqrt{T}}}}} = \frac{1}{\sqrt{T}} \left( -\frac{\eta'(\sigma; a)}{\eta(\sigma; a)} \right) \Big|_{\sigma = \varphi^{-1}(Tt) + \frac{\lambda}{\sqrt{T}}}. \end{aligned}$$

Since  $\varphi^{-1}(Tt) \searrow 1$  as  $T \rightarrow \infty$ , and thus  $\sigma = \varphi^{-1}(Tt) + \frac{\lambda}{\sqrt{T}} \searrow 1$ , Claim 6 tells us that

$$(\sigma - 1) \left( -\frac{\eta'(\sigma; a)}{\eta(\sigma; a)} \right) \rightarrow \rho.$$

On the other hand, by (8),

$$\varphi^{-1}(Tt) - 1 \sim \frac{1}{\sqrt{\varphi(\varphi^{-1}(Tt))}} = \frac{1}{\sqrt{T} \sqrt{t}},$$

and thus

$$\sqrt{T}(\sigma - 1) = \sqrt{T} \left( \varphi^{-1}(Tt) + \frac{\lambda}{\sqrt{T}} - 1 \right) = \lambda + \sqrt{T}(\varphi^{-1}(Tt) - 1) \rightarrow \lambda + \frac{1}{\sqrt{t}}.$$

Combining two convergences above, we have

$$\frac{1}{\sqrt{T}} \left( -\frac{\eta'(\sigma; a)}{\eta(\sigma; a)} \right) = \frac{1}{\sqrt{T}(\sigma - 1)} \cdot (\sigma - 1) \left( -\frac{\eta'(\sigma; a)}{\eta(\sigma; a)} \right) \rightarrow \frac{\rho}{\lambda + \frac{1}{\sqrt{t}}}.$$

Next, by (9),

$$\begin{aligned} \iint_{\substack{0 < s \leq t \\ u > 0}} e^{-\lambda u} u n^{(\rho)}(dsdu) &= \frac{\rho}{2} \int_0^t s^{-\frac{3}{2}} ds \int_0^\infty u e^{-(\frac{1}{\sqrt{s}} + \lambda)u} du \\ &= \frac{\rho}{2} \int_0^t s^{-\frac{3}{2}} \frac{ds}{(\frac{1}{\sqrt{s}} + \lambda)^2} \\ &\quad [\text{since } \int_0^\infty u e^{-\mu u} du = \frac{1}{\mu^2} \ (\mu > 0)] \\ &= \frac{\rho}{\lambda + \frac{1}{\sqrt{t}}}. \end{aligned}$$

□

**Claim 8.** As  $T \rightarrow \infty$ ,  $n_T(dsdu) \rightarrow n^{(\rho)}(dsdu)$  vaguely. That is, for  $\forall j \in C_c([0, \infty) \times (0, \infty))$ ,

$$\iint_{(0, \infty) \times (0, \infty)} j(s, u) n_T(dsdu) \rightarrow \iint_{(0, \infty) \times (0, \infty)} j(s, u) n^{(\rho)}(dsdu).$$

Here  $C_c([0, \infty) \times (0, \infty))$  is the set of all real-valued continuous functions on  $[0, \infty) \times (0, \infty)$  with compact support.

Proof. It is divided into 6 steps.

1° For  $\lambda \geq 0$ , let  $f_\lambda(u) = e^{-\lambda u}$ . Then

$$f_\lambda \in C_\infty([0, \infty)) \text{ if } \lambda > 0, \quad f_0 = 1 \in C_b([0, \infty)).$$

Here

$C_b([0, \infty))$  = the set of all real-valued, bounded, continuous functions on  $[0, \infty)$ ,

$C_\infty([0, \infty)) = \{f \in C_b([0, \infty)); \lim_{u \rightarrow \infty} f(u) = 0\}$ ,

$C_c([0, \infty)) = \{f \in C_b([0, \infty)); \text{supp } f \text{ is compact}\}$ .

Let  $[0, \infty]$  be the one-point compactification of  $[0, \infty)$ . If, at point  $\infty$ , we define

$$f_\lambda(\infty) := \begin{cases} 0 & \text{if } \lambda > 0, \\ 1 & \text{if } \lambda = 0, \end{cases}$$

then  $f_\lambda \in C([0, \infty])^2$ . Letting  $\mathcal{A} \subset C([0, \infty])$  be the set of all linear combinations of  $f_\lambda$ ,  $\lambda \geq 0$ , we can check that

- $\mathcal{A}$  is an algebra,
- $\mathcal{A}$  separates points on  $[0, \infty]$ ,
- $\mathcal{A}$  vanishes at no point of  $[0, \infty]$ .

Thus, by the Stone-Weierstrass theorem (cf. [9, Theorem 7.32]),  $\overline{\mathcal{A}} = C([0, \infty])$ . Particularly, for  $\forall f \in C_c([0, \infty))$  and  $\forall \varepsilon > 0$ ,

$$\exists g \in \mathcal{A} \text{ s.t. } \sup_{0 \leq u < \infty} |f(u) - g(u)| < \varepsilon.$$

---

<sup>2</sup>The extension of  $f_\lambda$  to  $[0, \infty]$  is denoted by the same symbol  $f_\lambda$ .

2° For  $\forall t > 0$  and  $\forall f \in C_c([0, \infty))$ ,

$$\lim_{T \rightarrow \infty} \iint_{\substack{0 < s \leq t \\ u > 0}} f(u)un_T(dsdu) = \iint_{\substack{0 < s \leq t \\ u > 0}} f(u)un^{(\rho)}(dsdu).$$

Proof. Fix  $t > 0$  and  $f \in C_c([0, \infty))$ . By 1°,  $f$  can be approximated by a sequence  $\{g_k\}$  of  $\mathcal{A}$ . Since, by Lemma 1,

$$\lim_{T \rightarrow \infty} \iint_{\substack{0 < s \leq t \\ u > 0}} g_k(u)un_T(dsdu) = \iint_{\substack{0 < s \leq t \\ u > 0}} g_k(u)un^{(\rho)}(dsdu),$$

it follows in a routine way that

$$\lim_{T \rightarrow \infty} \iint_{\substack{0 < s \leq t \\ u > 0}} f(u)un_T(dsdu) = \iint_{\substack{0 < s \leq t \\ u > 0}} f(u)un^{(\rho)}(dsdu).$$

□

3° For  $\forall t > 0$ ,

$$\lim_{\lambda \searrow 0} \limsup_{T \rightarrow \infty} \iint_{\substack{0 < s \leq t \\ u \geq \frac{1}{\lambda}}} un_T(dsdu) = 0.$$

Proof. Fix  $t > 0$ . Noting that for  $\lambda > 0$  and  $u > 0$ ,

$$1 - e^{-\lambda u} = \int_0^u (-e^{-\lambda v})' dv = \int_{(0, \infty)} \mathbf{1}_{v \leq u} \lambda e^{-\lambda v} dv,$$

we obtain the following lower estimate:

$$\begin{aligned} & \iint_{\substack{0 < s \leq t \\ u > 0}} un_T(dsdu) - \iint_{\substack{0 < s \leq t \\ u > 0}} e^{-\lambda u} un_T(dsdu) \\ &= \iint_{\substack{0 < s \leq t \\ u > 0}} (1 - e^{-\lambda u}) un_T(dsdu) \\ &= \iint_{\substack{0 < s \leq t \\ u > 0}} un_T(dsdu) \int_{(0, \infty)} \mathbf{1}_{v \leq u} \lambda e^{-\lambda v} dv \\ &= \int_{(0, \infty)} \lambda e^{-\lambda v} dv \iint_{\substack{0 < s \leq t \\ u \geq v}} un_T(dsdu) \\ &= \int_{(0, \infty)} e^{-w} dw \iint_{\substack{0 < s \leq t \\ u \geq \frac{w}{\lambda}}} un_T(dsdu) \quad [\text{by the change of variables } \lambda v = w] \\ &\geq \int_0^1 e^{-w} dw \iint_{\substack{0 < s \leq t \\ u \geq \frac{1}{\lambda}}} un_T(dsdu) \quad [\text{since } [\frac{w}{\lambda}, \infty) \supset [\frac{1}{\lambda}, \infty) \text{ for } 0 < w \leq 1] \end{aligned}$$

$$= (1 - e^{-1}) \iint_{\substack{0 < s \leq t \\ u \geq \frac{1}{\lambda}}} un_T(dsdu).$$

By Lemma 1,


$$\begin{aligned} \limsup_{T \rightarrow \infty} \iint_{\substack{0 < s \leq t \\ u \geq \frac{1}{\lambda}}} un_T(dsdu) &\leq \frac{1}{1 - e^{-1}} \lim_{T \rightarrow \infty} \left( \iint_{\substack{0 < s \leq t \\ u > 0}} un_T(dsdu) - \iint_{\substack{0 < s \leq t \\ u > 0}} e^{-\lambda u} un_T(dsdu) \right) \\ &= \frac{\rho}{1 - e^{-1}} \frac{\lambda t}{1 + \lambda \sqrt{t}} \rightarrow 0 \quad \text{as } \lambda \searrow 0. \end{aligned}$$

□

4° For  $\forall t > 0$  and  $\forall f \in C_b([0, \infty))$ ,

$$\lim_{T \rightarrow \infty} \iint_{\substack{0 < s \leq t \\ u > 0}} f(u)un_T(dsdu) = \iint_{\substack{0 < s \leq t \\ u > 0}} f(u)un^{(\rho)}(dsdu).$$

Proof. Fix  $t > 0$  and  $f \in C_b([0, \infty))$ . For each  $m \in \mathbb{N}$ , set  $h_m \in C_c([0, \infty))$  by



Note that  $f \cdot h_m \in C_c([0, \infty))$  and

$$(21) \quad |f(u) - (f \cdot h_m)(u)| = |f(u)|(1 - h_m(u)) \leq \|f\|_{\infty} \mathbf{1}_{[m, \infty)}(u).$$

By 2°,

$$\lim_{T \rightarrow \infty} \iint_{\substack{0 < s \leq t \\ u > 0}} (f \cdot h_m)(u)un_T(dsdu) = \iint_{\substack{0 < s \leq t \\ u > 0}} (f \cdot h_m)(u)un^{(\rho)}(dsdu).$$

Also, by (21) and 3°,

$$\limsup_{T \rightarrow \infty} \iint_{\substack{0 < s \leq t \\ u > 0}} |(f \cdot h_m)(u) - f(u)|un_T(dsdu) \rightarrow 0 \quad \text{as } m \rightarrow \infty.$$

Thus, it follows in the usual way that

$$\lim_{T \rightarrow \infty} \iint_{\substack{0 < s \leq t \\ u > 0}} f(u)un_T(dsdu) = \iint_{\substack{0 < s \leq t \\ u > 0}} f(u)un^{(\rho)}(dsdu).$$

□

5° For  $\forall h \in C_c([0, \infty) \times [0, \infty))$ ,

$$\lim_{T \rightarrow \infty} \iint_{(0, \infty) \times (0, \infty)} h(s, u)un_T(dsdu) = \iint_{(0, \infty) \times (0, \infty)} h(s, u)un^{(\rho)}(dsdu).$$

Proof. Fix  $h \in C_c([0, \infty) \times [0, \infty))$ . Since  $h$  is uniformly continuous on  $[0, \infty) \times [0, \infty)$ ,

$$(22) \quad \forall \varepsilon > 0, \exists \delta > 0 \text{ s.t. } |s - s'| < \delta, |u - u'| < \delta \Rightarrow |h(s, u) - h(s', u')| < \varepsilon.$$

Also, since  $\text{supp } h$  is compact,

$$\exists t > 0 \text{ s.t. } \text{supp } h \subset [0, t] \times [0, t].$$

Take a large  $n \in \mathbb{N}$  such that  $\frac{t}{n} < \delta$ , and rewrite

$$\begin{aligned} \iint_{(0, \infty) \times (0, \infty)} h(s, u) un_T(dsdu) &= \iint_{(0, t] \times (0, t]} h(s, u) un_T(dsdu) \\ &= \sum_{k=1}^n \iint_{\substack{\frac{k-1}{n}t < s \leq \frac{k}{n}t \\ u > 0}} h\left(\frac{k-1}{n}t, u\right) un_T(dsdu) \\ &\quad + \sum_{k=1}^n \iint_{(\frac{k-1}{n}t, \frac{k}{n}t] \times (0, t]} \left(h(s, u) - h\left(\frac{k-1}{n}t, u\right)\right) un_T(dsdu), \\ \iint_{(0, \infty) \times (0, \infty)} h(s, u) un^{(\rho)}(dsdu) &= \sum_{k=1}^n \iint_{\substack{\frac{k-1}{n}t < s \leq \frac{k}{n}t \\ u > 0}} h\left(\frac{k-1}{n}t, u\right) un^{(\rho)}(dsdu) \\ &\quad + \sum_{k=1}^n \iint_{(\frac{k-1}{n}t, \frac{k}{n}t] \times (0, t]} \left(h(s, u) - h\left(\frac{k-1}{n}t, u\right)\right) un^{(\rho)}(dsdu). \end{aligned}$$

Then, by (22),

$$\begin{aligned} &\left| \iint_{(0, \infty) \times (0, \infty)} h(s, u) un_T(dsdu) - \iint_{(0, \infty) \times (0, \infty)} h(s, u) un^{(\rho)}(dsdu) \right| \\ &\leq \left| \sum_{k=1}^n \left( \iint_{\substack{0 < s \leq \frac{k}{n}t \\ u > 0}} h\left(\frac{k-1}{n}t, u\right) un_T(dsdu) - \iint_{\substack{0 < s \leq \frac{k}{n}t \\ u > 0}} h\left(\frac{k-1}{n}t, u\right) un^{(\rho)}(dsdu) \right. \right. \\ &\quad \left. \left. - \left( \iint_{\substack{0 < s \leq \frac{k-1}{n}t \\ u > 0}} h\left(\frac{k-1}{n}t, u\right) un_T(dsdu) - \iint_{\substack{0 < s \leq \frac{k-1}{n}t \\ u > 0}} h\left(\frac{k-1}{n}t, u\right) un^{(\rho)}(dsdu) \right) \right) \right| \\ &\quad + \sum_{k=1}^n \left( \iint_{(\frac{k-1}{n}t, \frac{k}{n}t] \times (0, t]} \left| h(s, u) - h\left(\frac{k-1}{n}t, u\right) \right| un_T(dsdu) \right. \\ &\quad \left. + \iint_{(\frac{k-1}{n}t, \frac{k}{n}t] \times (0, t]} \left| h(s, u) - h\left(\frac{k-1}{n}t, u\right) \right| un^{(\rho)}(dsdu) \right) \\ &\leq \left| \sum_{k=1}^n \left( \iint_{\substack{0 < s \leq \frac{k}{n}t \\ u > 0}} h\left(\frac{k-1}{n}t, u\right) un_T(dsdu) - \iint_{\substack{0 < s \leq \frac{k}{n}t \\ u > 0}} h\left(\frac{k-1}{n}t, u\right) un^{(\rho)}(dsdu) \right) \right| \end{aligned}$$

$$\begin{aligned}
 & - \left( \iint_{\substack{0 < s \leq \frac{k-1}{n}t \\ u > 0}} h\left(\frac{k-1}{n}t, u\right) un_T(dsdu) - \iint_{\substack{0 < s \leq \frac{k-1}{n}t \\ u > 0}} h\left(\frac{k-1}{n}t, u\right) un^{(\rho)}(dsdu) \right) \\
 & + \varepsilon \left( \iint_{\substack{0 < s \leq t \\ u > 0}} un_T(dsdu) + \iint_{\substack{0 < s \leq t \\ u > 0}} un^{(\rho)}(dsdu) \right)
 \end{aligned}$$

=: the first term + the second term.

By 4°,  $\lim_{T \rightarrow \infty}$  the first term = 0 since  $h(\frac{k-1}{n}t, \cdot) \in C_b([0, \infty))$ , and by Lemma 1,  $\lim_{T \rightarrow \infty}$  the second term =  $\varepsilon \frac{2\rho}{1/\sqrt{t}} \rightarrow 0$  as  $\varepsilon \searrow 0$ . Thus we have the assertion of 5°.  $\square$

6° For  $\forall j \in C_c([0, \infty) \times (0, \infty))$ , set

$$h(s, u) := \begin{cases} \frac{1}{u} j(s, u) & \text{if } u > 0, \\ 0 & \text{if } u = 0. \end{cases}$$

Then  $h \in C_c([0, \infty) \times [0, \infty))$  and  $j(s, u) = h(s, u)u$ . By 5°,

$$\begin{aligned}
 \lim_{T \rightarrow \infty} \iint_{(0, \infty) \times (0, \infty)} j(s, u) n_T(dsdu) &= \lim_{T \rightarrow \infty} \iint_{(0, \infty) \times (0, \infty)} h(s, u) un_T(dsdu) \\
 &= \iint_{(0, \infty) \times (0, \infty)} h(s, u) un^{(\rho)}(dsdu) \\
 &= \iint_{(0, \infty) \times (0, \infty)} j(s, u) n^{(\rho)}(dsdu).
 \end{aligned}$$

$\square$


**Claim 9.** For  $\forall t > 0$ ,

$$\lim_{k \rightarrow \infty} \limsup_{T \rightarrow \infty} \iint_{\substack{0 < s \leq t \\ u \in (0, \infty) \setminus [\frac{1}{k}, k]}} un_T(dsdu) = 0.$$

Proof. Fix  $t > 0$ . First

$$\begin{aligned}
 \iint_{\substack{0 < s \leq t \\ u \in (0, \infty) \setminus [\frac{1}{k}, k]}} un_T(dsdu) &= \iint_{\substack{0 < s \leq t \\ 0 < u < \frac{1}{k}}} un_T(dsdu) + \iint_{\substack{0 < s \leq t \\ u > k}} un_T(dsdu) \\
 &\leq \iint_{\substack{0 < s \leq t \\ u > 0}} j_k(u) un_T(dsdu) + \iint_{\substack{0 < s \leq t \\ u \geq \frac{1}{k}}} un_T(dsdu).
 \end{aligned}$$

Here  $j_k \in C_c([0, \infty))$  is as follows:



By 2° and 3° in the proof of Claim 8,

$$\begin{aligned} & \limsup_{T \rightarrow \infty} \iint_{\substack{0 < s \leq t \\ u \in (0, \infty) \setminus [\frac{1}{k}, k]}} un_T(dsdu) \\ & \leq \iint_{\substack{0 < s \leq t \\ u > 0}} j_k(u)un^{(\rho)}(dsdu) + \limsup_{T \rightarrow \infty} \iint_{\substack{0 < s \leq t \\ u \geq \frac{1}{k}}} un_T(dsdu) \rightarrow 0 \quad \text{as } k \rightarrow \infty. \end{aligned}$$

□

**Corollary 1.** For each  $t > 0$ ,

the distribution of  $\frac{1}{\sqrt{T}}Z(\varphi^{-1}(Tt); a) \rightarrow \mathbf{1}_{(0, \infty)}(x) \frac{1}{\Gamma(\rho)}(\sqrt{t})^{-\rho} x^{\rho-1} e^{-\frac{x}{\sqrt{t}}} dx$  as  $T \rightarrow \infty$ .

(The limiting distribution is the gamma distribution with parameters  $\rho, \sqrt{t}$ .) In particular, letting  $t = 1$  and  $\sigma = \varphi^{-1}(T)$  and then noting that

$$T \rightarrow \infty \Leftrightarrow \sigma \searrow 1, \quad (\sigma - 1)^2 \varphi(\sigma) \rightarrow 1 \quad \text{as } \sigma \searrow 1$$

tell us that

the distribution of  $(\sigma - 1)Z(\sigma; a) \rightarrow \mathbf{1}_{(0, \infty)}(x) \frac{1}{\Gamma(\rho)} x^{\rho-1} e^{-x} dx$  as  $\sigma \searrow 1$ .

Proof. Fix  $\lambda > 0$  and  $t > 0$ . By Theorem 1,

$$\begin{aligned} \lim_{T \rightarrow \infty} E[e^{-\lambda \frac{1}{\sqrt{T}} Z(\varphi^{-1}(Tt); a)}] &= E[e^{-\lambda \int_0^{t+} \int_{(0, \infty)} un^{(\rho)}(dsdu)}] \\ &= \exp\left\{- \iint_{\substack{0 < s \leq t \\ u > 0}} (1 - e^{-\lambda u}) n^{(\rho)}(dsdu)\right\}. \end{aligned}$$

Here, by Lemma 1,

$$\begin{aligned} \iint_{\substack{0 < s \leq t \\ u > 0}} (1 - e^{-\lambda u}) n^{(\rho)}(dsdu) &= \iint_{\substack{0 < s \leq t \\ u > 0}} \int_0^1 e^{-a\lambda u} \lambda u da n^{(\rho)}(dsdu) \\ &= \lambda \int_0^1 \frac{\rho}{a\lambda + \frac{1}{\sqrt{t}}} da = \rho \log(1 + \sqrt{t}\lambda). \end{aligned}$$

Substituting this into the last right-hand side of the preceding expression, we have

$$\begin{aligned} \lim_{T \rightarrow \infty} E[e^{-\lambda \frac{1}{\sqrt{t}} Z(\varphi^{-1}(Tt); a)}] &= \exp\{-\rho \log(1 + \sqrt{t}\lambda)\} \\ &= (1 + \sqrt{t}\lambda)^{-\rho} \\ &= \frac{1}{\Gamma(\rho)} \int_0^\infty e^{-(1+\sqrt{t}\lambda)y} y^{\rho-1} dy \\ &\quad [\text{since } \mu^{-\rho} = \frac{1}{\Gamma(\rho)} \int_0^\infty e^{-\mu y} y^{\rho-1} dy \quad (\mu > 0)] \\ &= \frac{1}{\Gamma(\rho)} \int_0^\infty e^{-\lambda x} (\sqrt{t})^{-\rho} x^{\rho-1} e^{-\frac{x}{\sqrt{t}}} dx \\ &\quad [\text{by the change of variables } x = \sqrt{t}y], \end{aligned}$$

which shows the assertion of the corollary. □

### 3. Examples of arithmetical function $a(\cdot)$

EXAMPLE 1. Let a sequence  $(a(p))_p$  be nonnegative, i.e.,  $a(p) \geq 0$  ( $\forall p$ ). If  $a(p) \rightarrow c \in [0, \infty)$  as  $p \rightarrow \infty$ , then

$$\sum_{p \leq x} \frac{a(p) \log p}{p} = (c + o(1)) \log x \quad \text{as } x \rightarrow \infty.$$

Thus the condition (4) holds with  $\tau = c$ .

Proof. By Mertens' first theorem (6),

$$\begin{aligned} &\left| \frac{1}{\log x} \sum_{p \leq x} \frac{a(p) \log p}{p} - c \right| \\ &= \left| \frac{1}{\log x} \sum_{p \leq x} \frac{c \log p}{p} + \frac{1}{\log x} \sum_{p \leq x} \frac{(a(p) - c) \log p}{p} - c \right| \\ &= \left| c \frac{\log x + O(1)}{\log x} - c \right. \\ &\quad \left. + \frac{1}{\log x} \sum_{p \leq y} \frac{(a(p) - c) \log p}{p} + \frac{1}{\log x} \sum_{y < p \leq x} \frac{(a(p) - c) \log p}{p} \right| \\ &\quad [\text{where we fix } y \in (2, x) \text{ arbitrarily}] \\ &\leq \frac{c|O(1)|}{\log x} + \frac{1}{\log x} \left| \sum_{p \leq y} \frac{(a(p) - c) \log p}{p} \right| + \left( \sup_{p > y} |a(p) - c| \right) \left( 1 + \frac{|O(1)|}{\log x} \right). \end{aligned}$$

By letting  $x \rightarrow \infty$ ,

$$\limsup_{x \rightarrow \infty} \left| \frac{1}{\log x} \sum_{p \leq x} \frac{a(p) \log p}{p} - c \right| \leq \sup_{p > y} |a(p) - c| \rightarrow 0 \quad \text{as } y \rightarrow \infty.$$

This shows the assertion of Example 1. □



EXAMPLE 2. For coprime  $a, m \in \mathbb{N}$ , we set  $E_{a,m} := \{p; p \equiv a \pmod{m}\}$ . Then

$$\sum_{p \leq x} \frac{\mathbf{1}_{E_{a,m}}(p) \log p}{p} = \left(\frac{1}{\phi(m)} + o(1)\right) \log x \quad \text{as } x \rightarrow \infty,$$

where  $\phi(\cdot)$  is Euler's function. Thus,  $(\mathbf{1}_{E_{a,m}}(p))_p$  satisfies the condition (4) with  $\tau = \frac{1}{\phi(m)}$ .

Proof. We use the prime number theorem for arithmetic progressions in the following form:

$$(23) \quad \vartheta_{a,m}(x) := \sum_{p \leq x} \mathbf{1}_{E_{a,m}}(p) \log p = \left(\frac{1}{\phi(m)} + o(1)\right)x \quad \text{as } x \rightarrow \infty.$$

$\vartheta_{a,m}(\cdot)$  is non-decreasing, right-continuous and  $\vartheta_{a,m}(t) = 0$  ( $\forall t < 2$ ). Noting that

$$\vartheta_{a,m}(dt) = \sum_p \mathbf{1}_{E_{a,m}}(p) (\log p) \delta_p(dt),$$

we compute that for  $0 < \varepsilon < 1$ ,

$$\begin{aligned} \sum_{p \leq x} \frac{\mathbf{1}_{E_{a,m}}(p) \log p}{p} &= \int_{(2-\varepsilon, x]} \frac{1}{t} \vartheta_{a,m}(dt) \\ &= \int_{(2-\varepsilon, x]} \frac{1}{t} d\left(\frac{t}{\phi(m)} + t\delta(t)\right) \\ &\quad [\text{where } \delta(t) := \vartheta_{a,m}(t)/t - 1/\phi(m) \ (t \geq 1)] \\ &= \frac{1}{\phi(m)} \int_{2-\varepsilon}^x \frac{dt}{t} + \int_{2-\varepsilon}^x \frac{\delta(t)}{t} dt + \delta(x) + \frac{1}{\phi(m)}. \end{aligned}$$

By letting  $\varepsilon \searrow 0$ ,

$$\begin{aligned} \sum_{p \leq x} \frac{\mathbf{1}_{E_{a,m}}(p) \log p}{p} &= \frac{1}{\phi(m)} \int_2^x \frac{dt}{t} + \int_2^x \frac{\delta(t)}{t} dt + \delta(x) + \frac{1}{\phi(m)} \\ &= (\log x) \left(\frac{1}{\phi(m)} \left(1 + \frac{-\log 2 + 1}{\log x}\right) + \int_{\frac{\log 2}{\log x}}^1 \delta(x^r) dr + \frac{\delta(x)}{\log x}\right) \\ &\quad [\text{by the change of variables } \frac{\log t}{\log x} = r]. \end{aligned}$$

Since, from  $\lim_{x \rightarrow \infty} \delta(x) = 0$  (cf. (23)) and the bounded convergence theorem,

$$\lim_{x \rightarrow \infty} \int_{\frac{\log 2}{\log x}}^1 \delta(x^r) dr = 0,$$

we have

$$\sum_{p \leq x} \frac{\mathbf{1}_{E_{a,m}}(p) \log p}{p} = \left(\frac{1}{\phi(m)} + o(1)\right) \log x \quad \text{as } x \rightarrow \infty.$$

□

EXAMPLE 3. If a sequence  $(a(p))_p$  with  $0 \leq a(p) \leq 1$  ( $\forall p$ ) satisfies that  $\sum_p \frac{1-a(p)}{p} < \infty$ , then

$$\sum_{p \leq x} \frac{a(p) \log p}{p} = (1 + o(1)) \log x \quad \text{as } x \rightarrow \infty.$$

Thus the condition (4) holds with  $\tau = 1$ .

Proof. For simplicity,

$$D(x) := \sum_{p \leq x} \frac{1 - a(p)}{p}, \quad x \in \mathbb{R}.$$

$D(\cdot)$  is non-decreasing, right-continuous,  $D(\infty) < \infty$  and  $D(x) = 0$  ( $\forall x < 2$ ). Since, by Mertens' first theorem (6),

$$\begin{aligned} \sum_{p \leq x} \frac{a(p) \log p}{p} &= \sum_{p \leq x} \frac{\log p}{p} - \sum_{p \leq x} \frac{1 - a(p)}{p} \log p \\ &= \log x + O(1) - \sum_{p \leq x} \frac{1 - a(p)}{p} \log p, \end{aligned}$$

it suffices to show that

$$\sum_{p \leq x} \frac{1 - a(p)}{p} \log p = o(\log x).$$

$D(dt) = \sum_p \frac{1 - a(p)}{p} \delta_p(dt)$  and integration by parts tell us that for  $x \geq 2$ ,

$$\begin{aligned} &\sum_{p \leq x} \frac{1 - a(p)}{p} \log p \\ &= \int_{(2-\varepsilon, x]} (\log t) D(dt) \quad [\text{where } 0 < \varepsilon < 1] \\ &= \int_{(2-\varepsilon, x]} \left( d(D(t) \log t) - D(t) \frac{dt}{t} \right) \\ &= D(x) \log x - \int_1^x \frac{D(t)}{t} dt \quad [\text{since } D(t) = 0 \quad (\forall t < 2)] \\ &= (\log x) \int_0^1 (D(x) - D(x^r)) dr \quad [\text{by the change of variables } \frac{\log t}{\log x} = r]. \end{aligned}$$

By noting that for each  $r \in (0, 1]$ ,

$$0 \leq D(x) - D(x^r) \leq D(x) \leq D(\infty) < \infty \quad (\forall x \geq 2), \quad \lim_{x \rightarrow \infty} (D(x) - D(x^r)) = 0,$$

it follows from the bounded convergence theorem that

$$\lim_{x \rightarrow \infty} \frac{1}{\log x} \sum_{p \leq x} \frac{1 - a(p)}{p} \log p = 0.$$

□

REMARK 2. When  $0 \leq a(p) \leq 1$  ( $\forall p$ ), there is no implication between the condition for  $(a(p))_p$  in Example 1 and that in Example 3.

- (i) Let a subset  $E \subset \{p \text{ is prime}\}$  be such that<sup>3</sup>  $\#E = \infty$ ,  $\sum_{p \in E} \frac{1}{p} < \infty$ . Then  $(a(p) = 1 - \mathbf{1}_E(p))_p$  satisfies  $\sum_p \frac{1-a(p)}{p} < \infty$ , but  $\limsup_p a(p) = 1$  and  $\liminf_p a(p) = 0$ . This  $(a(p))_p$  is in Example 3, but not in Example 1.
- (ii) Let  $0 \leq a(p) \leq 1$  be such that  $a(p) = 1 - \frac{1}{\log \log p}$  ( $p \gg 1$ ). Then, clearly  $a(p) \rightarrow 1$ , but  $\sum_p \frac{1-a(p)}{p} = \infty$ . This  $(a(p))_p$  is in Example 1, but not in Example 3.

Proof. (i) Since  $E$  is an infinite set by assumption,  $a(p) = 0$  i.o. Since  $\sum_{p \notin E} \frac{1}{p} = \infty$  by  $\sum_p \frac{1}{p} = \infty$  and assumption, and thus, since  $\{p \text{ is prime}\} \setminus E$  is also an infinite set,  $a(p) = 1$  i.o.

(ii) We use the prime number theorem in the following form (cf. (23)):

$$\vartheta(x) := \sum_{p \leq x} \log p = (1 + o(1))x \quad \text{as } x \rightarrow \infty.$$

$\vartheta(\cdot)$  is non-decreasing, right-continuous and  $\vartheta(t) = 0$  ( $\forall t < 2$ ).  $\delta(t) := \frac{\vartheta(t)}{t} - 1$  ( $t > 0$ ) is of bounded variation on every bounded closed interval of  $(0, \infty)$ , and  $\lim_{t \rightarrow \infty} \delta(t) = 0$ . Take a prime  $q_0$  large enough such that  $1 - a(p) = \frac{1}{\log \log p}$  ( $p > q_0$ ). Then

$$\begin{aligned} \sum_{q_0 < p \leq x} \frac{1 - a(p)}{p} &= \sum_{q_0 < p \leq x} \frac{\log p}{p \log p \log \log p} \\ &= \int_{(q_0, x]} \frac{\vartheta(dt)}{t \log t \log \log t} \\ &= \int_{(q_0, x]} \frac{1}{t \log t \log \log t} d(t + t\delta(t)) \\ &= \int_{q_0}^x \frac{1}{(\log \log t)(\log t)} \frac{dt}{t} + \int_{q_0}^x \frac{\delta(t)}{(\log \log t)(\log t)} \frac{dt}{t} \\ &\quad + \int_{(q_0, x]} \frac{\delta(dt)}{\log t \log \log t} \\ &= \int_{q_0}^x \frac{1}{(\log \log t)(\log t)} \frac{dt}{t} + \int_{q_0}^x \frac{\delta(t)}{(\log \log t)(\log t)} \frac{dt}{t} \\ &\quad + \int_{(q_0, x]} \left( d\left(\frac{\delta(t)}{\log t \log \log t}\right) - \delta(t) d\left(\frac{1}{\log t \log \log t}\right) \right) \\ &\quad \text{[by integration by parts]} \\ &= \int_{q_0}^x \frac{1}{(\log \log t)(\log t)} \frac{dt}{t} + \int_{q_0}^x \frac{\delta(t)}{(\log \log t)(\log t)} \frac{dt}{t} \\ &\quad + \int_{q_0}^x \frac{\delta(t)(1 + \log \log t)}{(\log t \log \log t)^2} \frac{dt}{t} \\ &\quad + \frac{\delta(x)}{\log x \log \log x} - \frac{\delta(q_0)}{\log q_0 \log \log q_0} \\ &= \log \log \log x \left( 1 - \frac{\log \log \log q_0}{\log \log \log x} + \int_{\frac{\log \log \log q_0}{\log \log \log x}}^1 \delta(e^{e^v \log \log \log x}) dv \right) \end{aligned}$$

---

<sup>3</sup>Such an  $E$  exists. For example, take a sequence  $\{q_i\}_{i=1}^{\infty}$  of prime numbers such that  $q_1 = 2$ ,  $q_{i+1} > q_i^2$  ( $i \geq 1$ ), and set  $E := \{q_i; i \geq 1\}$ . Then this  $E$  clearly satisfies the above conditions.

$$\begin{aligned}
 &+ \int_{\log \log q_0}^{\log \log x} \delta(e^{e^r}) \left(1 + \frac{1}{r}\right) e^{-r} \frac{dr}{r} \\
 &+ \frac{\delta(x)}{\log x \log \log x} - \frac{\delta(q_0)}{\log q_0 \log \log q_0} \\
 &\quad \text{[by the change of variables } \log \log t = u].
 \end{aligned}$$

This implies that

$$\frac{1}{\log \log \log x} \sum_{q_0 < p \leq x} \frac{1 - a(p)}{p} \rightarrow 1 \quad \text{as } x \rightarrow \infty,$$

and thus

$$\sum_p \frac{1 - a(p)}{p} = \infty.$$

□

REMARK 3.  $E_{a,m}$  is an infinite set. By Mertens' first theorem (6) and Example 2,

$$\begin{aligned}
 \sum_{p \leq x} \frac{(1 - \mathbf{1}_{E_{a,m}}(p)) \log p}{p} &= \sum_{p \leq x} \frac{\log p}{p} - \sum_{p \leq x} \frac{\mathbf{1}_{E_{a,m}}(p) \log p}{p} \\
 &= \left(1 - \frac{1}{\phi(m)} + o(1)\right) \log x \quad \text{as } x \rightarrow \infty.
 \end{aligned}$$

Since  $\phi(m) \geq 2$  for  $m \geq 3$ , and thus  $1 - \frac{1}{\phi(m)} > 0$ ,  $\{p \text{ is prime}\} \setminus E_{a,m}$  is also an infinite set. Therefore  $\limsup_p \mathbf{1}_{E_{a,m}}(p) = 1$  and  $\liminf_p \mathbf{1}_{E_{a,m}}(p) = 0$ . This tells us that for  $m \geq 3$ ,  $(\mathbf{1}_{E_{a,m}}(p))_p$  is not in Example 1.

#### 4. Behavior of $Z(\sigma; a)$ as $\sigma \searrow 1$ for more general $a(\cdot)$

Roughly speaking, the aim of this section is as follows:

In case  $\tau + \#\{p; a(p) = p\} = 0$  in (5), how does  $Z(\sigma; a)$  behave as  $\sigma \searrow 1$ ?

To this end, for a nonnegative, completely multiplicative arithmetical function  $a(\cdot)$ , we consider, instead of (3) and (4), the following conditions:

$$(24) \quad \sup_p a(p) < \infty, \quad \sup_p \frac{a(p)}{p} < 1,$$

$$(25) \quad \sum_p \frac{a(p) \log p}{p} = \infty.$$

In the case where  $\tau + \#\{p; a(p) = p\} = 0 \Leftrightarrow \tau = 0$  and  $a(p) \neq p \ (\forall p)$ , (3) becomes (24). But (4) does not always imply (25). In this paper, let us consider this convenient condition for us.

We begin with the following claim, which states that Claim 5 is valid even under the slightly weak condition (24):

**Claim 10.** (i) For  $\sigma \in \mathbb{R} \setminus \{1\}$ ,

$$\sum_p \frac{a(p)}{p^\sigma} \begin{cases} = \infty & \text{if } -\infty < \sigma < 1, \\ < \infty & \text{if } \sigma > 1. \end{cases}$$

(ii) For  $\sigma > 1$  and  $t \in \mathbb{R}$ ,  $\prod_p \frac{1}{1 - \frac{a(p)}{p^{\sigma+it}}}$  is convergent and  $\sum_{n=1}^{\infty} \frac{a(n)}{n^{\sigma+it}}$  is absolutely convergent, and these coincide with each other.

(iii) For  $\sigma \in \mathbb{R} \setminus \{1\}$ ,

$$\sum_{n=1}^{\infty} \frac{a(n)}{n^\sigma} \begin{cases} = \infty & \text{if } -\infty < \sigma < 1, \\ < \infty & \text{if } \sigma > 1. \end{cases}$$

For  $\sigma = 1$ ,

$$\sum_{n=1}^{\infty} \frac{a(n)}{n} = \infty \Leftrightarrow \sum_p \frac{a(p)}{p} = \infty.$$

Proof. (i) From an inequality  $x > \log x$  ( $x > 0$ ), the following implication is seen: For  $\sigma \in \mathbb{R}$  and prime  $p$ ,

$$\begin{aligned} p^{1-\sigma} > \log p^{1-\sigma} &= (1-\sigma) \log p \Rightarrow \frac{1}{p^\sigma} > (1-\sigma) \frac{\log p}{p} \\ &\Rightarrow \frac{a(p)}{p^\sigma} \geq (1-\sigma) \frac{a(p) \log p}{p}. \end{aligned}$$

In the case where  $\sigma \in (-\infty, 1)$ ,  $1 - \sigma > 0$  and (25) imply

$$\sum_p \frac{a(p)}{p^\sigma} \geq (1-\sigma) \sum_p \frac{a(p) \log p}{p} = \infty;$$

In the case where  $\sigma \in (1, \infty)$ , (24) implies

$$\sum_p \frac{a(p)}{p^\sigma} \leq \sum_p \frac{\sup_q a(q)}{p^\sigma} < \sup_q a(q) \sum_{n=1}^{\infty} \frac{1}{n^\sigma} < \infty.$$

(ii) Let  $\sigma \geq 1$  and  $t \in \mathbb{R}$ . By (24),

$$\left| \frac{a(p)}{p^{\sigma+it}} \right| = \frac{a(p)}{p^\sigma} \leq \frac{a(p)}{p} \leq \sup_q \frac{a(q)}{q} < 1,$$

so that

$$\frac{1}{1 - \frac{a(p)}{p^{\sigma+it}}} = e^{\frac{a(p)}{p^{\sigma+it}}} \exp \left\{ \frac{a(p)^2}{p^{2\sigma+2it}} \int_0^1 \frac{s}{1 - \frac{a(p)}{p^{\sigma+it}} s} ds \right\}.$$

Multiplication in  $p \leq x$  yields that

$$\prod_{p \leq x} \frac{1}{1 - \frac{a(p)}{p^{\sigma+it}}} = \exp \left\{ \sum_{p \leq x} \frac{a(p)}{p^{\sigma+it}} \right\} \exp \left\{ \sum_{p \leq x} \frac{a(p)^2}{p^{2\sigma+2it}} \int_0^1 \frac{s}{1 - \frac{a(p)}{p^{\sigma+it}} s} ds \right\}.$$

Here, by noting that

$$\sum_p \left| \frac{a(p)^2}{p^{2\sigma+2t}} \int_0^1 \frac{s}{1 - \frac{a(p)}{p^{\sigma+it}} s} ds \right| \leq \frac{(\sup_q a(q))^2}{2(1 - \sup_q \frac{a(q)}{q})} \sum_p \frac{1}{p^2} < \infty,$$

the convergence of  $\prod_p \frac{1}{1 - \frac{a(p)}{p^{\sigma+it}}}$  is reduced to that of  $\sum_p \frac{a(p)}{p^{\sigma+it}}$ . Since, in the case where  $\sigma > 1$ ,

$$\sum_p \left| \frac{a(p)}{p^{\sigma+it}} \right| = \sum_p \frac{a(p)}{p^\sigma} < \infty$$

by (i),  $\prod_p \frac{1}{1 - \frac{a(p)}{p^{\sigma+it}}}$  ( $\sigma > 1, t \in \mathbb{R}$ ) is convergent.

Let  $\sigma \geq 1$  and  $t \in \mathbb{R}$  again. First, from the proof of Claim 5, note that

$$\begin{aligned} \sum_{n \in \mathbb{N}_L} \frac{a(n)}{n^{\sigma+it}} &= \left( \prod_{j=1}^L \frac{1}{1 - \frac{a(p_j)}{p_j^{\sigma+it}}} \right) \prod_{j=1}^L \left( 1 - \left( \frac{a(p_j)}{p_j^{\sigma+it}} \right)^{L+1} \right), \\ \prod_{j=1}^L \left( 1 - \left( \frac{a(p_j)}{p_j^{\sigma+it}} \right)^{L+1} \right) &= \exp \left\{ \sum_{j=1}^L \left( - \left( \frac{a(p_j)}{p_j^{\sigma+it}} \right)^{L+1} \right) \int_0^1 \frac{1}{1 - \left( \frac{a(p_j)}{p_j^{\sigma+it}} \right)^{L+1} s} ds \right\}, \\ \left| \sum_{j=1}^L \left( - \left( \frac{a(p_j)}{p_j^{\sigma+it}} \right)^{L+1} \right) \int_0^1 \frac{1}{1 - \left( \frac{a(p_j)}{p_j^{\sigma+it}} \right)^{L+1} s} ds \right| \\ &\leq L \left( \sup_p \frac{a(p)}{p} \right)^{L+1} \frac{1}{1 - \left( \sup_p \frac{a(p)}{p} \right)^{L+1}} \rightarrow 0 \quad \text{as } L \rightarrow \infty. \end{aligned}$$

In the case where  $\sigma > 1$ ,

$$\lim_{L \rightarrow \infty} \sum_{n \in \mathbb{N}_L} \frac{a(n)}{n^{\sigma+it}} = \prod_p \frac{1}{1 - \frac{a(p)}{p^{\sigma+it}}}.$$

When  $t = 0$ , the monotone convergence theorem tells us that

$$\sum_{n=1}^{\infty} \frac{a(n)}{n^\sigma} = \sum_{n \in \mathbb{N}} \frac{a(n)}{n^\sigma} = \lim_{L \rightarrow \infty} \sum_{n \in \mathbb{N}_L} \frac{a(n)}{n^\sigma} < \infty.$$

Thus we have the absolute convergence of  $\sum_{n=1}^{\infty} \frac{a(n)}{n^{\sigma+it}}$  and

$$\sum_{n=1}^{\infty} \frac{a(n)}{n^{\sigma+it}} = \prod_p \frac{1}{1 - \frac{a(p)}{p^{\sigma+it}}}.$$

(iii) In the case where  $\sigma < 1$ ,

$$\sum_{n=1}^{\infty} \frac{a(n)}{n^\sigma} \geq \sum_p \frac{a(p)}{p^\sigma} = \infty$$

by (i). In the case where  $\sigma > 1$ ,

$$\sum_{n=1}^{\infty} \frac{a(n)}{n^\sigma} < \infty$$

by (ii).

From the proof of (ii), it follows that

$$\sum_{n=1}^{\infty} \frac{a(n)}{n} = \lim_{L \rightarrow \infty} \sum_{n \in \mathbb{N}_L} \frac{a(n)}{n} = \lim_{L \rightarrow \infty} \prod_{i=1}^L \frac{1}{1 - \frac{a(p_i)}{p_i}} = \prod_p \frac{1}{1 - \frac{a(p)}{p}},$$

$$\prod_p \frac{1}{1 - \frac{a(p)}{p}} < \infty \Leftrightarrow \sum_p \frac{a(p)}{p} < \infty.$$

Combining these, we have

$$\sum_{n=1}^{\infty} \frac{a(n)}{n} < \infty \Leftrightarrow \sum_p \frac{a(p)}{p} < \infty.$$

□

DEFINITION 6. (i) By virtue of Claim 10, for  $s = \sigma + it$  ( $\sigma \in (1, \infty), t \in \mathbb{R}$ ), we define

$$\eta(s; a) := \sum_{n=1}^{\infty} \frac{a(n)}{n^s} = \prod_p \frac{1}{1 - \frac{a(p)}{p^s}}.$$

And, for  $\sigma \in (1, \infty)$ , we set

$$\eta_{\sigma}(t; a) := \frac{\eta(\sigma + it; a)}{\eta(\sigma; a)}, \quad t \in \mathbb{R}.$$

As before (cf. Definition 4), let  $\mu_{\sigma}(dx; a)$  be a 1-dimensional probability measure corresponding to  $\eta_{\sigma}(\cdot; a)$ .

(ii) For a sequence  $\{Y_p\}_p$  of independent geometric processes on some  $(\Omega, \mathcal{F}, P)$ , we define

$$Z(\sigma; a) := \sum_p Y_p \left( \frac{a(p)}{p^{\sigma}} \right) \log p, \quad \sigma \in (1, \infty).$$

Then  $(-Z(\sigma; a))_{1 < \sigma < \infty}$  is a backwards Lévy process whose marginal distribution at  $\sigma$  is  $\mu_{\sigma}(\cdot; a)$ . But, as compared with  $Z(\sigma; a)$  in Definition 5, there is the following difference: If  $\sum_p \frac{a(p)}{p} < \infty$ , then  $Z(1+; a) < \infty$  a.s., and if  $\sum_p \frac{a(p)}{p} = \infty$ , then  $Z(1+; a) = \infty$  a.s.

Our interest is the behavior of  $Z(\sigma; a)$  as  $\sigma \searrow 1$ . To see this, for a nonnegative, completely multiplicative arithmetical function  $a(\cdot)$  satisfying (24) and (25), we further suppose the following:

$$(26) \quad \begin{cases} \mathbb{R} \ni x \mapsto \sum_{p \leq e^x} \frac{a(p) \log p}{p} \in [0, \infty) \text{ is regularly varying at } \infty \text{ with exponent} \\ \gamma \in [0, \infty). \end{cases}$$

First of all, note that  $\gamma \leq 1$  from Mertens' first theorem. For, let  $L(\cdot)$ <sup>4</sup> be a slowly varying function at  $\infty$  defined as

$$(27) \quad \sum_{p \leq e^x} \frac{a(p) \log p}{p} = x^{\gamma} L(x),$$

then (24) and (6) tell us that

---

<sup>4</sup>In this paper, we call this  $L(\cdot)$  a slowly varying part of a regularly varying function  $x \mapsto \sum_{p \leq e^x} \frac{a(p) \log p}{p}$ .

$$x^\gamma L(x) \leq \sup_q a(q) \sum_{p \leq e^x} \frac{\log p}{p} = \sup_q a(q)(\log e^x + O(1)) = \sup_q a(q)(x + O(1)).$$

This implies

$$(28) \quad \limsup_{x \rightarrow \infty} x^{\gamma-1} L(x) \leq \sup_q a(q),$$

so that it must be that  $\gamma \leq 1$ .

We treat the following two cases:

Case 1  $\gamma < 1$

$$\text{or} \\ \gamma = 1 \text{ and } \int_2^\infty \frac{L(x)}{x} dx < \infty,$$

$$\text{Case 2 } \gamma = 1 \text{ and } \int_2^\infty \frac{L(x)}{x} dx = \infty.$$

**Theorem 2.** *In Case 1,  $\sum_p \frac{a(p)}{p} < \infty$ , and thus  $Z(1+; a) < \infty$  a.s. In Case 2,  $\sum_p \frac{a(p)}{p} = \infty$ , and thus  $Z(1+; a) = \infty$  a.s.*

Proof. First, by (10) and (27),

$$(29) \quad C(e^x) = x^\gamma L(x).$$

From (11), it follows that for  $x \geq 2$ ,

$$\begin{aligned} \sum_{p \leq x} \frac{a(p)}{p} &= \sum_{p \leq x} \frac{a(p) \log p}{p} \frac{1}{\log p} \\ &= \int_{(2-\varepsilon, x]} \frac{C(dt)}{\log t} \quad [\text{where } 0 < \varepsilon < 1] \\ &= \int_{(2-\varepsilon, x]} \left( d\left(\frac{C(t)}{\log t}\right) + C(t) \frac{1}{(\log t)^2} \frac{dt}{t} \right) \quad [\text{by integration by parts}] \\ &= \frac{C(x)}{\log x} + \int_2^x \frac{C(t)}{(\log t)^2} \frac{dt}{t} \quad [\text{since } C(t) = 0 \ (\forall t < 2)] \\ &= \frac{C(x)}{\log x} + \int_{\log 2}^{\log x} \frac{C(e^s)}{s^2} ds \quad [\text{by the change of variables } \log t = s]. \end{aligned}$$

By (29), this is rewritten as

$$\sum_{p \leq e^x} \frac{a(p)}{p} = \frac{L(x)}{x^{1-\gamma}} + \int_{\log 2}^x \frac{L(s)}{s^{2-\gamma}} ds.$$

In the case where  $\gamma < 1$ ,  $1 - \gamma > 0$  and the slow variation of  $L(\cdot)$  at  $\infty$  yield that

$$\begin{aligned} \lim_{x \rightarrow \infty} \frac{L(x)}{x^{1-\gamma}} &= 0, \\ \int_2^\infty \frac{L(s)}{s^{2-\gamma}} ds &= \int_2^\infty \frac{L(s)}{s^{1+1-\gamma}} ds < \infty, \end{aligned}$$

from which, it follows that



$$\sum_p \frac{a(p)}{p} < \infty.$$

In the case where  $\gamma = 1$  and  $\int^\infty \frac{L(x)}{x} dx < \infty$ , (28) implies

$$\begin{aligned} \sum_p \frac{a(p)}{p} &= \lim_{x \rightarrow \infty} \sum_{p \leq e^x} \frac{a(p)}{p} = \lim_{x \rightarrow \infty} \left( L(x) + \int_{\log 2}^x \frac{L(s)}{s} ds \right) \\ &\leq \sup_q a(q) + \int_{\log 2}^\infty \frac{L(s)}{s} ds < \infty. \end{aligned}$$

In the case where  $\gamma = 1$  and  $\int^\infty \frac{L(x)}{x} dx = \infty$ ,

$$\sum_p \frac{a(p)}{p} = \lim_{x \rightarrow \infty} \sum_{p \leq e^x} \frac{a(p)}{p} \geq \lim_{x \rightarrow \infty} \int_{\log 2}^x \frac{L(s)}{s} ds = \int_{\log 2}^\infty \frac{L(s)}{s} ds = \infty.$$

□

**Claim 11.** In Case 2,  $-\frac{\eta'(\sigma; a)}{\eta(\sigma; a)} \sim \frac{1}{\sigma - 1} L\left(\frac{1}{\sigma - 1}\right)$  as  $\sigma \searrow 1$ .

Proof. By 1° in the proof of Claim 6,

$$-\frac{\eta'(\sigma; a)}{\eta(\sigma; a)} = \sum_p \frac{a(p)}{p^\sigma} \log p + \sum_p \frac{a(p)^2}{p^{2\sigma}} \frac{\log p}{1 - \frac{a(p)}{p^\sigma}}.$$

Since the second term is convergent as  $\sigma \searrow 1$ , we may investigate the asymptotics of the first term as  $\sigma \searrow 1$ .

By (11) and (29),

$$\begin{aligned} (30) \quad \sum_p \frac{a(p)}{p^\sigma} \log p &= \sum_p \frac{a(p) \log p}{p} \frac{1}{p^{\sigma-1}} \\ &= \int_{(2-\varepsilon, \infty)} \frac{C(dx)}{x^{\sigma-1}} \quad [\text{where } 0 < \varepsilon < 1] \\ &= \int_{(2-\varepsilon, \infty)} \left( d\left(\frac{C(x)}{x^{\sigma-1}}\right) - C(x)(1 - \sigma)x^{-\sigma} dx \right) \quad [\text{by integration by parts}] \\ &= (\sigma - 1) \int_1^\infty \frac{C(x)}{x^\sigma} dx \quad \left[ \begin{array}{l} \text{since } C(t) = 0 \ (\forall t < 2) \text{ and by (29),} \\ \frac{C(x)}{x^{\sigma-1}} = \frac{C(e^{\log x})}{x^{\sigma-1}} = \frac{\log x L(\log x)}{x^{\sigma-1}} \rightarrow 0 \text{ as } x \rightarrow \infty \end{array} \right] \\ &= (\sigma - 1) \int_0^\infty e^{-(\sigma-1)s} C(e^s) ds \quad [\text{by the change of variables } \log x = s] \\ &= (\sigma - 1) \int_0^\infty e^{-(\sigma-1)s} d\left(\int_0^s C(e^x) dx\right). \end{aligned}$$

Here, since, by (29),  $x \mapsto C(e^x)$  is regularly varying at  $\infty$  with exponent 1, Feller [4, Chapter VIII, Theorem 1] tells us that

$$\frac{sC(e^s)}{\int_0^s C(e^x) dx} \rightarrow 2 \quad \text{as } s \rightarrow \infty,$$

and thus

$$\int_0^s C(e^x)dx \sim \frac{1}{2}sC(e^s) = \frac{1}{2}s^2L(s) \quad \text{as } s \rightarrow \infty.$$

This, by Feller [4, Chapter XIII, Theorem 2], implies

$$(31) \quad \int_0^\infty e^{-(\sigma-1)s}d\left(\int_0^s C(e^x)dx\right) \sim (\sigma-1)^{-2}L\left(\frac{1}{\sigma-1}\right) \quad \text{as } \sigma \searrow 1.$$

Therefore we have

$$\sum_p \frac{a(p)}{p^\sigma} \log p \sim \frac{1}{\sigma-1}L\left(\frac{1}{\sigma-1}\right) \quad \text{as } \sigma \searrow 1,$$

which is the assertion of the claim. □

We divide Case 2 into three cases:

Case 2.1  $\lim_{x \rightarrow \infty} L(x) = \tau \in (0, \infty)$ ,

Case 2.2  $\lim_{x \rightarrow \infty} L(x) = 0$ ,

Case 2.3 Neither Case 2.1 nor Case 2.2.

Since Case 2.3 is hard to deal with, this case is excluded from our consideration. Case 2.1 is  $C(e^x) = x(\tau + o(1))$  as  $x \rightarrow \infty$ , i.e.,

$$\sum_{p \leq x} \frac{a(p) \log p}{p} = (\log x)(\tau + o(1)) \quad \text{as } x \rightarrow \infty.$$

This is just the condition (4), so the answer to Case 2.1 is given from Corollary 1 in the following way:

$$\text{the distribution of } (\sigma-1)Z(\sigma; a) \rightarrow \mathbf{1}_{(0, \infty)}(x) \frac{1}{\Gamma(\tau)} x^{\tau-1} e^{-x} dx \quad \text{as } \sigma \searrow 1.$$

**Theorem 3.** *In Case 2.2,  $\lim_{\sigma \searrow 1} E[e^{-\lambda(\sigma-1)Z(\sigma; a)}] = 1$  ( $\forall \lambda > 0$ ). Thus*

$$(\sigma-1)Z(\sigma; a) \rightarrow 0 \quad \text{i.p. as } \sigma \searrow 1.$$

**Proof.** Fix  $\lambda > 0$ . First, by (14),

$$\begin{aligned} E[e^{-\lambda(\sigma-1)Z(\sigma; a)}] &= \prod_p E\left[e^{-\lambda(\sigma-1)(\log p)Y_p\left(\frac{a(p)}{p^\sigma}\right)}\right] \\ &= \prod_p \frac{1 - \frac{a(p)}{p^\sigma}}{1 - \frac{a(p)}{p^\sigma} \frac{1}{p^{\lambda(\sigma-1)}}} \\ &= \prod_p \exp\left\{-\frac{a(p)}{p^\sigma} - \frac{a(p)^2}{p^{2\sigma}} \int_0^1 \frac{s}{1 - \frac{a(p)}{p^\sigma} s} ds \right. \\ &\quad \left. + \frac{a(p)}{p^\sigma} \frac{1}{p^{\lambda(\sigma-1)}} + \frac{a(p)^2}{p^{2\sigma}} \frac{1}{p^{2\lambda(\sigma-1)}} \int_0^1 \frac{s}{1 - \frac{a(p)}{p^\sigma} \frac{s}{p^{\lambda(\sigma-1)}}} ds\right\} \\ &= \exp\left\{-\sum_p \frac{a(p)}{p^\sigma} \left(1 - \frac{1}{p^{\lambda(\sigma-1)}}\right) - \sum_p \frac{a(p)^2}{p^{2\sigma}} \int_0^1 \frac{s}{1 - \frac{a(p)}{p^\sigma} s} ds\right\} \end{aligned}$$

$$+ \sum_p \frac{a(p)^2}{p^{2\sigma}} \frac{1}{p^{2\lambda(\sigma-1)}} \int_0^1 \frac{s}{1 - \frac{a(p)}{p^\sigma} \frac{s}{p^{\lambda(\sigma-1)}}} ds \Big\}.$$

Since, as  $\sigma \searrow 1$ ,

$$\begin{aligned} \sum_p \frac{a(p)^2}{p^{2\sigma}} \int_0^1 \frac{s}{1 - \frac{a(p)}{p^\sigma} s} ds &\rightarrow \sum_p \frac{a(p)^2}{p^2} \int_0^1 \frac{s}{1 - \frac{a(p)}{p} s} ds, \\ \sum_p \frac{a(p)^2}{p^{2\sigma}} \frac{1}{p^{2\lambda(\sigma-1)}} \int_0^1 \frac{s}{1 - \frac{a(p)}{p^\sigma} \frac{s}{p^{\lambda(\sigma-1)}}} ds &\rightarrow \sum_p \frac{a(p)^2}{p^2} \int_0^1 \frac{s}{1 - \frac{a(p)}{p} s} ds, \end{aligned}$$

it suffices to show that

$$(32) \quad \sum_p \frac{a(p)}{p^\sigma} \left(1 - \frac{1}{p^{\lambda(\sigma-1)}}\right) \rightarrow 0.$$

Rewriting  $1 - \frac{1}{p^{\lambda(\sigma-1)}}$  as

$$1 - \frac{1}{p^{\lambda(\sigma-1)}} = \lambda(\sigma - 1) \int_0^1 \frac{1}{p^{\lambda(\sigma-1)t}} \log p \, dt$$

and then using (30) yield that

$$(33) \quad \begin{aligned} \sum_p \frac{a(p)}{p^\sigma} \left(1 - \frac{1}{p^{\lambda(\sigma-1)}}\right) \\ = \lambda(\sigma - 1) \int_0^1 (\sigma - 1)(1 + \lambda t) dt \int_0^\infty e^{-(\sigma-1)(1+\lambda t)s} d\left(\int_0^s C(e^x) dx\right). \end{aligned}$$

Here, by (31),

$$\kappa \int_0^\infty e^{-\kappa s} d\left(\int_0^s C(e^x) dx\right) \sim \frac{1}{\kappa} L\left(\frac{1}{\kappa}\right) \quad \text{as } \kappa \searrow 0,$$

and thus, for  $0 < \forall \varepsilon < 1$ ,

$$\exists \delta > 0 \text{ s.t. } 0 < \kappa < \delta \Rightarrow 1 - \varepsilon < \frac{\kappa \int_0^\infty e^{-\kappa s} d\left(\int_0^s C(e^x) dx\right)}{\frac{1}{\kappa} L\left(\frac{1}{\kappa}\right)} < 1 + \varepsilon.$$

Since, for  $1 < \sigma < 1 + \frac{\delta}{1+\lambda}$ ,

$$0 < (\sigma - 1)(1 + \lambda t) \leq (\sigma - 1)(1 + \lambda) < \delta \quad (0 \leq \forall t \leq 1),$$

it follows that

$$\frac{(\sigma - 1)(1 + \lambda t) \int_0^\infty e^{-(\sigma-1)(1+\lambda t)s} d\left(\int_0^s C(e^x) dx\right)}{\frac{1}{(\sigma-1)(1+\lambda t)} L\left(\frac{1}{(\sigma-1)(1+\lambda t)}\right)} < 1 + \varepsilon \quad (0 \leq \forall t \leq 1).$$

Using this estimate in (33), we have

$$0 \leq \sum_p \frac{a(p)}{p^\sigma} \left(1 - \frac{1}{p^{\lambda(\sigma-1)}}\right) \leq (1 + \varepsilon) \int_0^1 \frac{\lambda}{1 + \lambda t} L\left(\frac{1}{(\sigma - 1)(1 + \lambda t)}\right) dt.$$

Finally, noting that by  $\lim_{x \rightarrow \infty} L(x) = 0$  and the bounded convergence theorem,

$$\lim_{\sigma \searrow 1} \int_0^1 \frac{\lambda}{1 + \lambda t} L\left(\frac{1}{(\sigma - 1)(1 + \lambda t)}\right) dt = 0,$$

we obtain (32). □

To investigate the behavior of  $Z(\sigma; a)$  as  $\sigma \searrow 1$  in Case 2.2 in more detail, we suppose the following:

$$(34) \quad u \mapsto L(e^u) \text{ is regularly varying at } \infty \text{ with exponent } \delta.$$

Then note that  $-1 \leq \delta \leq 0$ . For, let  $l(\cdot)$  be its slowly varying part, then

$$\begin{aligned} \infty &= \int_0^\infty \frac{L(x)}{x} dx \quad [\text{cf. Case 2}] \\ &= \int_0^\infty L(e^u) du \quad [\text{by the change of variables } \log x = u] \\ &= \int_0^\infty u^\delta l(u) du, \\ 0 &= \lim_{x \rightarrow \infty} L(x) \quad [\text{cf. Case 2.2}] = \lim_{u \rightarrow \infty} L(e^u) = \lim_{u \rightarrow \infty} u^\delta l(u). \end{aligned}$$

These convergences imply neither  $\delta < -1$  nor  $\delta > 0$ , i.e.,  $-1 \leq \delta \leq 0$ .

We divide Case 2.2 into the following cases:

Case 2.2.1  $-1 < \delta \leq 0$ ,

Case 2.2.2  $\delta = -1$ ,

Case 2.2.2.1  $\lim_{u \rightarrow \infty} l(u) = \infty$ ,

Case 2.2.2.2  $\lim_{u \rightarrow \infty} l(u) = \kappa \in (0, \infty)$ ,

Case 2.2.2.3  $\lim_{u \rightarrow \infty} l(u) = 0$ .

**Theorem 4.** (i) *In Case 2.2.1 or Case 2.2.2.1,  $\lim_{\sigma \searrow 1} E[e^{-\lambda(\sigma-1)^\Delta Z(\sigma; a)}] = 0$  ( $\forall \lambda > 0, 0 \leq \Delta < 1$ ). Thus*

$$(\sigma - 1)^\Delta Z(\sigma; a) \rightarrow \infty \text{ i.p. as } \sigma \searrow 1, \quad 0 \leq \Delta < 1.$$

(ii) *In Case 2.2.2.2,  $\lim_{\sigma \searrow 1} E[e^{-\lambda(\sigma-1)^\Delta Z(\sigma; a)}] = \Delta^\kappa$  ( $\forall \lambda > 0, 0 < \Delta \leq 1$ ). Thus,  $(\sigma - 1)^\Delta Z(\sigma; a)$  being regarded as a  $[0, \infty]$ -valued random variable,*

$$\begin{aligned} &\text{the distribution of } (\sigma - 1)^\Delta Z(\sigma; a) \\ &\rightarrow \Delta^\kappa \delta_0 + (1 - \Delta^\kappa) \delta_\infty \text{ as } \sigma \searrow 1, \quad 0 < \Delta \leq 1. \end{aligned}$$

(iii) *In Case 2.2.2.3,  $\lim_{\sigma \searrow 1} E[e^{-\lambda(\sigma-1)^\Delta Z(\sigma; a)}] = 1$  ( $\forall \lambda > 0, 0 < \Delta \leq 1$ ). Thus*

$$(\sigma - 1)^\Delta Z(\sigma; a) \rightarrow 0 \text{ i.p. as } \sigma \searrow 1, \quad 0 < \Delta \leq 1.$$

Proof. Fix  $\lambda > 0$  and  $0 < \Delta < 1$ . First, from the proof of Theorem 3, it is seen that as  $\sigma \searrow 1$ ,

$$E[e^{-\lambda(\sigma-1)^\Delta Z(\sigma; a)}] \sim \exp\left\{-\sum_p \frac{a(p)}{p^\sigma} \left(1 - \frac{1}{p^{\lambda(\sigma-1)^\Delta}}\right)\right\},$$

$$\begin{aligned} & \sum_p \frac{a(p)}{p^\sigma} \left(1 - \frac{1}{p^{\lambda(\sigma-1)^\Delta}}\right) \\ &= \lambda(\sigma-1)^\Delta \int_0^1 (\sigma-1 + \lambda(\sigma-1)^\Delta t) dt \int_0^\infty e^{-(\sigma-1 + \lambda(\sigma-1)^\Delta t)s} d\left(\int_0^s C(e^x) dx\right) \\ &\sim \lambda(\sigma-1)^\Delta \int_0^1 \frac{1}{\sigma-1 + \lambda(\sigma-1)^\Delta t} L\left(\frac{1}{\sigma-1 + \lambda(\sigma-1)^\Delta t}\right) dt \\ &= \int_{\log \frac{1}{\sigma-1 + \lambda(\sigma-1)^\Delta}}^{\log \frac{1}{\sigma-1}} L(e^u) du \quad [\text{by the change of variables } \log \frac{1}{\sigma-1 + \lambda(\sigma-1)^\Delta t} = u] \\ &= \int_{\log \frac{1}{\sigma-1 + \lambda(\sigma-1)^\Delta}}^{\log \frac{1}{\sigma-1}} u^\delta l(u) du. \end{aligned}$$

In the case where  $-1 < \delta \leq 0$ , we take  $\varepsilon > 0$  such that  $-1 < \delta - \varepsilon < \delta \leq 0$ . Since  $u^{-\varepsilon} < l(u) < u^\varepsilon$  ( $u \gg 1$ ) by the slow variation of  $l(\cdot)$  at  $\infty$ , and  $\delta - \varepsilon + 1 > 0$ , it follows that

$$\begin{aligned} \int_{\log \frac{1}{\sigma-1 + \lambda(\sigma-1)^\Delta}}^{\log \frac{1}{\sigma-1}} u^\delta l(u) du &\geq \int_{\log \frac{1}{\sigma-1 + \lambda(\sigma-1)^\Delta}}^{\log \frac{1}{\sigma-1}} u^{\delta-\varepsilon} du \\ &= \frac{1}{\delta - \varepsilon + 1} \left(\log \frac{1}{\sigma-1}\right)^{\delta-\varepsilon+1} \left(1 - \left(\Delta + \frac{\log \frac{1}{\lambda + (\sigma-1)^{1-\Delta}}}{\log \frac{1}{\sigma-1}}\right)^{\delta-\varepsilon+1}\right) \\ &\rightarrow \infty \quad \text{as } \sigma \searrow 1. \end{aligned}$$

Thus

$$\lim_{\sigma \searrow 1} E[e^{-\lambda(\sigma-1)^\Delta Z(\sigma;a)}] = 0.$$

In the case where  $\delta = -1$ ,

$$\begin{aligned} \int_{\log \frac{1}{\sigma-1 + \lambda(\sigma-1)^\Delta}}^{\log \frac{1}{\sigma-1}} u^{-1} l(u) du &= \int_{\log \log \frac{1}{\sigma-1 + \lambda(\sigma-1)^\Delta}}^{\log \log \frac{1}{\sigma-1}} l(e^v) dv \quad [\text{by the change of variables } \log u = v] \\ &= \int_{\alpha(\sigma)}^{\beta(\sigma)} f(v) dv \quad \left[ \begin{array}{l} \text{where, for simplicity} \\ f(v) := l(e^v), \\ \alpha(\sigma) := \log \log \frac{1}{\sigma-1 + \lambda(\sigma-1)^\Delta}, \\ \beta(\sigma) := \log \log \frac{1}{\sigma-1} \end{array} \right] \\ &= \int_0^{\beta(\sigma) - \alpha(\sigma)} \frac{f(w + \alpha(\sigma))}{f(\alpha(\sigma))} dw f(\alpha(\sigma)). \end{aligned}$$

Here, note that as  $\sigma \searrow 1$ ,

$$\beta(\sigma) - \alpha(\sigma) = \log \frac{1}{\Delta + \frac{\log \frac{1}{\lambda + (\sigma-1)^{1-\Delta}}}{\log \frac{1}{\sigma-1}}} \rightarrow \log \frac{1}{\Delta},$$

$$\alpha(\sigma) \rightarrow \infty,$$

$$\frac{f(w + \alpha(\sigma))}{f(\alpha(\sigma))} = \frac{l(e^w e^{\alpha(\sigma)})}{l(e^{\alpha(\sigma)})} \xrightarrow{c} 1 \quad [\text{by the slow variation of } l(\cdot) \text{ at } \infty].$$

From these, it follows that

$$\int_0^{\beta(\sigma)-\alpha(\sigma)} \frac{f(w + \alpha(\sigma))}{f(\alpha(\sigma))} dw \rightarrow \log \frac{1}{\Delta} \quad \text{as } \sigma \searrow 1,$$

so that

$$\lim_{\sigma \searrow 1} \int_{\log \frac{1}{\sigma-1+\lambda(\sigma-1)^\Delta}}^{\log \frac{1}{\sigma-1}} u^{-1} l(u) du = \begin{cases} \infty & \text{if } \lim_{u \rightarrow \infty} l(u) = \infty, \\ \kappa \log \frac{1}{\Delta} = -\log \Delta^\kappa & \text{if } \lim_{u \rightarrow \infty} l(u) = \kappa \in (0, \infty), \\ 0 & \text{if } \lim_{u \rightarrow \infty} l(u) = 0. \end{cases}$$

Thus

$$\lim_{\sigma \searrow 1} E[e^{-\lambda(\sigma-1)^\Delta Z(\sigma; a)}] = \begin{cases} 0 & \text{if } \lim_{u \rightarrow \infty} l(u) = \infty, \\ \Delta^\kappa & \text{if } \lim_{u \rightarrow \infty} l(u) = \kappa \in (0, \infty), \\ 1 & \text{if } \lim_{u \rightarrow \infty} l(u) = 0. \end{cases}$$

□

Before closing this paper, we give some examples of  $a(\cdot)$ . For this, we need the following lemma:

**Lemma 2.** *Let  $t_0 > 0$ , and  $f : (t_0, \infty) \rightarrow (0, \infty)$  be of class  $C^1$  and ultimately non-increasing, i.e.,  $\exists t_1 > t_0$  s.t.  $f' \leq 0$  on  $[t_1, \infty)$ . Then, for  $q_0 := \min\{p:\text{prime}; t_0 < p\}$  and  $0 < \varepsilon < (q_0 - t_0) \wedge (q_0 - 1)$ ,*

$$\sum_{t_0 < p \leq x} \frac{\log p}{p} f(p) = \int_{q_0 - \varepsilon}^x \frac{f(t)}{t} dt + O(1) \quad \text{as } x \rightarrow \infty.$$

Proof. For simplicity, we set

$$M(x) = \sum_{p \leq x} \frac{\log p}{p}, \quad x \in \mathbb{R}.$$

Clearly  $M(\cdot)$  is non-decreasing, right-continuous,  $M(x) = 0$  ( $\forall x < 2$ ), and

$$M(dx) = \sum_p \frac{\log p}{p} \delta_p(dx).$$

Note that  $\sup_{x \geq 1} |M(x) - \log x| < \infty$  by Mertens' first theorem (6). By integration by parts,

$$\begin{aligned} \sum_{t_0 < p \leq x} \frac{\log p}{p} f(p) &= \sum_{q_0 - \varepsilon < p \leq x} \frac{\log p}{p} f(p) \\ &= \int_{(q_0 - \varepsilon, x]} f(t) M(dt) \\ &= \int_{(q_0 - \varepsilon, x]} (d(f(t)M(t)) - M(t)f'(t)dt) \\ &= f(x)(\log x + \eta(x)) - f(q_0 - \varepsilon)(\log(q_0 - \varepsilon) + \eta(q_0 - \varepsilon)) \\ &\quad - \int_{q_0 - \varepsilon}^x (\log t + \eta(t))f'(t)dt \quad [\text{where } \eta(x) := M(x) - \log x] \end{aligned}$$

$$\begin{aligned} &= \int_{q_0-\varepsilon}^x \frac{f(t)}{t} dt + f(x)\eta(x) - f(q_0 - \varepsilon)\eta(q_0 - \varepsilon) - \int_{q_0-\varepsilon}^x \eta(t)f'(t)dt \\ &= \int_{q_0-\varepsilon}^x \frac{f(t)}{t} dt + O(1) - \int_{q_0-\varepsilon}^x \eta(t)f'(t)dt \quad \text{as } x \rightarrow \infty. \end{aligned}$$

Here, since  $-f' \geq 0$  on  $[t_1, \infty)$ ,

$$\begin{aligned} \left| - \int_{q_0-\varepsilon}^x \eta(t)f'(t)dt \right| &= \left| - \int_{q_0-\varepsilon}^{t_1 \vee 1} \eta(t)f'(t)dt + \int_{t_1 \vee 1}^x \eta(t)(-f'(t))dt \right| \\ &\leq \left| \int_{q_0-\varepsilon}^{t_1 \vee 1} \eta(t)f'(t)dt \right| + \int_{t_1 \vee 1}^x |\eta(t)|(-f'(t))dt \\ &\leq \left| \int_{q_0-\varepsilon}^{t_1 \vee 1} \eta(t)f'(t)dt \right| + \left( \sup_{t \geq 1} |\eta(t)| \right) (-f(x) + f(t_1 \vee 1)) \\ &= O(1) \quad \text{as } x \rightarrow \infty. \end{aligned}$$

This, together with the preceding, implies the assertion of the lemma. □

Letting  $f(t) = \frac{1}{(\log t)^b}$  or  $\frac{1}{(\log \log t)^c}$  or  $\frac{1}{(\log \log t)^c (\log \log \log t)^d}$  or  $\frac{1}{(\log \log \log t)^d}$  in Lemma 2 yields the following example, whose details are omitted:

EXAMPLE 4. (i) For  $b > 0$ ,

$$\sum_{p \leq x} \frac{\log p}{p} \frac{1}{(\log p)^b} = \begin{cases} O(1) & \text{if } b > 1, \\ \log \log x + O(1) & \text{if } b = 1, \\ \frac{1}{1-b} (\log x)^{1-b} + O(1) & \text{if } 0 < b < 1 \end{cases} \quad \text{as } x \rightarrow \infty.$$

Thus

$$\sum_p \frac{\log p}{p} \frac{1}{(\log p)^b} \begin{cases} < \infty & \text{if } b > 1, \\ = \infty & \text{if } 0 < b \leq 1. \end{cases}$$

In the latter case,  $x \mapsto \sum_{p \leq e^x} \frac{\log p}{p} \frac{1}{(\log p)^b}$  is regularly varying at  $\infty$  with exponent  $1-b \in [0, 1)$ .

(ii) For  $c > 0$ ,

$$\sum_{e < p \leq x} \frac{\log p}{p} \frac{1}{(\log \log p)^c} \sim \frac{\log x}{(\log \log x)^c} \quad \text{as } x \rightarrow \infty.$$

Thus  $\sum_{p > e} \frac{\log p}{p} \frac{1}{(\log \log p)^c} = \infty$ , and  $x \mapsto \sum_{e < p \leq e^x} \frac{\log p}{p} \frac{1}{(\log \log p)^c}$  is regularly varying at  $\infty$  with exponent 1 and its slowly varying part  $L(x) \sim \frac{1}{(\log x)^c}$  as  $x \rightarrow \infty$ , so that

$$\lim_{x \rightarrow \infty} L(x) = 0, \quad \int^{\infty} \frac{L(x)}{x} dx \begin{cases} < \infty & \text{if } c > 1, \\ = \infty & \text{if } 0 < c \leq 1. \end{cases}$$

In the latter case,  $u \mapsto L(e^u)$  is regularly varying at  $\infty$  with exponent  $-c \in [-1, 0)$ .

(iii) For  $c > 0$  and  $d \in \mathbb{R}$ ,

$$\sum_{e^c < p \leq x} \frac{\log p}{p} \frac{1}{(\log \log p)^c (\log \log \log p)^d} \sim \frac{\log x}{(\log \log x)^c (\log \log \log x)^d} \quad \text{as } x \rightarrow \infty.$$

Thus  $\sum_{p>e^e} \frac{\log p}{p} \frac{1}{(\log \log p)^c (\log \log \log)^d} = \infty$ , and  $x \mapsto \sum_{e^e < p \leq e^x} \frac{\log p}{p} \frac{1}{(\log \log p)^c (\log \log \log p)^d}$  is regularly varying at  $\infty$  with exponent 1 and its slowly varying part  $L(x) \sim \frac{1}{(\log x)^c (\log \log x)^d}$  as  $x \rightarrow \infty$ , so that

$$\lim_{x \rightarrow \infty} L(x) = 0, \quad \int \frac{L(x)}{x} dx \begin{cases} < \infty & \text{if } c > 1 \text{ or } c = 1 \text{ and } d > 1, \\ = \infty & \text{if } 0 < c < 1 \text{ or } c = 1 \text{ and } d \leq 1. \end{cases}$$

In the latter case,  $u \mapsto L(e^u)$  is regularly varying at  $\infty$  with exponent  $-c \in [-1, 0)$  and its slowly varying part  $l(u) \sim \frac{1}{(\log u)^d}$  as  $u \rightarrow \infty$ , so that in the case where  $c = 1$  and  $d \leq 1$ ,

$$\lim_{u \rightarrow \infty} l(u) = \begin{cases} \infty & \text{if } d < 0, \\ 1 & \text{if } d = 0, \\ 0 & \text{if } 0 < d \leq 1. \end{cases}$$

(iv) For  $d > 0$ ,

$$\sum_{e^e < p \leq x} \frac{\log p}{p} \frac{1}{(\log \log \log p)^d} \sim \frac{\log x}{(\log \log \log x)^d} \quad \text{as } x \rightarrow \infty.$$

Thus  $\sum_{p>e^e} \frac{\log p}{p} \frac{1}{(\log \log \log)^d} = \infty$ , and  $x \mapsto \sum_{e^e < p \leq e^x} \frac{\log p}{p} \frac{1}{(\log \log \log p)^d}$  is regularly varying at  $\infty$  with exponent 1 and its slowly varying part  $L(x) \sim \frac{1}{(\log \log x)^d}$  as  $x \rightarrow \infty$ , so that

$$\lim_{x \rightarrow \infty} L(x) = 0, \quad \int \frac{L(x)}{x} dx = \infty,$$

$u \mapsto L(e^u)$  is regularly varying at  $\infty$  with exponent 0 (= slowly varying at  $\infty$ ).

---

### References

- [1] P. Billingsley: *Convergence of Probability Measures*, 2nd edition, Wiley-Interscience Publ., 1999.
- [2] W. Ehm: *A Riemann zeta stochastic process*, C. R. Acad. Sci. Paris **345** (2007), 279–282.
- [3] W. Ehm: *Functional limits of zeta type processes*, Acta Sci. Math. (Szeged) **74** (2008), 381–398.
- [4] W. Feller: *An Introduction to Probability Theory and Its Applications*, vol. II, 2nd edition, John Wiley & Sons, 1971.
- [5] G.H. Hardy and E.M. Wright: *An Introduction to the Theory of Numbers*, 6th edition, Oxford Univ. Press, 2008.
- [6] K. Itô: *Stochastic Processes*, Springer, 2004.
- [7] Y. Kasahara and S. Watanabe: *Limit theorems for point processes and their functionals*, J. Math. Soc. Japan, **38** (1986), 543–574.
- [8] T. Lindvall: *Weak convergence of probability measures and random functions in the function space  $D[0, \infty)$* , J. Appl. Prob., **10** (1973), 109–121.
- [9] W. Rudin: *Principles of Mathematical Analysis*, 3rd edition, McGraw-Hill, 1976.
- [10] G. Tenenbaum: *Introduction to Analytic and Probabilistic Number Theory*, Cambridge Univ. Press, 1995.



Faculty of Mathematics and Physics  
Institute of Science and Engineering  
Kanazawa University  
Kakuma-machi  
Kanazawa, Ishikawa 920-1192  
Japan  
e-mail: [takanob@staff.kanazawa-u.ac.jp](mailto:takanob@staff.kanazawa-u.ac.jp)