



Title	A concentration phenomenon around a shrinking hole for solutions of mean field equations
Author(s)	Ohtsuka, Hiroshi
Citation	Osaka Journal of Mathematics. 2002, 39(2), p. 395-407
Version Type	VoR
URL	https://doi.org/10.18910/7371
rights	
Note	

The University of Osaka Institutional Knowledge Archive : OUKA

<https://ir.library.osaka-u.ac.jp/>

The University of Osaka

A CONCENTRATION PHENOMENON AROUND A SHRINKING HOLE FOR SOLUTIONS OF MEAN FIELD EQUATIONS

HIROSHI OHTSUKA

(Received June 26, 2000)

1. Introduction

Let Ω be a bounded smooth domain in \mathbf{R}^2 . In this paper, we consider the following mean field equation in statistical mechanics of point vortices; see [6, 7, 15]:

$$(P) \quad \begin{aligned} -\Delta u &= \rho \frac{e^u}{\int_{\Omega} e^u} \quad \text{in } \Omega, \quad \rho > 0 \\ u &= 0 \quad \text{on } \partial\Omega. \end{aligned}$$

We note that the problem (P) for $\rho < 0$ is treated in [14]; see also [6, 7]. Analogous problems under Neumann boundary conditions are considered in relation to stationary problems of the Keller-Segel system of chemotaxis in [28]. Analogous problems on two-dimensional manifolds are also considered in relation to the prescribed Gauss curvature problem or Chern-Simons-Higgs gauge theory; see [12, 17, 26, 29] and references therein.

It should be also remarked that the following non-linear eigenvalue problem called the Gel'fand problem (see, for example, [3, 32]) also relates to our problem (P):

$$(G) \quad \begin{aligned} -\Delta u &= \lambda e^u \quad \text{in } \Omega, \quad \lambda > 0 \\ u &= 0 \quad \text{on } \partial\Omega. \end{aligned}$$

Indeed, every solution of (G) corresponds to the solution of (P) for $\rho = \int_{\Omega} \lambda \exp u \, dx$. (P) is the Euler-Lagrange equation of the following functional:

$$J_{\rho}(u) = \frac{1}{2} \int_{\Omega} |\nabla u|^2 - \rho \log \int_{\Omega} e^u \quad \text{for } u \in H_0^1(\Omega).$$

Caglioti et al. show the following facts on (P):

Facts 1.1 ([6]; see also [7]).

(1) From the Moser-Trudinger inequality [23],

$$\inf_{u \in H_0^1(\Omega)} J_\rho(u) > -\infty \quad \text{for } 0 < \rho \leq 8\pi.$$

Moreover, the problem (P) for $0 < \rho < 8\pi$ has a solution that minimizes J_ρ .

(2) The disks admit no solution of (P) for every $\rho \geq 8\pi$. More generally, let Ω be a strictly star-shaped domain, that is, there exists a constant $\alpha_0 > 0$ such that $(x \cdot \nu)(\int_{\partial\Omega} d\sigma)^{-1} \geq \alpha_0$ on $\partial\Omega$, where ν is the exterior unit outer normal vector field to $\partial\Omega$ and $d\sigma$ is the arclength measure on $\partial\Omega$. Then (P) admits no solutions if $\rho \geq 4/\alpha_0$ from the Pohožaev identity [27]. We note that $\alpha_0 = 1/(2\pi)$ when Ω is a disk.

(3) Each annulus admits the unique radial solution for every $\rho \in \mathbf{R}$.

It should be remarked that parts of Fact 1.1 are already known as results on (G). Indeed, Bandle [3, Theorem 4.16] and Suzuki and Nagasaki [35, Lemma 3] obtained similar conclusions to Fact 1.1 (2) for (G) from the Pohožaev identity (see also [3, p. 201]). The existence of radial solutions of (G) on annuli was proved by Nagasaki and Suzuki [24] (see also [30, 32, 34]) and independently by Lin [19]. Their studies on the solutions are sufficient to obtain Fact 1.1 (3) for $\rho > 0$. We note that they also studied, in different ways, the existence of non-radial solutions of (G) on annuli. It should be also remarked that, in the course of the study of (G), Suzuki proved the unique existence of solutions of (P) when Ω is simply connected and $0 < \rho < 8\pi$ [33] (see also [32, p. 263]).

We note that, on general domains other than disks and annuli, it is not clear whether a solution of (P) for $\rho \geq 8\pi$ exists. Caglioti et al. proved the existence of a minimizer of $J_{8\pi}(\cdot)$, that is, a solution of (P) for $\rho = 8\pi$ when Ω is sufficiently *thin* by analyzing the dual functional to $J_{8\pi}(\cdot)$ [6, p. 523]. In this case, supposing additionally that Ω is strictly star-shaped and admits the unique solution of (P) for $\rho = 8\pi$, they also proved the existence of a sequence $\rho_n \rightarrow 8\pi + 0$ such that (P) for ρ_n has at least two solutions [7, Theorem 7.1]. On the other hand, when Ω is simply connected and satisfies some additional conditions, we know the existence of the Weston branch of large solutions (λ, u_λ) of (G) for sufficiently small λ [36], which blows up at one point in Ω as $\lambda \rightarrow 0$. We note that Moseley [22] and subsequently Suzuki [31] (see also [32, Section 3.4]) reduced some sufficient conditions on Ω to construct the branch. Suzuki and Nagasaki proved that the Weston branch satisfies

$$\int_{\Omega} \lambda e^{u_\lambda} dx = 8\pi + C\lambda + o(\lambda) \quad \text{as } \lambda \rightarrow 0,$$

where C is a constant determined by a conformal mapping $B_1(0)$ onto Ω [35, Appendix I] (see also [32, Proposition 4.36]). This formula indicates that, on the domains satisfying $C > 0$, the solutions of (P) for $\rho > 8\pi$ and sufficiently close to 8π

exist. Moreover, Mizoguchi and Suzuki proved that the Weston branch and the trivial solution $(\lambda, u) = (0, 0)$ of (G) are connected under the additional conditions on Ω [21, Theorem 13]. This result indicates additionally the existence of solutions of (P) for $\rho = 8\pi$ as well as for $\rho > 8\pi$ and sufficiently close to 8π on the appropriate domains, an example of which is given in [21, pp. 207–208]. We note that this example is also *thin* in some sense. It should be remarked that Nagasaki and Suzuki [25] (see also [32, Section 3.3]) proved that, when a family of solutions $\{(\lambda_n, u_n)\}$ of (G) on a general domain (not necessarily a simply connected one) satisfies $\lambda_n \rightarrow 0$ and $\int_{\Omega} \lambda_n \exp u_n dx \rightarrow \Sigma_0$ as $n \rightarrow \infty$, the limit Σ_0 must be $8\pi m$ for some $m \in \{0, \infty\} \cup \mathbf{N}$. They also proved that, when $m \in \mathbf{N}$, the solution u_n of (G) blows up at distinct m points in Ω as $n \rightarrow \infty$ and obtained several necessary conditions of the limiting function of u_n . We note that this result resembles the later results of Brezis and Merle [5] and Li and Shafrir [18], which we refer as Fact 2.5 in this paper. Recently, Baraket and Pacard [4] considered the converse problem to this result of Nagasaki and Suzuki [25]. Baraket and Pacard gave, for each $m \in \mathbf{N}$, a sufficient condition of limiting functions that enables us to construct a one-parameter family of solutions $\{(\lambda, u_{\lambda})\}$ of (G) satisfying that $\int_{\Omega} \lambda \exp u_{\lambda} dx \rightarrow 8\pi m$ and u_{λ} converges to such a limiting function as $\lambda \rightarrow 0$. This result also suggests the existence of solutions of (P) at least near $\rho = 8\pi m$ on the appropriate domains for each $m \in \mathbf{N}$.

Recently, a new proof of the existence of a solution of (P) for $\rho > 8\pi$ appeared. Ding et al. proved the following fact by the minimax variational method:

Fact 1.2 ([12]). On every smooth bounded domain whose complement contains a bounded region, that is, on every smooth bounded domain with a hole, the mean field equation (P) has a solution for all $\rho \in (8\pi, 16\pi)$.

The purpose of this paper is to investigate the behavior of this solution as the hole of the domain shrinks to a point. To simplify the presentation, assuming that $0 \in \Omega$, we study the behavior of solutions of (P) on $\Omega_{\varepsilon} = \Omega \setminus \overline{B_{\varepsilon}(0)}$ as $\varepsilon \rightarrow 0$, where $B_{\varepsilon}(0) = \{x \in \mathbf{R}^2 : |x| < \varepsilon\}$. We refer (P) for Ω_{ε} as (P_{ε}) and the functional $J_{\rho}(\cdot)$ on $H_0^1(\Omega_{\varepsilon})$ for (P_{ε}) as $J_{\rho}^{\varepsilon}(\cdot)$.

Here we recall the minimax method used in [12] for the case (P_{ε}) . Let D_{ρ}^{ε} be a family of continuous functions $h : B_1(0) = \{(r, \theta) : 0 \leq r < 1, \theta \in [0, 2\pi)\} \rightarrow H_0^1(\Omega_{\varepsilon})$ satisfying

$$(1.1) \quad \lim_{r \rightarrow 1} J_{\rho}^{\varepsilon}(h(r, \theta)) \rightarrow -\infty$$

and

$$(1.2) \quad \lim_{r \rightarrow 1} m_{\Omega_{\varepsilon}}(h(r, \cdot)) \quad \text{is a continuous curve enclosing} \quad B_{\varepsilon}(0),$$

where

$$m_{\Omega_\varepsilon}(u) = \int_{\Omega_\varepsilon} x \frac{e^{u(x)}}{\int_{\Omega_\varepsilon} e^{u(x)}}.$$

Ding et al. proved that for every $\rho \in (8\pi, 16\pi)$ the minimax value

$$\alpha_\rho^\varepsilon = \inf_{h \in D_\rho^\varepsilon} \sup_{u \in h(B_1(0))} J_\rho^\varepsilon(u)$$

is achieved by a critical point of J_ρ^ε in $H_0^1(\Omega_\varepsilon)$, which is a solution of (P_ε) .

In the following, we assume each element of $H_0^1(\Omega_\varepsilon)$ to be an element of $H_0^1(\Omega)$ by extending it by 0 on $B_\varepsilon(0)$. Our result is stated as follows:

Main Theorem. *Fix $\rho \in (8\pi, 16\pi)$, a sequence $\varepsilon_n \downarrow 0$ as $n \rightarrow \infty$, and a solution u_n of (P_{ε_n}) that attains the minimax value $\alpha_\rho^{\varepsilon_n}$. Then,*

$$\frac{e^{u_n}}{\int_{\Omega} e^{u_n}} \rightharpoonup \delta_0 \quad \text{weakly * in } M(\bar{\Omega}) \quad \text{as } n \rightarrow \infty,$$

where $M(\bar{\Omega}) = C(\bar{\Omega})^*$ denotes the space of signed Radon measures over the compact space $\bar{\Omega}$ and δ_0 denotes the Dirac measure supported at the origin $0 \in \bar{\Omega}$.

We note that Lewandowski [16] obtained a concentration phenomenon similar to our Main Theorem in the following higher dimensional problem with the critical Sobolev exponent:

$$(P') \quad \begin{aligned} -\Delta u &= u^{(N+2)/(N-2)}, & u > 0 & \text{in } \Omega \subset \mathbf{R}^N \quad \text{for } N \geq 5, \\ & u = 0 & \text{on } \partial\Omega. \end{aligned}$$

Assuming that Ω is a smooth bounded star-shaped domain and $0 \in \Omega$, Lewandowski considered (P') also on the domain $\Omega_\varepsilon = \Omega \setminus \overline{B_\varepsilon(0)}$. We note that Coron [9] proved that Ω_ε admits a solution of (P') for sufficiently small ε ; see also [2] for the more general existence result for (P') . Let u_ε be a solution of (P') on Ω_ε satisfying the appropriate conditions. Then Lewandowski proved that $|\nabla u_\varepsilon|^2 \rightarrow (S_N)^{N/2} \delta_0$ as $\varepsilon \rightarrow 0$, where S_N is the best constant in the Sobolev inequality, that is, $S_N = \inf\{\int_{\mathbf{R}^N} |\nabla u|^2 : u \in H^1(\mathbf{R}^N), \|u\|_{L^{2N/(N-2)}(\mathbf{R}^N)} = 1\}$.

In contrast to our results, Lewandowski proved more on the behavior of u_ε as $\varepsilon \rightarrow 0$. Indeed, let $w_\varepsilon(x)$ be a blow-up around an appropriate point $a_\varepsilon \in \mathbf{R}^N$, that is, $w_\varepsilon(x) = t_\varepsilon^{(N-2)/2} u_\varepsilon(t_\varepsilon(x + a_\varepsilon))$ for appropriately chosen $t_\varepsilon \in (0, 1)$ satisfying $t_\varepsilon \rightarrow 0$. Then $w_\varepsilon(x)$ converges to a solution of (P') for $\Omega = \mathbf{R}^N$ in an appropriate topology.

Thus, also for our problem, it is natural to ask more precise behavior of u_n itself. It seems interesting to study the behavior of u_n by the blow-up analysis for (P) developed by Li and Shafir [18], though the author now thinks that it seems difficult.

2. Proof of Main Theorem

The key of the proof of Main Theorem is the following estimate on the minimax value α_ρ^ε :

Proposition 2.1. *For every $\rho \in (8\pi, 16\pi)$,*

$$\alpha_\rho^\varepsilon \longrightarrow -\infty \quad \text{as} \quad \varepsilon \longrightarrow 0.$$

Assuming this proposition, which we prove in Section 3, we prove Main Theorem in this section.

Set

$$\mu_n(x) = \frac{e^{\mu_n(x)}}{\int_{\Omega} e^{\mu_n(x)} dx}.$$

We regard $\{\mu_n\}$ as a bounded set in $M(\bar{\Omega})$. Thus, choosing a subsequence if necessary, we may assume that

$$\mu_n \rightharpoonup \mu_\infty \quad \text{weakly * in} \quad M(\bar{\Omega}) \quad \text{as} \quad n \longrightarrow \infty$$

for some $\mu_\infty \in M(\bar{\Omega})$. In the rest of this section, we prove μ_∞ is always δ_0 , which implies that $\mu_n \longrightarrow \delta_0$ without choosing a subsequence, that is, we obtain Main Theorem.

We prove $\mu_\infty = \delta_0$ by the following three steps:

STEP 1. $\mu_\infty = \delta_{x_\infty}$ for some $x_\infty \in \bar{\Omega}$.

STEP 2. $x_\infty \notin \partial\Omega$.

STEP 3. $x_\infty \notin \Omega \setminus \{0\}$, that is, $x_\infty = 0$.

We start from recalling the improved Moser-Trudinger inequality:

Fact 2.2 ([12, Lemma 2.2]; see also [1, Théorème 4] and [8, Theorem 2.1]). Let S_1 and S_2 be two subsets of $\bar{\Omega}$ satisfying $\text{dist}(S_1, S_2) \geq \delta_0 > 0$ and let γ_0 be a number satisfying $\gamma_0 \in (0, 1/2)$. Then for any $\varepsilon > 0$, there exists a constant $c = c(\varepsilon, \delta_0, \gamma_0) > 0$ such that

$$(2.1) \quad \int_{\Omega} e^u \leq c \exp \left\{ \frac{1}{32\pi - \varepsilon} \int_{\Omega} |\nabla u|^2 + c \right\}$$

holds for all $u \in H_0^1(\Omega)$ satisfying

$$\frac{\int_{S_1} e^u}{\int_{\Omega} e^u} \geq \gamma_0 \quad \text{and} \quad \frac{\int_{S_2} e^u}{\int_{\Omega} e^u} \geq \gamma_0.$$

From Fact 2.2, we obtain the following lemma:

Lemma 2.3. *Suppose that a sequence $\{u_n\} \subset H_0^1(\Omega)$ satisfies*

$$J_\rho(u_n) \longrightarrow -\infty \quad \text{and} \quad m_\Omega(u_n) \left(= \int_\Omega x \mu_n = \int_\Omega x \frac{e^{u_n}}{\int_\Omega e^{u_n}} \right) \longrightarrow x_\infty \quad \text{as} \quad n \longrightarrow \infty$$

for some $\rho \in (8\pi, 16\pi)$ and some $x_\infty \in \mathbf{R}^2$. Then $x_\infty \in \bar{\Omega}$ and

$$\mu_n \longrightarrow \delta_{x_\infty} \quad \text{weakly * in} \quad M(\bar{\Omega}).$$

Although we are able to prove this lemma easily by similar argument to the proof of [12, Lemma 2.3], we give a proof of Lemma 2.3 in Appendix for convenience.

Proof of Step 1. It is obvious that $J_\rho(u_n) \leq J_\rho^{\varepsilon_n}(u_n)$ because $u_n \equiv 0$ in $B_{\varepsilon_n}(0)$. Thus $J_\rho(u_n) (\leq J_\rho^{\varepsilon_n}(u_n) = \alpha_\rho^{\varepsilon_n}) \longrightarrow -\infty$ as $n \longrightarrow \infty$ from Proposition 2.1. On the other hand, there exists a subsequence of $m_\Omega(u_n)$ that converges because Ω is bounded. Using Lemma 2.3, we obtain the conclusion because $\mu_n \longrightarrow \mu_\infty$. \square

To make the next step, we recall the following fact from [10]:

Fact 2.4 ([10, p. 51 (8')]; see also [20, p. 628]). Let $N \subset \mathbf{R}^2$ be a neighborhood of $\partial\Omega$ (not $\partial\Omega_\varepsilon$) and set $\omega_0 = \bar{\Omega} \cap N$. Then, there exist positive constants ε , γ , and C depending on $\partial\Omega$ and ω_0 satisfying the following properties: $\omega = \{x \in \bar{\Omega} : \text{dist}(x, \partial\Omega) < \varepsilon\}$ is a subset of ω_0 and, for all $x \in \omega$, there exists a measurable set I_x such that

- (1) $\text{meas}(I_x) \geq \gamma$,
- (2) $I_x \subset \{y \in \omega_0 : \text{dist}(y, \partial\Omega) \geq \varepsilon/2\}$,
- (3) $u(x) \leq Cu(\xi)$ for all $\xi \in I_x$,

where u is any $C^2(\omega_0)$ function satisfying

$$\begin{aligned} -\Delta u &= f(u) \quad \text{and} \quad u > 0 \quad \text{in} \quad \omega_0 \cap \Omega (\subset \Omega), \\ u &= 0 \quad \text{on} \quad \omega_0 \cap \partial\Omega (= \partial\Omega) \end{aligned}$$

for some locally Lipschitz function $f : \mathbf{R} \longrightarrow \mathbf{R}$.

We note that Fact 2.4 is proved by the moving plane method established in [13].

Proof of Step 2. Fix a neighborhood $N \subset \mathbf{R}^2$ of $\partial\Omega$ such that $\Omega_{\varepsilon_n} \cap N$ is independent of n . Applying Fact 2.4 to this N , we obtain $\omega \subset \bar{\Omega}$ satisfying the several properties stated in Fact 2.4. We prove below that $\sup_n \|u_n\|_{L^\infty(\omega)} < \infty$, which prevents $x_\infty \in \partial\Omega$ since $\int_\Omega e^{u_n} \longrightarrow \infty$ as $n \longrightarrow \infty$ from Proposition 2.1 and Fact 1.1 (1).

Let $\omega_1 = \bigcup_{x \in \omega} I_x \subset \Omega$. Then we obtain that

$$0 \leq u_n(x) \leq \frac{C}{\gamma} \int_{I_x} u_n(y) dy \leq \frac{C}{\gamma} \|u_n\|_{L^1(\omega_1)} \leq \frac{C}{\gamma} \|u_n\|_{L^1(\Omega)} \quad \text{for every } x \in \omega,$$

that is,

$$\sup_n \|u_n\|_{L^\infty(\omega)} \leq \frac{C}{\gamma} \sup_n \|u_n\|_{L^1(\Omega)}.$$

It is rather standard to estimate $\sup_n \|u_n\|_{L^1(\Omega)}$. Indeed, let

$$\psi_n(x) = \int_{\Omega_{\varepsilon_n}} \left(\frac{1}{2\pi} \log |x-y|^{-1} - \frac{1}{2\pi} \log [\text{diam}(\Omega)]^{-1} \right) \rho \mu_n(y) dy.$$

It is obvious that

$$\begin{aligned} -\Delta \psi_n &= \rho \mu_n = \rho \frac{e^{u_n}}{\int_{\Omega_{\varepsilon_n}} e^{u_n}} \quad \text{in } \Omega_{\varepsilon_n}, \\ \psi_n &\geq 0 \quad \text{on } \partial \Omega_{\varepsilon_n}. \end{aligned}$$

We note that $\psi_n - u_n$ is harmonic in Ω_ε and non-negative on $\partial \Omega_{\varepsilon_n}$. Applying the maximum principle of harmonic functions to $\psi_n - u_n$, we obtain $\psi_n - u_n \geq 0$, that is, $\psi_n \geq u_n (\geq 0)$ in Ω_{ε_n} . Using the Young inequality for convolutions, we obtain

$$\begin{aligned} \|u_n\|_{L^1(\Omega)} &= \|u_n\|_{L^1(\Omega_{\varepsilon_n})} \leq \|\psi_n\|_{L^1(\Omega_{\varepsilon_n})} \\ &\leq \frac{\rho}{2\pi} \|\log |\cdot|^{-1}\|_{L^1(B_{\text{diam}(\Omega)}(0))} \cdot \|\mu_n\|_{L^1(\Omega_{\varepsilon_n})} + C' \\ &\leq C'' < \infty \end{aligned}$$

for some constants C' and C'' independent of n because $\|\mu_n\|_{L^1(\Omega_{\varepsilon_n})} \equiv 1$. \square

To make the final step, we recall the results of [5] and [18] concerning the solutions of $-\Delta u = V(x) \exp u$. Combining their results, we obtain the following fact:

Fact 2.5 ([5, Theorem 3] and [18, Theorem]). Let Ω be a bounded domain in \mathbf{R}^2 and let $\{w_n\} \subset C(\Omega)$ be a sequence of solutions of

$$-\Delta w = \rho e^w \quad \text{in } \mathcal{D}'(\Omega)$$

for some $\rho > 0$ such that $\sup_n \int_{\Omega} e^{w_n} < \infty$. Then, there exists a subsequence $\{u_{n_k}\}$ satisfying one of the following alternatives:

- (1) $\{u_{n_k}\}$ is bounded in $L_{\text{loc}}^\infty(\Omega)$,
- (2) $u_{n_k} \rightarrow -\infty$ uniformly on compact subsets of Ω ,
- (3) there exists a finite non-empty blow-up set $S = \{a_1, \dots, a_m\} \subset \Omega$ such that, for any $i = 1, \dots, m$, there exists $\{x_{n_k}\} \subset \Omega$ satisfying $x_{n_k} \rightarrow a_i$, $u_{n_k}(x_{n_k}) \rightarrow \infty$, and $u_{n_k}(x) \rightarrow -\infty$ uniformly on compact subsets of $\Omega \setminus S$. Moreover, $\rho \exp(u_{n_k}) \rightarrow \sum_{i=1}^m 8\pi m_i \delta_{a_i}$ weakly in the sense of measures on Ω , where m_i is a positive integer for all $i = 1, \dots, m$.

It should be remarked that, prior to [5] and [18], an analogous result to Facts 2.5 for the asymptotic behavior of the solutions of (G) as $\lambda \rightarrow 0$ exists [25], which we mentioned in Section 1.

Proof of Step 3. Suppose $x_\infty \in \Omega \setminus \{0\}$. Then, we are able to take $R > 0$ such that $\Omega_{\varepsilon_n} \supset B_R(x_\infty)$ for sufficient large n . Let $w_n(x) = u_n(x) - \log \int_{\Omega_{\varepsilon_n}} u_n(x) dx$. Then this $\{w_n(x)\}$ satisfies the assumptions of Fact 2.5 on the bounded domain $B_R(x_\infty)$. Since $\rho \exp(w_n) = \rho \mu_n \rightarrow \rho \delta_{x_\infty}$, only the alternative (3) is able to occur with $S = \{x_\infty\}$ and ρ must be $8\pi m$ for some positive integer m . Nevertheless, $\rho \in (8\pi, 16\pi)$ from the hypothesis. This is a contradiction and we obtain $x_\infty = 0$. \square

3. Estimate of the minimax value

To prove Proposition 2.1, it is enough to construct $h_\varepsilon \in D_\rho^\varepsilon$ such that

$$(3.1) \quad \sup_{u \in h_\varepsilon(B_1(0))} J_\rho^\varepsilon(u) \rightarrow -\infty \quad \text{as } \varepsilon \rightarrow 0.$$

Fix $s_0 > 0$ and set

$$u_s(t) = \begin{cases} 4 \log \frac{s_0}{s} & 0 \leq t \leq s, \\ 4 \log \frac{s_0}{t} & s \leq t \leq s_0, \\ 0 & s_0 \leq t. \end{cases}$$

We use $u_s(t)$ to construct h_ε . It is obvious that $u_{s,p}(x) = u_s(|x - p|) \in H_0^1(\Omega_\varepsilon) \subset H_0^1(\Omega)$ if $B_{s_0}(p) \subset \Omega_\varepsilon$. Moreover, we are able to obtain the following estimates:

Proposition 3.1. *Suppose $B_{s_0}(p) \subset \Omega_\varepsilon$. Then we obtain*

$$(3.2) \quad \int_{\Omega_\varepsilon} |\nabla u_{s,p}|^2 = \int_{B_{s_0}(0) \setminus B_s(0)} |\nabla(u_s(|x|))|^2 = 32\pi \log \frac{s_0}{s},$$

$$(3.3) \quad \int_{\Omega_\varepsilon} e^{u_{s,p}} \geq \int_{B_{s_0}(0) \setminus B_s(0)} e^{u_s(|x|)} = \frac{1}{s^2} \pi s_0^4 \left[1 - \left(\frac{s}{s_0} \right)^2 \right]$$

for every $0 < s < s_0$. Especially, we have

$$(3.4) \quad \frac{e^{u_{s,p}}}{\int_{\Omega_\varepsilon} e^{u_{s,p}}} \rightharpoonup \delta_p \quad \text{weakly } * \quad \text{in } M(\bar{\Omega}) \quad \text{as } s \rightarrow 0,$$

$$(3.5) \quad J_\rho^\varepsilon(u_{s,p}) \leq -2(\rho - 8\pi) \log \frac{1}{s} + O(1) \rightarrow -\infty \quad \text{as } s \rightarrow 0,$$

where $O(1)$ is independent of ε and p .

Since we obtain Proposition 3.1 by elementary calculations, we omit the proof.

We are able to take positive numbers R and $s_0 \leq R$ such that $B_{4R}(0) \setminus B_{2R}(0) \subset \Omega_\varepsilon$ for sufficiently small ε and $B_{4R+s_0}(0) \subset \Omega$. Take $s = s(\varepsilon) \rightarrow 0$ as $\varepsilon \rightarrow 0$ such that $\varepsilon \leq s \leq s_0$, which we specify later. We define

$$h_\varepsilon^0(r, \theta)(x) := \begin{cases} u_{s, p(4Rr, \theta)}(x) & 0 \leq r \leq \frac{1}{2}, \\ u_{2(1-r)s, p(4Rr, \theta)}(x) & \frac{1}{2} \leq r < 1, \end{cases}$$

where $p(r, \theta) = (r \cos \theta, r \sin \theta) \in \mathbf{R}^2$. From (3.4) and (3.5), it is easy to see that $h_\varepsilon^0(r, \theta)(\cdot)$ satisfies (1.1) and (1.2), though $h_\varepsilon^0(r, \theta)(\cdot) \notin H_0^1(\Omega_\varepsilon)$ if r is small, that is, $h_\varepsilon^0(\cdot) \notin D_\rho^\varepsilon$ yet.

We introduce the following logarithmic cut-off function, which is also used in [11]:

$$\eta_\varepsilon(t) := \begin{cases} 0, & 0 \leq t \leq \varepsilon \\ -\frac{2 \log(t/\varepsilon)}{\log \varepsilon}, & \varepsilon \leq t \leq \sqrt{\varepsilon}, \\ 1, & \sqrt{\varepsilon} \leq t. \end{cases}$$

Let

$$h_\varepsilon(r, \theta)(x) := \eta_\varepsilon(|x|)h_\varepsilon^0(r, \theta)(x).$$

This h_ε obviously belongs to D_ρ^ε and we are able to prove the following fact:

Proposition 3.2. *For every $\delta > 0$, if we take sufficiently small positive number $\sigma < 1/2$ and set $s = \varepsilon^\sigma (\geq \sqrt{\varepsilon} \geq \varepsilon)$, we obtain*

$$\sup_{(r, \theta) \in B_1(0)} J_\rho^\varepsilon(h_\varepsilon(r, \theta)(x)) \leq -2\sigma\{\rho - (1 + \delta)8\pi\} \log \frac{1}{\varepsilon} + O(1) \quad \text{as } \varepsilon \rightarrow 0.$$

Proof. We note that $h_\varepsilon(r, \theta)(x) \equiv h_\varepsilon^0(r, \theta)(x)$ if $1/2 \leq r < 1$. From (3.5), we obtain that

$$\begin{aligned} J_\rho^\varepsilon(h_\varepsilon(r, \theta)) &= J_\rho^\varepsilon(u_{2(1-r)s, p(4Rr, \theta)}) \leq -2(\rho - 8\pi) \log \frac{1}{2(1-r)s} + O(1) \\ (3.6) \quad &\leq -2(\rho - 8\pi) \log \frac{1}{s} + O(1) \quad \text{as } s \rightarrow 0 \quad \text{if } \frac{1}{2} \leq r < 1. \end{aligned}$$

For every $r \leq 1/2$ and every $\delta > 0$, we obtain

$$\begin{aligned} (3.7) \quad &\int_{\Omega_\varepsilon} |\nabla h_\varepsilon(r, \theta)|^2 \\ &\leq \left(1 + \frac{\delta}{2}\right) \int_{\Omega_\varepsilon} |\nabla h_\varepsilon^0(r, \theta)|^2 + C(\delta) \left(\sup_{x \in \Omega_\varepsilon} |h_\varepsilon^0(r, \theta)(x)|\right)^2 \int_{\Omega_\varepsilon} |\nabla(\eta_\varepsilon(|x|))|^2, \end{aligned}$$

where $C(\delta)$ is a constant depending only on δ . We note that $h_\varepsilon^0(r, \theta)(x)$ is a translation of $u_s(|x|)$ if $0 \leq r \leq 1/2$ and $\text{supp } h_\varepsilon^0(1/2, \theta) = B_{s_0}(p(2R, \theta)) \subset \Omega_\varepsilon$. Thus we obtain from (3.2) that

$$(3.8) \quad \begin{aligned} \int_{\Omega_\varepsilon} |\nabla h_\varepsilon^0(r, \theta)|^2 &\leq \int_{\Omega_\varepsilon} \left| (\nabla h_\varepsilon^0) \left(\frac{1}{2}, \theta \right) \right|^2 \\ &= \int_{\Omega_\varepsilon} |\nabla u_{s, p(2R, \theta)}|^2 = 32\pi \log \frac{1}{s} + O(1) \quad \text{as } s \rightarrow 0. \end{aligned}$$

It is easy to see that

$$(3.9) \quad \sup_{x \in \Omega_\varepsilon} |h_\varepsilon^0(r, \theta)(x)| = \sup_t |u_s(t)| = 4 \log \frac{s_0}{s}$$

and

$$(3.10) \quad \int_{\Omega_\varepsilon} |\nabla(\eta_\varepsilon(|x|))|^2 = \frac{4\pi}{\log(1/\varepsilon)}.$$

Combining (3.7–10) and choosing $s = \varepsilon^\sigma$ for sufficiently small $\sigma \in (0, 1/2)$, we obtain

$$(3.11) \quad \begin{aligned} \int_{\Omega_\varepsilon} |\nabla h_\varepsilon(r, \theta)|^2 &\leq 32\pi \left(1 + \frac{\delta}{2} \right) \log \frac{1}{s} + C(\delta)' \frac{\log s}{\log \varepsilon} \log \frac{1}{s} + O(1), \\ &\leq 32\pi\sigma(1 + \delta) \log \frac{1}{\varepsilon} + O(1) \quad \text{as } \varepsilon \rightarrow 0, \end{aligned}$$

where $C(\delta)'$ is a constant independent of ε .

On the other hand, we obtain from (3.3) that

$$(3.12) \quad \begin{aligned} \int_{\Omega_\varepsilon} e^{h_\varepsilon(r, \theta)} &\geq \int_{\Omega_\varepsilon} e^{h_\varepsilon(0, \theta)} \geq \int_{B_{s_0}(0) \setminus B_s(0)} e^{u_s(|x|)} \\ &\geq \frac{1}{s^2} \pi s_0^4 \left[1 - \left(\frac{s}{s_0} \right)^2 \right] = \frac{1}{\varepsilon^{2\sigma}} \pi s_0^4 \left[1 - \left(\frac{\varepsilon^\sigma}{s_0} \right)^2 \right]. \end{aligned}$$

Combining (3.11–12), we obtain

$$(3.13) \quad J_\rho^\varepsilon(h_\varepsilon(r, \theta)) \leq -2\sigma\{\rho - (1 + \delta)8\pi\} \log \frac{1}{\varepsilon} + O(1) \quad \text{as } \varepsilon \rightarrow 0 \quad \text{if } 0 \leq r \leq \frac{1}{2}.$$

Thus we obtain the conclusion from (3.6) and (3.13). \square

Proof of Proposition 2.1. As we assumed that $\rho > 8\pi$, we are able to take a sufficiently small $\delta > 0$ such that $\rho - (1 + \delta)8\pi > 0$. Then h_ε satisfies required property (3.1). \square

Appendix. Proof of Lemma 2.3

It is enough to see that, for every sufficiently small $r > 0$, there exists $x_{r,n} \in \bar{\Omega}$ such that

$$(A.1) \quad \int_{\Omega \cap B_r(x_{r,n})} \mu_n \geq 1 - r$$

if n is sufficiently large.

(2.1) is equivalent to the inequality

$$(A.2) \quad \frac{\rho - 16\pi + (\varepsilon/2)}{32\pi - \varepsilon} \int_{\Omega} |\nabla u|^2 + J_{\rho}(u) \geq -\rho \log c - \rho c.$$

Since we assumed that $\rho < 16\pi$, we are able to take a sufficiently small ε such that $\rho - 16\pi + (\varepsilon/2) < 0$. Then (A.2) with this ε does not hold for u_n with sufficiently large n because $J_{\rho}(u_n) \rightarrow -\infty$. Accordingly, for every $\delta_0 > 0$, every two subsets S_1 and S_2 of $\bar{\Omega}$ satisfying $\text{dist}(S_1, S_2) \geq \delta_0 > 0$, and every $\gamma_0 \in (0, 1/2)$, we obtain

$$(A.3) \quad \min \left(\frac{\int_{S_1} e^{\mu_n}}{\int_{\Omega} e^{\mu_n}}, \frac{\int_{S_2} e^{\mu_n}}{\int_{\Omega} e^{\mu_n}} \right) = \min \left(\int_{S_1} \mu_n, \int_{S_2} \mu_n \right) < \gamma_0$$

if n is sufficiently large.

Let $Q_n(r)$ be the concentration function of μ_n , that is,

$$Q_n(r) = \sup_{x \in \Omega} \int_{\Omega \cap B_r(x)} \mu_n.$$

For every $r > 0$, take $x_{r,n} \in \bar{\Omega}$ such that $\int_{\Omega \cap B_{r/2}(x_{r,n})} \mu_n = Q_n(r/2)$. Applying (A.3) for $\delta_0 = r/2$, $S_1 = \Omega \cap B_{r/2}(x_{r,n})$, and $S_2 = \Omega \setminus B_r(x_{r,n})$, we obtain that, for every $\gamma_0 \in (0, 1/2)$,

$$(A.4) \quad \min \left(\int_{S_1} \mu_n, \int_{S_2} \mu_n \right) = \min \left(Q_n \left(\frac{r}{2} \right), 1 - \int_{\Omega \cap B_r(x_{r,n})} \mu_n \right) < \gamma_0$$

if n is sufficiently large.

Since $\int_{\Omega} \mu_n \equiv 1$, it is easy to see that there exists a constant C independent of n such that

$$Q_n(r) \geq Cr^2 \quad \text{for every } 0 < r \leq \text{diam}(\Omega).$$

Taking sufficiently small γ_0 such that $Q_n(r/2) \geq \gamma_0 > 0$, we obtain (A.1) from (A.4). \square

ACKNOWLEDGEMENT. I would like to thank Professor Atsushi Inoue and Professor Takashi Suzuki for their useful suggestions.

References

- [1] T. Aubin: *Meilleures constantes dans le théorème d'inclusion de Sobolev et un théorème de Fredholm non linéaire pour la transformation conforme de la courbure scalaire*, J. Funct. Anal. **32** (1979), 148–174.
- [2] A. Bahri and J.M. Coron: *On a nonlinear elliptic equation involving the critical Sobolev exponent: The effect of the topology of the domain*, Comm. Pure Appl. Math. **41** (1988), 253–294.
- [3] C. Bandle: *Isoperimetric inequalities and applications*, Pitman, Boston, 1980.
- [4] S. Baraket and F. Pacard: *Construction of singular limits for a semilinear elliptic equation in dimension 2*, Calc. Var. Partial Differential Equations **6** (1998), 1–38.
- [5] H. Brezis and F. Merle: *Uniform estimates and blow-up behavior for solutions of $-\Delta = V(x)e^u$ in two dimensions*, Comm. in Partial Differential Equations **16** (1991), 1223–1253.
- [6] E. Caglioti, P.L. Lions, C. Marchioro and M. Pulvirenti: *A special class of stationary flows for two-dimensional Euler equations: A statistical mechanics description*, Comm. Math. Phys. **143** (1992), 501–525.
- [7] E. Caglioti, P.L. Lions, C. Marchioro and M. Pulvirenti: *A special class of stationary flows for two-dimensional Euler equations: A statistical mechanics description. Part II*, Comm. Math. Phys. **174** (1995), 229–260.
- [8] W. Chen and C. Li: *Prescribing Gaussian Curvatures on Surfaces with Conical Singularities*, J. Geom. Anal. **1** (1991), 359–372.
- [9] J.M. Coron: *Topologie et cas limite des injections de Sobolev*, C. R. Acad. Sci. Paris Math. **299** (1984), 209–212.
- [10] D.G. de Figueiredo, P.-L. Lions and R.D. Nussbaum: *A priori estimates and existence of positive solutions of semilinear elliptic equations*, J. Math. Pures Appl. **61** (1982), 41–63.
- [11] W. Ding: *Positive solutions of $\Delta u + u^{(n+2)/(n-2)} = 0$ on contractible domains*, J. Partial Differential Equations **2** (1989), 83–88.
- [12] W. Ding, J. Jost, J. Li and G. Wang: *Existence results for mean field equations*, Ann. Inst. H. Poincaré Anal. Non Linéaire **16** (1999), 653–666.
- [13] B. Gidas, W.-M. Ni and L. Nirenberg: *Symmetry and related properties via the maximum principle*, Commun. Math. Phys. **68** (1979), 209–243.
- [14] B. Gogny and P.L. Lions: *Sur les états d'équilibre pour les densités électroniques dans les plasmas*, RAIRO Modél. Math. Anal. Numér. **23** (1989), 137–153.
- [15] M.K.-H. Kiessling: *Statistical mechanics of classical particles with logarithmic interactions*, Comm. Pure Appl. Math. **46** (1993), 27–56.
- [16] R. Lewandowski: *Little holes and convergence of solutions of $-\Delta u = u^{(N+2)/(N-2)}$* , Nonlinear Anal. **14** (1990), 873–888.
- [17] Y.Y. Li: *Harnack type inequality: the method of moving planes*, Comm. Math. Phys. **200** (1999), 421–444.
- [18] Y.Y. Li and I. Shafrir: *Blow-up analysis for solutions of $-\Delta u = Ve^u$ in dimension two*, Indiana Univ. Math. J. **43** (1994), 1255–1270.
- [19] S.-S. Lin: *On non-radially symmetric bifurcation in the annulus*, J. Differential Equations **80** (1989), 251–279.
- [20] S.-S. Lin: *Semilinear elliptic equations on singularly perturbed domains*, Comm. Partial Differential Equations **16** (1991), 617–645.
- [21] N. Mizoguchi and T. Suzuki: *Equations of gas combustion: S-shaped bifurcation and mushrooms*, J. Differential Equations **134** (1997), 183–215.
- [22] J.L. Moseley: *Asymptotic solutions for a Dirichlet problem with exponential nonlinearity*, SIAM J. Math. Anal. **14** (1983), 719–735.
- [23] J. Moser: *A sharp form of an inequality by N. Trudinger*, Indiana Univ. Math. J. **20** (1971), 1077–1092.
- [24] K. Nagasaki and T. Suzuki: *Radial and nonradial solutions for the nonlinear eigenvalue problem $\Delta u + \lambda e^u = 0$ on annuli in \mathbf{R}^2* , J. Differential Equations **87** (1990), 144–168.
- [25] K. Nagasaki and T. Suzuki: *Asymptotic analysis for two-dimensional elliptic eigenvalue prob-*

lems with exponentially dominated nonlinearities, *Asymptot. Anal.* **3** (1990), 173–188.

[26] M. Nolasco and G. Tarantello: *On a sharp Sobolev-type inequality on two-dimensional compact manifolds*, *Arch. Ration. Mech. Anal.* **145** (1998), 161–195.

[27] S.I. Pohožaev: *Eigenfunctions of the equation $\Delta u + \lambda f(u) = 0$* , *Soviet Math. Dokl.* **6** (1965), 1408–1411.

[28] T. Senba and T. Suzuki: *Some structures of the solution set for a stationary system of chemotaxis*, *Adv. Math. Sci. Appl.* **10** (2000), 191–224.

[29] M. Struwe and G. Tarantello: *On multivortex solutions in Chern-Simons gauge theory*, *Boll. Unione Mat. Ital. Sez. B Artic. Ric. Mat.* **1** (1998), 109–121.

[30] T. Suzuki: *Radial and non-radial solutions for semilinear elliptic equations on circular domains*, *Geometry of solutions to partial differential equations*, Academic Press, London, G. Talenti ed. *Sympos. Math.* **30** (1989), 153–174.

[31] T. Suzuki: *Some remarks about singular perturbed solutions for Emden-Fowler equation with exponential nonlinearity*, *Functional analysis and related topics*, Springer, New York, H. Komatsu ed. *Lecture Notes in Math.* **1540** (1993), 341–360.

[32] T. Suzuki: *Semilinear elliptic equations*, Gakkōtoshō, Tokyo, 1994.

[33] T. Suzuki: *Global analysis for a two-dimensional elliptic eigenvalue problem with the exponential nonlinearity*, *Ann. Inst. H. Poincaré Anal. Non Linéaire* **9** (1992), 367–398.

[34] T. Suzuki and K. Nagasaki: *Lifting of local subdifferentiations and elliptic boundary value problems on symmetric domains. II*, *Proc. Japan Acad. Ser. A Math. Sci.* **64** (1988), 29–32.

[35] T. Suzuki and K. Nagasaki: *On the nonlinear eigenvalue problem $\Delta u + \lambda e^u = 0$* , *Trans. Amer. Math. Soc.* **309** (1988), 591–608.

[36] V.H. Weston: *On the asymptotic solution of a partial differential equation with an exponential nonlinearity*, *SIAM J. Math. Anal.* **9** (1978), 1030–1053.

Department of Mathematics
 Faculty of Science
 Tokyo Institute of Technology
 2-12-1 Oh-okayama Meguro-ku Tokyo
 152-8551, Japan
 e-mail: ohtsuka@math.titech.ac.jp

Current address:
 Department of Natural Science Education
 Kisarazu National College of Technology
 2-11-1 Kiyomidai-higashi Kisarazu-shi Chiba
 292-0041, Japan
 e-mail: ohtsuka@nebula.n.kisarazu.ac.jp