



Title	New examples of compact minitwistor spaces and their moduli space
Author(s)	Honda, Nobuhiro
Citation	Osaka Journal of Mathematics. 2010, 47(3), p. 717–730
Version Type	VoR
URL	https://doi.org/10.18910/7377
rights	
Note	

The University of Osaka Institutional Knowledge Archive : OUKA

<https://ir.library.osaka-u.ac.jp/>

The University of Osaka

NEW EXAMPLES OF COMPACT MINITWISTOR SPACES AND THEIR MODULI SPACE

NOBUHIRO HONDA

(Received July 16, 2008, revised March 27, 2009)

Abstract

In the paper [5] we obtained explicit examples of Moishezon twistor spaces of some compact self-dual four-manifolds admitting a non-trivial Killing field, and also determined their moduli space. In this note we investigate minitwistor spaces associated to these twistor spaces. We determine their structure, minitwistor lines and also their moduli space, by using a double covering structure of the twistor spaces. In particular, we find that these minitwistor spaces have different properties in many respects, compared to known examples of minitwistor spaces. Especially, we show that the moduli space of the minitwistor spaces is identified with the configuration space of different 4 points on a circle divided by the standard $PSL(2, \mathbf{R})$ -action.

1. Introduction

In [6] P.E. Jones and K.P. Tod established a reduction theory for self-dual 4-manifolds with a non-trivial Killing field. We briefly recall their results. Suppose that a self-dual metric g on a 4-manifold M admits a free isometric $U(1)$ -action. Then the quotient 3-manifold $M/U(1)$ is naturally equipped with so called a *Weyl structure*, which is a pair of a conformal structure (associated to the natural Riemannian metric on $M/U(1)$) and an affine connection compatible with the conformal structure. As a consequence of the self-duality of g , a curvature of the affine connection satisfies a kind of Einstein condition and the pair becomes *Einstein–Weyl structure* in the sense of N.J. Hitchin [3]. Moreover, the function on $M/U(1)$ obtained by associating the length of each $U(1)$ -orbits (with respect to g) satisfies certain linear equation, which is called a *monopole equation*. Thus, the Einstein–Weyl condition and the monopole equation can be thought as a non-linear and linear part of the self-duality equation respectively. This construction is invertible. Namely, if a 3-manifold N is equipped with an Einstein–Weyl structure, and if $M \rightarrow N$ is a principal $U(1)$ -bundle equipped with a (positive) solution of the monopole equation, a conformal structure on M is naturally constructed and it becomes self-dual. One can also refer to [8] for the details.

The last inversion of the reduction theory already produces non-trivial self-dual metrics even if one takes the flat Euclidean 3-space (with the natural Einstein–Weyl struc-

2000 Mathematics Subject Classification. 32L25, 32G05, 32G07, 53AA30, 53C25.

Partially supported by Research Fellowships of the Japan Society for the Promotion of Science for Young Scientists.

ture) and if one allows a certain kind of singularities for the $U(1)$ -bundle. Namely, the Eguchi–Hanson metric [1] on the cotangent bundle of \mathbf{CP}^1 and the Gibbons–Hawking metrics [2] on the minimal resolution of \mathbf{C}^2/Γ , where Γ is a cyclic subgroup of $SU(2)$, are obtained in this way. Later, C. LeBrun [7] successfully applies the inversion construction for the hyperbolic 3-space (again with the natural Einstein–Weyl structure) to realize explicit self-dual metrics on $n\mathbf{CP}^2$, the connected sum of n copies of the complex projective plane.

This reduction theory for self-dual metrics can be translated into that of twistor spaces [6] (see also [7, 8]). In particular, taking the quotient of a self-dual manifold by a $U(1)$ -action corresponds to taking the quotient of the associated twistor space by the natural holomorphic \mathbf{C}^* -action. Because \mathbf{C}^* -action can become pathological in general as seen in [4, §4], the quotient space of the \mathbf{C}^* -action does not necessarily possess a structure of a complex surface. But if the twistor space is Moishezon for example, such a pathology does not occur and the quotient space can have a natural structure of a complex surface (with singularities in general). In this case, the quotient complex surface is called a *minitwistor space*. For the Gibbons–Hawking metrics and the LeBrun metrics mentioned above, this is indeed the case; the resulting minitwistor spaces are the total space of the holomorphic tangent bundle of \mathbf{CP}^1 for the Gibbons–Hawking metrics, and the product $\mathbf{CP}^1 \times \mathbf{CP}^1$ for the LeBrun metrics. Here, an important feature in these two basic examples is that, while the self-dual (or hyperkähler) structure actually deforms, the corresponding Einstein–Weyl 3-manifolds, and hence their minitwistor spaces, do not deform.

In the paper [5] the author explicitly constructed a family of twistor spaces on $3\mathbf{CP}^2$ parametrized by a 3-dimensional connected space. The corresponding self-dual metrics have a non-trivial $U(1)$ -action but are not conformal to the LeBrun metrics. Moreover, the constructed family is complete in the sense that, on $3\mathbf{CP}^2$, every non-LeBrun self-dual metric with $U(1)$ -action is a member of this family, at least if the self-dual metric is supposed to have a positive scalar curvature. The purpose of this note is to investigate minitwistor spaces of these explicit twistor spaces of $3\mathbf{CP}^2$. Our first result is a determination of the structure of these minitwistor spaces; namely we show that the minitwistor spaces have a natural structure of a branched double covering of $\overline{\Sigma}_2$, the Hirzebruch surface of degree 2 with the (-2) -section contracted, and that the branching divisor is a smooth anticanonical curve which is an elliptic curve disjoint from the node of $\overline{\Sigma}_2$ (Theorem 2.3). Briefly speaking this is a reflection of the property that our twistor spaces of $3\mathbf{CP}^2$ have a structure of generically 2 to 1 covering of \mathbf{CP}^3 branched along a singular quartic surface that is bimeromorphic to an elliptic ruled surface. We will see that the branch elliptic curve of the minitwistor space is isomorphic to the base elliptic curve of the branch quartic surface. We also show that the isomorphism class of the branch elliptic curve uniquely determines the complex structure of the minitwistor space, and that the moduli space of our minitwistor spaces can be identified with the configuration space of different 4 points on $\mathbf{RP}^1 \simeq S^1$ modulo the

natural $PSL(2, \mathbf{R})$ -action (Theorem 2.7). In particular, our minitwistor spaces constitute a 1-dimensional moduli space, which contrasts with the case of the Gibbons–Hawking metrics and the LeBrun metrics where the moduli spaces are single points.

When investigating a twistor space, twistor lines are of fundamental significance. The images of twistor lines into minitwistor space (by the quotient map) are important as well and are called *minitwistor lines*. In Section 3, we investigate minitwistor lines in our minitwistor spaces. We prove that general minitwistor line is a nodal anticanonical curve of the minitwistor space, and that the natural morphism from a twistor line to the minitwistor line gives the normalization of the nodal curve. Thus the situation is quite different from the case of the Gibbons–Hawking metrics and the LeBrun metrics, since in these two cases, general minitwistor lines are smooth rational curves which are biholomorphic images of twistor lines. Geometrically, the appearance of the singularity of our minitwistor lines corresponds to the fact that for a general twistor line, there exists a unique \mathbf{C}^* -orbit intersecting the twistor line *twice*. Finally we give an account why such a situation occurs in our (mini)twistor spaces (Lemma 3.3), comparing with that of LeBrun in [7].

The author would like to express his gratitude to Professors Shin Nayatani for asking him what is the minitwistor space (or the Einstein–Weyl manifold) of the twistor spaces in [5], and also to Professor Takashi Nitta for useful conversations. He also thanks the referee for careful reading. Finally, he would like to thank Professor Akira Fujiki for his kind advise.

NOTATIONS. If Z is a twistor space of a self-dual 4-manifold, F denotes the canonical square root of the anticanonical line bundle $-K_Z$. Tensor product of line bundles is denoted additively.

2. The structure of minitwistor spaces and their moduli space

First we recall the main results of [5] which determine global structure of the moduli space of self-dual metrics on $3\mathbf{CP}^2 = \mathbf{CP}^2 \# \mathbf{CP}^2 \# \mathbf{CP}^2$ satisfying particular conditions.

Proposition 2.1 ([5]). *Let g be a self-dual metric on $3\mathbf{CP}^2$ satisfying the following three properties:*

- (i) *the scalar curvature of g is positive,*
- (ii) *g admits a non-zero killing field,*
- (iii) *g is not conformal to the self-dual metrics constructed by LeBrun [7].*

Let Z be the twistor space of $[g]$. Then there is a commutative diagram of holomorphic maps

$$(1) \quad \begin{array}{ccc} Z & \xrightarrow{\mu} & Z_0 \\ \Phi \downarrow & \nearrow \Phi_0 & \\ \mathbf{CP}^3 & & \end{array}$$

where Φ is a map associated to a linear system $|F|$ on Z , $\Phi_0: Z_0 \rightarrow \mathbf{CP}^3$ is a double covering whose branch locus is a quartic surface B with ordinary nodes, μ is a small resolution of the corresponding ordinary nodes of Z_0 . Moreover, the defining equation of B is given by

$$(2) \quad \{y_2y_3 + Q(y_0, y_1)\}^2 - y_0y_1(y_0 + y_1)(y_0 - ay_1) = 0,$$

where (y_0, y_1, y_2, y_3) are homogenous coordinates on \mathbf{CP}^3 , a is a positive real number, and $Q(y_0, y_1)$ is a homogeneous quadratic polynomial with real coefficients satisfying the following condition:

(*) as an equation on $\mathbf{CP}^1 = \{(y_0, y_1)\}$, the quartic equation

$$Q(y_0, y_1)^2 - y_0y_1(y_0 + y_1)(y_0 - ay_1) = 0$$

has a unique real double root.

Conversely, for any quartic surface (2) with Q and $a > 0$ satisfying the condition (*), the double covering of \mathbf{CP}^3 branched along B admits a small resolution such that the resulting manifold is a twistor space of a self-dual metric on $3\mathbf{CP}^3$ satisfying (i), (ii) and (iii).

Apriori the real double root in the condition (*) belongs to one of the two intervals $(-1, 0)$ and (a, ∞) on which the quartic $y_0y_1(y_0 + y_1)(y_0 - ay_1)$ is positive. But as showed in [5, §5.1] we can always suppose that the double root belongs to the latter interval (a, ∞) by applying a real projective transformation with respect to (y_0, y_1) , and in the following we always suppose this. We also recall that the quartic surface B defined by (2) satisfying the condition (*) has exactly three singular points

$$P_\infty := (0, 0, 0, 1), \quad \bar{P}_\infty := (0, 0, 1, 0), \quad P_0 := (\lambda_0, 1, 0, 0),$$

where $(\lambda_0, 1) \in \mathbf{CP}^1$ is the real double root (so that $\lambda_0 > a$ by the above normalization).

The killing field appeared in Proposition 2.1 generates an isometric $U(1)$ -action of the self-dual metric. This $U(1)$ -action naturally lifts and gives a holomorphic $U(1)$ -action on the twistor space Z . Taking the complexification of the last $U(1)$ -action, we obtain a holomorphic \mathbf{C}^* -action on Z . This \mathbf{C}^* -action then descends on $\mathbf{CP}^3 = \mathbf{P}H^0(F)^\vee$, which was shown to be of the form [5, Proposition 2.1]

$$(3) \quad (y_0, y_1, y_2, y_3) \mapsto (y_0, y_1, ty_2, t^{-1}y_3), \quad t \in \mathbf{C}^*.$$

Of course this \mathbf{C}^* -action leaves the quartic surface B invariant. Note that any orbit of this action is contained in a plane belonging to the pencil $\langle y_0, y_1 \rangle$, and that the closure of general orbits is a conic in these planes. On the other hand, the anti-holomorphic involution of \mathbf{CP}^3 naturally induced from the real structure on Z is explicitly given by,

again as shown in [5, Proposition 2.1],

$$(4) \quad (y_0, y_1, y_2, y_3) \mapsto (\bar{y}_0, \bar{y}_1, \bar{y}_3, \bar{y}_2),$$

where we are using the homogeneous coordinate in Proposition 2.1. Of course, as long as $t \in \mathrm{U}(1)$, the $\mathrm{U}(1)$ -action (3) and the real structure (4) commute.

With these preliminary results, we begin to investigate quotient spaces of the twistor spaces with respect to the \mathbf{C}^* -action. First we consider a quotient space of the \mathbf{C}^* -action (3) on \mathbf{CP}^3 . The rational map $\psi: \mathbf{CP}^3 \rightarrow \mathbf{CP}^3$ defined by

$$(5) \quad \psi: (y_0, y_1, y_2, y_3) \mapsto (z_0, z_1, z_2, z_3) = (y_0^2, y_1^2, y_0 y_1, y_2 y_3),$$

which is the rational map associated to a linear system formed by \mathbf{C}^* -invariant quadratic polynomials, can be regarded as a quotient map of the \mathbf{C}^* -action, since general fibers of the map is the closure of general orbits. The indeterminacy locus of ψ consists of the two points P_∞ and \bar{P}_∞ , which constitute a conjugate pair of points. The image of ψ is easily seen to be a quadratic cone $\bar{\Sigma}_2 := \{z_0 z_1 = z_2^2\}$ which has $(0, 0, 0, 1)$ as the vertex. Of course, $\bar{\Sigma}_2 \setminus \{(0, 0, 0, 1)\}$ is isomorphic to the total space of $\mathcal{O}(2)$, where the isomorphism is explicitly given by

$$(6) \quad \bar{\Sigma}_2 \setminus \{(0, 0, 0, 1)\} \ni (z_0, z_1, z_2, z_3) \mapsto (u, \zeta) = \left(\frac{z_2}{z_1}, \frac{z_3}{z_1} \right) \in \mathcal{O}(2),$$

where u is an affine coordinate on the base space of the bundle $\mathcal{O}(2)$ and ζ is a fiber coordinate valid there. Then by (2), (3), (5) and (6), it is immediate to see the following

Lemma 2.2. *Let ψ be as in (5) and B the quartic surface in Proposition 2.1. Then under the isomorphism (6), the image $\mathcal{B} := \psi(B)$ is explicitly given by*

$$(7) \quad \{\zeta + Q(u, 1)\}^2 - u(u+1)(u-a) = 0.$$

It is obvious from (7) that the projection $\mathcal{B} \rightarrow \mathbf{CP}^1$ (given by $(u, \zeta) \mapsto u$) is a double covering which has $u = 0, -1, a$ as simple branch points. Further, since (7) is an equation taking values in the line bundle $\mathcal{O}(4)$, $u = \infty$ is also a simple branched point. Also it is obvious that these are all branch points. Therefore, \mathcal{B} is a smooth elliptic curve whose complex structure is determined by a . Also we note that \mathcal{B} does not go through the node of $\bar{\Sigma}_2$, and it belongs to the anticanonical class on $\bar{\Sigma}_2$.

The following result describes a structure of quotient spaces of the twistor spaces in Proposition 2.1 with respect to the \mathbf{C}^* -action:

Theorem 2.3. *Let g , Z , Φ and B as in Proposition 2.1 and ψ , \mathcal{B} as in Lemma 2.2. Then there exists a commutative diagram of meromorphic maps:*

$$(8) \quad \begin{array}{ccc} Z & \xrightarrow{\psi} & \mathcal{T} \\ \Phi \downarrow & & \downarrow \phi \\ \mathbf{CP}^3 & \xrightarrow{\psi} & \overline{\Sigma}_2 \end{array}$$

where \mathcal{T} is a normal rational surface, Ψ is a surjective rational map, ϕ is a finite double covering map whose branch locus is the curve \mathcal{B} . Moreover, all fibers of Ψ are \mathbf{C}^* -invariant and general fibers are the closures of orbits of the \mathbf{C}^* -actions on Z .

By the last property, \mathcal{T} can be regarded as a quotient space of the \mathbf{C}^* -action on Z . Hence we call the normal rational surface \mathcal{T} as a minitwistor space associated to the twistor space in Proposition 2.1.

Proof of Theorem 2.3. Since ψ has indeterminacy at P_∞ and \overline{P}_∞ , the composition $\psi \circ \Phi$ also has indeterminacy along $\Phi^{-1}(P_\infty)$ and $\Phi^{-1}(\overline{P}_\infty)$. As seen in [5], both of the last two sets are chains of 3 smooth rational curves in Z . Let $Z' \rightarrow Z$ be a sequence of blow-ups which resolves the indeterminacy of the rational map $\psi \circ \Phi$. We may suppose that the image of the exceptional divisors are contained in $\Phi^{-1}(P_\infty) \cup \Phi^{-1}(\overline{P}_\infty)$. Let $\Phi_0: Z_0 \rightarrow \mathbf{CP}^3$ be the double covering whose branch is B as before. If $x \in \overline{\Sigma}_2 \setminus \mathcal{B}$, then $\psi^{-1}(x)$ is the closure of a \mathbf{C}^* -invariant conics which are not contained in B . Hence $\Phi_0^{-1}(\psi^{-1}(x))$ splits into the closure of two orbits of the \mathbf{C}^* -action on Z_0 . On the other hand, if $x \in \mathcal{B}$, then $\psi^{-1}(x)$ is a \mathbf{C}^* -invariant conic contained in B . Hence $\Phi_0^{-1}(\psi^{-1}(x))$ is biholomorphic to the conic $\psi^{-1}(x)$. Therefore if $Z' \rightarrow \mathcal{T} \rightarrow \overline{\Sigma}_2$ is a Stein factorization of the morphism $Z' \rightarrow \overline{\Sigma}_2$, the latter $\mathcal{T} \rightarrow \overline{\Sigma}_2$ is a double covering whose branch is exactly \mathcal{B} . Hence we obtained the commutative diagram (8). The statement about fibers of Ψ is obvious from the above argument. Further, the singular locus of \mathcal{T} is exactly the pre-image of the node of $\overline{\Sigma}_2$ since the branch curve \mathcal{B} does not go through the node. Hence \mathcal{T} is normal. Finally, the rationality of \mathcal{T} is an immediate consequence of the fact that the pre-images of the lines on the cone $\overline{\Sigma}_2$ gives a pencil of rational curves on \mathcal{T} . \square

Since the \mathbf{C}^* -action on the twistor space is compatible with the real structure, the minitwistor spaces also have real structures. It is explicitly described as follows:

Proposition 2.4. *Let \mathcal{T} be the minitwistor space in Theorem 2.3 and $\mathcal{B} \subset \overline{\Sigma}_2$ the branch elliptic curve of the double covering $\phi: \mathcal{T} \rightarrow \overline{\Sigma}_2$. Consider a real structure on $\overline{\Sigma}_2$ given by*

$$(9) \quad (u, \zeta) \mapsto (\overline{u}, \overline{\zeta}),$$

on the complement $\mathcal{O}(2)$ of the node of $\bar{\Sigma}_2$. Then \mathcal{B} is invariant under this real structure, and the real structure on \mathcal{T} covers the real structure (9) through the double covering $\mathcal{T} \rightarrow \bar{\Sigma}_2$.

Proposition 2.4 can be readily deduced by using (4)–(6) and we omit the proof.

Another realization of the minitwistor space \mathcal{T} is given by the following

Proposition 2.5. *Let \mathcal{T} be the minitwistor space in Theorem 2.3. Then as a complex surface, \mathcal{T} is obtained from $\mathbf{CP}^1 \times \mathbf{CP}^1$ in the following way: Let $p: \mathbf{CP}^1 \times \mathbf{CP}^1 \rightarrow \mathbf{CP}^1$ be (any) one of the projections, and A_1 and A_2 any two different sections of p whose self-intersection numbers are zero. Locate 4 points on $A_1 \cup A_2$ in such a way that 2 points are on A_1 and the remaining 2 points are on A_2 , and that the image of the 4 points under p is equivalent to $\{-1, 0, \infty, a\}$ under a projective transformation of \mathbf{CP}^1 . Next blowup $\mathbf{CP}^1 \times \mathbf{CP}^1$ at these 4 points. Then the strict transforms of A_1 and A_2 become (-2) -curves. Finally blow down these two curves, to obtain a surface having two ordinary nodes. This surface is biholomorphic to the minitwistor space \mathcal{T} .*

Proof. Let $v: \Sigma_2 \rightarrow \bar{\Sigma}_2$ be the minimal resolution and write $\mathcal{B}' = v^{-1}(\mathcal{B})$ which is isomorphic to \mathcal{B} . Let $\mathcal{T}' \rightarrow \Sigma_2$ be the double covering branched along \mathcal{B}' . Then \mathcal{T}' is the minimal resolution of \mathcal{T} . Consider the composition $\mathcal{T}' \rightarrow \Sigma_2 \rightarrow \mathbf{CP}^1$, where $\Sigma_2 \rightarrow \mathbf{CP}^1$ is a projection of a ruling. Since \mathcal{B}' is 2 to 1 over \mathbf{CP}^1 , general fiber of the above composition map is \mathbf{CP}^1 . Further, since \mathcal{B}' has 4 branched points, the composition map has precisely 4 singular fibers, all of which are two (-1) curves intersecting transversally. If we choose four (-1) -curves among eight ones in such a way that just two of them intersect one of the exceptional curves of the minimal resolution $\mathcal{T}' \rightarrow \mathcal{T}$, and that the other two of them intersect another exceptional curve of $\mathcal{T}' \rightarrow \mathcal{T}$, and if we blow them down, then we obtain a (relatively) minimal surface which must be $\mathbf{CP}^1 \times \mathbf{CP}^1$. This implies the claim of the proposition. \square

We note that although Proposition 2.5 gives an explicit construction of the minitwistor space as a complex surface, its real structure can never be obtained through this construction. More precisely, the blowing-down $\mathcal{T}' \rightarrow \mathbf{CP}^1 \times \mathbf{CP}^1$ in the above proof does not preserve the real structure. This can be seen, by going back to the twistor space, as follows. Consider singular fibers of $\mathcal{T}' \rightarrow \mathbf{CP}^1$ in the above proof, which are pairs of (-1) -curves intersecting transversally at a point. Then each of these singular fibers is the image of a reducible member of the linear system $|\Phi^* \mathcal{O}(1)| = |F|$, where $\Phi: Z \rightarrow \mathbf{CP}^3$ is the generically 2 to 1 covering as in Proposition 2.1. Namely, the (-1) -curves are the images of the irreducible components, by the quotient map. Since the real structure of Z exchanges the irreducible components, the two (-1) -curves in \mathcal{T}' must be a conjugate pair. Since the blowing-down $\mathcal{T}' \rightarrow \mathbf{CP}^1 \times \mathbf{CP}^1$ contracts just one of the (-1) -curves for each reducible fiber, it cannot preserve the real structure.

The complex structure of our twistor spaces in Proposition 2.1 depends not only on $a > 0$ but also on the coefficients of $Q(y_0, y_1)$ in the defining equation (2) of the branch quartic surface B . We next show that the complex structure of the minitwistor spaces does not depend on $Q(y_0, y_1)$. Namely we show the following

Proposition 2.6. *Let \mathcal{T} be the minitwistor space in Theorem 2.3. Then the complex structure of \mathcal{T} is uniquely determined by $a > 0$ in the equation (2). In other words, the complex structure of \mathcal{T} does not depend on the quadratic polynomial $Q(y_0, y_1)$ in (2).*

Proof. Fix $a > 0$ and let $Q_1 = Q_1(y_0, y_1)$ and $Q_2 = Q_2(y_0, y_1)$ be two real homogeneous quadratic polynomials satisfying the condition $(*)$ in Proposition 2.1. Let B_1 and B_2 be the quartic surfaces determined by (Q_1, a) and (Q_2, a) by the equation (2) respectively. Then we can write $Q_1(u, 1) - Q_2(u, 1) = d_0 + d_1u + d_2u^2$ for $d_0, d_1, d_2 \in \mathbf{R}$. Using these d_0, d_1, d_2 we consider a map

$$(10) \quad (u, \zeta) \mapsto (u, \zeta + d_0 + d_1u + d_2u^2).$$

Viewing (u, ζ) as a holomorphic coordinate on the total space of $\mathcal{O}(2)$ as in the proof of Theorem 2.3, this map is easily seen to be a holomorphic automorphism of the Hirzebruch surface Σ_2 . Moreover, by (7), the automorphism (10) maps \mathcal{B}_1 to \mathcal{B}_2 , where \mathcal{B}_1 and \mathcal{B}_2 are the images of B_1 and B_2 under the quotient map from \mathbf{CP}^3 to $\overline{\Sigma}_2$. Thus we have concretely obtained an isomorphism of the pair $(\overline{\Sigma}_2, \mathcal{B}_1)$ and $(\overline{\Sigma}_2, \mathcal{B}_2)$. Thus the double cover \mathcal{T}_1 and \mathcal{T}_2 whose branches are \mathcal{B}_1 and \mathcal{B}_2 respectively are mutually biholomorphic, as desired. \square

Note that the isomorphism (10) between the pairs $(\Sigma_2, \mathcal{B}_1)$ and $(\Sigma_2, \mathcal{B}_2)$ given in the above proof commutes with the real structure $(u, \zeta) \mapsto (\bar{u}, \bar{\zeta})$, since d_0, d_1 and d_2 in (10) are real. Thus the minitwistor space \mathcal{T} is uniquely determined by a not only as a complex surface but also as a complex surface with real structure.

By Proposition 2.6 we can determine the moduli space of our minitwistor spaces. Let \mathcal{M} be the moduli space of isomorphism classes of twistor spaces in Proposition 2.1. As showed in [5], \mathcal{M} is naturally identified with \mathbf{R}^3/G , where G is a reflection of \mathbf{R}^3 having 2-dimensional fixed locus. Let \mathcal{N} be the moduli space of isomorphism classes of the associated minitwistor spaces, where the isomorphism is required to commute with the real structures. We have a natural surjective map $\mathcal{M} \rightarrow \mathcal{N}$ sending each isomorphism class of a twistor space Z to the isomorphism class of minitwistor space \mathcal{T} . Then it is immediate from Proposition 2.6 to obtain the following

Theorem 2.7. *Let \mathcal{N} be the moduli space of isomorphism classes of our minitwistor spaces as explained above. Then \mathcal{N} is naturally identified with the configuration space of different 4 points on a circle, divided by the usual $\mathrm{PSL}(2, \mathbf{R})$ -action on the circle.*

In particular our minitwistor space has a non-trivial moduli, which contrasts with the known examples such as Gibbons–Hawking’s [2] and LeBrun’s [7, 8]; in these two cases the minitwistor spaces are the total space of $\mathcal{O}(2)$ and a quadratic surface $\mathbf{CP}^1 \times \mathbf{CP}^1$ respectively and therefore do not deform, although the corresponding self-dual (or hyperkähler) metrics on 4-manifolds constitute non-trivial moduli spaces.

3. Description of minitwistor lines

As showed in the previous section, our minitwistor space \mathcal{T} is a rational surface with two ordinary double points. In this section we investigate minitwistor lines in \mathcal{T} ; namely the images of twistor lines by the (rational) quotient map $\Psi: Z \rightarrow \mathcal{T}$. We investigate these minitwistor lines by using the diagram (8).

A basic fact about twistor lines in our twistor space Z was that, the image of general twistor line by the map $\Phi: Z \rightarrow \mathbf{CP}^3$ is a very special kind of conic, called *touching conic*, meaning that the conic is tangent to the branch quartic surface B at any intersection points which consist of 4 points in general [5, Definition 3.1 and Proposition 3.2]. Hence we first study the images of these touching conics by the (rational) quotient map $\psi: \mathbf{CP}^3 \rightarrow \overline{\Sigma}_2$:

Lemma 3.1. *Let $\psi: \mathbf{CP}^3 \rightarrow \overline{\Sigma}_2$ be as in Lemma 2.2. Then the image of general conics in \mathbf{CP}^3 under ψ are anticanonical curves on $\overline{\Sigma}_2$ with a unique node. Further, this is true even for general touching conics of B , and their images are nodal anticanonical curves which touch the smooth anticanonical curve \mathcal{B} at 4 points.*

Proof. Since any conic in \mathbf{CP}^3 is contained in some plane, we first study the restriction of ψ onto a general plane. Since general orbits of our \mathbf{C}^* -action (3) are conics, the restriction $\psi|_H$ is 2 to 1 for general plane H . Further, by elementary calculations, we can readily see that $\psi|_H$ can be identified with a quotient map of $H = \mathbf{CP}^2$ by a reflection with respect to some line in H , where the line is exactly the set of tangents points of \mathbf{C}^* -orbits. Further, the unique isolated fixed point of the reflection is mapped to the node of $\overline{\Sigma}_2$.

In the following we say that a conic in a plane H is *symmetric* if it is invariant under the reflection. Then in the complete linear system $|\mathcal{O}(2)|$ on H , symmetric conics in H form a codimension 2 linear subsystem. It is easily seen that if a conic C is not symmetric, its image is an anticanonical curve in $\overline{\Sigma}_2$ which has a unique node corresponding to the pair of intersection points of C and its image in H by the reflection which are not on the line. (In contrast, the image of symmetric conic becomes linearly equivalent to the branch curve of the map $H \rightarrow \overline{\Sigma}_2$.) Thus we have seen that the image of a conic C by ψ is a nodal anticanonical curve in $\overline{\Sigma}_2$, as long as C is not symmetric. This shows the first claim of the proposition.

In order to show that the claim is still true for general touching conics of B , it suffices to show that in general for a smooth quartic B_H on a plane H and for any one

of the 63 one-dimensional families of touching conics of B_H (cf. [5, Proposition 3.10]), there exists no real line in H for which all of the touching conics in the family become symmetric. In the sequel we prove this by contradiction.

Let \mathcal{C} be any one of the families of touching conics of B_H and suppose that there is a line l in H for which all members of \mathcal{C} are symmetric. Let $\Phi_H: S_H \rightarrow H$ be the double covering whose branch is B_H , and put $A := \Phi_H^{-1}(l)$. Since $\Phi_H^* \mathcal{O}(1) \simeq -K_{S_H}$, the curve A is an anticanonical curve of S_H . By [5, Lemma 3.9 (ii)] we know that there are precisely 6 reducible members of \mathcal{C} , all of which are of course pairs of bitangents. Since these reducible members also must be symmetric with respect to l by the assumption, the intersection points of each pair of bitangents must belong to l .

For any irreducible member $C \in \mathcal{C}$, the inverse image $\Phi_H^{-1}(C)$ splits into a sum of two smooth rational curves F_1 and F_2 satisfying $F_1^2 = F_2^2 = 0$ on S_H ([5, Lemma 3.8]). On the curves F_1 and F_2 , Φ_H is isomorphic to their images. Let F be any one of F_1 and F_2 and consider the pencil $|F|$. Let $h: S_H \rightarrow \mathbf{CP}^1$ be the morphism associated to $|F|$. In the next paragraph we show that the restriction $h|_A: A \rightarrow \mathbf{CP}^1$ is a double covering ramified at least 6 points, and that all these ramifications are simple (namely the map can locally be written as $z \mapsto z^2$ in a neighborhood of the ramification point.)

For this, let $l_1 + l_2$ be any reducible member of \mathcal{C} , so that l_1 and l_2 are bitangents of B_H . Then the inverse image $\Phi_H^{-1}(l_1)$ is a sum of two (-1) -curves f_1 and f'_1 intersecting transversally at two points (over the two tangent points). Similarly we can write $\Phi_H^{-1}(l_2) = f_2 + f'_2$. We may suppose that f_1 and f_2 intersect transversally. Then f'_1 and f'_2 intersect transversally, and $f_1 \cap f'_2 = f'_1 \cap f_2 = \emptyset$. Since l is supposed to pass the intersection point $l_1 \cap l_2$, the curve A passes the two points $f_1 \cap f_2$ and $f'_1 \cap f'_2$. Moreover, since Φ_H is locally biholomorphic in a neighborhood of these two points (since $l_1 \cap l_2 \notin B_H$), all the intersection of A and the four (-1) -curves f_1, f'_1, f_2, f'_2 are transversal. Furthermore, either $f_1 + f_2 \in |F|$ or $f'_1 + f'_2 \in |F|$ holds by the choice of F . We may suppose the former holds. Then by what we argued in this paragraph, on the surface S_H , we have

$$F \cdot A = (f_1 + f_2) \cdot A = f_1 \cdot A + f_2 \cdot A = 1 + 1 = 2.$$

This means that the restriction $h|_A$ is a double covering and the intersection point $A \cap (f_1 \cup f_2)$ is a simple ramification point of $h|_A$. Therefore, since the family \mathcal{C} has exactly 6 reducible members, and since the above argument works for arbitrary such member, we conclude that the double covering map $h|_A$ has at least 6 simple ramification points.

Now since the curve A is known to be irreducible (and reduced), its geometric genus makes sense. Then as the double covering $h|_A$ has at least 6 ramification points, the geometric genus of A is at least two. This is a contradiction since the geometric genus of A is at most one as it is an anticanonical curve of a smooth complex surface.

Thus we have shown that there exists no line on H with respect to which all members of \mathcal{C} are symmetric.

Combined with what we have proved in the first paragraph of this proof, it follows that $\psi(C)$ is an anticanonical curve having a unique node, for a general touching conic C . Since $\psi|_H$ is locally isomorphic outside the symmetric line on H , it follows that $\psi(C)$ still touches the image $\mathcal{B} = \psi(B)$ at four points. Thus we have proved all the claims of the proposition. \square

Using Lemma 3.1 we show the following

Proposition 3.2. *Let $\Psi: Z \rightarrow \mathcal{T}$ be the (rational) quotient map by the \mathbf{C}^* -action on Z as in Theorem 2.3. Then the image of a general twistor line in Z under Ψ is a real anticanonical curve of \mathcal{T} which has a unique node.*

In particular, general minitwistor lines in our minitwistor space \mathcal{T} are not smooth. This contrasts with the case for LeBrun's metrics, since in LeBrun twistor spaces, since in LeBrun's case, minitwistor space is $\mathbf{CP}^1 \times \mathbf{CP}^1$ and a general minitwistor line is a real (irreducible) curve of bidegree $(1, 1)$, so that always non-singular.

Proof of Proposition 3.2. As is already mentioned, $C = \Phi(L)$ is a touching conic of B for a general twistor line L . By Lemma 3.1 the image $\Gamma := \psi(C)$ is a nodal anticanonical curve of $\bar{\Sigma}_2$. Then by the diagram (8) the minitwistor line $\mathcal{L} := \Psi(L)$ is an irreducible component of $\phi^{-1}(\Gamma)$. As before let $\mu: \mathcal{T}' \rightarrow \mathcal{T}$ and $\nu: \Sigma_2 \rightarrow \bar{\Sigma}_2$ be the minimal resolutions of \mathcal{T} and $\bar{\Sigma}_2$ respectively. Let $\phi': \mathcal{T}' \rightarrow \Sigma_2$ be the natural lift of ϕ . Define a line in \mathbf{CP}^3 by $l_\infty := \{y_0 = y_1 = 0\}$ in the coordinate of Proposition 2.1. Then l_∞ is exactly the fiber of $\psi: \mathbf{CP}^3 \rightarrow \bar{\Sigma}_2$ over the node. The branch locus of ϕ' is a smooth anticanonical curve $\mathcal{B}' = \nu^{-1}(\mathcal{B})$. If L is chosen so as to satisfy $\Phi(L) \cap l_\infty = \emptyset$, then $\Gamma = \Psi(\Phi(L))$ does not go through the node. Hence $\Gamma' := \nu^{-1}(\Gamma)$ is a nodal anticanonical curve of Σ_2 which is tangent to the branch curve \mathcal{B}' at 4 points.

To prove the proposition, we have to look at irreducible components of the curve $(\phi')^{-1}(\Gamma')$. It is immediate to see that $(\phi')^*(-K_{\Sigma_2}) \simeq -2K_{\mathcal{T}'}$. Moreover, since Γ' is tangent to the branch curve \mathcal{B}' at every intersection points, $(\phi')^{-1}(\Gamma')$ splits into two irreducible curves \mathcal{L}_1 and \mathcal{L}_2 . There are two possible situations:

- (a) both \mathcal{L}_1 and \mathcal{L}_2 are smooth and the morphisms $\mathcal{L}_1 \rightarrow \Gamma'$ and $\mathcal{L}_2 \rightarrow \Gamma'$ (which are the restrictions of ϕ') are the normalizations of the nodal curve Γ' ; or
- (b) both \mathcal{L}_1 and \mathcal{L}_2 remain nodal curves and the morphisms $\mathcal{L}_1 \rightarrow \Gamma'$ and $\mathcal{L}_2 \rightarrow \Gamma'$ are isomorphic.

We now show that (a) cannot occur for general twistor lines by contradiction. To this end, recall first that \mathcal{T}' is realized as 4 points blown-up of $\mathbf{CP}^1 \times \mathbf{CP}^1$ as in Proposition 2.5. In particular we have $(-K_{\mathcal{T}'})^2 = 4$ on \mathcal{T}' . On the other hand, as is seen in the proof of Proposition 2.5, the restriction of the projection $\Sigma_2 \rightarrow \mathbf{CP}^1$ onto \mathcal{B}' has 4 branch points and consequently the composition $\mathcal{T}' \rightarrow \Sigma_2 \rightarrow \mathbf{CP}^1$ has precisely

4 singular fibers, all of which are the sum of two smooth rational curves intersecting transversally. Let $E_i + E'_i$, $1 \leq i \leq 4$, be these 4 reducible fibers. Then the blowing-down $\alpha: \mathcal{T}' \rightarrow \mathbf{CP}^1 \times \mathbf{CP}^1$ is obtained by appropriately choosing one of E_i and E'_i for each $1 \leq i \leq 4$ and then blowing them down. After possible renaming, we suppose that E_i , $1 \leq i \leq 4$, are blown-down by α . Then we can write

$$(11) \quad \mathcal{L}_i = \alpha^* \mathcal{O}(a_i, b_i) - \sum_{j=1}^4 n_{ij} E_j, \quad i = 1, 2.$$

Since $\mathcal{L}_1 + \mathcal{L}_2 = -2K_{\mathcal{T}'}$, we have $a_1 + a_2 = b_1 + b_2 = 4$ and $n_{1j} + n_{2j} = 2$ for $1 \leq j \leq 4$. Moreover obviously we have $(a_i, b_i) = (1, 3), (2, 2)$ or $(3, 1)$ for $i = 1, 2$.

Now in order to exclude the situation (a), we show that all n_{ij} in (11) must be 1. To see this, recall that the linear system $|F|$ on the twistor space has precisely 4 reducible members $\{D_i + \bar{D}_i\}_{i=1}^4$ and that all of them are \mathbf{C}^* -invariant. Since inverse images of fibers of $\mathcal{T} \rightarrow \mathbf{CP}^1$ by the quotient map Ψ are \mathbf{C}^* -invariant members of $|F|$, it follows that the 4 reducible fibers $E_i + E'_i$, $1 \leq i \leq 4$, are the images of $D_i + \bar{D}_i$. On the other hand, because $D_i \cdot L = \bar{D}_i \cdot L = 1$, general twistor lines intersect transversally with both of D_i and \bar{D}_i . Hence $\mathcal{L} = \Psi(L)$ intersects both E_i and \bar{E}_i ($1 \leq i \leq 4$) for general L . Thus combining with $n_{1j} + n_{2j} = 2$, we have $n_{ij} = 1$ for all i and j . Once this is proved, it readily follows $(a_1, b_1) = (a_2, b_2) = (2, 2)$, since the blown-up 4 points of α are located in the way described in Proposition 2.5, and in particular there are two sections of the projection with self-intersection zero on which 2 of the 4 blown-up points lie. Thus we have

$$(12) \quad \mathcal{L}_i = \alpha^* \mathcal{O}(2, 2) - E_1 - E_2 - E_3 - E_4 \quad \text{for } i = 1, 2.$$

Therefore the above situation (a) cannot occur and (b) must hold for a general L . Moreover, it is now obvious from (12) that \mathcal{L}_1 and \mathcal{L}_2 are anticanonical curves of \mathcal{T}' . The reality of minitwistor lines is clear since the quotient map Ψ preserve the real structure. Thus we obtain all claims of the proposition. \square

Finally we give another proof of the property that general minitwistor line in \mathcal{T} has a node (Proposition 3.2), and compare the case of LeBrun twistor spaces:

Lemma 3.3. *Consider the natural real structure on \mathcal{T} which is induced from that on Z (cf. Proposition 2.4). Then the real locus on \mathcal{T} consists of two disjoint 2-dimensional spheres. Moreover, exactly one of the sphere parametrizes \mathbf{C}^* -orbits (in Z) whose closures are \mathbf{C}^* -invariant twistor lines.*

Proof. As in Proposition 2.4, the real structure on the total space of $\mathcal{O}(2)$ (which is the smooth locus of $\bar{\Sigma}_2$) is given by $(u, \zeta) \mapsto (\bar{u}, \bar{\zeta})$. Therefore the real locus of $\bar{\Sigma}_2$ consists of the closure of the set $\{(u, \zeta) \mid u \in \mathbf{R}, \zeta \in \mathbf{R}\}$, which form a pinched torus,

where the pinched point is the node of $\overline{\Sigma}_2$. As is already seen, the branch locus \mathcal{B} of the double covering $\phi: \mathcal{T} \rightarrow \overline{\Sigma}_2$ is defined by

$$(13) \quad \{\xi + Q(u, 1)\}^2 - u(u+1)(u-a) = 0.$$

Hence the real locus on \mathcal{B} is a union of the two sets given by

$$(14) \quad \{(u, \xi) \in \mathcal{O}(2) \mid -1 \leq u \leq 0, \xi = \pm\sqrt{u(u+1)(u-a)}\}$$

and

$$(15) \quad \{(u, \xi) \in \mathcal{O}(2) \mid a \leq u \leq \infty, \xi = \pm\sqrt{u(u+1)(u-a)}\}$$

where in the last condition we regard $\xi = 0$ if $u = \infty$.

The sets (14) and (15) are smooth circles in \mathcal{B} . The sign of the left-hand side of (13) changes across these circles. Since the double covering map $\mathcal{T} \rightarrow \overline{\Sigma}_2$ preserves the real structure, the real locus of \mathcal{T} lies over the real locus of $\overline{\Sigma}_2$. These mean that the real locus of \mathcal{T} is either the inverse image of the two closed disks bounded by the circles (14) and (15), or the inverse image of the complement of the last two disks. But since the two points over the node of $\overline{\Sigma}_2$ (which is clearly outside the two circles) are a conjugate pair of points, the former must hold. These two double covers of the closed disks are smooth spheres.

Next we see that exactly one of the two spheres parametrizes \mathbf{C}^* -orbits in Z whose closures are \mathbf{C}^* -invariant twistor lines. From the above description, the two spheres are over two intervals $[-1, 0]$ and $[a, \infty]$ respectively, and every corresponding \mathbf{C}^* -orbits lie over a \mathbf{C}^* -invariant planes (determined by u). Then as is shown in [5, Proposition 5.22], if $-1 \leq u \leq 0$, then every real \mathbf{C}^* -orbits lying on the plane $y_0 = uy_1$ must be an image of \mathbf{C}^* -invariant twistor lines. Thus the sphere in \mathcal{T} lying over $[-1, 0]$ parametrizes \mathbf{C}^* -invariant twistor lines. On the other hand, real orbits lying on a plane $y_0 = uy_1$ with $u \geq a$ are not the image of twistor lines [5, Proposition 5.22]. This proves all the claims of the lemma. \square

By using Lemma 3.3 we now give another explanation as to why general minitwistor lines in the minitwistor space \mathcal{T} become singular. Let \mathcal{T}_2^σ and \mathcal{T}_4^σ be the connected components of the real locus \mathcal{T}^σ of \mathcal{T} , where the former and latter lie over the interval $[-1, 0]$ and $[a, \infty]$ in $\mathbf{RP}^1 \subset \mathbf{CP}^1$ respectively. (The subscripts 2 and 4 come from the notations in [5], where we wrote $I_2 = [-1, 0]$ and $I_4 = [a, \infty]$.) As above, both \mathcal{T}_2^σ and \mathcal{T}_4^σ are 2-spheres smoothly embedded in \mathcal{T} . As explained in the final part of the proof of Lemma 3.3, \mathcal{T}_2^σ parametrizes \mathbf{C}^* -orbits in Z whose closures are \mathbf{C}^* -invariant twistor lines, while \mathbf{C}^* -orbits parametrized by \mathcal{T}_4^σ are not (contained in) twistor lines. Let $O \in \mathcal{T}_4^\sigma$ be any point and think O as a real \mathbf{C}^* -orbit in Z . Consider a twistor line $L \subset Z$ which intersects O . Then by what we have explained above, O is not contained

in L and it follows by reality that $L \cap O$ consists of even number of points. The last number is 2 since O is contained in some $S \in |F|$ and $S \cdot L = 2$. Then because the orbit map $\Psi: Z \rightarrow \mathcal{T}$ identifies these conjugate pair of points, the image $\Psi(L)$ in \mathcal{T} must have a singular point at $O \in \mathcal{T}$. This is precisely the node of minitwistor line as stated in Proposition 3.2. For each $O \in \mathcal{T}_4^\sigma$, there are obviously 2-dimensional family of twistor lines intersecting L . Moreover, \mathcal{T}_4^σ to which O belongs, is also real 2-dimensional. Thus there are real 4-dimensional family of twistor lines intersecting real orbit in \mathcal{T}_4^σ . This means that the image of general twistor line must have a node.

In contrast with the situation described in Lemma 3.3, the real locus of the minitwistor space of LeBrun twistor spaces on $n\mathbf{CP}^2$ consists of a unique sphere, and it parametrizes \mathbf{C}^* -orbits whose closures are \mathbf{C}^* -invariant twistor lines. This is a reason why the image of general twistor lines by the orbit map is non-singular for LeBrun's twistor spaces.

References

- [1] T. Eguchi and A.J. Hanson: *Asymptotically flat solutions to Euclidean gravity*, Phys. Lett. **74B** (1978), 249–251.
- [2] G.W. Gibbons and S.W. Hawking: *Gravitational multi-instantons*, Phys. Lett. **78B** (1978), 430–432.
- [3] N.J. Hitchin: *Complex manifolds and Einstein's equations*; in Twistor Geometry and Nonlinear Systems (Primorsko, 1980), Lecture Notes in Math. **970**, Springer, Berlin, 73–99, 1982.
- [4] N. Honda: *Equivariant deformations of meromorphic actions on compact complex manifolds*, Math. Ann. **319** (2001), 469–481.
- [5] N. Honda: *Self-dual metrics and twenty-eight bitangents*, J. Differential Geom. **75** (2007), 175–258.
- [6] P.E. Jones and K.P. Tod: *Minitwistor spaces and Einstein–Weyl spaces*, Classical Quantum Gravity **2** (1985), 565–577.
- [7] C. LeBrun: *Explicit self-dual metrics on $\mathbf{CP}_2 \sharp \dots \sharp \mathbf{CP}_2$* , J. Differential Geom. **34** (1991), 223–253.
- [8] C. LeBrun: *Self-dual manifolds and hyperbolic geometry*; in Einstein Metrics and Yang–Mills Connections (Sanda, 1990), Lecture Notes in Pure and Appl. Math. **145**, Dekker, New York, 99–131, 1993.

Department of Mathematics
 Graduate School of Science and Engineering
 Tokyo Institute of Technology
 2-12-1, O-okayama, Meguro, 152–8551
 Japan

Current address:
 Mathematical Institute
 Tohoku University
 Sendai, 980–8578, Japan
 e-mail: honda@math.tohoku.ac.jp