



Title	J-groups of suspensions of stunted lens spaces mod 4
Author(s)	Kono, Susumu; Tamamura, Akie
Citation	Osaka Journal of Mathematics. 1989, 26(2), p. 319-345
Version Type	VoR
URL	https://doi.org/10.18910/7432
rights	
Note	

The University of Osaka Institutional Knowledge Archive : OUKA

<https://ir.library.osaka-u.ac.jp/>

The University of Osaka

J-GROUPS OF SUSPENSIONS OF STUNTED LENS SPACES MOD 4

Dedicated to Professor Shôrô Araki on his 60th birthday

SUSUMU KÔNO AND AKIE TAMAMURA

(Received August 5, 1988)

1. Introduction

Let $L^n(q) = S^{2n+1}/\mathbf{Z}_q$ be the $(2n+1)$ -dimensional standard lens space mod q . As defined in [7], we set

$$(1.1) \quad \begin{aligned} L_q^{2n+1} &= L^n(q), \\ L_q^{2n} &= \{[z_0, \dots, z_n] \in L^n(q) \mid z_n \text{ is real } \geq 0\}. \end{aligned}$$

In the previous paper [15], we determined the KO -groups $\widetilde{KO}(S^j(L_q^m/L_q^n))$ of the suspensions of the stunted lens spaces L_q^m/L_q^n for $j \equiv 1 \pmod{2}$. For primes p , the J -groups $\tilde{J}(S^j(L_p^m/L_p^n))$ have been determined (cf. [11] for $p=2$ and [12] for odd primes p). The purpose of this paper is to determine the KO - and J -groups of suspensions of stunted lens spaces mod 4.

This paper is organized as follows. In section 2 we state the main theorems: the structures of $\tilde{J}(S^j(L_{2q}^m/L_{2q}^n))$ for $j \equiv 1 \pmod{2}$ are given in Theorem 1, the proof of which is similar to that for the case $q=1$ (cf. [11]) and omitted, the structures of $\widetilde{KO}(S^j(L_4^m/L_4^n))$ and $\tilde{J}(S^j(L_4^m/L_4^n))$ for $j \equiv 0 \pmod{2}$ are given in Theorems 2 and 3 respectively. In section 3 we prepare some lemmas and recall known results in [8], [10] and [13]. By virtue of the results in [8], the proofs of Theorem 2 and 3 for the case $j \equiv 0 \pmod{4}$ are given in section 4. Applying the method used in the corresponding parts of [8], we prove Theorems 2 and 3 for the case $j \equiv 2 \pmod{4}$ in the final section.

The authors would like to express their gratitude to Professor T. Kobayashi and Professor H. Ôshima for helpful suggestions.

2. Statement of results

Let $\nu_p(s)$ denote the exponent of the prime p in the prime power decomposition of s , and $m(s)$ the function defined on positive integers as follows (cf. [3]):

$$\nu_p(m(s)) = \begin{cases} 0 & (p \neq 2 \text{ and } s \not\equiv 0 \pmod{p-1}) \\ 1 + \nu_p(s) & (p \neq 2 \text{ and } s \equiv 0 \pmod{p-1}) \\ 1 & (p = 2 \text{ and } s \not\equiv 0 \pmod{2}) \\ 2 + \nu_2(s) & (p = 2 \text{ and } s \equiv 0 \pmod{2}) \end{cases}.$$

Let \mathbf{Z}/k denote the cyclic group $\mathbf{Z}/k\mathbf{Z}$ of order k . For an integer n , $A(n)$ denotes the group defined by

$$(2.1) \quad A(n) = \begin{cases} \mathbf{Z}/2 \oplus \mathbf{Z}/2 & (n \equiv 0 \pmod{8}) \\ \mathbf{Z}/2 & (n \equiv 1 \text{ or } 7 \pmod{8}) \\ 0 & (\text{otherwise}). \end{cases}$$

If $j \equiv 1 \pmod{2}$, then we have

$$\widetilde{KO}(S^j(L_{2q}^m/L_{2q}^n)) \cong \widetilde{KO}(S^j(RP(m)/RP(n)))$$

(cf. [15, Remark 4]), and the proof of the following theorem is similar to that for the corresponding part of the theorem in [11].

Theorem 1. *Let q, j, m and n be non-negative integers with $q \geq 1$ and $m \geq n+2$.*

(1) *If $j \equiv 1 \pmod{4}$, then we have*

$$\tilde{J}(S^j(L_{2q}^m/L_{2q}^n)) \cong \begin{cases} \mathbf{Z}/m((m+j)/2) \oplus A(n+j) & (m \equiv 3 \pmod{4}) \\ A(n+j) & (\text{otherwise}). \end{cases}$$

(2) *If $j \equiv 3 \pmod{4}$, then we have*

$$\tilde{J}(S^j(L_{2q}^m/L_{2q}^n)) \cong \begin{cases} \mathbf{Z}/m((m+j)/2) & (m \equiv 1 \pmod{4}) \\ \mathbf{Z}/2 \oplus \mathbf{Z}/2 & (m+j \equiv 2 \pmod{8}) \\ \mathbf{Z}/2 & (m+j \equiv 1 \text{ or } 3 \pmod{8}) \\ 0 & (\text{otherwise}). \end{cases}$$

REMARK. (1) In the case $m=n+1$, $S^j(L_q^{n+1}/L_q^n)$ is homeomorphic to the sphere S^{n+j+1} , and J -groups of the spheres are well-known:

$$\tilde{J}(S^k) \cong \begin{cases} \mathbf{Z}/m(k/2) & (k \equiv 0 \pmod{4}) \\ \mathbf{Z}/2 & (k \equiv 1 \text{ or } 2 \pmod{8}) \\ 0 & (\text{otherwise}). \end{cases}$$

(2) If $j \equiv 1 \pmod{2}$, then the above theorem and [11] imply

$$\tilde{J}(S^j(L_{2q}^m/L_{2q}^n)) \cong \tilde{J}(S^j(RP(m)/RP(n)))$$

for any q .

In order to state the next theorem, we prepare functions h_1 , h_2 , a_1 and b_1 defined by

$$(2.2) \quad \begin{cases} h_1(n) = [n/4] + [(n+7)/8] + [(n+4)/8] \\ h_2(n) = [n/8] + [(n+6)/8] . \end{cases}$$

$$(2.3) \quad \begin{cases} a_1(m, n) = h_1(m) - [(n+1)/4] - [(n+1)/8] - [(n+6)/8] \\ b_1(m, n) = h_2(m) - [(n+7)/8] - [(n+5)/8] . \end{cases}$$

We denote the direct sum $\mathbf{Z}/n_1 \oplus \cdots \oplus \mathbf{Z}/n_i$ by (n_1, \dots, n_i) , and \mathbf{Z} by (∞) .

Theorem 2. Let j, m and n be non-negative integers with $m > n$.

(1) Suppose $j \equiv 0 \pmod{4}$.

i) If $n \not\equiv 3 \pmod{4}$, then we have

$$\widetilde{KO}(S^j(L_4^m/L_4^n)) \cong \begin{cases} \mathbf{Z}/2^{a_1(m+j, n+j)} \oplus \mathbf{Z}/2^{b_1(m+j, n+j)} & (b_1(m+j, n+j) \geq 0) \\ 0 & (b_1(m+j, n+j) < 0) . \end{cases}$$

ii) If $n \equiv 3 \pmod{4}$, then we have

$$\widetilde{KO}(S^j(L_4^m/L_4^n)) \cong \begin{cases} \mathbf{Z} \oplus \mathbf{Z}/2^{a_1(m+j, n+j)} \oplus \mathbf{Z}/2^{b_1(m+j, n+j)} & (b_1(m+j, n+j) \geq 0) \\ \mathbf{Z} & (b_1(m+j, n+j) < 0) . \end{cases}$$

(2) Suppose $j \equiv 2 \pmod{4}$.

i) If $m \geq n+9$, then we have

$$\widetilde{KO}(S^j(L_4^m/L_4^n)) \cong \mathbf{Z}/2^{[(m+j)/4] - [(n+j+1)/4]} \oplus A(m+j-1) \oplus B(n+j) ,$$

where $A(m)$ is the group defined by (2.1), and $B(n)$ is the group defined by

$$B(n) = \begin{cases} \mathbf{Z} & (n \equiv 3 \pmod{4}) \\ \mathbf{Z}/2 \oplus \mathbf{Z}/2 & (n \equiv 1 \pmod{8}) \\ \mathbf{Z}/2 & (n \equiv 0 \text{ or } 2 \pmod{8}) \\ 0 & (\text{otherwise}) . \end{cases}$$

ii) If $n+8 \geq m > n$, then the groups $\widetilde{KO}(S^j(L_4^m/L_4^n))$ are isomorphic to the corresponding groups in the following table :

$m-n$ $n+j \pmod{8}$	1	2	3	4	5	6	7	8
0	(2)	(2, 2)	(2)	(2, 2)	(2, 2)	(2, 2)	(2, 2)	(4, 2, 2)
1	(2)	(2)	(4, 2)	(4, 2)	(4, 2)	(4, 2)	(4, 4, 2)	(4, 4, 2, 2)
2	0	(4)	(4)	(4)	(4)	(4, 4)	(4, 4, 2)	(4, 2, 2)
3	(∞)	(∞)	(∞)	(∞)	($\infty, 4$)	($\infty, 4, 2$)	($\infty, 2, 2$)	($\infty, 2$)
4	0	0	0	(4)	(4, 2)	(2, 2)	(2)	(4)
5	0	0	(4)	(4, 2)	(2, 2)	(2)	(4)	(4)
6	0	(4)	(4, 2)	(2, 2)	(2)	(4)	(4)	(4)
7	(∞)	($\infty, 2$)	($\infty, 2$)	(∞)	($\infty, 2$)	($\infty, 2$)	($\infty, 2$)	($\infty, 2$)

REMARK. (1) Combining this theorem with [15, Theorem 2], we obtain the complete results for the groups $\widehat{KO}(S^j(L_4^m/L_4^n))$.

(2) The partial results for the case $n=0$ of this theorem have been obtained in [8].

In order to state the next theorem, we set

$$(2.4) \quad \begin{cases} a(j, m, n) = \begin{cases} a_1(m, n) & (j=0) \\ \min \{v_2(j)+1, a_1(m+j, n+j)\} & (j>0) \end{cases} \\ b(j, m, n) = \begin{cases} b_1(m, n) & (j=0) \\ \min \{v_2(j)+1, b_1(m+j, n+j)\} & (j>0) \end{cases} \end{cases}.$$

Main result is the following theorem.

Theorem 3. *Let j, m and n be non-negative integers with $m>n$.*

(1) *Suppose $j \equiv 0 \pmod{4}$.*

i) *If $n \not\equiv 3 \pmod{4}$, then we have*

$$\tilde{J}(S^j(L_4^m/L_4^n)) \cong \begin{cases} \mathbf{Z}/2^{a(j,m,n)} \oplus \mathbf{Z}/2^{b(j,m,n)} & (b(j, m, n) \geq 0) \\ 0 & (b(j, m, n) < 0) \end{cases}.$$

ii) *In the case $n \equiv 3 \pmod{4}$, we have*

$$\tilde{J}(S^j(L_4^m/L_4^n)) \cong \begin{cases} \mathbf{Z}/m((n+j+1)/2) \cdot 2^c \oplus \mathbf{Z}/2^{d+i} \oplus \mathbf{Z}/2^k & (b(j, m, n) \geq 0) \\ \mathbf{Z}/m((n+j+1)/2) & (b(j, m, n) < 0) \end{cases},$$

where i, k, c and d are integers defined by

$$(2.5) \quad \begin{cases} i = \begin{cases} \min \{v_2(n+1)-1, a(j, m, n)\} & (n+j \equiv 7 \pmod{8}) \\ \min \{v_2(n+1), a(j, m, n)\} & (n+j \equiv 3 \pmod{8}) \end{cases} \\ k = \min \{v_2(n+1)-1, b(j, m, n)\} \\ c = \max \{a(j, m, n)-i, b(j, m, n)-k\} \\ d = \min \{a(j, m, n)-i, b(j, m, n)-k\} \end{cases}.$$

(2) *Suppose $j \equiv 2 \pmod{4}$.*

i) *If $m \geq n+9$, then we have*

$$\tilde{J}(S^j(L_4^m/L_4^n)) \cong A(m+j-1) \oplus C(n+j),$$

where $A(m)$ is the group defined by (2.1), and $C(n)$ is the group defined by

$$C(n) = \begin{cases} \mathbf{Z}/2m((n+1)/2) \oplus \mathbf{Z}/2 & (n \equiv 3 \pmod{4}) \\ \mathbf{Z}/4 \oplus \mathbf{Z}/2 \oplus \mathbf{Z}/2 & (n \equiv 1 \pmod{8}) \\ \mathbf{Z}/4 \oplus \mathbf{Z}/2 & (n \equiv 0 \text{ or } 2 \pmod{8}) \\ \mathbf{Z}/4 & (\text{otherwise}). \end{cases}$$

ii) If $n+8 \geq m > n$, then the groups $\mathcal{J}(S^j(L_4^m/L_4^n))$ are isomorphic to the corresponding groups in the following table, where M denotes the integer $m((n+j+1)/2)$:

$m-n$ $n+j \pmod{8}$	1	2	3	4	5	6	7	8
0	(2)	(2, 2)	(2)	(2, 2)	(2, 2)	(2, 2)	(2, 2)	(4, 2, 2)
1	(2)	(2)	(4, 2)	(4, 2)	(4, 2)	(4, 2)	(4, 4, 2)	(4, 4, 2, 2)
2	0	(4)	(4)	(4)	(4)	(4, 4)	(4, 4, 2)	(4, 2, 2)
3	(M)	(M)	(M)	(M)	(M, 4)	(M, 4, 2)	(M, 2, 2)	(M, 2)
4	0	0	0	(4)	(4, 2)	(2, 2)	(2)	(4)
5	0	0	(4)	(4, 2)	(2, 2)	(2)	(4)	(4)
6	0	(4)	(4, 2)	(2, 2)	(2)	(4)	(4)	(4)
7	(M)	(M, 2)	(M, 2)	(M)	(M, 2)	(M, 2)	(M, 2)	(M, 2)

REMARK. (1) Combining this theorem with Theorem 1, we obtain the complete results for the groups $\mathcal{J}(S^j(L_4^m/L_4^n))$.

(2) The partial results for the case $j=n=0$ of this theorem have been obtained in [9].

3. Preliminaries

In this section we prepare some lemmas and recall known results which are needed to prove Theorems 2 and 3.

Lemma 3.1. *Let j be a positive integer with $j \equiv 0 \pmod{2}$ and k be an odd integer. Then we have*

$$k^j - 1 \equiv (k^2 - 1)(j/2) \pmod{2^{\nu_2(j)+4}}.$$

Proof. Since $k^2 \equiv 1 \pmod{8}$, we have

$$\begin{aligned} k^j - 1 &= (k^2 - 1)((k^2)^{(j/2)-1} + (k^2)^{(j/2)-2} + \dots + 1) \\ &\equiv (k^2 - 1)(j/2) \pmod{2^6}. \end{aligned}$$

This proves the lemma for the case $\nu_2(j)=1$. Assume that

$$k^j - 1 \equiv (k^2 - 1)(j/2) \pmod{2^{\nu_2(j)+4}}.$$

Then we have

$$\begin{aligned} k^{2j} - 1 &= (k^j - 1)(k^j + 1) \\ &\equiv (k^2 - 1)(j/2)(k^j + 1) \pmod{2^{\nu_2(j)+5}} \\ &\equiv (k^2 - 1)(j/2)(2 + (k^2 - 1)(j/2)) \pmod{2^{2\nu_2(j)+6}} \\ &\equiv (k^2 - 1)(2j/2) \pmod{2^{2\nu_2(j)+4}} \end{aligned}$$

Since $\nu_2(j) \geq 1$, this implies

$$k^{2j} - 1 \equiv (k^2 - 1)(2j/2) \pmod{2^{\nu_2(2j)+4}}$$

Thus the lemma is proved by the induction with respect to $\nu_2(j)$. q.e.d.

Considering the $\mathbf{Z}/4$ -action on $S^{2n+1} \times \mathbf{C}$ given by

$$\exp(2\pi\sqrt{-1}/4)(z, u) = (z \cdot \exp(2\pi\sqrt{-1}/4), u \cdot \exp(2\pi\sqrt{-1}/4))$$

for $(z, u) \in S^{2n+1} \times \mathbf{C}$, we have a complex line bundle

$$\eta: (S^{2n+1} \times \mathbf{C})/(\mathbf{Z}/4) \rightarrow L_4^{2n+1}.$$

Then we have the following elements

$$(3.2) \quad \begin{cases} \sigma = \eta - 1 \in \tilde{K}(L_4^{2n+1}) \\ \sigma(1) = \eta^2 - 1 \in \tilde{K}(L_4^{2n+1}). \end{cases}$$

The following proposition is well known.

Proposition 3.3. *If $m \geq 2$, then we have*

(1) (Mahammed [13]) *The ring $K(L_4^m)$ is isomorphic to the truncated polynomial ring*

$$\mathbf{Z}[\sigma]/(\sigma^{[m/2]+1}, (\sigma+1)^4-1),$$

where $(\sigma^{[m/2]+1}, (\sigma+1)^4-1)$ means the ideal of $\mathbf{Z}[\sigma]$ generated by $\sigma^{[m/2]+1}$ and $(\sigma+1)^4-1$.

(2) (Kobayashi and Sugawara [10]) *The group $\tilde{K}(L_4^m)$ is isomorphic to the direct sum of cyclic groups of order $2^{[m/2]+1}$, $2^{[m/4]}$ and $2^{[(m-2)/4]}$ generated by σ , $\sigma(1)+2^{[m/4]+1}\sigma$ and $\sigma(1)\sigma+2^{[(m+2)/4]+1}\sigma$ respectively. That is,*

$$\tilde{K}(L_4^m) \cong \langle \sigma, \sigma(1), \sigma(1)\sigma \rangle / \langle \{X_1, X_2, X_3\} \rangle,$$

where $X_1 = 2^{[m/2]+1}\sigma$, $X_2 = 2^{[m/4]}\sigma(1) + 2^{[m/4]+1}\sigma$ and $X_3 = 2^{[(m-2)/4]}\sigma(1)\sigma + 2^{[(m+2)/4]}\sigma$.

The following lemma is obtained by the above proposition.

Lemma 3.4. *Let u be a positive integer. Then, in $K(L_4^m)$,*

$$\sigma^u = a_u \sigma + b_u \sigma(1) + c_u \sigma(1)\sigma,$$

where a_u , b_u and c_u are integers defined by

$$a_u = (-2)^{u-1},$$

$$b_u = \begin{cases} 2(-4)^{(u/4)-1} & (u \equiv 0 \pmod{4}) \\ 0 & (u \equiv 1 \pmod{4}) \\ (-4)^{(u-2)/4} & (u \equiv 2 \pmod{4}) \\ -2(-4)^{(u-3)/4} & (u \equiv 3 \pmod{4}) \end{cases}$$

and

$$c_u = \begin{cases} -2^{u-2} & (u \equiv 0 \pmod{4}) \\ 2^{u-2} + 2(-4)^{(u-5)/4} & (u \equiv 1 \pmod{4}) \\ -2^{u-2} + (-4)^{(u-2)/4} & (u \equiv 2 \pmod{4}) \\ 2^{u-2} - (-4)^{(u-3)/4} & (u \equiv 3 \pmod{4}). \end{cases}$$

Proof. By making use of the relation $(\sigma+1)^4=1$, we obtain equalities

$$\begin{aligned} a_{u+1} &= -2a_u, \\ b_{u+1} &= a_u - 2c_u \end{aligned}$$

and

$$c_{u+1} = b_u - 2c_u,$$

where $a_1=1$, $b_1=0$, and $c_1=0$. Thus the lemma is proved by the induction with respect to u . q.e.d.

For each integer n with $0 \leq n < m$, we denote the inclusion map of L_4^n into L_4^m by i_n^m , and denote the kernel of the homomorphism

$$(i_n^m)^!: \tilde{K}(L_4^m) \rightarrow \tilde{K}(L_4^n)$$

by V_n . Then by Proposition 3.3 and Lemma 3.4, we obtain the following lemma.

Lemma 3.5. *Let u be a positive integer with $2u < m$. Then we have*

$$\sigma^u \equiv \begin{cases} \sigma & (u=1) \\ \sigma(1) - 2\sigma & (u=2) \\ (-1)^{(u-1)/2} (2^{(u-3)/2} \sigma(1) \sigma + 2^{u-1} \sigma) & (u \equiv 1 \pmod{2} \text{ and } u > 1) \\ (-1)^{(u-2)/2} (2^{(u-2)/2} \sigma(1) + 2^{u-1} \sigma) & (u \equiv 0 \pmod{2} \text{ and } u > 2) \end{cases}$$

modulo the subgroup V_{2u} .

Considering the $\mathbf{Z}/4$ -action on $S^{2n+1} \times \mathbf{R}$ given by

$$\exp(2\pi\sqrt{-1}/4)(z, v) = (z \cdot \exp(2\pi\sqrt{-1}/4), -v)$$

for $(z, v) \in S^{2n+1} \times \mathbf{R}$, we have a real line bundle

$$\nu: (S^{2n+1} \times \mathbf{R})/(\mathbf{Z}/4) \rightarrow L_4^{2n+1}.$$

We set

$$\kappa = \nu - 1 \in \widetilde{KO}(L_4^{2n+1}).$$

It is easy to see that

$$(3.6) \quad \begin{cases} c(\kappa) = \sigma(1) \\ r(\sigma(1)) = 2\kappa, \end{cases}$$

where $c: KO \rightarrow K$ is the complexification and $r: K \rightarrow KO$ is the real restriction. Let

$$I: \tilde{K}(X) \rightarrow \tilde{K}(S^2 X)$$

and

$$I_R: \widetilde{KO}(X) \rightarrow \widetilde{KO}(S^8 X)$$

be the Bott periodicity isomorphisms for K - and KO -theory respectively. Then we have the following proposition.

Proposition 3.7. (1) (Kobayashi and Sugawara [10]) *If $j \equiv 0 \pmod{8}$ and $m \geq 2$, then $\widetilde{KO}(S^j(L_4^m))$ is isomorphic to the direct sum of the cyclic groups of order $2^{h_1(m)}$ and $2^{h_2(m)}$ generated by $r(I^{j/2}(\sigma))$ and $I_R^{j/8}(\kappa) + 2^{\lfloor m/4 \rfloor} r(I^{j/2}(\sigma))$ respectively. That is,*

$$\widetilde{KO}(S^j(L_4^m)) \cong \langle \{r(I^{j/2}(\sigma)), I_R^{j/8}(\kappa)\} \rangle / \langle \{Y_1, Y_2\} \rangle,$$

where $Y_1 = 2^{h_1(m)} r(I^{j/2}(\sigma))$ and $Y_2 = 2^{h_2(m)} I_R^{j/8}(\kappa) + 2^{h_2(m) + \lfloor m/4 \rfloor} r(I^{j/2}(\sigma))$.

In the case $j \equiv 0 \pmod{8}$ and $m = 1$, the group $\widetilde{KO}(S^j(L_4^1)) \cong \widetilde{KO}(S^{j+1})$ is isomorphic to $\mathbb{Z}/2$ generated by $I_R^{j/8}(\kappa)$.

(2) (Kobayashi [8]) *If $j \equiv 4 \pmod{8}$ and $m \geq 4$, then the group $\widetilde{KO}(S^j(L_4^m))$ is isomorphic to the direct sum of the cyclic groups of order $2^{h_1(m+4)-2}$ and $2^{h_2(m+4)-2}$ generated by $r(I^{j/2}(\sigma))$ and $r(I^{j/2}(\sigma(1) + 2^{\lfloor m/4 \rfloor + 1} \sigma))$ respectively. That is,*

$$\widetilde{KO}(S^j(L_4^m)) \cong \langle \{r(I^{j/2}(\sigma)), r(I^{j/2}(\sigma(1)))\} \rangle / \langle \{Y_1, Y_2\} \rangle,$$

where $Y_1 = 2^{h_1(m+4)-2} r(I^{j/2}(\sigma))$ and

$$Y_2 = 2^{h_2(m+4)-2} r(I^{j/2}(\sigma(1))) + 2^{h_2(m+4) + \lfloor m/4 \rfloor - 1} r(I^{j/2}(\sigma)).$$

If $j \equiv 4 \pmod{8}$ and $1 \leq m < 4$, then we have $\widetilde{KO}(S^j(L_4^m)) \cong 0$.

4. Proof for the case $j \equiv 0 \pmod{4}$

In this section we prove the parts (1) of Theorems 2 and 3. Throughout this section, j denotes a non-negative integer with $j \equiv 0 \pmod{4}$.

We consider the elements y_1 and y_2 of $\widetilde{KO}(S^j L_4^m)$ defined by

$$(4.1) \quad \begin{cases} y_1 = r(I^{j/2}(\sigma)) \\ y_2 = \begin{cases} I_R^{j/8}(\kappa) & (j \equiv 0 \pmod{8}) \\ r(I^{j/2}(\sigma(1))) & (j \equiv 4 \pmod{8}). \end{cases} \end{cases}$$

According to [1] and [4], we have the following lemma.

Lemma 4.2. *The Adams operations are given by the following formulae.*

$$(1) \quad \psi^k(y_1) = \begin{cases} k^{j/2}y_1 & (k \equiv 1 \pmod{2}) \\ 2k^{j/2}y_2 & (j \equiv 0 \pmod{8} \text{ and } k \equiv 2 \pmod{4}) \\ k^{j/2}y_2 & (j \equiv 4 \pmod{8} \text{ and } k \equiv 2 \pmod{4}) \\ 0 & (k \equiv 0 \pmod{4}). \end{cases}$$

$$(2) \quad \psi^k(y_2) = \begin{cases} k^{j/2}y_2 & (k \equiv 1 \pmod{2}) \\ 0 & (k \equiv 0 \pmod{2}). \end{cases}$$

For each integer n with $0 \leq n < m$, we denote the kernel of the homomorphism

$$(i_n^m)^!: \widetilde{KO}(S^j L_4^m) \rightarrow \widetilde{KO}(S^j L_4^n)$$

by $VO_{m,n}^j$.

Lemma 4.3. *If $0 \leq n < m$, then we have*

$$VO_{m,n}^j \cong \begin{cases} \mathbb{Z}/2^{h_1(m+j)-h_2(n+j)-[(n+j)/4]} \oplus \mathbb{Z}/2^{h_2(m+j)-h_1(n+j)+[(n+j)/4]} & (h_2(m+j) \geq h_1(n+j) - [(n+j)/4]) \\ 0 & (h_2(m+j) < h_1(n+j) - [(n+j)/4]). \end{cases}$$

Proof. By Proposition 3.7, $VO_{m,n}^j$ is the subgroup of $\widetilde{KO}(S^j L_4^m)$ generated by Y_1 and Y_2 , where

$$Y_1 = \begin{cases} y_1 & (1 - 4[j/8] + j/2 \geq n) \\ 2^{h_1(n)}y_1 & (j \equiv 0 \pmod{8} \text{ and } n \geq 2) \\ 2^{h_1(n+4)-2}y_1 & (j \equiv 4 \pmod{8} \text{ and } n \geq 4) \end{cases}$$

and

$$Y_2 = \begin{cases} 2y_2 & (j \equiv 0 \pmod{8} \text{ and } n=1) \\ y_2 & (j \equiv 4 \pmod{8} \text{ and } 4 > n \geq 0) \\ 2^{h_2(n)}(y_2 + 2^{[n/4]}y_1) & (j \equiv 0 \pmod{8} \text{ and } n \neq 1) \\ 2^{h_2(n+4)-2}(y_2 + 2^{[n/4]+1}y_1) & (j \equiv 4 \pmod{8} \text{ and } n \geq 4). \end{cases}$$

Consider the case $h_2(m+j) \geq h_1(n+j) - [(n+j)/4]$. Suppose that $[(m+j)/4] + h_2(n+j) \geq h_1(n+j)$ and $m \geq 2$. Then we have the relations $A_i = 0$ ($i=1, 2$), where

$$A_1 = \begin{cases} 2^{h_1(m)}Y_1 & (j \equiv 0 \pmod{8} \text{ and } n=1) \\ 2^{h_1(m+4)-2}Y_1 & (j \equiv 4 \pmod{8} \text{ and } 4 > n \geq 0) \\ 2^{h_1(m+j)-h_1(n+j)}Y_1 & (\text{otherwise}) \end{cases}$$

and

$$A_2 = \begin{cases} 2^{h_2(m)}(y_2 + 2^{\lfloor m/4 \rfloor} y_1) & (j \equiv 0 \pmod{8}) \\ 2^{h_2(m+4)-2}(y_2 + 2^{\lfloor m/4 \rfloor + 1} y_1) & (j \equiv 4 \pmod{8}). \end{cases}$$

Setting

$$A_3 = \begin{cases} A_1 & (1 - 4\lfloor j/8 \rfloor + j/2 \geq n \geq 1 + 2\lfloor j/8 \rfloor - j/4) \\ A_1 + 2^{h_1(m+j) - h_2(m+j) - \lfloor (n+j)/4 \rfloor} A_2 & (\text{otherwise}), \end{cases}$$

$$u_1 = \begin{cases} Y_1 & (1 - 4\lfloor j/8 \rfloor + j/2 \geq n \geq 1 + 2\lfloor j/8 \rfloor - j/4) \\ Y_2 + 2^{h_2(n+j) - h_1(n+j) + \lfloor (m+j)/4 \rfloor} Y_1 & (\text{otherwise}) \end{cases}$$

and

$$u_2 = \begin{cases} Y_2 + 2^{\lfloor m/4 \rfloor + 1} Y_1 & (1 - 4\lfloor j/8 \rfloor + j/2 \geq n \geq 1 + 2\lfloor j/8 \rfloor - j/4) \\ (2^{\lfloor m/4 \rfloor - \lfloor n/4 \rfloor} - 1) Y_1 + 2^{h_1(n+j) - h_2(n+j) - \lfloor (n+j)/4 \rfloor} Y_2 & (\text{otherwise}), \end{cases}$$

we have

$$A_3 = 2^{h_1(m+j) - h_2(n+j) - \lfloor (n+j)/4 \rfloor} u_1$$

and

$$A_2 = 2^{h_2(m+j) - h_1(n+j) + \lfloor (n+j)/4 \rfloor} u_2.$$

Noting that

$$A_1 = \begin{cases} A_3 & (1 - 4\lfloor j/8 \rfloor + j/2 \geq n \geq 1 + 2\lfloor j/8 \rfloor - j/4) \\ A_3 - 2^{h_1(m+j) - h_2(m+j) - \lfloor (n+j)/4 \rfloor} A_2 & (\text{otherwise}), \end{cases}$$

$$Y_1 = \begin{cases} u_1 & (1 - 4\lfloor j/8 \rfloor + j/2 \geq n \geq 1 + 2\lfloor j/8 \rfloor - j/4) \\ 2^{h_1(n+j) - h_2(n+j) - \lfloor (n+j)/4 \rfloor} u_1 - u_2 & (\text{otherwise}) \end{cases}$$

and

$$Y_2 = \begin{cases} -2^{\lfloor m/4 \rfloor + 1} u_1 + u_2 & (1 - 4\lfloor j/8 \rfloor + j/2 \geq n \geq 1 + 2\lfloor j/8 \rfloor - j/4) \\ (1 - 2^{\lfloor m/4 \rfloor - \lfloor n/4 \rfloor}) u_1 + 2^{h_2(n+j) - h_1(n+j) + \lfloor (m+j)/4 \rfloor} u_2 & (\text{otherwise}), \end{cases}$$

we see that $VO_{m,n}^j$ is isomorphic to the group generated by u_1 and u_2 with relations $A_i = 0$ ($i=2, 3$). This implies the lemma for the case $\lfloor (m+j)/4 \rfloor + h_2(n+j) \geq h_1(n+j)$ and $m \geq 2$.

Suppose that $h_2(m+j) + \lfloor (n+j)/4 \rfloor \geq h_1(n+j) > \lfloor (m+j)/4 \rfloor + h_2(n+j)$ and $n \neq 1$. Then we have $n+j \equiv 1 \pmod{8}$, $n+2 \geq m > n$ and $VO_{m,n}^j \cong \mathbf{Z}/2$ generated by Y_2 . If $n=1$ and $2 \leq m \leq 3$, then we have $VO_{m,n}^j \cong \mathbf{Z}/2$ generated by Y_1 . If $n=0$, the lemma follows from Proposition 3.7. Thus the proof of the lemma for the case $h_2(m+j) \geq h_1(n+j) - \lfloor (n+j)/4 \rfloor$ is completed.

If $h_2(m+j) < h_1(n+j) - \lfloor (n+j)/4 \rfloor$, then we have $\lfloor (m+j)/8 \rfloor = \lfloor (n+j-4)/8 \rfloor$. This implies $h_1(m+j) = h_1(n+j)$, $h_2(m+j) = h_2(n+j)$ and $\lfloor (m+j)/4 \rfloor = \lfloor (n+j)/4 \rfloor$. Hence we have $VO_{m,n}^j \cong 0$.

Thus the proof of the lemma is completed.

q.e.d.

Suppose that $n \not\equiv 3 \pmod{4}$. Then we have

$$a_1(m+j, n+j) = h_1(m+j) - h_2(n+j) - [(n+j)/4]$$

and

$$b_1(m+j, n+j) = h_2(m+j) - h_1(n+j) + [(n+j)/4].$$

Thus the part i) of (1) of Theorem 2 is proved by making use of [15, Corollary 3] and Lemma 4.3.

Proof of the part i) of (1) of Theorem 3. We set

$$(4.4) \quad UO_{m,n}^j = \sum_k (\cap_e k^e (\psi^k - 1) VO_{m,n}^j).$$

Since the order of $VO_{m,n}^j$ is equal to a power of 2, we have

$$UO_{m,n}^j = \sum_{k: \text{odd}} (\psi^k - 1) VO_{m,n}^j = \sum_{k: \text{odd}} (k^{j/2} - 1) VO_{m,n}^j = 2^{\nu_2(j)+1} VO_{m,n}^j,$$

by Lemma 4.2 and Lemma 3.1. Since the order of $\widetilde{KO}(S^j(L_4^m/L_4^n))$ is finite, we have

$$\widetilde{J}(S^j(L_4^m/L_4^n)) \cong VO_{m,n}^j / UO_{m,n}^j = VO_{m,n}^j / 2^{\nu_2(j)+1} VO_{m,n}^j.$$

Thus the part i) of (1) of Theorem 3 is proved by making use of Lemma 4.3. q.e.d.

Now, we turn to the case $n \equiv 3 \pmod{4}$. In the rest of this section, n denotes a positive integer with $n \equiv 3 \pmod{4}$. It follows from [15] that we have the commutative diagram

$$(4.5) \quad \begin{array}{ccccccc} & & 0 & & 0 & & \\ & & \downarrow & & \downarrow & & \\ & & \widetilde{KO}(S^{j+n+1}) & = & \widetilde{KO}(S^{j+n+1}) & & \\ & & \downarrow \delta_1 & & \downarrow \delta_2 & & \\ 0 & \longrightarrow & VO_{m,n+1}^j & \xrightarrow{f_1} & \widetilde{KO}(S^j(L_4^m/L_4^n)) & \xrightarrow{f_2} & \widetilde{KO}(S^{j+n+1}) \longrightarrow 0 \\ & & \parallel & & \downarrow f_3 & & \downarrow \\ 0 & \longrightarrow & VO_{m,n+1}^j & \longrightarrow & VO_{m,n-1}^j & \xrightarrow{f_4} & \widetilde{KO}(S^j(L_4^{n+1}/L_4^{n-1})) \longrightarrow 0 \\ & & & & \downarrow & & \downarrow \\ & & & & 0 & & 0 \end{array}$$

of exact sequences. Since $\widetilde{KO}(S^{j+n+1})$ is isomorphic to \mathbf{Z} , the upper row of (4.5) splits. Choose $y \in \widetilde{KO}(S^j(L_4^m/L_4^n))$ such that $\beta = f_2(y)$ generates the group $\widetilde{KO}(S^{j+n+1})$. Then we have an isomorphism

$$f: VO_{m,n+1}^j \oplus \widetilde{KO}(S^{j+n+1}) \rightarrow \widetilde{KO}(S^j(L_4^m/L_4^n))$$

defined by $f(x, k\beta) = f_1(x) + ky$ for every $(x, k) \in VO_{m,n+1}^j \oplus \mathbf{Z}$. This proves the

part ii) of (1) of Theorem 2. Moreover, we have the following lemma.

Lemma 4.6. *If $j \equiv 0 \pmod{4}$ and $n \equiv 3 \pmod{4}$, then there is an element $y \in \widetilde{KO}(S^j(L_4^m/L_4^n))$ which satisfies the following conditions.*

- (1) $\beta = f_2(y)$ generates the group $\widetilde{KO}(S^{j+n+1})$.
- (2) $f_3(y) = \begin{cases} 2^{(n-1)/2}y_1 & (n+j+1 \equiv 4 \pmod{8}) \\ 2^{(n-3)/4}y_2 + 2^{(n-3)/2}y_1 & (n+1 \equiv j \equiv 0 \pmod{8}) \\ 2^{(n-7)/4}y_2 + 2^{(n-3)/2}y_1 & (n+1 \equiv j \equiv 4 \pmod{8} \text{ and } n > 3) \\ y_1 & (j \equiv 4 \pmod{8} \text{ and } n = 3). \end{cases}$

Proof. Suppose that $j \equiv 0 \pmod{8}$ and $n \equiv 7 \pmod{8}$. By the proof of Lemma 4.3, we have

$$VO_{m,n+1}^j = \langle \{2^{(n+1)/2}y_1, 2^{(n+1)/4}y_2\} \rangle$$

and

$$VO_{m,n-1}^j = \langle \{2^{(n-1)/2}y_1, 2^{(n-3)/4}y_2 + 2^{(n-3)/2}y_1\} \rangle.$$

Hence

$$\widetilde{KO}(S^j(L_4^{n+1}/L_4^{n-1})) \cong VO_{m,n-1}^j / VO_{m,n+1}^j \cong \mathbb{Z}/4$$

and the first group is generated by $f_4(2^{(n-3)/4}y_2 + 2^{(n-3)/2}y_1)$. It follows from the commutativity of the diagram (4.5) that the element y can be chosen to satisfy $f_3(y) = 2^{(n-3)/4}y_2 + 2^{(n-3)/2}y_1$. The proofs for the other cases are similar. q.e.d.

In the rest of this section, we fix an element $y \in \widetilde{KO}(S^j(L_4^m/L_4^n))$ which satisfies the conditions of Lemma 4.6.

Lemma 4.7. *If k is an odd integer, then the Adams operation ψ^k is given by*

$$\psi^k(y) = k^{(n+j+1)/2}y + ((k^{j/2} - k^{(n+j+1)/2})/4)f_1(4f_3(y)).$$

Proof. We necessarily have

$$\psi^k(y) = uy + f_1(x)$$

for some integer u and an element $x \in VO_{m,n+1}^j$. By using the ψ -map f_2 , we see that $u = k^{(n+j+1)/2}$. Under f_3 , $f_1(x)$ maps into x and y maps into $f_3(y)$, and we see that

$$\psi^k(f_3(y)) = k^{(n+j+1)/2}f_3(y) + x.$$

It follows from Lemma 4.2 that

$$k^{j/2}f_3(y) = k^{(n+j+1)/2}f_3(y) + x.$$

This implies that

$$x = ((k^{j/2} - k^{(n+j+1)/2})/4)(4f_3(y))$$

and

$$\begin{aligned}\psi^k(y) &= k^{(n+j+1)/2}y + f_1(x) \\ &= k^{(n+j+1)/2}y + ((k^{j/2} - k^{(n+j+1)/2})/4)f_1(4f_3(y)).\end{aligned}\quad \text{q.e.d.}$$

We now recall some definition in [3]. Set $Y = \widetilde{KO}(S^j(L_4^m/L_4^n))$ and let f be a function which assigns to each integer k a non-negative integer $f(k)$. Given such a function f , we define Y_f to be the subgroup of Y generated by

$$\{k^{f(k)}(\psi^k - 1)(y) \mid k \in \mathbf{Z}, y \in Y\};$$

that is,

$$Y_f = \langle \{k^{f(k)}(\psi^k - 1)(y) \mid k \in \mathbf{Z}, y \in Y\} \rangle.$$

Then the kernel of the homomorphism $J'': Y \rightarrow J''(Y)$ coincides with $\bigcap_f Y_f$, where the intersection runs over all functions f .

Suppose that f satisfies

$$(4.8) \quad f(k) \geq m + \max \{v_p(m((n+j+1)/2)) \mid p \text{ is a prime divisor of } k\}$$

for every $k \in \mathbf{Z}$. For each odd integer i , $N(i)$ denotes the integer chosen to satisfy the property

$$(4.9) \quad iN(i) \equiv 1 \pmod{2^m}.$$

In the following calculation we put $(n+j+1)/2 = u$ for the sake of simplicity. From Lemmas 3.1 and 4.7, we have

$$\begin{aligned}k^{f(k)}(\psi^k - 1)(y) &= k^{f(k)}(k^u - 1)y + k^{f(k)}((k^{j/2} - k^u)/4)f_1(4f_3(y)) \\ &= k^{f(k)}(k^u - 1)y + k^{f(k)}N(u/2^{v_2(u)})((u(k^{j/2} - 1) - u(k^u - 1))/2^{v_2(u)+2})f_1(4f_3(y)) \\ &\equiv k^{f(k)}(k^u - 1)y + k^{f(k)}N(u/2^{v_2(u)})((j/2)(k^u - 1) - u(k^u - 1))/2^{v_2(u)+2}f_1(4f_3(y)) \\ &\quad \pmod{f_1(UO_{m,n+1}^j)} \\ &= (k^{f(k)}(k^u - 1)/2^{v_2(u)+2})(2^{v_2(u)+2}y - N(u/2^{v_2(u)}((n+1)/2)f_1(4f_3(y)))\end{aligned}$$

By virtue of [3; II, Theorem (2.7) and Lemma (2.12)], we have

$$\begin{aligned}&\langle f_1(UO_{m,n+1}^j) \cup \{k^{f(k)}(\psi^k - 1)(y) \mid k \in \mathbf{Z}\} \rangle \\ &= \langle f_1(UO_{m,n+1}^j) \cup \{m(u)/2^{v_2(u)+2}(2^{v_2(u)+2}y - N(u/2^{v_2(u)}((n+1)/2)f_1(4f_3(y))))\} \rangle\end{aligned}$$

Therefore,

$$Y_f = \langle f_1(VO_{m,n+1}^j) \cup \{m((n+j+1)/2)y - Mf_1(4f_3(y))\} \rangle$$

where $M = (m((n+j+1)/2)/2^{v_2(n+j+1)+1})N((n+j+1)/2^{v_2(n+j+1)}((n+1)/2)$. Since this is true for every function f which satisfies (4.8), we have

$$(4.10) \quad J''(Y) \cong Y / \langle f_1(VO_{m,n+1}^j) \cup \{m((n+j+1)/2)y - Mf_1(4f_3(y))\} \rangle.$$

Suppose that $b(j, m, n) \geq 0$. It follows from the proof of Lemma 4.3 that $VO_{m,n+1}^j \cong \mathbf{Z}/2^{a_1(m+j,n+j)} \oplus \mathbf{Z}/2^{b_1(m+j,n+j)}$ is generated by

$$u_1 = \begin{cases} 2^{(n+1)/4}y_2 + (2^{\lfloor (m+n+1)/4 \rfloor} + 2^{(n+1)/2})y_1 & (j \equiv 0 \pmod{8}) \\ 2^{(n-3)/4}y_2 + (2^{\lfloor (m+n+1)/4 \rfloor} + 2^{(n+1)/2})y_1 & (j \equiv 4 \pmod{8}) \end{cases}$$

and

$$u_2 = \begin{cases} 2^{(n+5)/4}y_2 + 2^{\lfloor (m+n+5)/4 \rfloor}y_1 & (j \equiv n-3 \equiv 0 \pmod{8}) \\ 2^{(n+1)/4}y_2 + 2^{\lfloor (m+n+5)/4 \rfloor}y_1 & (j \equiv n-3 \equiv 4 \pmod{8}) \\ 2^{(n+1)/4}y_2 + 2^{\lfloor (m+n+1)/4 \rfloor}y_1 & (j \equiv n+1 \equiv 0 \pmod{8}) \\ 2^{(n-3)/4}y_2 + 2^{\lfloor (m+n+1)/4 \rfloor}y_1 & (j \equiv n+1 \equiv 4 \pmod{8}). \end{cases}$$

By Lemma 4.6, we have

$$4f_3(y) = \begin{cases} 2u_1 - u_2 & (n+j \equiv 3 \pmod{8}) \\ u_1 - u_2 & (j \equiv 4 \pmod{8} \text{ and } n=3) \\ (1 - 2^{\lfloor (m-n+3)/4 \rfloor})u_1 + (1 + 2^{\lfloor (m-n+3)/4 \rfloor})u_2 & (\text{otherwise}). \end{cases}$$

Therefore

$$J''(Y) \cong \langle y, u_1, u_2 \rangle / \langle \{M_0y + M_1u_1 + M_2u_2, 2^{a(j,m,n)}u_1, 2^{b(j,m,n)}u_2\} \rangle,$$

where

$$M_0 = m((n+j+1)/2),$$

$$M_1 = \begin{cases} -2M & (n+j \equiv 3 \pmod{8}) \\ -M & (j \equiv 4 \pmod{8} \text{ and } n=3) \\ -(1 - 2^{\lfloor (m-n+3)/4 \rfloor})M & (\text{otherwise}) \end{cases}$$

and

$$M_2 = \begin{cases} -(1 + 2^{\lfloor (m-n+3)/4 \rfloor})M & (n+j \equiv 7 \pmod{8} \text{ and } n>3) \\ M & (\text{otherwise}). \end{cases}$$

Set

$$i = \begin{cases} \min \{a(j, m, n), v_2(n+1)\} & (n+j \equiv 3 \pmod{8}) \\ \min \{a(j, m, n), v_2(n+1)-1\} & (n+j \equiv 7 \pmod{8}) \end{cases}$$

and

$$k = \min \{b(j, m, n), \nu_2(n+1)-1\}.$$

Since $\nu_2(M)=\nu_2(n+1)-1$, the greatest common divisor of M_1 and $2^{a(j,m,n)}$ is equal to 2^i , and the greatest common divisor of M_2 and $2^{b(j,m,n)}$ is equal to 2^k . Choose integers e_1, e_2, e_3 and e_4 with

$$e_1 2^{a(j,m,n)} + e_2 M_1 = 2^i$$

and

$$e_3 2^{b(j,m,n)} + e_4 M_2 = 2^k.$$

For the sake of simplicity, we put $a=a(j, m, n)$ and $b=b(j, m, n)$ in the following calculation. If $a-i \geq b-k$, then we have

$$A \begin{pmatrix} M_0 y + M_1 u_1 + M_2 u_2 \\ 2^a u_1 \\ 2^b u_2 \end{pmatrix} = \begin{pmatrix} 2^{a-i} M_0 y \\ 2^{b-k+i} ((e_2 M_0 / 2^i) y + u_1) \\ 2^k ((e_4 M_0 / 2^k) y + (e_4 M_1 / 2^k) u_1 + u_2) \end{pmatrix},$$

where

$$A = \begin{pmatrix} 2^{a-i} & -M_1/2^i & -(M_2/2^k) 2^{a-b-i+k} \\ e_2 2^{b-k} & e_1 2^{b-k} & -e_2 M_2/2^k \\ e_4 & 0 & e_3 \end{pmatrix}$$

and $\det A=1$. This implies that

$$J''(Y) \cong \mathbf{Z}/2^{a-i} M_0 \oplus \mathbf{Z}/2^{b-k+i} \oplus \mathbf{Z}/2^k.$$

On the other hand, if $b-k > a-i$, then we have

$$B \begin{pmatrix} M_0 y + M_1 u_1 + M_2 u_2 \\ 2^a u_1 \\ 2^b u_2 \end{pmatrix} = \begin{pmatrix} 2^{b-k} M_0 y \\ 2^a u_1 \\ 2^k ((e_4 M_0 / 2^k) y + (e_4 M_1 / 2^k) u_1 + u_2) \end{pmatrix},$$

where

$$B = \begin{pmatrix} 2^{b-k} & -(M_1/2^i) 2^{-a+b+i-k} & -M_2/2^k \\ 0 & 1 & 0 \\ e_4 & 0 & e_3 \end{pmatrix}$$

and $\det B=1$. This implies that

$$J''(Y) \cong \mathbf{Z}/2^{b-k} M_0 \oplus \mathbf{Z}/2^a \oplus \mathbf{Z}/2^k.$$

Thus we have

(4.11) If $j \equiv 0 \pmod{4}$, $n \equiv 3 \pmod{4}$ and $b(j, m, n) \geq 0$, then we have

$$\mathcal{J}(S^j(L_4^m/L_4^n)) \cong \mathbf{Z}/m((n+j+1)/2) \cdot 2^c \oplus \mathbf{Z}/2^{d+i} \oplus \mathbf{Z}/2^k,$$

where i, k, c and d are integers defined by (2.5).

Next suppose that $b(j, m, n) < 0$. It follows from Lemma 4.3 that we have $VO_{m,n+1}^j \cong 0$. This implies that the homomorphism f_2 in the diagram (4.5) is an isomorphism of ψ -groups. Thus we obtain

(4.12) If $j \equiv 0 \pmod{4}$, $n \equiv 3 \pmod{4}$ and $b(j, m, n) < 0$ then we have

$$\mathcal{J}(S^j(L_4^m/L_4^n)) \cong \mathbf{Z}/m((n+j+1)/2).$$

Now, combining (4.11) and (4.12) we obtain the part ii) of (1) of Theorem 3. Thus the proof for the case $j \equiv 0 \pmod{4}$ is completed.

5. Proof for the case $j \equiv 2 \pmod{4}$

In this section we prove the parts (2) of Theorems 2 and 3. Throughout this section j denotes a positive integer with $j \equiv 2 \pmod{4}$. Consider the elements x_1, x_2 and x_3 of $\tilde{K}(S^j L_4^m)$ defined by

$$(5.1) \quad \begin{cases} x_1 = I^{j/2} \sigma, \\ x_2 = I^{j/2} \sigma(1), \\ x_3 = I^{j/2} (\sigma(1) \sigma). \end{cases}$$

According to [1], we have the following lemma.

Lemma 5.2. *The Adams operations are given by the following formulae.*

$$\begin{aligned} (1) \quad \psi^k(x_1) &= \begin{cases} k^{j/2}(x_1 + x_2 + x_3) & (k \equiv 3 \pmod{4}) \\ k^{j/2} x_1 & (k \equiv 1 \pmod{4}) \\ k^{j/2} x_2 & (k \equiv 2 \pmod{4}) \\ 0 & (k \equiv 0 \pmod{4}). \end{cases} \\ (2) \quad \psi^k(x_2) &= \begin{cases} k^{j/2} x_2 & (k \equiv 1 \pmod{2}) \\ 0 & (k \equiv 0 \pmod{2}). \end{cases} \\ (3) \quad \psi^k(x_3) &= \begin{cases} k^{j/2}(-x_3 - 2x_2) & (k \equiv 3 \pmod{4}) \\ k^{j/2} x_3 & (k \equiv 1 \pmod{4}) \\ 0 & (k \equiv 0 \pmod{2}). \end{cases} \end{aligned}$$

Consider the elements X_1, X_2 and X_3 of $\tilde{K}(S^j L_4^m)$ defined by

$$(5.3) \quad \begin{cases} X_1 = \begin{cases} 2^{[(n+3)/2]} x_1 & (n \geq 1) \\ x_3 & (n = 0) \end{cases} \\ X_2 = \begin{cases} 2^{[(n+1)/4]} x_2 & (n \equiv 0 \text{ or } 3 \pmod{4}) \\ 2^{[(n+1)/4]} x_2 + 2^{[(n+1)/2]} x_1 & (n \equiv 1 \text{ or } 2 \pmod{4}) \end{cases} \\ X_3 = \begin{cases} 2^{[(n-1)/4]} x_3 & (n \equiv 1 \text{ or } 2 \pmod{4}) \\ 2^{[(n-1)/4]} x_3 + 2^{[(n+1)/2]} x_1 & (n \equiv 0 \text{ or } 3 \pmod{4} \text{ and } n \geq 3) \\ x_1 & (n = 0) \end{cases} \end{cases}$$

For each integer n with $0 \leq n \leq m$, we denote the kernel of the homomorphism

$$(i_n^m)^!: \tilde{K}(S^j L_4^m) \rightarrow \tilde{K}(S^j L_4^n)$$

by $V_{m,n}^j$. Then by Proposition 3.3, we have

$$(5.4) \quad V_{m,2[(n+1)/2]}^j = \langle \{X_1, X_2, X_3\} \rangle.$$

Consider the Bott exact sequence (cf. [5] and [6, (12.2)])

$$(5.5) \quad \rightarrow \widetilde{KO}(S^{j+2}X) \xrightarrow{c} \tilde{K}(S^{j+2}X) \xrightarrow{r \circ I^{-1}} \widetilde{KO}(S^jX) \xrightarrow{\partial} \widetilde{KO}(S^{j+1}X) \rightarrow$$

for $X = L_4^m/L_4^n$, where ∂ is the homomorphism defined by the exterior product with the generator of $\widetilde{KO}(S^1)$. Using the isomorphisms

$$VO_{m,2[(n+1)/2]}^{j+2} \cong \widetilde{KO}(S^{j+2}(L_4^m/L_4^{2[(n+1)/2]}))$$

and

$$V_{m,2[(n+1)/2]}^j \cong \tilde{K}(S^j(L_4^m/L_4^{2[(n+1)/2]})),$$

we obtain the exact sequence

$$(5.6) \quad \rightarrow VO_{m,2u}^{j+2} \xrightarrow{I^{-1} \circ c} V_{m,2u}^j \xrightarrow{r_1} \widetilde{KO}(S^j(L_4^m/L_4^{2u})) \xrightarrow{\partial} G \rightarrow 0,$$

where $u = [(n+1)/2]$ and

$$G = \begin{cases} \widetilde{KO}(S^{j+1}(L_4^m/L_4^{2u})) & (m+j \equiv 0, 1 \text{ or } 2 \pmod{8}) \\ 0 & (\text{otherwise}). \end{cases}$$

Consider the generators y_1 and y_2 of $\widetilde{KO}(S^{j+2}L_4^m)$ defined by (4.1).

Lemma 5.7. (1) $I^{-1} \circ c(y_1) = 2x_1 + x_2 + x_3$.

$$(2) \quad I^{-1} \circ c(y_2) = \begin{cases} x_2 & (j \equiv 6 \pmod{8}) \\ 2x_2 & (j \equiv 2 \pmod{8}). \end{cases}$$

Proof. (1) By (4.1), we have

$$\begin{aligned} I^{-1} \circ c(y_1) &= I^{-1}(c \circ r(I^{(j+2)/2}(\sigma))) = I^{j/2}((1+t)(\sigma)) \\ &= I^{j/2}(2\sigma + \sigma(1) + \sigma(1)\sigma) = 2x_1 + x_2 + x_3. \end{aligned}$$

(2) If $j \equiv 6 \pmod{8}$, then by (3.6) we have

$$I^{-1} \circ c(y_2) = I^{-1}(I^{(j+2)/2}(c(\kappa))) = I^{j/2}(\sigma(1)) = x_2.$$

If $j \equiv 2 \pmod{8}$, then we have

$$\begin{aligned} I^{-1} \circ c(y_2) &= I^{-1}(I^{(j+2)/2}(c \circ r(\sigma(1)))) = I^{j/2}((1+t)(\sigma(1))) \\ &= I^{j/2}(2\sigma(1)) = 2x_2. \end{aligned} \quad \text{q.e.d.}$$

5.1. Proof for the case $n \equiv 0 \pmod{2}$. By Proposition 3.7 and (5.4), we have

$$VO_{m,n}^{j+2} = \begin{cases} \langle \{2^{h_1(n)}y_1, 2^{h_2(n)}(y_2 + 2^{\lfloor n/4 \rfloor}y_1)\} \rangle & (j \equiv 6 \pmod{8}) \\ \langle \{2^{h_1(n+4)-2}y_1, 2^{h_2(n+4)-2}(y_2 + 2^{\lfloor n/4 \rfloor+1}y_1)\} \rangle & (j \equiv 2 \pmod{8} \text{ and } n \geq 4) \\ \langle \{y_1, y_2\} \rangle & (j \equiv 2 \pmod{8} \text{ and } 0 \leq n \leq 2) \end{cases}$$

and $V_{m,n}^j = \langle \{X_1, X_2, X_3\} \rangle$. Using Lemma 5.7, we obtain

(5.8) For the homomorphism r_1 in the exact sequence (5.6), we have

$$\text{Ker } r_1 = \begin{cases} \langle \{2X_2, (1-2^{n/4})X_1 + X_2 + 2^{(n+4)/4}X_3\} \rangle & (n+j \equiv 2 \pmod{8} \text{ and } n \geq 4) \\ \langle \{2X_2, X_1 + X_2 + 2X_3\} \rangle & (j \equiv 2 \pmod{8} \text{ and } n=0) \\ \langle \{X_2, (1-2^{n/4})X_1 + 2^{(n+4)/4}X_3\} \rangle & (n+j \equiv 6 \pmod{8} \text{ and } n \geq 4) \\ \langle \{X_2, X_1 + 2X_3\} \rangle & (j \equiv 6 \pmod{8} \text{ and } n=0) \\ \langle \{2X_2 - X_1, 2^{(n+2)/4}X_3 + 2X_2\} \rangle & (n+j \equiv 0 \pmod{8}) \\ \langle \{2X_2 - X_1, 2^{(n-2)/4}X_3 + X_2\} \rangle & (n+j \equiv 4 \pmod{8}). \end{cases}$$

If $m \geq n+2$, then $\text{Im } r_1$ is isomorphic to the group generated by $\{X_1, X_2, X_3\}$ with relations $A_i = 0$ ($1 \leq i \leq 5$), where

$$\begin{aligned} A_1 &= \begin{cases} 2X_2 & (n+j \equiv 2 \pmod{8}) \\ X_2 & (n+j \equiv 6 \pmod{8}) \\ 2X_2 - X_1 & (n \equiv 2 \pmod{4}), \end{cases} \\ A_2 &= \begin{cases} (1-2^{n/4})X_1 + X_2 + 2^{(n+4)/4}X_3 & (4 \leq n \equiv 0 \pmod{4}) \\ X_1 + X_2 + 2X_3 & (n=0) \\ 2^{(n+2)/4}X_3 + 2X_2 & (n+j \equiv 0 \pmod{8}) \\ 2^{(n-2)/4}X_3 + X_2 & (n+j \equiv 4 \pmod{8}), \end{cases} \end{aligned}$$

$$A_3 = \begin{cases} 2^{\lfloor (m-n+2)/4 \rfloor} X_3 + 2^{\lfloor (m-n-2)/4 \rfloor} (2^{\lfloor (m-n+2)/4 \rfloor} - 1) X_1 & (4 \leq n \equiv 0 \pmod{4}) \\ 2^{\lfloor (m-2)/4 \rfloor} X_1 + 2^{2\lfloor (m+2)/4 \rfloor} X_3 & (n=0) \\ 2^{\lfloor (m-n)/4 \rfloor} X_3 + 2^{2\lfloor (m-n)/4 \rfloor} X_1 & (n \equiv 2 \pmod{4}), \end{cases}$$

$$A_4 = \begin{cases} 2^{\lfloor (m-n)/4 \rfloor} X_2 + 2^{2\lfloor (m-n)/4 \rfloor} X_1 & (4 \leq n \equiv 0 \pmod{4}) \\ 2^{\lfloor m/4 \rfloor} X_2 + 2^{2\lfloor m/4 \rfloor + 1} X_3 & (n=0) \\ 2^{\lfloor (m-n-2)/4 \rfloor} (2X_2 + (2^{\lfloor (m-n+2)/4 \rfloor} - 1)X_1) & (n \equiv 2 \pmod{4}) \end{cases}$$

and

$$A_5 = \begin{cases} 2^{\lfloor (m+2)/2 \rfloor} X_3 & (n=0) \\ 2^{\lfloor (m-n)/2 \rfloor} X_1 & (\text{otherwise}). \end{cases}$$

Thus we obtain

(5.9) If $m+j-2 \geq n+j \equiv 2 \pmod{8}$ or $m+j-6 \geq n+j \equiv 2 \pmod{8}$, then we have

$$\begin{aligned} r_1(V_{m,n}^j) &\cong \langle \{X_1, X_2, X_3\} \rangle / \langle \{A_1, A_2, B_3\} \rangle \\ &\cong \begin{cases} \mathbf{Z}/2^{\lfloor (m+j)/4 \rfloor - \lfloor (n+j)/4 \rfloor} \oplus \mathbf{Z}/2 & (n+j \equiv 0 \text{ or } 2 \pmod{8}) \\ \mathbf{Z}/2^{\lfloor (m+j)/4 \rfloor - \lfloor (n+j)/4 \rfloor} & (n+j \equiv 4 \text{ or } 6 \pmod{8}), \end{cases} \end{aligned}$$

where $B_3 = 2^{\lfloor (m+j)/4 \rfloor - \lfloor (n+j)/4 \rfloor} X_3$,

$$A_1 = \begin{cases} 2X_2 & (n+j \equiv 2 \pmod{8}) \\ X_2 & (n+j \equiv 6 \pmod{8}) \\ 2X_2 - X_1 & (n \equiv 2 \pmod{8}) \end{cases}$$

and

$$A_2 = \begin{cases} 2^{(n+4)/4} X_3 + X_2 + (1 - 2^{n/4}) X_1 & (4 \leq n \equiv 0 \pmod{4}) \\ 2X_3 + X_2 + X_1 & (n=0) \\ 2^{(n+2)/4} X_3 + 2X_2 & (n+j \equiv 0 \pmod{8}) \\ 2^{(n-2)/4} X_3 + X_2 & (n+j \equiv 4 \pmod{8}). \end{cases}$$

If $n+j \equiv 2 \pmod{8}$ and $n+5 \geq m \geq n+2$, then we have

$$r_1(V_{m,n}^j) \cong \langle \{X_1, X_2, X_3\} \rangle / \langle \{B_1, X_2 - 2X_3, 4X_3\} \rangle \cong \mathbf{Z}/4,$$

where

$$B_1 = \begin{cases} X_1 + 2X_3 & (n \geq 4) \\ X_1 & (n=0). \end{cases}$$

In the case $m=n+1$, we have $r_1(V_{m,n}^j) \cong 0$.

By Lemma 5.2 and (5.8), we obtain the following.

(5.10) The Adams operations are given by the following formulae.

$$\begin{aligned}
(1) \quad \psi^k(r_1(X_3)) &= \begin{cases} k^{j/2} r_1(X_3) & (k \equiv 1 \pmod{4}) \\ -k^{j/2} r_1(X_3) & (k \equiv 3 \pmod{4}) \\ 0 & (k \equiv 0 \pmod{2}). \end{cases} \\
(2) \quad \psi^k(r_1(X_2)) &= \begin{cases} r_1(X_2) & (n \equiv 0 \pmod{4} \text{ and } k \equiv 1 \pmod{2}) \\ 0 & (n \equiv 0 \pmod{4} \text{ and } k \equiv 0 \pmod{2}). \end{cases} \\
(3) \quad \psi^k(r_1(2^{(n-2)/4} X_3 + X_2)) &= \begin{cases} r_1(2^{(n-2)/4} X_3 + X_2) & (n \equiv 2 \pmod{4} \text{ and } k \equiv 1 \pmod{2}) \\ 0 & (n \equiv 2 \pmod{4} \text{ and } k \equiv 0 \pmod{2}). \end{cases}
\end{aligned}$$

By Lemma 3.1, (5.6), (5.9) and (5.10), we obtain the results for the cases $j \equiv 2 \pmod{4}$, $n \equiv 0 \pmod{2}$ and $m+j \equiv 3, 4, 5, 6$ or $7 \pmod{8}$.

We now turn to the case $m+j \equiv 1 \pmod{8}$. Suppose that $m \geq n+3$, and consider the commutative diagram

$$\begin{array}{ccccccc}
& & 0 & & 0 & & \\
& & \uparrow & & \uparrow & & \\
& & V_{m-2,n}^j & \xrightarrow{r_1} & \widetilde{KO}(S^j(L_4^{m-2}/L_4^n)) & \longrightarrow & 0 \\
& & \uparrow & & \uparrow & & \\
& & V_{m,n}^j & \xrightarrow{r_1} & \widetilde{KO}(S^j(L_4^m/L_4^n)) & \xrightarrow{\partial} & \widetilde{KO}(S^{j+1}(L_4^m/L_4^n)) \longrightarrow 0 \\
& & \uparrow f & & \uparrow g & & \uparrow h \\
0 \rightarrow \tilde{K}(S^{m+j-1}) & \xrightarrow{r_2} & \widetilde{KO}(S^{m+j}) \oplus \widetilde{KO}(S^{m+j-1}) & \xrightarrow{\partial_1 \oplus \partial_2} & \widetilde{KO}(S^{m+j+1}) \oplus \widetilde{KO}(S^{m+j}) & \rightarrow & 0 \\
& & & & \uparrow & & \\
& & & & 0 & &
\end{array}$$

of exact sequences, where $\partial_1: \widetilde{KO}(S^{m+j}) \rightarrow \widetilde{KO}(S^{m+j+1})$ is an isomorphism. We denote the generators of $\widetilde{KO}(S^{m+j})$ and $\widetilde{KO}(S^{m+j+1})$ by ω_1 and ω_2 respectively. Since $\widetilde{KO}(S^{m+j}) \cong \mathbf{Z}/2$, Lemma 3.5 implies that $\tilde{K}(S^{m+j-1}) \cong \mathbf{Z}$ has a generator γ with

$$f(\gamma) = \begin{cases} 2^{(m-7)/4} x_3 + 2^{(m-3)/2} x_1 & (m \geq 7) \\ x_1 & (m=3) \end{cases}$$

and $r_2(\gamma) = 2\beta$, where β is a generator of the group $\widetilde{KO}(S^{m+j-1}) \cong \mathbf{Z}$. It follows from (5.9) that we have

$$\begin{aligned}
2g(\beta) &= r_1(f(\gamma)) \\
&= \begin{cases} r_1(2^{(m-7)/4} x_3 + 2^{(m-3)/2} x_1) & (m \geq 7) \\ r_1(x_1) & (m=3) \end{cases}
\end{aligned}$$

$$= \begin{cases} 2^{(m-n-3)/4} r_1(X_3) + 2^{(m-n-7)/4} r_1(X_2) & (n+j \equiv 2 \pmod{8}) \\ 2^{(m-n-3)/4} r_1(X_3) & (n+j \equiv 6 \pmod{8}) \\ 2^{(m-n-5)/4} r_1(X_3) & (n \equiv 2 \pmod{4}). \end{cases}$$

If $m \geq n+7$, we set $\alpha = g(\beta) - 2^{((m-7)/4) - [(n+2)/4]} r_1(X_3)$. Then we have $\partial(\alpha) = h(\omega_1)$, and

$$2\alpha = \begin{cases} 0 & (m \geq n+9) \\ r_1(X_2) & (m = n+7). \end{cases}$$

By (5.10) and the fact $4g(\beta) = 0$, we have

$$\begin{aligned} \psi^k(\alpha) &= k^{(m+j-1)/2} g(\beta) - \psi^k(2^{((m-7)/4) - [(n+2)/4]} r_1(X_3)) \\ &= \begin{cases} \alpha & (k: \text{odd}) \\ 0 & (k: \text{even}). \end{cases} \end{aligned}$$

According to [3, II], we have

$$\psi^k(\omega_i) = \begin{cases} \omega_i & (k: \text{odd}) \\ 0 & (k: \text{even}) \end{cases}$$

($i=1, 2$). If $m \geq n+9$, then the short exact sequence

$$0 \rightarrow r_1(V_{m,n}^j) \rightarrow \widetilde{KO}(S^j(L_4^m/L_4^n)) \rightarrow \widetilde{KO}(S^{j+1}(L_4^m/L_4^n)) \rightarrow 0$$

of ψ -groups splits. Hence

$$\widetilde{KO}(S^j(L_4^m/L_4^n)) \cong r_1(V_{m,n}^j) \oplus \mathbf{Z}/2 \oplus \mathbf{Z}/2$$

and

$$\mathcal{J}(S^j(L_4^m/L_4^n)) \cong \mathcal{J}''(r_1(V_{m,n}^j)) \oplus \mathbf{Z}/2 \oplus \mathbf{Z}/2.$$

If $m = n+7$, then we have

$$\widetilde{KO}(S^j(L_4^m/L_4^n)) = \langle r_1(V_{m,n}^j) \cup \{\alpha, g(\omega_1)\} \rangle = \langle \{r_1(X_3), \alpha, g(\omega_1)\} \rangle.$$

Since $\text{ord } \widetilde{KO}(S^j(L_4^m/L_4^n)) = 32$ by [15], $\text{ord } \langle r_1(X_3) \rangle = \text{ord } \langle \alpha \rangle = 4$ and $\text{ord } \langle g(\omega_1) \rangle = 2$, we have

$$\widetilde{KO}(S^j(L_4^m/L_4^n)) \cong \mathcal{J}(S^j(L_4^m/L_4^n)) \cong \mathbf{Z}/4 \oplus \mathbf{Z}/4 \oplus \mathbf{Z}/2.$$

If $m = n+5$ or $n+3$, then we have

$$\widetilde{KO}(S^j(L_4^m/L_4^n)) = \langle \{g(\beta), g(\omega_1)\} \rangle.$$

Since $\text{ord } \widetilde{KO}(S^j(L_4^m/L_4^n)) = 8$ by [15], $\text{ord } \langle g(\beta) \rangle = 4$ and $\text{ord } \langle g(\omega_1) \rangle = 2$, we have

$$\widetilde{KO}(S^j(L_4^m/L_4^n)) \cong \mathcal{J}(S^j(L_4^m/L_4^n)) \cong \mathbf{Z}/4 \oplus \mathbf{Z}/2.$$

Thus we obtain the results for the case $j \equiv 2 \pmod{4}$, $n \equiv 0 \pmod{2}$ and $m+j \equiv 1 \pmod{8}$.

The proof for the case $j \equiv 2 \pmod{4}$, $n \equiv 0 \pmod{2}$ and $m+j \equiv 0 \pmod{8}$ is similar to that for the above case, so we omit it.

Finally we consider the case $m+j \equiv 2 \pmod{8}$. Inspect the commutative diagram

$$\begin{array}{ccccccc}
 & & V_{m-2,n}^j & \xrightarrow{r_1} & \widetilde{KO}(S^j(L_4^{m-2}/L_4^n)) & & 0 \\
 & \uparrow & & & \uparrow & & \uparrow \\
 & V_{m,n}^j & \xrightarrow{r_1} & \widetilde{KO}(S^j(L_4^m/L_4^n)) & \xrightarrow{\partial} & \widetilde{KO}(S^{j+1}(L_4^m/L_4^n)) & \longrightarrow 0 \\
 & \uparrow & & \uparrow & & \uparrow & \\
 \tilde{K}(S^j(L_4^m/L_4^{m-2})) & \xrightarrow{r} & \widetilde{KO}(S^j(L_4^m/L_4^{m-2})) & \xrightarrow{\partial} & \widetilde{KO}(S^{j+1}(L_4^m/L_4^{m-2})) & \longrightarrow 0 & \\
 & & & & & & \uparrow \\
 & & & & & & 0
 \end{array}$$

of exact sequences. Since

$$\widetilde{KO}(S^j(L_4^m/L_4^{m-2})) \cong \widetilde{KO}(S^{j+m-2}L_4^2) \cong \mathbf{Z}/2 \oplus \mathbf{Z}/2$$

by Proposition 3.7, and

$$r(\tilde{K}(S^j(L_4^m/L_4^{m-2}))) \cong \widetilde{KO}(S^{j+1}(L_4^m/L_4^{m-2})) \cong \mathbf{Z}/2,$$

the short exact sequence

$$0 \rightarrow r(\tilde{K}(S^j(L_4^m/L_4^{m-2}))) \rightarrow \widetilde{KO}(S^j(L_4^m/L_4^{m-2})) \rightarrow \widetilde{KO}(S^{j+1}(L_4^m/L_4^{m-2})) \rightarrow 0$$

splits. The Adams operations on $\widetilde{KO}(S^j(L_4^m/L_4^{m-2}))$ or $\widetilde{KO}(S^{j+1}(L_4^m/L_4^{m-2}))$ are given by

$$\psi^k = \begin{cases} 1 & (k: \text{odd}) \\ 0 & (k: \text{even}). \end{cases}$$

Hence the short exact sequence

$$0 \rightarrow r_1(V_{m,n}^j) \rightarrow \widetilde{KO}(S^j(L_4^m/L_4^n)) \rightarrow \widetilde{KO}(S^{j+1}(L_4^m/L_4^n)) \rightarrow 0$$

of ψ -groups splits. Thus we obtain the result for the case $j \equiv 2 \pmod{4}$, $n \equiv 0 \pmod{2}$ and $m+j \equiv 2 \pmod{8}$.

Thus the proof for the case $j \equiv 2 \pmod{4}$ and $n \equiv 0 \pmod{2}$ is completed.

5.2. Proof for the case $n \equiv 3 \pmod{4}$. Consider the following commutative diagram, in which the row is exact.

$$\begin{array}{ccccccc}
0 & \longrightarrow & V_{m,n+1}^j & \xrightarrow{f_1} & \tilde{K}(S^j(L_4^m/L_4^n)) & \xrightarrow{f_2} & \tilde{K}(S^{n+j+1}) \longrightarrow 0 \\
& & \parallel & & \downarrow f_3 & & \\
& & V_{m,n+1}^j & \hookrightarrow & \tilde{K}(S^j(L_4^m)) & &
\end{array}$$

By Lemma 3.5, we can choose an element $x \in \tilde{K}(S^j(L_4^m/L_4^n))$ such that $f_2(x)$ generates the group $\tilde{K}(S^{n+j+1}) \cong \mathbb{Z}$ and

$$f_3(x) = 2^{(n-1)/2} x_1 + 2^{(n-3)/4} x_2 + 2^{(n-3)/2} x_3.$$

Applying the method used in the proof of Lemma 4.7 to x , we obtain the following result by Lemma 5.2.

(5.11) *The Adams operations are given by*

$$\psi^k(x) = \begin{cases} k^u x + ((k^{j/2} - k^u)/4) f_1(4f_3(x)) & (k \equiv 1 \pmod{2}) \\ k^u x - (k^u/4) f_1(4f_3(x)) & (k \equiv 0 \pmod{4}) \\ k^u x + f_1(k^{j/2} 2^{(n-3)/4} X_2 - k^u f_3(x)) & (k \equiv 2 \pmod{4}), \end{cases}$$

where $u = (n+j+1)/2$.

This implies that $c \circ r(x) = (1 + \psi^{-1})(x) = 0$. By (5.8), we have

$$r_1(4f_3(x)) = r_1((1 - 2^{(n+1)/4})X_1 + 2X_2 + 2^{(n+5)/4}X_3) = r_1(X_2).$$

Thus we obtain

$$(5.12) \quad (1) \quad 2r(x) = r(c \circ r(x)) = 0.$$

$$(2) \quad \psi^k(r(x)) = k^{(n+j+1)/2} r(x) = \begin{cases} r(x) & (k \equiv 1 \pmod{2}) \\ 0 & (k \equiv 0 \pmod{2}). \end{cases}$$

Inspect the following commutative diagram

$$\begin{array}{ccccccc}
0 & \longrightarrow & V_{m,n+1}^j & \xrightarrow{f_1} & \tilde{K}(S^j(L_4^m/L_4^n)) & \xrightarrow{f_2} & \tilde{K}(S^{n+j+1}) \longrightarrow 0 \\
& & \downarrow r_1 & & \downarrow r & & \downarrow r \\
0 & \longrightarrow & \widetilde{KO}(S^j(L_4^m/L_4^{n+1})) & \longrightarrow & \widetilde{KO}(S^j(L_4^m/L_4^n)) & \longrightarrow & \widetilde{KO}(S^{n+j+1}) \longrightarrow 0 \\
& & & & & & \downarrow \\
& & & & & & 0
\end{array}$$

of exact sequences. Since

$$\widetilde{KO}(S^{n+j+1}) \cong \begin{cases} \mathbb{Z}/2 & (n+j \equiv 1 \pmod{8}) \\ 0 & (n+j \equiv 5 \pmod{8}), \end{cases}$$

using (5.12) we see that the short exact sequence

$$0 \rightarrow \widetilde{KO}(S^j(L_4^m/L_4^{n+1})) \rightarrow \widetilde{KO}(S^j(L_4^m/L_4^n)) \rightarrow \widetilde{KO}(S^{n+j+1}) \rightarrow 0$$

of ψ -groups splits. This implies that

$$\widetilde{KO}(S^j(L_4^m/L_4^n)) \cong \widetilde{KO}(S^j(L_4^m/L_4^{n+1})) \oplus \widetilde{KO}(S^{n+j+1})$$

and

$$\widetilde{J}(S^j(L_4^m/L_4^n)) \cong \widetilde{J}(S^j(L_4^m/L_4^{n+1})) \oplus \widetilde{J}(S^{n+j+1}).$$

Thus, results of the case $j \equiv 2 \pmod{4}$ and $n \equiv 3 \pmod{4}$ follow from those of the case $j \equiv 2 \pmod{4}$ and $n \equiv 0 \pmod{4}$.

5.3. Proof for the case $n \equiv 1 \pmod{4}$. Consider the following commutative diagram, in which the row is exact.

$$\begin{array}{ccccccc} 0 & \longrightarrow & V_{m,n+1}^j & \xrightarrow{f_1} & \tilde{K}(S^j(L_4^m/L_4^n)) & \xrightarrow{f_2} & \tilde{K}(S^{n+j+1}) \longrightarrow 0 \\ & & \parallel & & \downarrow f & & \\ & & V_{m,n+1}^j & \hookrightarrow & \tilde{K}(S^j(L_4^m)) & & \end{array}$$

By Lemma 3.5, we can choose an element $x \in \tilde{K}(S^j(L_4^m/L_4^n))$ such that $f_2(x)$ generates the group $\tilde{K}(S^{n+j+1}) \cong \mathbb{Z}$ and

$$f_3(x) = \begin{cases} 2^{(n-5)/4}x_3 + 2^{(n-1)/2}x_1 & (n \geq 5) \\ x_1 & (n = 1). \end{cases}$$

Applying the method used in the proof of Lemma 4.7 to x , we obtain the following result by Lemma 5.2.

(5.13) *The Adams operations are given by*

$$\psi^k(x) = \begin{cases} k^u x + ((k^{j/2} - k^u)/4)f_1(4f_3(x)) & (k \equiv 1 \pmod{4}) \\ k^u x - ((k^{j/2} + k^u)/4)f_1(4f_3(x)) + k^{j/2}f_1(2^{(n-5)/4}(2X_2 + 2X_3 - X_1) + X_1 - X_2) & (k \equiv 3 \pmod{4} \text{ and } n \geq 5) \\ k^u x - ((k^{j/2} + k^u)/4)f_1(4f_3(x)) + k^{j/2}f_1(X_2 + X_3) & (k \equiv 3 \pmod{4} \text{ and } n = 1) \\ k^u x - (k^u/4)f_1(4f_3(x)) + (k^{j/2}/2)2^{(n-1)/4}f_1(2X_2 - X_1) & (k \equiv 2 \pmod{4}) \\ k^u x - (k^u/4)f_1(4f_3(x)) & (k \equiv 0 \pmod{4}), \end{cases}$$

where $u = (n+j+1)/2$.

Inspect the following commutative diagram

$$\begin{array}{ccccccc}
0 & & 0 & & \widetilde{KO}(S^{n+j+2}) & & 0 \\
\downarrow & & \downarrow & & \downarrow & & \downarrow \\
VO_{m,n+1}^{j+2} & \xrightarrow{I^{-1} \circ c} & V_{m,n+1}^j & \xrightarrow{r_1} & \widetilde{KO}(S^j(L_4^m/L_4^{n+1})) & \xrightarrow{\partial} & \widetilde{KO}(S^{j+1}(L_4^m/L_4^{n+1})) \\
\downarrow & & \downarrow f_1 & & \downarrow g_1 & & \downarrow h_1 \\
VO_{m,n}^{j+2} & \xrightarrow{\quad} & \widetilde{K}(S^j(L_4^m/L_4^n)) & \xrightarrow{r_2} & \widetilde{KO}(S^j(L_4^m/L_4^n)) & \xrightarrow{\partial} & \widetilde{KO}(S^{j+1}(L_4^m/L_4^n)) \\
\downarrow & & \downarrow f_2 & & \downarrow g_2 & & \downarrow h_2 \\
\widetilde{KO}(S^{n+j+3}) & \xrightarrow{I^{-1} \circ c} & \widetilde{K}(S^{n+j+1}) & \xrightarrow{r} & \widetilde{KO}(S^{n+j+1}) & \xrightarrow{\partial} & \widetilde{KO}(S^{n+j+2}) \longrightarrow 0 \\
\downarrow & & \downarrow & & \downarrow & & \\
0 & & 0 & & \widetilde{KO}(S^{j-1}(L_4^m/L_4^{n+1})) & &
\end{array}$$

of exact sequences. By Proposition 3.7, we have

$$VO_{m,n}^{j+2} = \begin{cases} \langle \{2^{(n+1)/2}y_1, 2^{(n-5)/4}y_2 + 2^{(n-1)/2}y_1\} \rangle & (j \equiv 2 \pmod{8} \text{ and } n \geq 5) \\ \langle \{2^{(n+1)/2}y_1, 2^{(n-1)/4}y_2 + 2^{(n-1)/2}y_1\} \rangle & (j \equiv 6 \pmod{8} \text{ and } n \geq 5) \\ \langle \{y_1, 2y_2\} \rangle & (j \equiv 6 \pmod{8} \text{ and } n = 1) \\ \langle \{y_1, y_2\} \rangle & (j \equiv 2 \pmod{8} \text{ and } n = 1). \end{cases}$$

Using Lemma 5.7, we obtain

$$(5.14) \quad \text{Ker } r_2 = \langle \{f_1(2X_2 - X_1), f_1(2^{(n-1)/4}X_3 + X_2)\} \rangle.$$

If $m \geq n+3$, then we have

$$\begin{aligned} \text{Coker } g_2 &\cong \widetilde{KO}(S^{n+j+2}) \\ &\cong \begin{cases} \mathbf{Z}/2 & (n+j \equiv 7 \pmod{8}) \\ 0 & (n+j \equiv 3 \pmod{8}), \end{cases} \end{aligned}$$

and hence

$$\begin{aligned} r(\widetilde{K}(S^{n+j+1})) &= g_2(\widetilde{KO}(S^j(L_4^m/L_4^n))) \\ &= \begin{cases} 2\widetilde{KO}(S^{n+j+1}) & (n+j \equiv 7 \pmod{8}) \\ \widetilde{KO}(S^{n+j+1}) & (n+j \equiv 3 \pmod{8}). \end{cases} \end{aligned}$$

Since h_1 is a monomorphism, we have $\text{Ker } g_1 \subset r_1(V_{m,n+1}^j)$. Thus we obtain a split short exact sequence

$$0 \rightarrow \widetilde{KO}(S^j(L_4^m/L_4^{n+1}))/\text{Ker } g_1 \xrightarrow{\bar{g}_1} \widetilde{KO}(S^j(L_4^m/L_4^n)) \xrightarrow{\bar{g}_2} \mathbf{Z} \rightarrow 0,$$

where

$$\text{Ker } g_1 = \langle r_1(2^{(n-1)/4}X_3 + X_2) \rangle.$$

By (5.9), we obtain

$$(5.15) \quad \text{If } m \geq n+3, \text{ then we have}$$

$$r_1(V_{m,n+1}^j)/\text{Ker } g_1 \cong \langle \{X_1, X_2, X_3\} \rangle / \langle \{A_1, B_2, B_3\} \rangle,$$

where $A_1 = 2X_2 - X_1$, $B_2 = 2^{(n-1)/4}X_3 + X_2$ and $B_3 = 2^{[(m-n-1)/4]}X_3$.

Thus the group $\widetilde{KO}(S^j(L_4^m/L_4^n))$ is determined by using results of the case $j \equiv 2 \pmod{4}$ and $n \equiv 2 \pmod{4}$. In order to determine the group $\widetilde{J}(S^j(L_4^m/L_4^n))$, we use the following fact which is obtained from (5.13) and (5.14).

(5.16) *The Adams operations are given by*

$$\psi^k(r_2(x)) = \begin{cases} k^{(n+j+1)/2}r_2(x) + ((k^{j/2} - k^{(n+j+1)/2})/4)r_2(f_1(4f_3(x))) & (k \equiv 1 \pmod{4}) \\ k^{(n+j+1)/2}r_2(x) - ((k^{j/2} + k^{(n+j+1)/2})/4)r_2(f_1(4f_3(x))) & (k \equiv 3 \pmod{4}) \\ k^{(n+j+1)/2}r_2(x) - (k^{(n+j+1)/2}/4)r_2(f_1(4f_3(x))) & (k \equiv 0 \pmod{2}) \end{cases}$$

Set $U = \sum_{k: \text{odd}} (\psi^k - 1) \widetilde{KO}(S^j(L_4^m/L_4^{n+1}))$. By Lemma 3.1 and (5.9), we have $U = \langle 4r_1(X_3) \rangle$. If $k \equiv \varepsilon \pmod{4}$ ($\varepsilon = \pm 1$), then we have

$$\begin{aligned} & ((\varepsilon k^{j/2} - k^{(n+j+1)/2})/4)r_2(f_1(4f_3(x))) \\ & \equiv ((\varepsilon k^{j/2} - k^{(n+j+1)/2})/2)g_1(r_1(X_3)) \quad (\text{mod } g_1(U)) \\ & \equiv ((k - \varepsilon)/2)g_1(r_1(X_3)) \quad (\text{mod } g_1(U)) \\ & \equiv ((k^{(n+j+1)/2} - 1)/2^{v_2(n+j+1)})g_1(r_1(X_3)) \quad (\text{mod } g_1(U)). \end{aligned}$$

Thus we have $\widetilde{J}(S^j(L_4^m/L_4^n)) \cong \widetilde{KO}(S^j(L_4^m/L_4^n))/U_1$, where U_1 is the subgroup of $\widetilde{KO}(S^j(L_4^m/L_4^n))$ generated by $4g_1(r_1(X_3))$ and $m((n+j+1)/2)r_2(x) - 2g_1(r_1(X_3))$.

Suppose $m+j \equiv 3, 4, 5, 6$ or $7 \pmod{8}$. Then we have

$$\widetilde{J}(S^j(L_4^m/L_4^n)) \cong \langle \{r_2(x), g_1(r_1(X_3))\} \rangle / \langle \{A_1, A_2\} \rangle,$$

where $A_1 = m((n+j+1)/2)r_2(x) - 2g_1(r_1(X_3))$ and

$$A_2 = \begin{cases} 4g_1(r_1(X_3)) & (m \geq n+9) \\ 2g_1(r_1(X_3)) & (n+8 \geq m \geq n+5) \\ g_1(r_1(X_3)) & (n+4 \geq m \equiv n+3). \end{cases}$$

Thus we obtain the results of the cases $j \equiv 2 \pmod{4}$, $n \equiv 1 \pmod{4}$ and $m+j \equiv 3, 4, 5, 6$ or $7 \pmod{8}$.

Since $\text{Ker } g_1 = r_1(\langle 2^{(n-1)/4}X_3 + X_2 \rangle)$, the rest of the proof is similar to that for the case $j \equiv 2 \pmod{4}$ and $n \equiv 2 \pmod{4}$.

References

- [1] J.F. Adams: *Vector fields on spheres*, Ann. of Math. **75** (1962), 603–632.
- [2] J.F. Adams: *On the groups $J(X)$* -I, Topology **2** (1963), 181–195.
- [3] J.F. Adams: *On the groups $J(X)$* -II, -III, Topology **3** (1965), 137–171, 193–222.

- [4] J.F. Adams and G. Walker: *On complex Stiefel manifolds*, Proc. Camb. Phil. Soc. **61** (1965), 81–103.
- [5] D.W. Anderson: *A new cohomology theory*, Thesis, Univ. of California, Berkeley, 1964.
- [6] R. Bott: *Lectures on $K(X)$* , Benjamin, 1969.
- [7] K. Fujii, T. Kobayashi and M. Sugawara: *Stable homotopy types of stunted lens spaces*, Mem. Fac. Sci. Kochi Univ. (Math.) **3** (1982), 21–27.
- [8] T. Kobayashi: *KO-cohomology of the lens space mod 4*, Mem. Fac. Sci. Kochi Univ. (Math.) **6** (1985), 45–63.
- [9] T. Kobayashi and M. Sugawara: *On stable homotopy types of stunted lens spaces*, Hiroshima Math. J. **1** (1971), 287–304.
- [10] T. Kobayashi and M. Sugawara: *K_{Λ} -rings of the lens spaces $L^n(4)$* , Hiroshima Math. J. **1** (1971), 253–271.
- [11] S. Kôno and A. Tamamura: *On J -groups of $S^l(RP(t-l)/RP(n-l))$* , Math. J. Okayama Univ. **24** (1982), 45–51.
- [12] S. Kôno and A. Tamamura: *J -groups of the suspensions of the stunted lens spaces mod p* , Osaka J. Math. **24** (1987), 481–498.
- [13] N. Mahammed: *A propos de la K -théorie des espaces lenticulaires*, C.R. Acad. Sc. Paris **271** (1970), 639–642.
- [14] D. Quillen: *The Adams conjecture*, Topology **10** (1971), 67–80.
- [15] A. Tamamura and S. Kôno: *On the KO-cohomologies of the stunted lens spaces*, Math. J. Okayama Univ. **29** (1987), 233–244.

Susumu Kôno
 Department of Mathematics
 Osaka University
 Toyonaka, Osaka 560, Japan

Akie Tamamura
 Department of Applied Mathematics
 Okayama University of Science
 Ridai, Okayama 700, Japan

