

Title	TROPICALIZATION OF 1-TACNODAL CURVES ON TORIC SURFACES
Author(s)	Takahashi, Takuhiro
Citation	Osaka Journal of Mathematics. 57(2) P.451-P.491
Issue Date	2020-04
Text Version	publisher
URL	https://doi.org/10.18910/75921
DOI	10.18910/75921
rights	
Note	

Osaka University Knowledge Archive : OUKA

<https://ir.library.osaka-u.ac.jp/>

Osaka University

TROPICALIZATION OF 1-TACNODAL CURVES ON TORIC SURFACES

TAKUHIRO TAKAHASHI

(Received October 25, 2017, revised January 28, 2019)

Abstract

A degeneration of a singular curve on a toric surface, called a tropicalization, was constructed by E. Shustin. He classified the degeneration of 1-cuspidal curves using polyhedral complexes called tropical curves. In this paper, we define a tropical version of a 1-tacnodal curve, that is, a curve having exactly one singular point whose topological type is A_3 , and by applying the tropicalization method, we classify tropical curves which correspond to 1-tacnodal curves.

Contents

1. Introduction	451
2. Preliminaries	453
3. Tropical 1-tacnodal curves	459
4. Main Result	477
References	490

1. Introduction

Tropical geometry is a modern study area on a polyhedral complex, which can be obtained as the non-linear locus of a polynomial over the max-plus algebra. Among previous studies on tropical geometry, the most famous result is an application to the enumerative geometry on toric surfaces by G. Mikhalkin [5]. T. Nishinou and B. Siebert [6] also showed that the enumerative problem on toric varieties equals to the enumeration of a certain type of tropical curves. These results are obtained by connecting tropical geometry with degeneration of nodal curves.

In order to apply tropical geometry to general singular curves, E. Shustin [8] presented a degeneration of a curve, called a *tropicalization*, and showed that the tropicalization of a curve which has only one singular point whose topological type is A_2 (he called such a curve a *1-cuspidal curve* for simplicity) is related to a certain tropical curve, called a tropical 1-cuspidal curve. Furthermore, using the theory of patchworking, he showed that the enumeration of 1-cuspidal curves reduced into that of the tropical 1-cuspidal curves.

In this paper, we apply the tropicalization method to 1-tacnodal curves, that is, curves which have exactly one singular point whose topological type is A_3 , on a toric surface, and

classify them using tropical curves.

To state our result, we prepare some terminology in tropical geometry. Let F be a polynomial in two variables over the field of convergent Puiseux series over \mathbb{C} , denoted by $K := \mathbb{C}\{\{t\}\}$. Then we can define a valuation $\text{val} : K^* := K \setminus \{0\} \rightarrow \mathbb{R}$ as follows. For a given element $b(t) \in K^*$, take the minimal exponent q of $b(t)$ in t , then define $\text{val}(b(t)) := -q$. We set

$$\text{Val} : (K^*)^2 \rightarrow \mathbb{R}^2; (z, w) \mapsto (\text{val}(z), \text{val}(w)).$$

We call the closure

$$T_F := \text{Closure}(\text{Val}(\{F = 0\}) \cap (K^*)^2) \subset \mathbb{R}^2$$

of the curve defined by F in $(K^*)^2$ the *tropical amoeba* defined by F .

Each tropical amoeba T has a positive integer $\text{rk}(T)$ called a *rank*, which, roughly speaking, is the dimension of the space of tropical curves which are combinatorially same as T . The formal definition of the rank will be given in Subsection 2.1. We will also give the definition of a *tropical 1-tacnodal curve* in Definition 3.1 as a tropical analogy of a classical 1-tacnodal curve.

The following statement is the main result in this paper.

Theorem 4.1. *Let $F \in K[z, w]$ be a polynomial which defines an irreducible 1-tacnodal curve. If the rank of the tropical amoeba T_F defined by F is more than or equal to the number of the lattice points of the Newton polytope of F minus four, then T_F is a tropical 1-tacnodal curve.*

In [8], Shustin proved the 1-cuspidal version of this theorem and the statement that “for each tropical 1-cuspidal curve, we can calculate the number of classical curves degenerated into the 1-cuspidal curve by using the patchworking method”, which means that the enumeration of 1-cuspidal curves on toric surfaces can be carried out by using tropical 1-cuspidal curves. The original aim of this paper is to enumerate 1-tacnodal curves on toric surfaces by the same method. However, it does not work unlike the studies for nodal and 1-cuspidal curves because the criterion of patchworking developed by Shustin [7, 8] cannot be used in this case. We will discuss this in Remark 4.12 below. Note that, in [8], the rank of tropical amoeba of A_2 -curve passing through “good” generic points, whose number is equal to the dimension of the space of A_2 -curves, is more than or equal to the number of lattice points of the Newton polytope minus three. In our case, it becomes “minus four” since the singularity is A_3 instead of A_2 (See Corollary 2.5 of this paper). This setting seems to be natural though there is no concrete observation about it.

We organize this paper as follows. In Subsection 2.1, we define some basic terminology on tropical geometry such as the dual subdivision and the rank of a tropical curve, and introduce a lemma on the rank of the tropical curves proved by E. Shustin. In Subsection 2.2, we consider a necessary and sufficient condition for a complex curve to have a tacnode, and estimate the dimension of the space of the 1-tacnodal curves on a toric surface. In Subsection 2.3, we summarize the tropicalization and its refinement. In Section 3, we define tropical 1-tacnodal curves and discuss polytopes appearing in their dual subdivisions. We also study a reduced curve associated with a tropical 1-tacnodal curve. In Subsections 4.1 and 4.2, before the proof of Theorem 4.1, we prepare some definition and lemmata on relation between

singular curves and their Newton polytopes. The proof of Theorem 4.1 is carried out from Subsections 4.3 to 4.5.

2. Preliminaries

A set in \mathbb{R}^2 is a (*lattice*) *polyhedron* if it is the intersection of a finite number of half-spaces in \mathbb{R}^2 whose vertices are contained in the lattice $\mathbb{Z}^2 \subset \mathbb{R}^2$. A set is a *polytope* in \mathbb{R}^2 if it is a compact polyhedron, that is, the convex hull of a finite number of lattice points. We call a facet of a polytope an *edge*. Similarly, we call the 0-dimensional sub-polytope obtained as the corner of a polytope a *vertex*. The *boundary* $\partial\Delta$ is the union of all facets of Δ . The *interior* $\text{Int}\Delta$ is defined by $\Delta \setminus \partial\Delta$.

A polytope is said to be *parallel* if the opposite edges have the same directional vector (up to orientation) and the same lattice length. A polytope is called an *m-gon* if the number of its edges is *m*.

Let $\Delta \subset \mathbb{R}^2$ be a polytope. We denote the set of lattice points in Δ , $\text{Int}\Delta$ and $\partial\Delta$ as $\Delta_{\mathbb{Z}}$, $\text{Int}\Delta_{\mathbb{Z}}$ and $\partial\Delta_{\mathbb{Z}}$, respectively. That is,

$$\Delta_{\mathbb{Z}} := \Delta \cap \mathbb{Z}^2, \quad \text{Int}\Delta_{\mathbb{Z}} := \text{Int}\Delta \cap \mathbb{Z}^2, \quad \partial\Delta_{\mathbb{Z}} := \partial\Delta \cap \mathbb{Z}^2.$$

For a polytope $\Delta \subset \mathbb{R}^2$, we can construct a polarized toric surface associated with Δ over \mathbb{C} , denoted by $(X(\Delta), D(\Delta))$, where $D(\Delta)$ is the polarization on $X(\Delta)$ associated with Δ .

2.1. Basics of tropical plane curves. Throughout this paper, $K := \mathbb{C}\{\{t\}\}$ represents the field of convergent Puiseux series over \mathbb{C} . The field K admits a non-Archimedean valuation

$$\text{val} : K^* \rightarrow \mathbb{Q}; \sum_{k=k_0}^{\infty} b_k t^{\frac{k}{N}} \mapsto -\frac{k_0}{N},$$

where $b_{k_0} \neq 0$. For a polynomial

$$F(z, w) = \sum_{(i,j) \in \Delta_{\mathbb{Z}}} c_{ij} z^i w^j \in K[z, w],$$

the sets

$$\text{Supp}(F) := \{(i, j) \in \mathbb{R}^2; c_{ij} \neq 0\} \quad \text{and} \quad N_F := \text{Conv}(\text{Supp}(F)) \subset \mathbb{R}^2$$

are called the *support* of F and the *Newton polytope* of F , respectively, where $\text{Conv}(A)$ is the convex hull of A in \mathbb{R}^2 . In this paper, we always assume that the dimension of any polytope is 2. Then the map Val is defined by

$$\text{Val} : (K^*)^2 \rightarrow \mathbb{R}^2; (z, w) \mapsto (\text{val}(z), \text{val}(w)).$$

The set T_F is defined by

$$T_F := \text{Closure}(\text{Val}(\{p \in (K^*)^2; F(p) = 0\})) \subset \mathbb{R}^2,$$

where $\text{Closure}(A)$ is the closure of A with usual topology on \mathbb{R}^2 . The set T_F is called the *tropical amoeba* defined by F .

On the other hand, for F , the *tropical polynomial* τ_F is defined by

$$\tau_F(x, y) := \max\{\text{val}(c_{ij}) + ix + jy; (i, j) \in \text{Supp}(F)\}$$

over the max-plus algebra. The non-linear locus of a polynomial over the max-plus algebra in two variables is called a *tropical plane curve*, which is a 1-dimensional polyhedral complex in \mathbb{R}^2 . It is known as Kapranov's Theorem [3] that the tropical amoeba T_F coincides with the tropical plane curve of the tropical polynomial τ_F . Hence, T_F has the structure of a 1-dimensional polyhedral complex in \mathbb{R}^2 . A 1-simplex and a 0-simplex of T_F are called an *edge* and a *vertex* of T_F , respectively.

An edge E of T_F corresponds to the intersection of two linearity domains of τ_F . If one of the linearity domain is defined by a term $a_{ij} + ix + jy$ of τ_F and the other is defined by a term $a_{i'j'} + i'x + j'y$ of τ_F , then the *weight* $w(E)$ of the edge E is defined as the greatest common divisor of $i - i'$ and $j - j'$.

Now we introduce a subdivision of the Newton polytope N_F which is dual to the tropical amoeba T_F . Let $\nu_F : N_F \rightarrow \mathbb{R}$ be the discrete Legendre transform of τ_F , which is a continuous concave PL-function (see, for example, [1, Chapter 1.5]). Then we obtain a subdivision of N_F consisting of the following three kinds of polytopes from ν_F :

- linearity domains of ν_F : $\Delta_1, \dots, \Delta_N$,
- 1-dimensional polytopes: $\sigma_{ij} := \Delta_i \cap \Delta_j \neq \emptyset, \neq \{\text{pt}\}$,
- 0-dimensional polytopes: $\Delta_{i_1} \cap \Delta_{i_2} \cap \Delta_{i_3} \neq \emptyset$.

These polytopes give a subdivision of N_F , which we denote by S_F . We call a 1-dimensional and a 0-dimensional sub-polytope of N_F contained in S_F an *edge* and a *vertex*, respectively.

The following claim is in [5, Proposition 3.11].

Theorem 2.1 (Duality Theorem). *There exists a correspondence between a tropical curve T_F with the weight $w(E)$ on each edge $E \subset T_F$ and the corresponding subdivision S_F of N_F in the following sense:*

- (1) *the components of $\mathbb{R}^2 \setminus T_F$ are in 1-to-1 correspondence with the vertices of the subdivision S_F ,*
- (2) *the edges of T_F are in 1-to-1 correspondence with the edges of S_F so that an edge $E \subset T_F$ is dual to an orthogonal edge of S_F having the lattice length equal to $w(E)$*
- (3) *the vertices of T_F are in 1-to-1 correspondence with the polytopes $\Delta_1, \dots, \Delta_N$ of S_F so that the valency of a vertex of T_F is equal to the number of sides of the corresponding polytope.*

We call the set S_F the *dual subdivision* of T_F . By Theorem 2.1, we can study any plane tropical curve by using the dual subdivision of the corresponding Newton polytope.

Next, we discuss the dimension of the space of tropical curves. For a given polytope Δ , let $\mathfrak{T}(\Delta)$ denote the set of tropical curves which are defined by polynomials in two variables over the max-plus algebra with Newton polytope Δ . Let S be the dual subdivision of $T \in \mathfrak{T}(\Delta)$ and define the *rank* of the tropical curve T (or of S) as

$$\text{rk}(T) := \text{rk}(S) := \dim\{T' \in \mathfrak{T}(\Delta); S = S'\},$$

where S' is the dual subdivision of T' . By [5, Lemma 3.14], the set $\{T' \in \mathfrak{T}(\Delta); S = S'\}$ is a polyhedron in \mathbb{R}^M for some positive integer M . Thus, the definition of the rank is well-defined.

Let $\Delta_1, \dots, \Delta_N$ be the 2-dimensional polytopes of S . According to [8], we define the *expected rank* of the tropical curve T (or of S) as

$$\text{rk}_{\text{exp}}(T) := \text{rk}_{\text{exp}}(S) := \#V(S) - 1 - \sum_{k=1}^N (\#V(\Delta_k) - 3),$$

where $V(S)$ is the set of vertices of S and $V(\Delta_k)$ is the set of vertices of Δ_k .

DEFINITION 2.2. A lattice subdivision of a polytope is a *TP-subdivision* if the subdivision consists of only triangles and parallelograms.

We remark that, this definition is same as the definition of the *nodal subdivision* in [8, Subsection 3.1] except the condition on the boundary $\partial\Delta$.

For any subdivision S , we denote the number of ℓ -gons and the number of parallel $(2m)$ -gons contained in S as N_ℓ and N'_{2m} , respectively.

The following statement is in [8, Lemma 2.2].

Lemma 2.3 (Shustin [8]). *The difference*

$$d(T) := \text{rk}(T) - \text{rk}_{\text{exp}}(T)$$

of a tropical curve T satisfies $d(T) \geq 0$. Moreover, for the dual subdivision S of T , the difference $d(T)$ satisfies

- $d(T) = 0$ if S is a TP-subdivision and
- $0 \leq 2d(T) \leq \mathcal{N}_S$ otherwise,

where

$$\begin{aligned} \mathcal{N}_S &:= \sum_{m \geq 2} ((2m - 3)N_{2m} - N'_{2m}) + \sum_{m \geq 2} (2m - 2)N_{2m+1} - 1 \\ &= \sum_{\ell \geq 3} (\ell - 3)N_\ell - \sum_{m \geq 2} N'_{2m} - 1. \end{aligned}$$

2.2. Some remarks on 1-tacnodal curves. In this paper, a curve on a projective surface is called a *1-tacnodal curve* if the curve has exactly one singular point at a smooth point of the surface whose topological type is A_3 . The term ‘‘tacnode’’ means A_3 -singularity. In this subsection we prepare some lemmata related to 1-tacnodal curves.

For a polynomial f and $p \in \mathbb{C}^2$, we use the notations $f_x(p) = \frac{\partial f}{\partial x}(p)$, $f_y(p) = \frac{\partial f}{\partial y}(p)$ and so on. We set $\text{Hess}(f)(p) = f_{xx}(p)f_{yy}(p) - f_{xy}(p)^2$ and

$$\begin{aligned} K(f)(p) &:= -f_{xy}(p)^3 f_{xxx}(p) + 3f_{xx}(p)f_{xy}(p)^2 f_{xxy}(p) \\ &\quad - 3f_{xx}(p)^2 f_{xy}(p)f_{xyy}(p) + f_{xx}(p)^3 f_{yyy}(p). \end{aligned}$$

Lemma 2.4. *Suppose that a polynomial $f \in \mathbb{C}[x, y]$ satisfies $f_{xx}(p) \neq 0$. Then the curve $\{f = 0\} \subset \mathbb{C}^2$ has a tacnode at p if and only if f satisfies*

- (1) $f(p) = f_x(p) = f_y(p) = 0$,
- (2) $\text{Hess}(f)(p) = 0$,
- (3) $K(f)(p) = 0$,
- (4) $a_{12}(p)^2 - 4f_{xx}(p)a_{04}(p) \neq 0$,

where

$$a_{12}(p) := f_{xy}(p)^2 f_{xxx}(p) - 2f_{xx}(p)f_{xy}(p)f_{xxy}(p) + f_{xx}(p)^2 f_{xyy}(p),$$

$$a_{04}(p) := f_{xy}(p)^4 f_{xxx}(p) - 4f_{xx}(p)f_{xy}(p)^3 f_{xxxy}(p) \\ + 6f_{xx}(p)^2 f_{xy}(p)^2 f_{xxyy}(p) - 4f_{xx}(p)^3 f_{xy}(p)f_{xyyy}(p) + f_{xx}(p)^4 f_{yyyy}(p).$$

Proof. For simplicity, we assume that p is the origin $(0, 0)$ of \mathbb{C}^2 . First, if the origin is a singular point then we can represent f as

$$f = Ax^2 + Bxy + Cy^2 + (\text{higher terms}),$$

where $(A, B, C) = (f_{xx}(0, 0)/2, f_{xy}(0, 0), f_{yy}(0, 0)/2)$. If $\text{Hess}(f)(0, 0) \neq 0$, then the origin is an A_1 -singularity of $\{f = 0\}$. Therefore $\text{Hess}(f)(0, 0) = 0$ for the origin to be an A_3 -singularity. Then we can rewrite f as

$$f = \frac{1}{4A}(2Ax + By)^2 + (\text{higher terms}).$$

The tangent line of $\{f = 0\}$ at the origin is defined by

$$f_{xx}(0, 0)x + f_{xy}(0, 0)y = 0.$$

Now we define new coordinates (u, v) as

$$\begin{pmatrix} u \\ v \end{pmatrix} = \begin{pmatrix} f_{xx}(0, 0) & f_{xy}(0, 0) \\ 0 & 1 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix}$$

and set

$$\hat{f}(u, v) := f(x(u, v), y(u, v)).$$

Note that the condition $f(0, 0) = f_x(0, 0) = f_y(0, 0) = \text{Hess}(f)(0, 0) = 0$ is equivalent to $\hat{f}(0, 0) = \hat{f}_u(0, 0) = \hat{f}_v(0, 0) = \text{Hess}(\hat{f})(0, 0) = 0$.

By direct computation, we obtain the equalities:

$$\begin{aligned} \hat{f}_{uu}(0, 0) &= \frac{1}{f_{xx}(0, 0)}, \\ \hat{f}_{uv}(0, 0) &= 0, \\ \hat{f}_{vv}(0, 0) &= \frac{1}{f_{xx}(0, 0)} \text{Hess}(f)(0, 0), \\ (*) \quad \hat{f}_{uuv}(0, 0) &= \frac{1}{f_{xx}(0, 0)^3} a_{12}(0, 0), \\ \hat{f}_{uvv}(0, 0) &= \frac{1}{f_{xx}(0, 0)^3} K(f)(0, 0), \\ \hat{f}_{vvv}(0, 0) &= \frac{1}{f_{xx}(0, 0)^4} a_{04}(0, 0). \end{aligned}$$

From the properties of the Newton diagram of a plane curve singularity [4], the condition that the singularity at the origin is A_3 can be rewritten as

$$\hat{f}_{uv}(0, 0) = \hat{f}_{vv}(0, 0) = \hat{f}_{uvv}(0, 0) = 0, \quad \hat{f}_{uu}(0, 0) \neq 0$$

and

$$\hat{f}_{uuv}(0, 0)^2 - 4\hat{f}_{uu}(0, 0)\hat{f}_{vvv}(0, 0) \neq 0$$

on the new coordinate system. By (*), these conditions coincide with the conditions in the

assertion. □

For $\mu = 1, 3$, let $U(\Delta, A_\mu)$ denote a locally closed subvariety in the complete linear system $|D(\Delta)|$ of $D(\Delta)$ which parametrizes the set of curves having exactly one singular point, whose topological type is A_μ . Let $V(\Delta, A_\mu)$ be the closure of $U(\Delta, A_\mu)$ in $|D(\Delta)|$.

Corollary 2.5. *If $V(\Delta, A_3)$ is non-empty then $\dim V(\Delta, A_3) \geq \#\Delta_{\mathbb{Z}} - 4$.*

Proof. For $\mu = 1, 3$, we set

$$\Sigma(\Delta, A_\mu) := \{(C, p); p \text{ is a singular point of } C\} \subset U(\Delta, A_\mu) \times X(\Delta) \subset |D(\Delta)| \times X(\Delta).$$

For a curve $C \in V(\Delta, A_\mu)$, we choose a local coordinate system (x, y) of $X(\Delta)$ around the singular point $p = (x_0, y_0) \in C$. Let f be a defining polynomial of C . By Lemma 2.4, $\Sigma(\Delta, A_3)$ is locally defined by

$$(**) \quad f(x_0, y_0) = f_x(x_0, y_0) = f_y(x_0, y_0) = \text{Hess}(f)(x_0, y_0) = K(f)(x_0, y_0) = 0.$$

Note that, by [2, Theorem (1.49)], the dimension of the Severi variety $V(\Delta, A_1)$ satisfies

$$\dim V(\Delta, A_1) = \dim \Sigma(\Delta, A_1) = \#\Delta_{\mathbb{Z}} - 1 - 1$$

and $\Sigma(\Delta, A_1)$ is defined by the first three equations of (**). Therefore, we obtain

$$\dim V(\Delta, A_3) \geq \dim \Sigma(\Delta, A_3) \geq \#\Delta_{\mathbb{Z}} - 1 - 3 = \#\Delta_{\mathbb{Z}} - 4.$$

□

2.3. Tropicalization of curves. We briefly introduce the *tropicalization* of a curve and its refinement (see [8, Section 3] for more details).

Let $F \in K[z, w]$ be a reduced polynomial which defines a curve $C \subset X(N_F)$. Set $\Delta = N_F$ and let T_F be the tropical amoeba defined by F introduced in Section 2.1 and S_F be the dual subdivision of T_F . We consider the 3-dimensional unbounded polyhedron

$$\check{\Delta}_F := \text{Conv}\{(i, j, t) \in \mathbb{R}^2 \times \mathbb{R}; t \geq v_F(i, j)\} \subset \mathbb{R}^3.$$

We remark that a compact facet $\check{\Delta}_i$ of $\check{\Delta}_F$ corresponds to a 2-dimensional polytope Δ_i in S_F by the projection $\check{\Delta}_F \subset \mathbb{R}^2 \times \mathbb{R} \rightarrow \mathbb{R}^2$.

We then obtain a toric flat morphism $X(\check{\Delta}_F) = \mathfrak{X} \rightarrow \mathbb{C}$ from the toric 3-fold associated with $\check{\Delta}_F$ to the complex line, which is called a *toric degeneration*. A generic fiber \mathfrak{X}_t is isomorphic to $X(\Delta)$, and its central fiber \mathfrak{X}_0 is isomorphic to $\bigcup_{i=1, \dots, N} X(\Delta_i)$ (see [6, Section 3] for more details). Let $D \subset \mathbb{C}$ be a small disk centered at the origin. We regard the indeterminate t of K as the variable in $D^* := D \setminus \{0\}$. Then we can get an analytic function $F(t; z, w)$ in three variables. From this analytic function, we obtain an equisingular family on the toric surface $X(\Delta)$

$$\{C^{(t)} := \text{Closure}(\{F(t; z, w) = 0\})\}_{t \in D^*}.$$

The limit $C^{(0)}$ of this family is constructed as follows: For each $i = 1, \dots, N$, a complex polynomial $f_i \in \mathbb{C}[z, w]$ whose Newton polytope is $\Delta_i \in S_F$ is induced from the face function of F on $\check{\Delta}_i$ by the transformation induced by the projection from $\check{\Delta}_i$ to Δ_i . The union of these curves is the limit $C^{(0)}$, which is a curve on the central fiber \mathfrak{X}_0 of the toric degeneration.

The limit $C^{(0)}$ is called a *tropicalization* of C .

For each singular point z of C , there exists a continuous family of singular points $\{z_t\}$ for $t \in D^*$, where $z_t \in C^{(t)}$, and this family defines a section $s : D^* \rightarrow X(\check{\Delta}_F)$. If the limit $s(0) = \lim_{t \rightarrow 0} s(t)$ does not belong to the intersection lines $\bigcup_{i \neq j} X(\Delta_i \cap \Delta_j)$ and bears just one singular point of $C^{(t)}$, the point $s(0)$ is called a *regular singular point*. Otherwise it is called an *irregular singular point*. Note that if $s(0)$ is a regular singular point then it is topologically equivalent to the original singularity.

If the singular point $s(0)$ is irregular, additional information can be obtained by the *refinement* of the tropicalization, see Figure 1. In the rest of this section, we explain this method briefly. See [8, Subsection 3.5] for the details of the refinement.

Hereafter, we assume that F defines a 1-tacnodal curve in $X(\Delta)$. Let $\Delta_1 \in S_F$ and $\Delta_2 \in S_F$ be polytopes which have a common edge σ of length $m \geq 2$ and we observe the case where an irregular singularity degenerates into the subvariety $X(\sigma)$ of \mathfrak{X}_0 . For each $i = 1, 2$, let f_i be a polynomial whose Newton polytope is Δ_i such that the union of curves $C_1 \cap C_2 \subset C^{(0)}$ defined by $f_1 = f_2 = 0$ intersects $X(\sigma)$ at $z \in \mathfrak{X}_0$. In this paper, by later discussion, we can assume that, for each $i = 1, 2$, the polynomial f_i has an isolated singularity at $z \in X(\sigma)$ and their Newton boundary intersects the x - and y -axes at $(m_i, 0)$ and $(0, m)$, respectively, where the y -axis corresponds to $X(\sigma)$.

Find an automorphism $M_\sigma \in \text{Aff}(\mathbb{Z}^2)$ such that $M_\sigma(\Delta)$ is contained in the right half-plane of \mathbb{R}^2 and $M_\sigma(\sigma) =: \sigma'$ is a horizontal segment, see Figure 1. The automorphism M_σ induces a transformation $(x, y) \mapsto (x', y')$, by which we obtain a new polynomial $F'(x', y')$ from F . We can assume that $F' \in K[x', y']$ by multiplying a monomial. We remark that the point z corresponds to a root $\xi \neq 0$ of the truncation polynomial $F'^{\sigma'}(x', y')$ of F' on σ' . Here the truncation polynomial F^σ of a polynomial F on a facet σ of N_F is the sum of the terms of F corresponding to the lattice points on σ .

Then we choose an element $\tau \in K$ such that the coefficient of \tilde{x}^{m-1} in $\tilde{F}(\tilde{x}, \tilde{y}) = F'(\tilde{x} + \xi + \tau, \tilde{y})$ is zero. Moreover, the dual subdivision of the tropical amoeba defined by \tilde{F} contains a subdivision of the triangle $\Delta_z := \text{Conv}\{(m, 0), (0, m_1), (0, -m_2)\}$. In this paper, we call the polytope Δ_z the *exceptional polytope* for the irregular singularity $z \in \mathfrak{X}_0$. We remark that, the exceptional polytope is the union of the complements of the Newton diagrams of the polynomials f_1 and f_2 at $z \in X(\sigma)$ in the first quadrant of \mathbb{R}^2 . Making the exceptional polytope Δ_z by the translation is an operation similar to a blowing-up of the 3-fold \mathfrak{X} . We can restore the topological type of the irregular singularity z in $X(\Delta_z)$ by this operation.




Fig. 1. A refinement of a tropicalization

DEFINITION 2.6. For each $i = 1, 2$, let f_i be a polynomial which defines C_i such that $f_1^\sigma = f_2^\sigma$, and ϕ_i denote the composition of f_i and the translation which maps z to the origin of \mathbb{C}^2 . Set

$$\hat{\sigma}_i := \Delta_z \cap N_{\phi_i} \subset \Delta_z,$$

where N_{ϕ_i} is the Newton polytope of ϕ_i . We assume that $\hat{\sigma}_i$ is an edge of Δ_z . We call a polynomial ϕ whose Newton polytope is Δ_z and that satisfies

- (a) the coefficient of x^{m-1} is zero, and
- (b) the truncation polynomial $\phi^{\hat{\sigma}_i}$ is equal to ϕ_i for each edge $\hat{\sigma}_i$ of Δ_z .

a deformation pattern compatible with given data (f_1, f_2, z) .

We remark that, by the same reason as in [8, Subsection 3.5], except case (E), if the curve defined by F has only one singular point which is an irregular singularity and there does not exist a deformation pattern compatible with the irregular singularity which defines a 1-tacnodal curve, then F does not define a 1-tacnodal curve. We will discuss what happen in case (E) in Subsection 3.4.

3. Tropical 1-tacnodal curves

3.1. Definition of tropical 1-tacnodal curves. In this subsection, we define a tropical 1-tacnodal curve. We can think of it as a tropical version of a 1-tacnodal curve, which is the main theorem (Theorem 4.1) in this paper.

Set

$$\begin{aligned} \Delta_I &:= \text{Conv}\{(0, 7), (1, 0), (2, 0)\}, \quad \Delta_{II} := \text{Conv}\{(0, 7), (2, 0), (3, 0)\}, \\ \Delta_{III} &:= \text{Conv}\{(0, 0), (2, 0), (1, 3)\}, \quad \Delta_{IV} := \text{Conv}\{(0, 0), (2, 0), (1, 2)\} \\ \Delta_V &:= \text{Conv}\{(0, 0), (4, 0), (0, 1)\}, \quad \Delta_{VI} := \text{Conv}\{(1, 0), (2, 0), (0, 3), (1, 3)\}, \\ \Delta_{VII} &:= \text{Conv}\{(0, 0), (1, 0), (2, 1), (0, 1), (1, 2)\}, \\ \Delta_{VIII} &:= \text{Conv}\{(0, 0), (1, 0), (0, 1), (3, 3)\}, \quad \Delta_{IX} := \text{Conv}\{(0, 0), (1, 0), (0, 1), (4, 2)\}, \\ \Delta_E &:= \text{Conv}\{(0, 0), (2, 0), (0, 1), (1, 2)\}, \end{aligned}$$

see Figure 2.

We say that a polytope $P \subset \mathbb{R}^2$ is $\text{Aff}(\mathbb{Z}^2)$ -equivalent to (or simply, equivalent to) P' if there exists an affine isomorphism $A \in \text{Aff}(\mathbb{Z}^2)$ such that $A(P) = P'$, and denote it as $P \simeq P'$.

DEFINITION 3.1. A tropical curve T is said to be *tropical 1-tacnodal* if the dual subdivision S of T contains one of the following polytopes or unions of polytopes:

- (I) a triangle equivalent to Δ_I ,
- (II) a triangle equivalent to Δ_{II} ,
- (III) the union of a triangle equivalent to Δ_{III} and a triangle with edges of lattice length 1, 1 and 2 and without interior lattice point glued in such a way that they share the edge of lattice length 2,
- (IV) the union of two triangles equivalent to Δ_{IV} which share the edge of lattice length 2,
- (V) the union of two triangles equivalent to Δ_V which share the edge of lattice length 4,
- (VI) a parallelogram equivalent to Δ_{VI} ,

- (VII) a pentagon equivalent to Δ_{VII} ,
- (VIII) a quadrangle equivalent to Δ_{VIII} ,
- (IX) a quadrangle equivalent to Δ_{IX} ,
- (E) the union of a quadrangle equivalent to Δ_{E} and a triangle with edges of lattice length 1, 1 and 2 and without interior lattice point which share the edge of lattice length 2, and the rest of S consists of triangles of area $1/2$.




Fig. 2. Polytopes in Definition 3.1. The notation Δ means a lattice point on the boundary which is not a vertex and the notation \star means an interior lattice point.

3.2. Polytopes corresponding to tropical 1-tacnodal curves. In this subsection, we mention some remark on polytopes appearing in Definition 3.1.

We denote an m -gon which has edges of lattice lengths ℓ_1, \dots, ℓ_m and I interior lattice points by

$$\Delta_m(I; \ell_1, \dots, \ell_m).$$

Similarly, we denote a parallel $2m$ -gon which has m pairs of antipodal parallel edges of lattice length ℓ_1, \dots, ℓ_m by

$$\Delta_{2m}^{\text{par}}(I; \ell_1, \dots, \ell_m).$$

When we consider polytopes of the same type $(I; \ell_1, \dots, \ell_m)$ simultaneously, we denote one as $\Delta_m(I; \ell_1, \dots, \ell_m)$ and the others as $\Delta'_m(I; \ell_1, \dots, \ell_m)$, $\Delta''_m(I; \ell_1, \dots, \ell_m)$ and so on. Note that in these notations, the order of lengths is not relevant to the cyclic order of edges.

Lemma 3.2. *The following holds up to $\text{Aff}(\mathbb{Z}^2)$ -equivalence:*

- (1) A triangle $\Delta_3(3; 1, 1, 1)$ is either Δ_{I} or Δ_{II} .
- (2) A triangle $\Delta_3(2; 2, 1, 1)$ is Δ_{III} .

- (3) A triangle $\Delta_3(1; 2, 1, 1)$ is Δ_{IV} .
- (4) A triangle $\Delta_3(0; 4, 1, 1)$ is Δ_V .
- (5) A parallelogram $\Delta_4^{\text{par}}(2; 1, 1)$ is Δ_{VI} .
- (6) A pentagon $\Delta_5(1; 1, 1, 1, 1)$ is Δ_{VII} .
- (7) A non-parallel quadrangle $\Delta_4(2; 1, 1, 1, 1)$ is equivalent to one of the following polytopes:

$$\Delta_{VIII}, \Delta_{IX}, \text{Conv}\{(1, 0), (0, 1), (2, 1), (1, 3)\}.$$

Proof. (1) We can take $A \in \text{Aff}(\mathbb{Z}^2)$ which maps $\Delta_3(3; 1, 1, 1)$ to

$$\hat{\Delta}_n := \text{Conv}\{(0, q), (n, 0), (n + 1, 0)\}$$

for some q, n . By Pick's formula, we obtain $q = 7$. We remark that, $\hat{\Delta}_n$ and $\hat{\Delta}_{n+7}$ are equivalent by

$$(***) \quad \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}.$$

Moreover we do not have to discuss the cases $n = 0$ and $n = 6$ since they have an edge of lattice length more than 1.

We get the isomorphisms

$$\hat{\Delta}_1 \simeq \hat{\Delta}_5, \quad \hat{\Delta}_2 \simeq \hat{\Delta}_4$$

by the reflection, and $\hat{\Delta}_1 \simeq \hat{\Delta}_3$ by

$$\begin{pmatrix} 3 & 1 \\ -7 & -2 \end{pmatrix}.$$

Because of the configuration of interior lattice points, we can show that $\hat{\Delta}_1 = \Delta_I$ and $\hat{\Delta}_2 = \Delta_{II}$ are not isomorphic.

- (2) For any $\Delta_3(2; 2, 1, 1)$, there exists $A \in \text{Aff}(\mathbb{Z}^2)$ such that $\Delta_3(2; 2, 1, 1)$ maps to

$$\text{Conv}\{(p, 0), (p + 2, 0), (0, q)\}$$

for some $p, q \in \mathbb{N}$. Then we have $q = 3$ by Pick's formula, and we may assume $p = 0, 1, 2$ by the isomorphism (***). But the cases $p = 0, 1$ do not satisfy the conditions of lattice length. Hence we get $p = 2$. This triangle is equivalent to Δ_{III} .

The claims (3), (4), (5) and (6) can be proved by the same method.

- (7) We can split $P := \Delta_4(2; 1, 1, 1, 1)$ into two triangles which satisfies one of the following:

- $\Delta_3(1; 2, 1, 1)$ and $\Delta_3(0; 2, 1, 1)$ such that their intersection is a segment of length 2,
- $\Delta_3(2; 1, 1, 1)$ and $\Delta_3(0; 1, 1, 1)$ such that their intersection is a segment of length 1,
- $\Delta_3(0; 3, 1, 1)$ and $\Delta'_3(0; 3, 1, 1)$ such that their intersection is a segment of length 3,
- $\Delta_3(1; 1, 1, 1)$ and $\Delta'_3(1; 1, 1, 1)$ such that their intersection is a segment of length 1.

In the first case, $\Delta_3(1; 2, 1, 1)$ is uniquely determined as $\text{Conv}\{(0, 0), (2, 0), (1, 2)\}$, so P has two descriptions

$$\hat{P}_1 := \text{Conv}\{(0, 0), (2, 0), (1, 2), (0, -1)\}, \quad \hat{P}_2 := \text{Conv}\{(0, 0), (2, 0), (1, 2), (1, -1)\}.$$

In the second case, by [8, Lemma 4.1], any triangle $\Delta_3(2; 1, 1, 1)$ is isomorphic to

$$Q := \text{Conv}\{(0, 0), (3, 2), (2, 3)\}.$$

We denote the other triangle, which is $\Delta_3(0; 1, 1, 1)$, by R . We can easily check that Q is equivalent to

$$Q_1 := \text{Conv}\{(1, 0), (2, 0), (0, 5)\}, \quad Q_2 := \text{Conv}\{(2, 0), (3, 0), (0, 5)\}.$$

If the intersection of Q with R is $\text{Conv}\{(0, 0), (3, 2)\} \subset Q$ or $\text{Conv}\{(0, 0), (2, 3)\} \subset Q$, then we can assume that the intersection is the bottom edge of Q_1 . Similarly, if the intersection is $\text{Conv}\{(2, 3), (3, 2)\} \subset Q$, then we can assume that R shares the bottom edge of Q_2 . Thus, the polytope P is equivalent to either

$$\hat{P}_3 := \text{Conv}\{(1, 0), (2, 0), (0, 5), (2, -1)\} \quad \text{or} \quad \hat{P}_4 := \text{Conv}\{(2, 0), (3, 0), (0, 5), (3, -1)\}.$$

In the third and fourth cases, we obtain the following polytopes in the same way as above:

$$\hat{P}_5 := \text{Conv}\{(0, 0), (0, 1), (1, -1), (3, 0)\}, \quad \hat{P}_6 := \text{Conv}\{(0, 0), (0, 1), (2, -1), (3, 0)\}.$$

Between the polytopes $\hat{P}_1, \dots, \hat{P}_6$, we have the following isomorphisms:

$$\hat{P}_1 \simeq \hat{P}_3 \quad \text{by} \quad \begin{pmatrix} -1 & 0 \\ 3 & -1 \end{pmatrix}, \quad \hat{P}_5 \simeq \hat{P}_4 \quad \text{by} \quad \begin{pmatrix} -1 & -1 \\ 2 & 1 \end{pmatrix}, \quad \hat{P}_6 \simeq \hat{P}_2 \quad \text{by} \quad \begin{pmatrix} 0 & -1 \\ 1 & 1 \end{pmatrix}.$$

Notice that, the polytope \hat{P}_2 is the translation of $\text{Conv}\{(1, 0), (0, 1), (2, 1), (1, 3)\}$. Also, the polytopes \hat{P}_3 and \hat{P}_4 are equivalent to Δ_{IX} and Δ_{VIII} by

$$\begin{pmatrix} 1 & 1 \\ -1 & 0 \end{pmatrix}: \mathbb{Z}^2 \rightarrow \mathbb{Z}^2,$$

respectively.

Furthermore, by the configuration of interior lattice points and vertices, we obtain $\Delta_{\text{VIII}} \neq \Delta_{\text{IX}}$, $\Delta_{\text{IX}} \neq \text{Conv}\{(1, 0), (0, 1), (2, 1), (1, 3)\}$, and $\text{Conv}\{(1, 0), (0, 1), (2, 1), (1, 3)\} \neq \Delta_{\text{VIII}}$. \square

Lemma 3.3. *A quadrangle $\Delta_4(1; 2, 1, 1, 1)$ is Δ_{E} .*

Proof. We can split $P = \Delta_4(1; 2, 1, 1, 1)$ into two polytopes Q, R along a line passing through the mid point of the edge of length two. Then there are three cases:

(3-1) $Q = \Delta_3(0; 1, 1, 1), R = \Delta_4(1; 1, 1, 1, 1)$ and these polytopes share an edge of length 1,

(3-2) $Q = \Delta_3(0; 2, 1, 1), R = \Delta_4(0; 2, 1, 1, 1)$ and these polytopes share the edge of length 2,

(3-3) $Q = \Delta_3(1; 1, 1, 1), R = \Delta_4(0; 1, 1, 1, 1)$ and these polytopes share an edge of length 1.

(3-1) If R is a parallelogram, then we can assume that R is

$$\text{Conv}\{(1, 0), (2, 0), (0, 2), (1, 2)\}$$

by [8, Lemma 4.1] and the common edge of R with Q is its bottom edge. Hence, we get

$$Q = \text{Conv}\{(1, 0), (2, 0), (2, -1)\},$$

by Pick's formula, but their union does not satisfy the condition of P .

If R is not a parallelogram, then we can assume that R is

$$\text{Conv}\{(0, 0), (1, 0), (0, 1), (2, 2)\}$$

by [8, Lemma 4.1] and the common edge with Q is either

$$\text{Conv}\{(0, 0), (1, 0)\} \text{ or } \text{Conv}\{(1, 0), (2, 2)\}.$$

In the former case, Q is uniquely determined as

$$\text{Conv}\{(0, 0), (1, 0), (0, -1)\}$$

and the union $Q \cup R = \text{Conv}\{(0, -1), (1, 0), (0, 1), (2, 2)\}$ is isomorphic to P . In the latter case, we can assume that R is

$$\text{Conv}\{(1, 0), (2, 0), (0, 2), (0, 3)\}$$

and the common edge is the bottom edge. Then Q must be

$$\text{Conv}\{(1, 0), (2, 0), (2, 1)\},$$

but the union $Q \cup R$ does not satisfy the condition of P .

(3-2) We can assume that R is

$$\text{Conv}\{(0, 0), (2, 0), (0, 1), (1, 1)\}$$

by [8, Lemma 4.1] and the common edge is the bottom edge. Then Q must be either

$$\text{Conv}\{(0, 0), (2, 0), (0, -1)\} \text{ or } \text{Conv}\{(0, 0), (2, 0), (3, -1)\}.$$

In both cases, the union $Q \cup R$ are isomorphic to Δ_E .

(3-3) We can assume that R is

$$\text{Conv}\{(0, 0), (1, 0), (0, 1), (1, 1)\},$$

by direct computation but any union with Q does not satisfy the condition of Δ_E . \square

3.3. Existence of 1-tacnodal curves for $\Delta_I, \dots, \Delta_{IX}$. For a polytope P , we set

$$\mathcal{F}(P) := \{f \in \mathbb{C}[x, y]; N_f = P\}.$$

We denote the plane curve defined by $f \in \mathcal{F}(P)$ in $X(P)$ as V_f . We remark that V_f is a member of $|D(P)|$. We consider the following two conditions:

- (S1) $V_f \subset X(P)$ is a 1-tacnodal curve whose singular point is contained in the maximal torus of $X(P)$,
- (S2) V_f intersects the toric boundary $X(\partial P)$ transversally.

In the rest of this section, except cases (III), (IV), and (V), we only consider polytopes whose edges are only of length one. Hence the condition (S2) is automatically satisfied except the three cases.

Lemma 3.4. *For each $k = \text{I, II}$ and given coefficients c_{ij} on the vertices $(i, j) \in V(P)$, there is a polynomial $f \in \mathcal{F}(\Delta_k)$ which has the fixed coefficients on the vertices and satisfies the conditions (S1), (S2). Furthermore, there is no polynomial $f \in \mathcal{F}(\Delta_k)$ that defines a curve with more complicated singularity than A_3 , i.e., the curve does not have an isolated*

singularity whose Milnor number is more than 3.

Proof. (I) We first show that we can assume that the coefficients on the vertices of Δ_I are 1. We transform the polynomial

$$f = c_{10}x + c_{20}x^2 + Axy + Bxy^2 + Cxy^3 + c_{07}y^7 \in \mathcal{F}(\Delta_I)$$

by substituting $x = X^{-1}, y = Y$ and multiplying X^2 . Then we get a new polynomial

$$\tilde{f} := c_{20} + c_{10}X + AXY + BXY^2 + CXY^3 + c_{07}X^2Y^7.$$

By multiplying suitable constants to the variables and the whole polynomial, we can assume that $c_{20} = c_{10} = c_{07} = 1$. Transforming \tilde{f} by $x = X^{-1}, y = Y$ again, we get

$$x + x^2 + A'xy + B'xy^2 + C'xy^3 + y^7.$$

We re-denote this polynomial by f .

For a polynomial

$$f = x + x^2 + Axy + Bxy^2 + Cxy^3 + y^7 \in \mathcal{F}(\Delta_I),$$

we apply Lemma 2.4 and eliminate the variables by the system $f = f_x = f_y = \text{Hess}(f) = K(f) = 0$. First, by $f = 0$, we can get A as

$$A = -\frac{x + x^2 + Bxy^2 + Cxy^3 + y^7}{xy}.$$

Therefore the system is reduced as

$$\begin{cases} (1) & x^2 - y^7 = 0, \\ (2) & -x - x^2 + Bxy^2 + 2Cxy^3 + 6y^7 = 0, \\ (3) & \text{substituting } A \text{ for } \text{Hess}(f) = 0, \\ (4) & \text{substituting } A \text{ for } K(f) = 0. \end{cases}$$

Secondly, by equation (2), we can get B as

$$B = \frac{x + x^2 - 2Cxy^3 - 6y^7}{xy^2}.$$

Then the system is reduced as

$$\begin{cases} (1') & x^2 - y^7 = 0, \\ (3') & 4x^3 + 4x^4 + 4Cx^3y^3 + 60x^2y^7 - 49y^{14} = 0, \\ (4') & 2Cx^3 + 7xy^4 + 77x^2y^4 + 7Cxy^7 - 42y^{11} = 0. \end{cases}$$

Thirdly, by equation (3'), we can get C as

$$C = \frac{-4x^3 - 4x^4 - 60x^2y^7 + 49y^{14}}{4x^3y^3}.$$

Then the system is reduced as

$$\begin{cases} x^2 - y^7 = 0, \\ 8x^5 + 8x^6 - 160x^4y^7 + 490x^2y^{14} - 343y^{21} = 0. \end{cases}$$

Hence we obtain $x = 8/5$ and the equation

$$(\text{****}) \quad y^7 - (8/5)^2 = 0.$$

Next, we check that the above f satisfies the condition (S1). Let y_0, y_1, \dots, y_6 be the solutions of equation (****) and, for each $i = 0, \dots, 6$, let $f^{(i)}$ denote the polynomial f with the solution $y = y_i$. By the above calculation, the curve $V_{f^{(i)}}$ defined by $f^{(i)}$ has a tacnode at $(8/5, y_i) \in (\mathbb{C}^*)^2$. Notice that the coefficients A, B and C of $f^{(i)}$ are determined by $x = 8/5$ and $y = y_i$. Let (s, t) be a singular point of $f^{(i)}$ on $V_{f^{(i)}}$. Solving $f_x^{(i)} = 0$, we obtain $s = s_i(t, y_i)$. Set

$$f_1(t, y_i) := f^{(i)}(s_i(t, y_i), t), \quad f_2(t, y_i) := f_{y_i}^{(i)}(s_i(t, y_i), t).$$

Eliminating y_i from f_1, f_2 by $y_i^7 - (8/5)^2 = 0$, we obtain two equations with variable t . We can check that their greatest common divisor is $t^7 - (5/8)^2$. Thus, the singularities of $f^{(i)}$ are only tacnodes.

The coefficient A of $f^{(i)}$ depends only on the solution y_i of (****) and we can check directly that the coefficients A for $y = y_i$ and $y = y_j$ are different if $i \neq j$. That is, the defining polynomials $f^{(i)}$ and $f^{(j)}$ are different for $i \neq j$. Therefore each $f^{(i)}$ satisfies the condition (S1).

(II) For the polynomial

$$f = x^2 + x^3 + Ax^2y + Bx^2y^2 + Cxy^4 + y^7 \in \mathcal{F}(\Delta_{II}),$$

we apply Lemma 2.4 and eliminate the variables by the system $f = f_x = f_y = \text{Hess}(f) = K(f) = 0$. First, by $f = 0$, we can get A as

$$A = -\frac{x^2 + x^3 + Bx^2y^2 + Cxy^4 + y^7}{x^2y}.$$

Therefore the system is reduced as

$$\left\{ \begin{array}{l} (1) \quad x^3 - Cxy^4 - 2y^7 = 0, \\ (2) \quad -x^2 - x^3 + Bx^2y^2 + 3Cxy^4 + 6y^7 = 0, \\ (3) \quad \text{substituting } A \text{ for } \text{Hess}(f) = 0, \\ (4) \quad \text{substituting } A \text{ for } K(f) = 0. \end{array} \right.$$

Secondly, by equation (2), we can get B as

$$B = \frac{x^2 + x^3 - 3Cxy^4 - 6y^7}{x^2y^2}.$$

Then the system is reduced as

$$\left\{ \begin{array}{l} (1') \quad x^3 - Cxy^4 - 2y^7 = 0, \\ (3') \quad 8x^5 + 8x^6 - 4Cx^3y^4 + 20Cx^4y^4 - 4x^2y^7 + 116x^3y^7 - 28C^2x^2y^8 - 184Cxy^{11} \\ \quad - 256y^{14} = 0, \\ (4') \quad \text{substituting } B \text{ for } (4) = 0. \end{array} \right.$$

Thirdly, by equation (1'), we can get C as

$$C = \frac{x^3 - 2y^7}{xy^4}.$$

Then the system is reduced as

$$\begin{cases} x^3 + y^7 + xy^7 = 0, \\ 4x^9 + 14x^6y^7 + 5x^7y^7 + 16x^3y^{14} + 11x^4y^{14} + 6y^{21} + 7xy^{21} = 0. \end{cases}$$

By direct computation, we can see that the solution of the above system is

(*****) $(x, y) = (y_0^7, y_0),$

where y_0 is a solution of $y^{14} + y^7 + 1 = 0$.

Next, we check that the above f satisfies the condition (S1). Notice that the curve V_f defined by f has a tacnode at $(y_0^7, y_0) \in (\mathbb{C}^*)^2$, where y_0 is a solution of (*****). Let $(s, t) \in (\mathbb{C}^*)^2$ be a singular point of V_f . Then, we can easily check that the system $f(s, t) = f_x(s, t) = f_y(s, t) = y_0^{14} + y_0^7 + 1 = 0$ implies $t = y_0$. After substituting $t = y_0$ for $f(s, t), f_x(s, t), f_y(s, t)$, we obtain $s - y_0^7$ as their greatest common divisor. That is, the singularities of V_f are only tacnodes. Moreover, we can easily check that for two different solutions y_0 and y'_0 of $y^{14} + y^7 + 1 = 0$, the triples (A, B, C) of the coefficients of the polynomial f , which are determined by y_0 and y'_0 , are different. Therefore, for each solution of $y^{14} + y^7 + 1 = 0$, the polynomial f satisfies the condition (S1). □

Lemma 3.5. *For each $k = \text{VI, VII, VIII, IX}$, and given coefficients c_{ij} on the vertices $(i, j) \in V(P)$, there is a polynomial $f \in \mathcal{F}(\Delta_k)$ which has the fixed coefficients on the vertices and satisfies (S1) and (S2) if and only if*

$$\begin{aligned} c_{03}c_{20} &= 64c_{10}c_{13}, \quad \text{if } k = \text{VI}, \\ c_{21}c_{00}^2 &= -4c_{01}c_{10}^2, \quad \text{and } c_{12}c_{00}^2 = -4c_{10}c_{01}^2, \quad \text{if } k = \text{VII}, \\ 8^6 c_{33}c_{00}^5 &= 5^5 c_{10}^3c_{01}^3, \quad \text{if } k = \text{VIII}, \\ 256c_{42}c_{00}^5 &= (41 + 38\sqrt{-1})c_{10}^4c_{01}^2, \quad \text{or } 256c_{42}c_{00}^5 = (41 - 38\sqrt{-1})c_{10}^4c_{01}^2, \quad \text{if } k = \text{IX}. \end{aligned}$$

Furthermore, there is no polynomial $f \in \mathcal{F}(\Delta_k)$ that defines a curve with more complicated singularity than A_3 .

Proof. (VI) We transform the polynomial

$$f := c_{10}x + c_{20}x^2 + Axy + Bxy^2 + c_{03}y^3 + c_{13}xy^3 \in \mathcal{F}(\Delta_{\text{VI}})$$

by substituting $x = X^{-1}, y = Y$ and multiplying X^2 . Then we get the new polynomial

$$\tilde{f} := c_{10}X + c_{20} + A'XY + B'XY^2 + c_{03}X^2Y^3 + c_{13}XY^3.$$

By multiplying suitable constants to the variables and the whole polynomial, we can rewrite \tilde{f} as

$$1 + X + A''XY + B''XY^2 + XY^3 + CX^2Y^3,$$

where

$$C = \frac{c_{03}c_{20}}{c_{10}c_{13}}.$$

For the polynomial

$$1 + x + Axy + Bxy^2 + xy^3 + Cx^2y^3$$

we apply Lemma 2.4 and eliminate the variables by the system $f = f_x = f_y = \text{Hess}(f) = K(f) = 0$. First, by $f = 0$, we can get A as

$$A = -\frac{1 + x + Bxy^2 + xy^3 + Cx^2y^3}{xy}.$$

Therefore the system is reduced as

$$\begin{cases} (1) & -1 + Cx^2y^3 = 0, \\ (2) & -1 - x + Bxy^2 + 2xy^3 + 2Cx^2y^3 = 0, \\ (3) & \text{substituting } A \text{ for } \text{Hess}(f) = 0, \\ (4) & \text{substituting } A \text{ for } K(f) = 0. \end{cases}$$

Secondly, by equation (1), we can get C as

$$C = \frac{1}{x^2y^3}.$$

Then the system is reduced as

$$\begin{cases} (2') & 1 - x + Bxy^2 + 2xy^3 = 0, \\ (3') & -4 + 8x - x^2 - 4Bxy^2 + 2Bx^2y^2 - 4xy^3 + 4x^2y^3 - B^2x^2y^4 - 4Bx^2y^5 - 4x^2y^6 = 0, \\ (4') & 48 - 144x + 36x^2 + 48Bxy^2 + 48xy^3 - 48Bx^2y^2 - 72x^2y^3 + 12B^2x^2y^4 + 24Bx^2y^5 \\ & = 0. \end{cases}$$

Thirdly, by equation (2'), we can get B as

$$B = -\frac{1 - x + 2xy^3}{xy^2}.$$

Then the system is reduced as

$$\begin{cases} 4x + 4xy^3 - 1 = 0, \\ 6x + 2xy^3 - 1 = 0. \end{cases}$$

The solution of the above system is

$$(\text{*****}) \quad (x, y) = (1/8, y_0),$$

where y_0 is a solution of $y^3 = 1$. Then we obtain

$$A = -9/y_0, \quad B = -9/y_0^2, \quad C = 1/x^2y_0^3 = 64.$$

Next, we check that the above f satisfies the condition (S1). Notice that the curve V_f defined by f has a tacnode at $(1/8, y_0) \in (\mathbb{C}^*)^2$, where y_0 is a solution of (*****). Let $(s, t) \in (\mathbb{C}^*)^2$ be a singular point of V_f . Then, we obtain $t^3 - 1 = 0$ and $s = 1/8$ from the system $f(s, t) = f_x(s, t) = f_y(s, t) = 0$ and the equation $y_0^3 - 1 = 0$. That is, the singularities of V_f are only tacnodes. Moreover, we can easily check that for two different solutions y_0 and y'_0 of $y^3 - 1 = 0$, the triples (A, B, C) of the coefficients of the polynomial f , which

are determined by y_0 and y'_0 , are different. Therefore, for each solution of $y^3 - 1 = 0$, the polynomial f satisfies the condition (S1).

(VII) We can rewrite the polynomial

$$f = c_{00} + c_{10}x + c_{01}y + Axy + c_{21}x^2y + c_{12}xy^2 \in \mathcal{F}(\Delta_{\text{VII}})$$

as

$$f = 1 + x + y + Axy + Bx^2y + Cxy^2$$

by the same manner as above, where

$$B = \frac{c_{21}c_{00}^2}{c_{01}c_{10}^2}, \quad C = \frac{c_{12}c_{00}^2}{c_{10}c_{01}^2}.$$

For the polynomial

$$f = 1 + x + y + Axy + Bx^2y + Cxy^2,$$

we apply Lemma 2.4 and eliminate the variables by the system $f = f_x = f_y = \text{Hess}(f) = K(f) = 0$. First, by $f = 0$, we can get A as

$$A = -\frac{1 + x + y + Bx^2y + Cxy^2}{xy}.$$

Therefore the system is reduced as

$$\left\{ \begin{array}{l} (1) \quad -1 - y + Bx^2y = 0, \\ (2) \quad -1 - x + Cxy^2 = 0, \\ (3) \quad \text{substituting } A \text{ for } \text{Hess}(f) = 0, \\ (4) \quad \text{substituting } A \text{ for } K(f) = 0. \end{array} \right.$$

Secondly, by equations (1) and (2), we can get B and C as

$$B = \frac{1 + y}{x^2y}, \quad C = \frac{1 + x}{xy^2},$$

respectively. Then the system is reduced as

$$\left\{ \begin{array}{l} (3') \quad 3 + 4x + 4y + 4xy = 0, \\ (4') \quad (1 + y)^2(1 + 2x) = 0. \end{array} \right.$$

The solution of the above system is

$$(x, y) = (-1/2, -1/2),$$

and we obtain

$$A = B = C = -4.$$

Next, we check that the above f satisfies the condition (S1). Notice that the curve V_f defined by f has a tacnode at $(-1/2, -1/2) \in (\mathbb{C}^*)^2$. Let $(s, t) \in (\mathbb{C}^*)^2$ be a singular point of V_f . Then, we can solve $f(s, t) = f_x(s, t) = f_y(s, t) = 0$, and the solution is $(s, t) = (-1/2, -1/2)$. That is, the singularity of f is only one point and is a tacnode. Therefore the

f satisfies the condition (S1).

(VIII) We can rewrite the polynomial

$$f = c_{00} + c_{10}x + c_{01}y + Axy + Bx^2y^2 + c_{33}x^3y^3 \in \mathcal{F}(\Delta_{\text{VIII}})$$

as

$$f = 1 + x + y + Axy + Bx^2y^2 + Cx^3y^3$$

by the same manner as above, where

$$C = \frac{c_{33}c_{00}^5}{c_{10}^3c_{01}^3}.$$

For the polynomial

$$f = 1 + x + y + Axy + Bx^2y^2 + Cx^3y^3,$$

we apply Lemma 2.4 and eliminate the variables by the system $f = f_x = f_y = \text{Hess}(f) = K(f) = 0$. First, by $f = 0$, we can get A as

$$A = -\frac{1 + x + y + Bx^2y^2 + Cx^3y^3}{xy}.$$

Therefore the system is reduced as

$$\begin{cases} (1) & -1 - y + Bx^2y^2 + 2Cx^3y^3 = 0, \\ (2) & -1 - x + Bx^2y^2 + 2Cx^3y^3 = 0, \\ (3) & \text{substituting } A \text{ for } \text{Hess}(f) = 0, \\ (4) & \text{substituting } A \text{ for } K(f) = 0. \end{cases}$$

Secondly, by equation (1), we can get B as

$$B = \frac{1 + y - 2Cx^3y^3}{x^2y^2}.$$

Then the system is reduced as

$$\begin{cases} (2') & x - y = 0, \\ (3') & 4 - x + 4y + 4Cx^3y^3 = 0, \\ (4') & \text{substituting } B \text{ for } (4) = 0. \end{cases}$$

Thirdly, by equation (3'), we can get C as

$$C = \frac{-4 + x - 4y}{4x^3y^3}.$$

Then the system is reduced as

$$\begin{cases} x - y = 0, \\ -8 + 3x - 8y = 0. \end{cases}$$

The solution of the above system is

$$(x, y) = (-8/5, -8/5),$$

and we also obtain

$$A = 75/64, \quad B = -5^4/2^{12}, \quad C = 5^5/8^6.$$

Next, we check that the above f satisfies the condition (S1). Notice that the curve V_f defined by f has a tacnode at $(-8/5, -8/5) \in (\mathbb{C}^*)^2$. Let $(s, t) \in (\mathbb{C}^*)^2$ be a singular point of V_f . Then, we can solve $f(s, t) = f_x(s, t) = f_y(s, t) = 0$, and the solution is $(s, t) = (-8/5, -8/5)$. That is, the singularity of f is only one point and is a tacnode. Therefore the f satisfies the condition (S1).

(IX) We can rewrite the polynomial

$$f = c_{00} + c_{10}x + c_{01}y + Axy + Bx^2y + c_{42}x^4y^2 \in \mathcal{F}(\Delta_{IX})$$

as

$$f = 1 + x + y + Axy + Bx^2y + Cx^4y^2$$

by the same manner as above, where

$$C = \frac{c_{42}c_{00}^5}{c_{10}^4c_{01}^2}.$$

For the polynomial

$$f = 1 + x + y + Axy + Bx^2y + Cx^4y^2,$$

we apply Lemma 2.4 and eliminate the variables by the system $f = f_x = f_y = \text{Hess}(f) = K(f) = 0$. First, by $f = 0$, we can get A as

$$A = -\frac{1 + x + y + Bx^2y + Cx^4y^2}{xy}.$$

Therefore the system is reduced as

$$\left\{ \begin{array}{l} (1) \quad -1 - y + Bx^2y + 3Cx^4y^2 = 0, \\ (2) \quad -1 - x + Cx^4y^2 = 0, \\ (3) \quad \text{substituting } A \text{ for } \text{Hess}(f) = 0, \\ (4) \quad \text{substituting } A \text{ for } K(f) = 0. \end{array} \right.$$

Secondly, by equations (1), we can get B as

$$B = \frac{1 + y - 3Cx^4y^2}{x^2y}.$$

Then the system is reduced as

$$\left\{ \begin{array}{l} (2') \quad -1 - x + Cx^4y^2 = 0, \\ (3') \quad 1 - 4Cx^2y^2 - 8Cx^3y^2 - 4Cx^2y^3 + 4C^2x^6y^4 = 0, \\ (4') \quad \text{substituting } B \text{ for } (4) = 0. \end{array} \right.$$

Thirdly, by equations (2'), we can get C as

$$C = \frac{1+x}{x^4 y^2}.$$

Then the system is reduced as

$$\begin{cases} 4x + 4y + 3x^2 + 4xy = 0, \\ (4 + 3x)(16x + 8y + 24x^2 + 22xy + 4y^2 + 9x^3 + 12x^2y + 5xy^2) = 0. \end{cases}$$

The solutions of the above system are

$$(x_0, y_0) = \left(-\frac{6}{5} + \frac{2}{5}\sqrt{-1}, \frac{2}{5} - \frac{4}{5}\sqrt{-1}\right),$$

$$(x_1, y_1) = \left(-\frac{6}{5} - \frac{2}{5}\sqrt{-1}, \frac{2}{5} + \frac{4}{5}\sqrt{-1}\right) \text{ and } (x_2, y_2) = \left(-\frac{4}{3}, 0\right),$$

and we obtain

$$C = -\frac{41}{256} + \frac{19}{128}\sqrt{-1} \text{ if } x = x_0, \quad C = \frac{41}{256} + \frac{19}{128}\sqrt{-1} \text{ if } x = x_1.$$

Here, we can ignore the case $x = x_2$ since $f(-4/3, 0) \neq 0$

Next, we check that the above f satisfies the condition (S1). Notice that the curve V_f defined by f has a tacnode at $(x_0, y_0) \in (\mathbb{C}^*)^2$. Let $(s, t) \in (\mathbb{C}^*)^2$ be a singular point of V_f . Then, we can solve $f(s, t) = f_x(s, t) = f_y(s, t) = 0$, and the solution is $(s, t) = (x_0, y_0)$. That is, the singularity of f is only one point and is a tacnode. Therefore the f satisfies the condition (S1). Also, we can check the condition (S1) for (x_1, y_1) by the same manner. \square

Lemma 3.6. *For each $k = \text{III}, \text{IV}, \text{V}$ and given coefficients c_{ij} on the vertices $(i, j) \in V(P)$, there is a polynomial $f \in \mathcal{F}(\Delta_k)$ which has the fixed coefficients on the vertices such that f defines a curve which has*

- (III) *an A_2 -singularity on the toric divisor corresponding to the edge of length 2,*
- (IV) *an A_1 -singularity on the toric divisor corresponding to the edge of length 2,*
- (V) *an intersection with the toric divisor corresponding to the edge of length 4 whose multiplicity is 4.*

Proof. (III) We set

$$f := 1 + Ax + x^2 + Bxy + Cxy^2 + xy^3 \in \mathcal{F}(\Delta_{\text{III}}).$$

Let $\sigma \subset \Delta_{\text{III}}$ be the edge of length 2. The intersection point of $X(\sigma)$ and the curve defined by f is an A_2 -singularity and this implies $A = \pm 2$.

We assume $A = 2$ and the singularity is at $(-1, 0)$. For $f = (1+x)^2 + Bxy + Cxy^2 + y^3$, the solution of $f(-1, 0) = f_x(-1, 0) = f_y(-1, 0) = \text{Hess}(f)(-1, 0) = 0$ is $B = C = 0$. Therefore we obtain the polynomial $f := 1 + 2x + x^2 + xy^3 \in \mathcal{F}(\Delta_{\text{III}})$.

(IV) We set

$$f := 1 + Ax + x^2 + Bxy + xy^2 \in \mathcal{F}(\Delta_{\text{IV}}).$$

Let $\sigma \subset \Delta_{\text{IV}}$ be the edge of length 2. The intersection point of $X(\sigma)$ and the curve defined by f is an A_1 -singularity and this implies $A = \pm 2$.

We assume $A = 2$ and the singularity is at $(-1, 0)$. For $f = (1+x)^2 + Bxy + xy^2$, the solution of $f(-1, 0) = f_x(-1, 0) = f_y(-1, 0) = 0$ is $B = 0$. Therefore we obtain the

polynomial $f := 1 + 2x + x^2 + xy^2 \in \mathcal{F}(\Delta_{IV})$.

(V) We can prove that the polynomial

$$f := (1 \pm x)^4 + y \in \mathcal{F}(\Delta_V)$$

satisfies the condition. □

Set

$$\hat{\Delta}_{III} := \text{Conv}\{(0, -1), (2, 0), (0, 3)\},$$

$$\hat{\Delta}_{IV} := \text{Conv}\{(0, -2), (2, 0), (0, 2)\},$$

$$\hat{\Delta}_V := \text{Conv}\{(0, -1), (4, 0), (0, 1)\}.$$




Fig. 3. Polytopes $\hat{\Delta}_{III}$, $\hat{\Delta}_{IV}$ and $\hat{\Delta}_V$. The notation Δ means a lattice point on the boundary which is not a vertex and the notation \star means an interior lattice point.

For the polytopes Δ_{III} and $\Delta_3(0; 2, 1, 1)$ appearing in Definition 3.1 (III), the polynomial on Δ_{III} obtained in Lemma 3.6 induces the polynomial on $\Delta_3(0; 2, 1, 1)$ as

$$1 + Ax + x^2 + y,$$

where $A = \pm 2$. Therefore the exceptional polytope in this case is $\hat{\Delta}_{III}$.

For the polytopes Δ_{IV} and $\Delta_3(1; 2, 1, 1)$ appearing in Definition 3.1 (IV), the polynomial on Δ_{IV} obtained in Lemma 3.6 induces the polynomial on $\Delta_3(1; 2, 1, 1)$ as

$$1 + Ax + x^2 + Bxy + xy^2,$$

where $A = \pm 2$. If $B = 0$, the exceptional polytope compatible with the data is $\hat{\Delta}_{IV}$. Note that, if $B \neq 0$, the exceptional polytope compatible with the data is

$$\text{Conv}\{(2, 0), (0, 2), (0, -1)\},$$

and it has no deformation pattern which defines an 1-tacnodal curve, see the discussion in Lemma 4.9.

For the polytopes Δ_V and $\Delta_3(0; 4, 1, 1)$ appearing in Definition 3.1 (V), the polynomial on Δ_V obtained in Lemma 3.6 induces the same polynomial on $\Delta_3(0; 4, 1, 1)$. Therefore, the exceptional polytope compatible with the data is $\hat{\Delta}_V$.

Lemma 3.7. *For each $k = \text{III}, \text{IV}, \text{V}$, there is a deformation pattern $\phi \in \mathcal{F}(\hat{\Delta}_k)$ compatible with given data in Lemma 3.6 which has the fixed coefficients on the vertices such that the curve defined by ϕ in $X(\hat{\Delta}_k)$ is a 1-tacnodal curve.*

Proof. (III) For the polynomial

$$\phi := 1 + Ay + x^2y + By^2 + Cxy^2 + Dy^3 + y^4 \in \mathcal{F}(\hat{\Delta}_{\text{III}})$$

we apply Lemma 2.4 and eliminate the variables by the system $\phi = \phi_x = \phi_y = \text{Hess}(\phi) = K(\phi) = 0$. Notice that y is nonzero. First, by $\phi_x = 0$, we can get C as

$$C = -\frac{2x}{y}.$$

Therefore the system is reduced as

$$\left\{ \begin{array}{l} (1) \quad 1 + Ay - x^2y + By^2 + Dy^3 + y^4 = 0, \\ (2) \quad A - 3x^2 + 2By + 3Dy^2 + 4y^3 = 0, \\ (3) \quad 4By - 12x^2 + 12Dy^2 + 24y^3 = 0, \\ (4) \quad -x^2 + Dy^2 + 4y^3 = 0. \end{array} \right.$$

Secondly, by equation (4), we obtain

$$x^2 = y^2(D + 4y).$$

Then the system is reduced as

$$\left\{ \begin{array}{l} (1') \quad 1 + Ay - 3y^4 + By^2 = 0, \\ (2') \quad A - 8y^3 + 2By = 0, \\ (3') \quad -B + 6y^2 = 0. \end{array} \right.$$

Thirdly, by equation (3'), we can get B as

$$B = 6y^2.$$

Then the system is reduced as

$$\left\{ \begin{array}{l} (1'') \quad 1 + Ay + 3y^4 = 0, \\ (2'') \quad A + 4y^3 = 0. \end{array} \right.$$

Hence we obtain $A = -4y^3$ and then the equation

$$y^4 - 1 = 0.$$

The solution is

$$(A, B, C, D, x, y) = (-4y_0^3, 6y_0^2, -2x_0/y_0, D, x_0, y_0),$$

where y_0 is a solution of $y^4 - 1 = 0$ and x_0 is a solution of $x^2 = y_0^2(D + 4y_0)$.

Next, we check that the above ϕ has only one singularity and it is a tacnode. Notice that the curve V_ϕ defined by ϕ has a tacnode at $(x_0, y_0) \in \mathbb{C}^2$. Let $(s, t) \in \mathbb{C}^2$ be a singular point of V_ϕ . Then we solve $\phi(s, t) = \phi_x(s, t) = \phi_y(s, t) = 0$ and we check that the solution is only $(s, t) = (x_0, y_0)$. That is, the singularity of ϕ is only one point and is a tacnode.

(IV) We consider the following polynomial

$$\phi := 1 + Ay + By^2 + Cy^3 + y^4 + c_{11}xy + c_{13}xy^3 + x^2y^2 \in \mathcal{F}(\hat{\Delta}_{IV}).$$

Note that c_{11}, c_{13} are both zero because of the form of the polynomials derived by Lemma 3.6 (IV).

For the polynomial

$$\phi := 1 + Ay + By^2 + Cy^3 + y^4 + x^2y^2 \in \mathcal{F}(\hat{\Delta}_{IV}),$$

we eliminate the variables by the system $\phi = \phi_x = \phi_y = \text{Hess}(\phi) = K(\phi) = 0$ by Lemma 2.4. Notice that y is nonzero. First, by $\phi_x = 0$, we obtain $x = 0$. Therefore the system is reduced as

$$\begin{cases} (1) & 1 + Ay + By^2 + Cy^3 + y^4 = 0, \\ (2) & A + 2By + 3Cy^2 + 4y^3 = 0, \\ (3) & B + 3Cy + 6y^2 = 0, \\ (4) & C + 4y = 0. \end{cases}$$

Secondly, by equation (4), we obtain

$$C = -4y.$$

Then the system is reduced as

$$\begin{cases} (1') & 1 + Ay + By^2 - 3y^4 = 0, \\ (2') & A + 2By - 8y^3 = 0, \\ (3') & B - 6y^2 = 0. \end{cases}$$

Thirdly, by equation (3'), we can get B as

$$B = 6y^2.$$

Then the system is reduced as

$$\begin{cases} (1'') & 1 + Ay + 3y^4 = 0, \\ (2'') & A + 4y^3 = 0. \end{cases}$$

Hence we obtain $A = -4y^3$ and then the equation

$$y^4 - 1 = 0.$$

The solution is

$$(A, B, C, x, y) = (-4y_0^3, 6y_0^2, -4y_0, 0, y_0),$$

where y_0 is a solution of $y^4 - 1 = 0$.

Next, we check that the above ϕ has only one singularity and it is a tacnode. Notice that the curve V_ϕ defined by ϕ has a tacnode at $(0, y_0) \in \mathbb{C}^2$. Let $(s, t) \in \mathbb{C}^2$ be a singular point of V_ϕ . Then, we solve $\phi(s, t) = \phi_x(s, t) = \phi_y(s, t) = 0$ and check that the solution is only $(s, t) = (0, y_0)$. That is, the singularity of ϕ is only one point and is a tacnode.

(V) In this case, in order to achieve $\phi_{xx} \neq 0$, we exchange the variables x and y in ϕ .

For the polynomial

$$\phi := 1 + Ax + Bxy + Cxy^2 + xy^4 + x^2 \in \mathcal{F}(\hat{\Delta}_V),$$

we eliminate the variables by the system $\phi = \phi_x = \phi_y = \text{Hess}(\phi) = K(\phi) = 0$ by Lemma 2.4. Notice that x is nonzero. First, by $\phi = 0$, we obtain

$$A = -\frac{1 + x^2 + Bxy + Cxy^2 + xy^4}{x}.$$

Therefore the system is reduced as

$$\left\{ \begin{array}{l} (1) \ (x - 1)(x + 1) = 0, \\ (2) \ B + 2Cy + 4y^3 = 0, \\ (3) \ \text{substituting } A \text{ for } \text{Hess}(\phi) = 0, \\ (4) \ \text{substituting } A \text{ for } K(\phi) = 0. \end{array} \right.$$

Secondly, by equation (2), we obtain

$$B = -2y(C + 2y).$$

Then the system is reduced as

$$\left\{ \begin{array}{l} (1') \ (x - 1)(x + 1) = 0, \\ (3') \ 4x(C + 6y^2) = 0, \\ (4') \ 192xy = 0. \end{array} \right.$$

Thirdly, by equation (3'), we can get C as

$$C = -6y^2.$$

The solution is

$$(A, B, C, x, y) = (\mp 2, 0, 0, \pm 1, 0).$$

Next, we check that the above ϕ has only one singularity and it is a tacnode. Suppose that the tacnode is at $(1, 0)$. Let $(s, t) \in \mathbb{C}^2$ be a singular point of V_ϕ . Then we solve $\phi(s, t) = \phi_x(s, t) = \phi_y(s, t) = 0$ and check that the solution is only $(s, t) = (1, 0)$. That is, the singularity of ϕ is only one point and is a tacnode. We can check the condition for the case where the tacnode is at $(-1, 0)$ by the same manner. \square

REMARK 3.8. Among the calculation in this section, there are finitely many polynomials which define 1-tacnodal curves except case (III) in Lemma 3.7. In case (III) in Lemma 3.7, we can get the conclusion without eliminating the variable D . This means that there exists a one-parameter family of deformation patterns which define 1-tacnodal curves.

3.4. Remarks on the polytope Δ_E . By the above discussion, for each tropical 1-tacnodal curve, except case (E), there is a “degenerate model of 1-tacnodal curve” whose tropical amoeba is the tropical 1-tacnodal curve. In this subsection, we discuss what happens in case (E).

Lemma 3.9. *There is NOT a polynomial $f \in \mathcal{F}(\Delta_E)$ which defines a 1-tacnodal curve on $X(\Delta_E)$.*

Proof. We assume that a polynomial

$$f := c_{00} + Ax + c_{20}x^2 + c_{01}y + Bxy + c_{12}xy^2$$

defines a 1-tacnodal curve. Then, since f_{xx} is non-zero, we can apply Lemma 2.4 and obtain $y = -B/2c_{12}$ from $K(f) = 0$. Substituting it for $f_y = c_{01} + Bx + 2c_{12}xy = 0$, we get $c_{01} = 0$, but this is a contradiction. \square

On the other hand, there is a polynomial $f \in \mathcal{F}(\Delta_E)$ which has a Newton degenerate singularity on $X(\sigma) \subset X(\Delta_E)$, where $\sigma \subset \Delta_E$ is the edge of length 2. Actually, we can calculate as follows: Set $P := \Delta_4(1; 2, 1, 1, 1)$, $Q := \Delta_3(0; 2, 1, 1)$. We consider the polynomial

$$f := c_{00} + Ax + c_{01}y + c_{20}x^2 + Bxy + c_{12}xy^2 \in \mathcal{F}(P).$$

By multiplying suitable constants to the variables and the whole polynomial, we can rewrite f as

$$1 + Ax + y + x^2 + Bxy + Cxy^2 \in \mathcal{F}(P).$$

If the curve $V_f \subset X(P)$ defined by f intersects $X(\sigma)$ at two points, we can easily check that these points are smooth points of V_f and the intersection $V_f \cap X(\sigma)$ is transversal. Therefore $V_f \cap X(\sigma)$ is exactly one point. Then, f can be rewritten as follows:

$$f = (\epsilon + x)^2 + y + Bxy + Cxy^2 \in \mathcal{F}(P),$$

where $\epsilon = \pm 1$. Set $(X, Y) := (x + \epsilon, y)$. Then f is rewritten as follows:

$$\tilde{f}(X, Y) := X^2 + BXY + (1 \mp B)Y + CXY^2 \mp CY^2.$$

Thus the most complicated isolated singular point defined by this polynomial at the origin (under the condition that the form of the polynomial does not change) is given as

$$X^2 \pm XY + \frac{1}{4}Y^2 + (\text{higher terms}).$$

More precisely, since the polynomial f is irreducible, the number of interior lattice points of Δ_E is two and the curve defined by f has no singularity more complicated than A_3 , the curve has only a cusp as the singularity.

Applying mechanically refinement arguments in this case, we find that the edge $\Delta_4(1; 2, 1, 1, 1) \cap \Delta_3(0; 2, 1, 1)$ does not correspond a 1-tacnodal curve as follows: By above discussion, the exceptional polytope in this case is $\hat{\Delta}_2$. We only consider the case of $\epsilon = 1$. The other case can be proved by the same argument. According to the explanation of a deformation pattern in Definition 2.6, we set

$$\phi := 1 + A'y + x^2y + B'y^2 + xy^2 + \frac{1}{4}y^3 \in \mathcal{F}(\hat{\Delta}_2).$$

By direct computation, we get $\phi_{xx} \neq 0$. Using Lemma 2.4, we obtain $8B'x = 0$. Both cases $x = 0$ and $B' = 0$ contradict $\phi = 0$.

In [8], it is assumed that each polynomial f_i has only semi-quasi-homogeneous singularity

since the paper only deals with the case of nodal or 1-cuspidal curves. This assumption may not be reasonable in the case of 1-tacnodal curves. Actually, when we list the possible polytopes for 1-tacnodal curves we cannot ignore case (E). This is the reason why this case is included in the definition of tropical 1-tacnodal curves. Note that, in fact, by the above discussion, there is no degenerate model of 1-tacnodal curve corresponding to case (E).

4. Main Result

The main theorem of this paper is the following:

Theorem 4.1. *Let $F \in K[z, w]$ be a polynomial which defines an irreducible 1-tacnodal curve. If the rank of the tropical amoeba T_F defined by F is more than or equal to the number of the lattice points of the Newton polytope of F minus four, then T_F is a tropical 1-tacnodal curve.*

Let F be a polynomial in the assertion, T_F be the tropical amoeba defined by F , whose rank satisfies

$$\#\Delta_{\mathbb{Z}} - 1 \geq \text{rk}(T_F) \geq \#\Delta_{\mathbb{Z}} - 4,$$

and S be the dual subdivision of T_F . We remark that, from the discussion in [8, Section 4], if $\#\Delta_{\mathbb{Z}} - 1 \geq \text{rk}(T_F) \geq \#\Delta_{\mathbb{Z}} - 3$, then T_F is smooth, nodal or 1-cuspidal. Thus, by Remark 4.6, we can assume that the rank of T_F is $\#\Delta_{\mathbb{Z}} - 4$.

From the discussion in [8, Subsection 3.3] and the equality $g(C^{(t)}) = \#\text{Int}\Delta_{\mathbb{Z}} - 2$, we can see that

$$\#\partial\Delta_{\mathbb{Z}} - \#\{V(S) \cap \partial\Delta\} = 0 \text{ or } 1.$$

We decompose the proof into four cases

- (A) S is a TP-subdivision and satisfies $\#\partial\Delta_{\mathbb{Z}} - \#\{V(S) \cap \partial\Delta\} = 0$,
- (B) S is a TP-subdivision and satisfies $\#\partial\Delta_{\mathbb{Z}} - \#\{V(S) \cap \partial\Delta\} = 1$,
- (C) S is NOT a TP-subdivision and satisfies $\#\partial\Delta_{\mathbb{Z}} - \#\{V(S) \cap \partial\Delta\} = 0$,
- (D) S is NOT a TP-subdivision and satisfies $\#\partial\Delta_{\mathbb{Z}} - \#\{V(S) \cap \partial\Delta\} = 1$.

For each case, we remove polytopes which cannot correspond to a 1-tacnodal curve and show that the remaining polytopes are exactly tropical 1-tacnodal curves in Definition 3.1.

To explain the removing process more precisely, we prepare some terminologies.

DEFINITION 4.2. A 2-dimensional polytope P is *1-tacnodal* if there is a polynomial $f \in \mathcal{F}(P)$ which defines a 1-tacnodal curve $V_f \in |D(P)|$ satisfying the conditions (S1) and (S2) in Subsection 3.3.

Let $\sigma := P_1 \cap P_2$ be an edge which is the intersection of 2-dimensional polytopes P_1 and P_2 . The edge σ is *1-tacnodal* if there is a pair of polynomials $(f_1, f_2) \in \mathcal{F}(P_1) \times \mathcal{F}(P_2)$ such that

- their truncation polynomials f_1^σ, f_2^σ on σ are same,
- each of the curves C_1 and C_2 defined by f_1 and f_2 has a smooth point or an isolated singular point at z in $X(\sigma)$,
- there exists a deformation pattern $\phi \in \mathcal{F}(\Delta_z)$ compatible with the above data which defines a 1-tacnodal curve in $X(\Delta_z)$.

It can be seen from the discussion in Subsection 3.3 that the polytopes and the pairs of polytopes appearing in Definition 3.1 are 1-tacnodal. To prove the theorem, for each of cases (A), (B), (C) and (D), we carry out the following arguments.

- (1) Remove configurations of edges and interior lattice points of polytopes which do not exist.
- (2) Classify polytopes that are not 1-tacnodal.
- (3) From the list in (2), remove polytopes which do not have 1-tacnodal edges.

In Subsection 4.1, we prepare lemmata for the non-existence of polytopes in (1), and then prove the theorem for case (A), (B), (C) and (D) in Subsection 4.2, 4.3, 4.4 and 4.5, respectively.

4.1. Auxiliary definitions and lemmata.

Lemma 4.3 (On interior lattice points). (1) *The number of interior lattice points of non-parallel quadrangle whose edges are length 1 is larger than 0.*

(2) *For an integer $m \geq 5$, the number of interior lattice points of an m -gon is larger than 0.*

Proof. (1) If a non-parallel $\Delta_4(0; 1, 1, 1, 1)$ exists, it can be decomposed into two triangles of area $1/2$. Thus, we can map this polytope to

$$\text{Conv}\{(0, 0), (1, 0), (0, 1), (p, q)\}$$

by some isomorphism. Then, from Pick's formula, we obtain

$$\frac{p+q}{2} = 1.$$

Hence $p = q = 1$. This is a parallelogram.

(2) It is obvious from the facts that the minimum pentagon is Δ_{VII} and any m -gon can be decomposed into polytopes including a pentagon. \square

Lemma 4.4 (Non-existence of some polytopes). (1) *Following polytopes do NOT exist:*

$$\Delta_3(1; 2, 2, 1), \quad \Delta_3(1; 3, 1, 1), \quad \Delta_3(0; 2, 2, 1), \quad \Delta_3(0; 3, 2, 1), \quad \Delta_5(0; 2, 1, 1, 1, 1).$$

(2) *There is NO non-parallel quadrangle $\Delta_4(0; 2, 2, 1, 1)$.*

Proof. (1) The first triangle is equivalent to

$$\text{Conv}\{(p, 0), (p+2, 0), (0, q)\}.$$

By Pick's formula, we obtain $q = 5/2$. But this contradicts $q \in \mathbb{Z}$. We can easily check the non-existence of the second, third and fourth triangles. If there exists a pentagon $\Delta_5(0; 2, 1, 1, 1, 1)$, we can split it into two quadrangles $\Delta_4(0; 1, 1, 1, 1)$ and $\Delta'_4(0; 1, 1, 1, 1)$. But these quadrangles are parallelograms by the fact that already proved in Lemma 4.3 (1). Thus the union can not be a pentagon.

(2) If it exists, then the edges of length 2 are either adjacent or in opposite sides. The former case can not occur since a triangle $\Delta_3(0; 2, 2, 1)$ does not exist. In the latter case, we can split it into two triangles $\Delta_3(0; 2, 1, 1)$ and $\Delta'_3(0; 2, 1, 1)$. We can assume that one of the triangles is isomorphic to $\text{Conv}\{(0, 0), (1, 0), (0, 2)\}$ and the common edge is the bottom edge. Then, by Pick's formula and the convexity, the last vertex of $\Delta_4(0; 2, 2, 1, 1)$ must be

(1, -2), but this does not satisfy the required conditions. \square

Lemma 4.5. *For the polytope*

$$P := \text{Conv}\{(0, 0), (2, 0), (0, 1), (2, 1)\},$$

the polynomial

$$f := c_{00} + Ax + c_{20}x^2 + c_{01}y + Bxy + c_{21}x^2y \in \mathcal{F}(P)$$

satisfies $f = f_x = f_y = \text{Hess}(f) = 0$ for some A, B and at some point if and only if

$$c_{21}c_{00} = c_{20}c_{01}.$$

Moreover, if the above condition holds, i.e., $V_f \subset X(\Delta_X)$ has a singularity more complicated than A_2 , then f has the form

$$(y + 1)(x \pm 1)^2$$

up to multiplication of a non-zero constant. In particular, the set of singularities of V_f is non-isolated.

Proof. By direct computation. \square

REMARK 4.6 (FACTS). The following facts can be found in [8, Lemma 4.1], or can be proved by easily computation.

(1) Let $I \geq 0$, $s, t, u \geq 1$ be integers such that

$$0 \leq I + (s - 1) + (t - 1) + (u - 1) \leq 2.$$

For each $(I; s, t, u)$, a triangle $\Delta_3(I; s, t, u)$ is uniquely determined up to $\text{Aff}(\mathbb{Z}^2)$ -equivalence as follows:

$$\begin{aligned} \Delta_3(2; 1, 1, 1) &\simeq \text{Conv}\{(0, 0), (3, 2), (2, 3)\}, \\ \Delta_3(1; 2, 1, 1) &\simeq \text{Conv}\{(0, 0), (2, 0), (1, 2)\}, \\ \Delta_3(1; 1, 1, 1) &\simeq \text{Conv}\{(0, 0), (1, 2), (2, 1)\}, \\ \Delta_3(0; 3, 1, 1) &\simeq \text{Conv}\{(0, 0), (3, 0), (0, 1)\}, \\ \Delta_3(0; 2, 1, 1) &\simeq \text{Conv}\{(0, 0), (2, 0), (0, 1)\}, \\ \Delta_3(0; 1, 1, 1) &\simeq \text{Conv}\{(0, 0), (1, 0), (0, 1)\}. \end{aligned}$$

(2) For integers $I \in \{0, 1\}$, $s, t \geq 1$ such that

$$0 \leq I + 2(s - 1) + 2(t - 1) \leq 2,$$

a parallelogram $\Delta_4^{\text{par}}(I; s, t)$ is uniquely determined up to $\text{Aff}(\mathbb{Z}^2)$ -equivalence as follows:

$$\begin{aligned} \Delta_4^{\text{par}}(1; 1, 1) &\simeq \text{Conv}\{(0, 0), (1, 0), (1, 2), (2, 2)\}, \\ \Delta_4^{\text{par}}(0; 2, 1) &\simeq \text{Conv}\{(0, 0), (2, 0), (0, 1), (2, 1)\}, \\ \Delta_4^{\text{par}}(0; 1, 1) &\simeq \text{Conv}\{(0, 0), (1, 0), (0, 1), (1, 1)\}. \end{aligned}$$

(3) The polytopes in this remark are not 1-tacnodal (By [8, Lemma 4.2] and Lemma 4.5, or

direct computation).

Lemma 4.7 (Describing some polytopes). (1) *Let $I \geq 0$, $s, t, u \geq 1$ be integers such that*

$$I + (s - 1) + (t - 1) + (u - 1) = 3.$$

For each $(I; s, t, u)$, a triangle $\Delta_3(I; s, t, u)$ has the following isomorphisms:

$$\Delta_3(3; 1, 1, 1) \simeq \Delta_I \text{ or } \Delta_{II},$$

$$\Delta_3(2; 2, 1, 1) \simeq \Delta_{III},$$

$$\Delta_3(0; 4, 1, 1) \simeq \Delta_V,$$

$$\Delta_3(0; 2, 2, 2) \simeq \text{Conv}\{(0, 0), (2, 0), (0, 2)\}.$$

(2) *A quadrangle $\Delta_4(0; 2, 1, 1, 1)$ is uniquely determined as $\text{Conv}\{(0, 0), (2, 0), (0, 1), (1, 1)\}$ up to $\text{Aff}(\mathbb{Z}^2)$ -equivalence.*

Proof. (1) These claims, except the last case, are the same as Lemma 3.2. We prove the last one. Without loss of generality, the polytope can be assumed to be

$$\text{Conv}\{(p, 0), (p + 2, 0), (0, q)\}.$$

From Pick's formula, we obtain $q = 2$ and $p = 2k$ for some $k \in \mathbb{Z}$. Thus, by the isomorphism

$$\begin{pmatrix} 1 & k \\ 0 & 1 \end{pmatrix} : \mathbb{Z}^2 \rightarrow \mathbb{Z}^2,$$

it is mapped to the polytope $\text{Conv}\{(0, 0), (2, 0), (0, 2)\}$.

(2) We can split $P = \Delta_4(0; 2, 1, 1, 1)$ into two polytopes Q, R which are either

- $Q = \Delta_3(0; 2, 1, 1)$, $R = \Delta_3(0; 1, 1, 1)$ and these polytopes share an edge of length 1, or
- $Q = \Delta_3(0; 1, 1, 1)$, $R = \Delta_4(0; 1, 1, 1, 1)$ and these polytopes share an edge of length 1.

In the former case, we can assume that Q is

$$\text{Conv}\{(0, 0), (1, 0), (0, 2)\}$$

and the common edge is its bottom edge. Then the last vertex of P must be $(1, -1)$. In the latter case, we can assume that R is

$$\text{Conv}\{(0, 0), (1, 0), (0, 1), (1, 1)\}$$

and the common edge is its bottom edge. Then the last vertex of P must be either $(0, -1)$, or $(1, -1)$. All of them are equivalent to

$$\text{Conv}\{(0, 0), (2, 0), (0, 1), (1, 1)\}.$$

□

Lemma 4.8 (Non-1-tacnodal polytopes). *The following polytopes are not 1-tacnodal polytopes:*

- (1) $\Delta_3(0; 2, 2, 2)$,
- (2) $\Delta_3(0; 4, 1, 1)$,
- (3) $\Delta_3(2; 2, 1, 1)$,

- (4) $\Delta_4(0; 2, 1, 1, 1)$,
 (5) $\text{Conv}\{(1, 0), (0, 1), (2, 1), (1, 3)\}$.

Proof. (1) This is by the fact that the Milnor number of an isolated singularity of a projective conic does not exceed 1.

(2) Notice that this polytope is isomorphic to $\text{Conv}\{(0, 0), (0, 1), (4, 0)\}$. Then a polynomial f with this Newton polytope has no singularity since f_y is a non-zero constant.

(3) We assume that a polynomial

$$f := 1 + Ax + x^2 + Bxy + Cxy^2 + xy^3 \in \mathcal{F}(\Delta_3(2; 2, 1, 1))$$

satisfies the condition (S1). Since the polynomial f satisfies $f_{xx} \neq 0$, the system $f = f_x = f_y = \text{Hess}(f) = K(f) = 0$ must have a solution. But, we obtain $K(f) = 48x$. This is a contradiction.

(4) Notice that this polytope is isomorphic to $\text{Conv}\{(0, 0), (2, 0), (0, 1), (1, 1)\}$. We set a polynomial f as

$$f := c_{00} + Ax + c_{20}x^2 + c_{01}y + c_{11}xy \in \mathcal{F}(\text{Conv}\{(0, 0), (2, 0), (0, 1), (1, 1)\}).$$

The Hessian of f is $-c_{11}^2 \neq 0$.

(5) We assume that a polynomial

$$f := c_{10}x + c_{01}y + Axy + c_{21}x^2y + Bxy^2 + c_{13}xy^3$$

defines a 1-tacnodal curve. Then, since f_{xx} is non-zero, we can apply Lemma 2.4 and obtain

$$4c_{01}x(c_{13}y^3 + c_{10}) = -4c_{01}c_{13}xy^3 = 0.$$

This equation and $f(x, y) = 0$ imply $(x, y) = (0, 0)$, but we have $f_x(0, 0) = c_{10} \neq 0$. This is a contradiction. \square

Set

$$\hat{\Delta}_1 = \text{Conv}\{(2, 0), (0, 1), (0, -1)\},$$

$$\hat{\Delta}_2 = \text{Conv}\{(2, 0), (0, 2), (0, -1)\},$$

$$\hat{\Delta}_3 = \text{Conv}\{(3, 0), (0, 1), (0, -1)\},$$

see Figure 4.

Lemma 4.9 (Non-1-tacnodal edges). *The following edges σ are not 1-tacnodal edges:*

- (1) the edge $\Delta_3(0; 2, 1, 1) \cap \Delta_3(0; 2, 1, 1)$ of length 2,
- (2) the edge $\Delta_3(1; 2, 1, 1) \cap \Delta_3(0; 2, 1, 1)$ of length 2 and the edge $\Delta_3(1; 2, 1, 1) \cap \Delta_4(0; 2, 1, 1, 1)$ of length 2,
- (3) the edge $\Delta_3(0; 3, 1, 1) \cap \Delta_3(0; 3, 1, 1)$ of length 3,
- (4) the edge $\Delta_4(0; 2, 1, 1, 1) \cap \Delta_3(0; 2, 1, 1)$ of length 2,
- (5) the edge $\Delta_4(0; 2, 1, 1, 1) \cap \Delta_4(0; 2, 1, 1, 1)$ of length 2,
- (6) the edge $\Delta_4^{\text{par}}(0; 2, 1) \cap \Delta_3(0; 2, 1, 1)$ of length 2 and the edge $\Delta_4^{\text{par}}(0; 2, 1) \cap \Delta_4(0; 2, 1, 1, 1)$ of length 2,
- (7) the edge $\Delta_3(0; 2, 2, 2) \cap \Delta_3(0; 2, 1, 1)$ of length 2.




Fig.4. Polytopes $\hat{\Delta}_1, \hat{\Delta}_2$ and $\hat{\Delta}_3$. The notation \triangle means a lattice point on the boundary which is not a vertex and the notation \star means an interior lattice point.

Proof. The assertion for cases (1), (3), (4) and (5) are already proved in [8, Lemma 3.9, 3.10 and 4.4]. Here we only prove (2), (6) and (7).

(2) Set $P := \Delta_3(1; 2, 1, 1), Q := \Delta_3(0; 2, 1, 1)$. It is easy to check that a curve in $|D(P)|$ cannot have an isolated singularity more complicated than A_1 . Also, we can easily check that if a curve V_f intersects $X(\sigma)$ at two points then the points are smooth points of V_f and those intersections are transversal. Therefore we can set

$$f := (x + \epsilon)^2 + Axy + xy^2 \in \mathcal{F}(P),$$

where $\epsilon = \pm 1$ and suppose that f defines a curve which has an A_1 -singularity on $X(\sigma) \subset X(P)$. With a simple calculation, we obtain $A = 0$. The polynomial corresponding to the polytope Q becomes

$$f' := (x + \epsilon)^2 + y \in \mathcal{F}(Q).$$

Then the exceptional polytope in this case is $\hat{\Delta}_2$. According to the explanation of a deformation pattern in Definition 2.6, we set

$$\phi := 1 + A'y + \epsilon x^2y + B'y^2 + y^3 \in \mathcal{F}(\hat{\Delta}_2).$$

In the case $\epsilon = 1$, we get $\phi_{xx} \neq 0$ by $y \neq 0$. Using Lemma 2.4, we obtain $48y^3 = 0$, but this is a contradiction. We also have a contradiction in the case $\epsilon = -1$.

(6) Set $P := \Delta_4^{\text{par}}(0; 2, 1), Q := \Delta_3(0; 2, 1, 1)$. For P , we set

$$f := (\epsilon + x)^2 + (1 + Ax + x^2)y \in \mathcal{F}(P),$$

where $\epsilon = \pm 1$. Then a polynomial corresponding to Q must be

$$f' := (\epsilon + x)^2 + y \in \mathcal{F}(Q).$$

Then the exceptional polytope in this case is $\hat{\Delta}_1$.

If $\epsilon = 1$, ϕ is given as

$$\phi := 1 + x^2y + y^2 + A'y \in \mathcal{F}(\hat{\Delta}_1),$$

and we can easily check that the solution of the system $\phi = \phi_x = \phi_y = \text{Hess}(\phi) = 0$ does not exist. The case $\epsilon = -1$ can be proved by the same argument.

(7) Set $P := \Delta_3(0; 2, 2, 2)$, $Q := \Delta_3(0; 2, 1, 1)$. Without loss of generality, we can assume that P and Q are

$$P = \text{Conv}\{(0, 0), (2, 0), (0, 2)\}, \quad Q = \text{Conv}\{(0, 0), (2, 0), (0, -1)\}.$$

For P , we set

$$f := 1 + 2\epsilon x + x^2 + By + y^2 + Cxy \in \mathcal{F}(P),$$

where $\epsilon = \pm 1$. Applying the new coordinates $(X, Y) := (x + \epsilon, y)$ for f , we obtain

$$f = X^2 + (B - C\epsilon)Y + CXY + Y^2.$$

Notice that, if $\text{Hess}(f) = C^2 - 4 = 0$, f defines a line of multiplicity 2, that is, f has non-isolated singularity. Therefore we may assume $C^2 - 4 \neq 0$. If $B - C\epsilon \neq 0$, the exceptional polytope in this case is $\hat{\Delta}_1$. If $B - C\epsilon = 0$, then $(\epsilon, 0) \in \mathbb{C}^2$ is an A_1 -singularity, i.e., f has the form $f = X^2 + CXY + Y^2$. Hence, the exceptional polytope in this case is $\hat{\Delta}_2$. The conclusion is derived by the same calculation as in (7) for the former case and in (2) for the latter case, respectively. □

REMARK 4.10 (ON AN EDGE OF LENGTH 1). Let Δ_1, Δ_2 be polytopes such that their intersection $\sigma := \Delta_1 \cap \Delta_2$ is an edge of length 1. The edge σ is NOT an 1-tacnodal edge. Actually, we can prove it as follows: For integers $m_1, m_2 > 0$ and the triangle

$$\hat{\Delta} := \text{Conv}\{(1, 0), (0, m_1), (0, -m_2)\},$$

a polynomial $\phi \in \mathcal{F}(\hat{\Delta})$ can be given as

$$\phi = 1 + \psi(y) + xy^{m_2},$$

where $\psi \in \mathbb{C}[y]$ is a polynomial in y which satisfies $\psi(0) = 0$. If the polynomial ϕ defines a singular curve, then $\phi = \phi_x = \phi_y = 0$ at the singular point. By $\phi_x = y^{m_2} = 0$, the singular point satisfies $y = 0$. However it satisfies $\phi(x, 0) \neq 0$ and this is a contradiction. Therefore, any deformation pattern cannot define a 1-tacnodal curve.

To prevent complication of the proof of the main theorem, we give the following auxiliary definition.

DEFINITION 4.11. The notation \mathbb{T}_{-1} means the set of polytopes equivalent to $\Delta_3(1; 1, 1, 1)$ and pairs of polytopes equivalent to the pair of $\Delta_3(0; 2, 1, 1)$ and $\Delta'_3(0; 2, 1, 1)$ such that their intersection $\Delta_3(0; 2, 1, 1) \cap \Delta'_3(0; 2, 1, 1)$ is a segment of length 2.

The notation \mathbb{T}_{-2} means the set of polytopes equivalent to $\Delta_3(2; 1, 1, 1)$ and pairs of polytopes equivalent to either

- the pair of $\Delta_3(1; 2, 1, 1)$ and $\Delta_3(0; 2, 1, 1)$ such that their intersection $\Delta_3(1; 2, 1, 1) \cap \Delta_3(0; 2, 1, 1)$ is a segment of length 2,
- the pair of $\Delta_3(0; 3, 1, 1)$ and $\Delta'_3(0; 3, 1, 1)$ such that their intersection $\Delta_3(0; 3, 1, 1) \cap \Delta'_3(0; 3, 1, 1)$ is a segment of length 3.

The triple $\Delta_3(0; 2, 2, 1)$, $\Delta_3(0; 2, 1, 1)$ and $\Delta'_3(0; 2, 1, 1)$ such that the intersections $\Delta_3(0; 2, 2, 1) \cap \Delta_3(0; 2, 1, 1)$ and $\Delta_3(0; 2, 2, 1) \cap \Delta'_3(0; 2, 1, 1)$ are segments of length 2 does

not exist by Lemma 4.4.

Note that, by Remark 4.6 and Lemma 4.9, these polytopes and their sharing edges are not 1-tacnodal.

4.2. Case (A). Let S be the dual subdivision of T_F . We assume that S is a TP-subdivision and satisfies $\#\partial\Delta_{\mathbb{Z}} - \#\!(V(S) \cap \partial\Delta) = 0$. Then $d(S) = 0$ by Lemma 2.3. Thus

$$\text{rk}(T_F) = \text{rk}_{\text{exp}}(T_F) = \#\Delta_{\mathbb{Z}} - 4.$$

By the definition of $\text{rk}_{\text{exp}}(T_F)$, we get

$$\begin{aligned} \#\Delta_{\mathbb{Z}} - 4 &= \#\!V(S) - 1 - \sum_{k=1}^N (\#\!V(\Delta_k) - 3) \\ &= \#\!V(S) - 1 - N'_4. \end{aligned}$$

Since $\#\!V(S) \leq \#\Delta_{\mathbb{Z}}$, we obtain $0 \leq N'_4 \leq 3$.

(A-0) If S satisfies $N'_4 = 0$, then it satisfies $\#\!V(S) = \#\Delta_{\mathbb{Z}} - 3$ and consists of triangles.

Then, the subdivision S must contain exactly one of the following polytopes:

- (i) $\Delta_3(3; 1, 1, 1)$,
- (ii) $\Delta_3(2; 1, 1, 1)$ with one of \mathbb{T}_{-1} ,
- (iii) $\Delta_3(2; 2, 1, 1)$ and $\Delta_3(0; 2, 1, 1)$ such that their intersection is a segment whose length is 2,
- (iv) $\Delta_3(1; 2, 1, 1)$ and $\Delta_3(1; 2, 1, 1)$ such that their intersection is a segment whose length is 2,
- (v) $\Delta_3(1; 2, 1, 1)$ and $\Delta_3(0; 2, 1, 1)$ such that their intersection is a segment whose length is 2 with one of \mathbb{T}_{-1} ,
- (vi) $\Delta_3(1; 2, 2, 1)$, $\Delta_3(0; 2, 1, 1)$ and $\Delta'_3(0; 2, 1, 1)$ such that their intersections $\Delta_3(1; 2, 2, 1) \cap \Delta_3(0; 2, 1, 1)$ and $\Delta_3(1; 2, 2, 1) \cap \Delta'_3(0; 2, 1, 1)$ are segments whose lengths are 2,
- (vii) $\Delta_3(0; 2, 2, 1)$, $\Delta_3(0; 2, 1, 1)$ and $\Delta_3(1; 2, 1, 1)$ such that their intersections $\Delta_3(0; 2, 2, 1) \cap \Delta_3(0; 2, 1, 1)$ and $\Delta_3(0; 2, 2, 1) \cap \Delta_3(1; 2, 1, 1)$ are segments whose lengths are 2,
- (viii) $\Delta_3(0; 2, 2, 2)$, $\Delta_3(0; 2, 1, 1)$, $\Delta'_3(0; 2, 1, 1)$ and $\Delta''_3(0; 2, 1, 1)$ such that their intersections $\Delta_3(0; 2, 2, 2) \cap \Delta_3(0; 2, 1, 1)$, $\Delta_3(0; 2, 2, 2) \cap \Delta'_3(0; 2, 1, 1)$ and $\Delta_3(0; 2, 2, 2) \cap \Delta''_3(0; 2, 1, 1)$ are segments whose lengths are 2,
- (ix) $\Delta_3(1; 3, 1, 1)$ and $\Delta_3(0; 3, 1, 1)$ such that their intersection $\Delta_3(1; 3, 1, 1) \cap \Delta_3(0; 3, 1, 1)$ is a segment whose length is 3,
- (x) $\Delta_3(0; 3, 1, 1)$ and $\Delta'_3(0; 3, 1, 1)$ such that their intersection $\Delta_3(0; 3, 1, 1) \cap \Delta'_3(0; 3, 1, 1)$ is a segment whose length is 3, with one of \mathbb{T}_{-1} ,
- (xi) $\Delta_3(0; 3, 2, 1)$, $\Delta_3(0; 3, 1, 1)$ and $\Delta_3(0; 2, 1, 1)$ such that their intersections $\Delta_3(0; 3, 2, 1) \cap \Delta_3(0; 3, 1, 1)$ and $\Delta_3(0; 3, 2, 1) \cap \Delta_3(0; 2, 1, 1)$ are segments whose lengths are 3 and 2, respectively,
- (xii) $\Delta_3(0; 4, 1, 1)$ and $\Delta'_3(0; 4, 1, 1)$ such that their intersection $\Delta_3(0; 4, 1, 1) \cap \Delta'_3(0; 4, 1, 1)$ is a segment whose length is 4,
- (xiii) three of \mathbb{T}_{-1} ,
- (xiv) one of \mathbb{T}_{-2} and one of \mathbb{T}_{-1} .

(A-1) If S satisfies $N'_4 = 1$, then it satisfies $\#V(S) = \#\Delta_{\mathbb{Z}} - 2$ and contains only one parallelogram in the following list and the rest of S consists of triangles:

- (i) $\Delta_4^{\text{par}}(2; 1, 1)$,
- (ii) $\Delta_4^{\text{par}}(0; 2, 1)$, $\Delta_3(0; 2, 1, 1)$ and $\Delta'_3(0; 2, 1, 1)$ such that their intersections $\Delta_4^{\text{par}}(0; 2, 1) \cap \Delta_3(0; 2, 1, 1)$ and $\Delta_4^{\text{par}}(0; 2, 1) \cap \Delta'_3(0; 2, 1, 1)$ are segments whose lengths are 2,
- (iii) $\Delta_4^{\text{par}}(1; 1, 1)$ with one of \mathbb{T}_{-1} ,
- (iv) $\Delta_4^{\text{par}}(0; 1, 1)$ with two of \mathbb{T}_{-1} ,
- (v) $\Delta_4^{\text{par}}(0; 1, 1)$ with one of \mathbb{T}_{-2} .

(A-2) If S satisfies $N'_4 = 2$, then it satisfies $\#V(S) = \#\Delta_{\mathbb{Z}} - 1$ and contains exactly two parallelograms in the following list and the rest of S consists of triangles:

- (i) $\Delta_4^{\text{par}}(1; 1, 1)$, $\Delta_4^{\text{par}}(0; 1, 1)$
- (ii) two $\Delta_4^{\text{par}}(0; 1, 1)$ with one of \mathbb{T}_{-1} .

(A-3) If S satisfies $N'_4 = 3$, then $\#V(S) = \#\Delta_{\mathbb{Z}}$ holds and S contains exactly three parallelograms. Thus S has three $\Delta_4^{\text{par}}(0; 1, 1)$ and the rest of S consists of triangles whose area is $1/2$.

In the above list, by Remark 4.6 and Lemma 4.4, cases (vi), (vii), (ix), (xi) in (A-0) does NOT occur. Furthermore, the following cases do NOT have a regular singularity by Lemma 4.8:

- (ii), (v), (viii), (x), (xiii), (xiv) in (A-0),
- (ii), (iii), (iv), (v) in (A-1),
- (i), (ii) in (A-2),
- (A-3).

Among them, the refinement of the following cases do NOT have an irregular singularity by Lemma 4.9 and Remark 4.10:

- (ii), (v), (viii), (x), (xiii), (xiv) in (A-0),
- (ii), (iii), (iv), (v) in (A-1),
- (i), (ii) in (A-2),
- (A-3).

The remaining cases are (i), (iii), (iv) and (xii) in (A-0) and (i) in (A-1), and they correspond to the polytopes “ Δ_I or Δ_{II} ”, Δ_{III} , Δ_{IV} , Δ_V and Δ_{VI} , respectively, by Lemma 3.2. Moreover, by Lemma 3.4, 3.5, 3.6 and 3.7, these polytopes are 1-tacnodal.

4.3. Case (B). We assume that S is a TP-subdivision and satisfies $\#\partial\Delta_{\mathbb{Z}} - \#(V(S) \cap \partial\Delta) = 1$. By the latter condition, S must have exactly one polytope $P \in S$ such that $P \cap \partial\Delta$ is a segment of length 2. By Lemma 2.3, we get

$$\text{rk}(T_F) = \text{rk}_{\text{exp}}(T_F) = \#\Delta_{\mathbb{Z}} - 4.$$

By the definition of $\text{rk}_{\text{exp}}(T_F)$, we obtain

$$\begin{aligned} \#\Delta_{\mathbb{Z}} - 4 &= \#V(S) - 1 - \sum_{k=1}^N (\#V(\Delta_k) - 3) \\ &= \#V(S) - 1 - N'_4. \end{aligned}$$

Since $\#V(S) \leq \#\Delta_{\mathbb{Z}} - 1$, we have $0 \leq N'_4 \leq 2$.

- (B-0) If S satisfies $N'_4 = 0$, then S satisfies $\#V(S) = \#\Delta_{\mathbb{Z}} - 3$ and consists of triangles. Let $P \in S$ be a polytope which intersects $\partial\Delta$ as a segment of length 2. Then S satisfies one of the following:
- (i) $P = \Delta_3(0; 2, 1, 1)$ and S contains two of \mathbb{T}_{-1} or one of \mathbb{T}_{-2} ,
 - (ii) $P = \Delta_3(1; 2, 1, 1)$ and S contains one of \mathbb{T}_{-1} ,
 - (iii) $P = \Delta_3(2; 2, 1, 1)$,
 - (iv) $P = \Delta_3(0; 2, 2, 2)$,
 - (v) $P = \Delta_3(0; 2, 2, 1)$, and S contains one of \mathbb{T}_{-1} ,
 - (vi) $P = \Delta_3(1; 2, 2, 1)$,
 - (vii) $P = \Delta_3(0; 2, 3, 1)$.
- (B-1) If S satisfies $N'_4 = 1$, then S satisfies $\#V(S) = \#\Delta_{\mathbb{Z}} - 2$. Let $P \in S$ be a polytope which intersects $\partial\Delta$ as a segment of length 2. Then S satisfies one of the following:
- (i) $P = \Delta_4^{\text{par}}(0; 2, 1)$ and $\Delta_3(0; 2, 1, 1)$ such that their intersection $P \cap \Delta_3(0; 2, 1, 1)$ is a segment of length 2,
 - (ii) $P = \Delta_3(0; 2, 1, 1)$ and S contains $\Delta_4^{\text{par}}(1; 1, 1)$,
 - (iii) $P = \Delta_3(1; 2, 1, 1)$ and S contains $\Delta_4^{\text{par}}(0; 1, 1)$,
 - (iv) $P = \Delta_3(0; 2, 2, 1)$ and S contains $\Delta_4^{\text{par}}(0; 1, 1)$,
 - (v) $P = \Delta_3(0; 2, 1, 1)$ and S contains $\Delta_4^{\text{par}}(0; 1, 1)$, and one of \mathbb{T}_{-1} .
- (B-2) If S satisfies $N'_4 = 2$, then S satisfies $\#V(S) = \#\Delta_{\mathbb{Z}}$ and contains exactly two parallelograms. Thus $P = \Delta_3(0; 2, 1, 1)$ and S contains two $\Delta_4^{\text{par}}(0; 1, 1)$.

In the above list, by Lemma 4.4, the following cases do NOT occur:

- (v), (vi), (vii) in (B-0),
- (iv) in (B-1).

Furthermore, the following cases do NOT have a regular singularity by Remark 4.6 and Lemma 4.8:

- (i), (ii), (iv) in (B-0),
- (i), (ii), (iii), (v) in (B-1),
- (B-2).

Among them, (iv) in (B-0) does NOT have an irregular singularity by Lemma 4.9 and the other polytopes except (iii) in (B-0) also do NOT have it since they have only one edge of length more than 1, which should be on the boundary $\partial\Delta$, and this edge cannot be a 1-tacnodal edge. The remaining case is (iii) in (B-0) and this corresponds to the polytope Δ_{III} by Lemma 3.2. Moreover, by Lemma 4.8 (3), this polytope is NOT 1-tacnodal.

4.4. Case (C). We assume that S is NOT a TP-subdivision. Then

$$\begin{aligned} d(S) &= \#\Delta_{\mathbb{Z}} - 4 - \{\#V(S) - 1 - \sum_{k=1}^N (\#V(\Delta_k) - 3)\} \\ &= \#\Delta_{\mathbb{Z}} - \#V(S) - 3 + \sum_{k=1}^N (\#V(\Delta_k) - 3) \\ &\geq -3 + \sum_{k=1}^N (\#V(\Delta_k) - 3) \end{aligned}$$

$$= \sum_{m \geq 3} (m - 3)N_m - 3.$$

By $0 \leq d(S) \leq \mathcal{N}_S/2$ due to Lemma 2.3, we get

$$\sum_{m \geq 3} (m - 3)N_m \leq - \sum_{m \geq 2} N'_{2m} + 5 \quad \text{and} \quad \sum_{m \geq 2} N'_{2m} \leq 2.$$

We decompose the proof into the following three cases:

- (C-0) $\sum_{m \geq 2} N'_{2m} = 0$ and $\sum_{m \geq 3} (m - 3)N_m \leq 5$,
 - (C-1) $\sum_{m \geq 2} N'_{2m} = 1$ and $\sum_{m \geq 3} (m - 3)N_m \leq 4$,
 - (C-2) $\sum_{m \geq 2} N'_{2m} = 2$ and $\sum_{m \geq 3} (m - 3)N_m \leq 3$.
- (C-0) In this case, since $N_4 + 2N_5 + 3N_6 + 4N_7 + 5N_8 \leq 5$ and $\sum_{m \geq 2} N'_{2m} = 0$, possible patterns are the following:
- (i) $(N_8, N'_8) = (1, 0)$,
 - (ii) $(N_7, N_4, N'_4) = (1, 0, 0)$ or $(1, 1, 0)$,
 - (iii) $N'_4 = N'_6 = 0$ and $(N_6, N_5, N_4) = (1, 0, 0), (1, 1, 0), (1, 0, 1)$ or $(1, 0, 2)$,
 - (iv) $(N_5, N_4, N'_4) = (2, 0, 0)$ or $(2, 1, 0)$,
 - (v) $(N_5, N_4, N'_4) = (1, N_4, 0)$ for $N_4 = 0, 1, 2, 3$,
 - (vi) $(N_4, N'_4) = (N_4, 0)$ for $N_4 = 1, 2, 3, 4, 5$.

In case (i), since $\mathcal{N}_S = 4$, we get $0 \leq d(S) \leq 2$. On the other hand, any octagon has two or more inner lattice points (Lemma 4.3), so

$$\begin{aligned} d(S) &= \#\Delta_{\mathbb{Z}} - 4 - \{\#V(S) - 1 - 5\} \\ &= \#\Delta_{\mathbb{Z}} - \#V(S) + 2 \\ &\geq 4. \end{aligned}$$

This is a contradiction. Therefore case (i) does not occur. We can prove that the above cases except the cases (v) with $N_4 = 0$ and (vi) with $N_4 = 1, 2$ do NOT occur by the same argument.

Next, we observe the remaining cases.

Case (v) with $N_4 = 0$. S has exactly one pentagon and the rest of S consists of triangles. Then $\text{rk}_{\text{exp}}(S) = \#V(S) - 3$ holds. Therefore, the set $(\Delta \cap \mathbb{Z}^2) \setminus V(S)$ is exactly one lattice point. By Lemma 4.3 (2), the pentagon is $\Delta_5(1; 1, 1, 1, 1, 1)$. This polytope is equivalent to Δ_{VII} by Lemma 3.2 (6). Moreover, by Lemma 3.5, the pentagon is a 1-tacnodal polytope.

Case (vi) with $N_4 = 1$. S has exactly one non-parallel quadrangle and the rest of S consists of triangles. Since $\text{rk}_{\text{exp}}(S) = \#V(S) - 2$, the set $(\Delta \cap \mathbb{Z}^2) \setminus V(S)$ consists of two lattice points. Therefore, a possible non-parallel quadrangle $\Delta_4(I; s, t, u, v)$ is one of the following list:

- (a) $\Delta_4(0; 1, 1, 1, 1)$,
- (b) $\Delta_4(0; 2, 1, 1, 1)$,
- (c) $\Delta_4(0; 2, 2, 1, 1)$,
- (d) $\Delta_4(1; 1, 1, 1, 1)$,
- (e) $\Delta_4(1; 2, 1, 1, 1)$,
- (f) $\Delta_4(2; 1, 1, 1, 1)$.

Cases (a) and (c) do NOT occur by Lemma 4.3 and Lemma 4.4, respectively. The polytopes in (b) and (e) are NOT 1-tacnodal polytopes by (4) of Lemma 4.8 and Lemma 3.9, respectively. Also the polytope in (d) is NOT a 1-tacnodal polytope by [8, Lemma 4.2 (i)]. Notice that, by Remark 4.10, the polytope in (d) does NOT have a 1-tacnodal edge.

By Lemma 3.2, the polytope (f) is equivalent to one of

$$\Delta_{\text{VIII}}, \Delta_{\text{IX}} \text{ and } \text{Conv}\{(1, 0), (0, 1), (2, 1), (1, 3)\}.$$

The polytopes $\Delta_{\text{VIII}}, \Delta_{\text{IX}}$ are 1-tacnodal polytopes by Lemma 3.5. On the other hand, the polytope $\text{Conv}\{(1, 0), (0, 1), (2, 1), (1, 3)\}$ is NOT a 1-tacnodal polytope by Lemma 4.8 (5) and does NOT have a 1-tacnodal edge by Remark 4.10.

If S contains the polytope in (b), since $\text{rk}_{\text{exp}}(S) = \sharp\Delta_{\mathbb{Z}} - 3$, the adjacent polytope which shares the edge of length 2 of $\Delta_4(1; 2, 1, 1, 1)$ must be either $\Delta_3(0; 2, 1, 1)$ or $\Delta_3(1; 2, 1, 1)$. Each of their intersection with $\Delta_4(1; 2, 1, 1, 1)$ is NOT a 1-tacnodal edge by (2) and (4) of Lemma 4.9. Therefore, any edge contained in S is NOT a 1-tacnodal edge.

If S contains the polytope (e), since $\text{rk}_{\text{exp}}(S) = \sharp\Delta_{\mathbb{Z}} - 4 = \text{rk}(S)$, the adjacent polytope which shares the edge of length 2 of $\Delta_4(1; 2, 1, 1, 1)$ must be $\Delta_3(0; 2, 1, 1)$. This is a dual subdivision of a tropical 1-tacnodal curve of type (E).

Case (vi) with $N_4 = 2$. S has exactly two non-parallel quadrangles and the rest of S consists of triangles. Since $\text{rk}_{\text{exp}}(S) = \sharp V(S) - 3$, the set $(\Delta \cap \mathbb{Z}^2) \setminus V(S)$ consists of exactly one lattice point. Therefore S contains $\Delta_4(0; 2, 1, 1, 1)$ and $\Delta'_4(0; 2, 1, 1, 1)$ such that their intersection is a segment whose length is 2. This is because a non-parallel quadrangle must satisfy either “the number of interior lattice points is non-zero” or “the polytope has an edge of length ≥ 2 ”, by Lemma 4.3. These polytopes are NOT 1-tacnodal polytopes by Lemma 4.8. Also their intersection is NOT a 1-tacnodal edge by Lemma 4.9 (5).

(C-1) In this case, since $N_4 + 2N_5 + 3N_6 + 4N_7 \leq 4$ and $\sum_{m \geq 2} N'_{2m} = 1$, the following patterns can occur:

- (i) $(N_6, N'_6, N_4, N'_4) = (1, 1, 0, 0)$ or $(1, 1, 1, 0)$,
- (ii) $(N_6, N'_6, N_4, N'_4) = (1, 0, 1, 1)$
- (iii) $(N_5, N_4, N'_4) = (1, 2, 1)$,
- (iv) $(N_5, N_4, N'_4) = (1, 1, 1)$,
- (v) $(N_4, N'_4) = (N_4, 1)$ for $N_4 = 2, 3, 4$.

However, we can check that the cases, except case (v) with $N_4 = 2$, are impossible by the same argument as in case (i) in (C-0).

We observe case (v) with $N_4 = 2$. S contains a non-parallel quadrangle P and a parallelogram Q , and the rest of S consists of triangles. Notice that, by Lemma 4.3, P must satisfy either “the number of interior lattice points is non-zero” or “the polytope has an edge of length ≥ 2 ”. Since $\text{rk}_{\text{exp}}(S) = \sharp V(S) - 3$, the set $(\Delta \cap \mathbb{Z}^2) \setminus V(S)$ consists of exactly one lattice point. Therefore P and Q must be either

- $P = \Delta_4(1; 1, 1, 1, 1)$ and $Q = \Delta_4^{\text{par}}(0; 1, 1)$, or
- $P = \Delta_4(0; 2, 1, 1, 1)$ and $Q = \Delta_4^{\text{par}}(0; 1, 1)$ such that the edge of length 2 of P intersects the triangle $\Delta_3(0; 2, 1, 1)$.

In both cases, the polytopes are not 1-tacnodal by Lemma 4.8, Lemma 4.9 and Remark 4.10.

(C-2) In this case, since $N_4 + 2N_5 + 3N_6 \leq 3$ and $\sum_{m \geq 2} N'_{2m} = 2$, any possible subdivision satisfies $N_4 = 3$ and $N'_4 = 2$. Since $\mathcal{N}_S = 0$, we get $d(S) = 0$. On the other hand, since $\#V(S) \leq \#\Delta_{\mathbb{Z}} - 1$ by Lemma 4.3,

$$d(S) = \#\Delta_{\mathbb{Z}} - \#V(S) \geq 1.$$

This is a contradiction.

4.5. Case (D). We assume that S is NOT a TP-subdivision and satisfies $\#\partial\Delta_{\mathbb{Z}} - \#(V(S) \cap \partial\Delta) = 1$. By the former condition, we can apply the same argument of (C) to case (D) and obtain the list of possible subdivisions as follows:

- (1) (v) with $N_4 = 0$ in (C-0),
- (2) (vi) with $N_4 = 1$ in (C-0),
- (3) (vi) with $N_4 = 2$ in (C-0),
- (4) (v) with $N_4 = 2$ in (C-1).

Case (1). S has exactly one pentagon and the rest of S consists of triangles. Then $\text{rk}_{\text{exp}}(S) = \#V(S) - 3$ holds. By the boundary condition, the set $(\Delta \cap \mathbb{Z}^2) \setminus V(S)$ is empty. If S contains a triangle P whose intersection with $\partial\Delta$ is an edge of length 2, then, by Lemma 4.3, S does NOT have a pentagon. Therefore, the possible pentagon is $\Delta_5(0; 2, 1, 1, 1, 1)$, whose intersection with $\partial\Delta$ is an edge of length 2. However, the pentagon does NOT exist by Lemma 4.4.

Case (2). S has exactly one non-parallel quadrangle and the rest of S consists of triangles. By $\text{rk}_{\text{exp}}(S) = \#V(S) - 2$ and the boundary condition, the set $(\Delta \cap \mathbb{Z}^2) \setminus V(S)$ consists of one lattice point. Therefore, possible non-parallel quadrangle $\Delta_4(I; s, t, u, v)$ is one of the following list:

- (a) $\Delta_4(0; 1, 1, 1, 1)$,
- (b) $\Delta_4(0; 2, 1, 1, 1)$,
- (c) $\Delta_4(1; 1, 1, 1, 1)$.

Case (a) does NOT occur by Lemma 4.3. The polytope in (b) is NOT a 1-tacnodal polytope by Lemma 4.8 (4). Also the polytope in (c) is NOT a 1-tacnodal polytope by [8, Lemma 4.2 (i)]. Notice that, by Remark 4.10, the polytope in (c) does NOT have a 1-tacnodal edge.

If S contains the polytope in (b), since $\text{rk}_{\text{exp}}(S) = \#\Delta_{\mathbb{Z}} - 4$, the intersection of the quadrangle $\Delta_4(0; 2, 1, 1, 1)$ and $\partial\Delta$ is an edge of length 2. Thus the edge is NOT a 1-tacnodal edge.

Case (3) and (4). S has exactly two non-parallel quadrangles and the rest of S consists of triangles. By $\text{rk}_{\text{exp}}(S) = \#V(S) - 3$ and the boundary condition, the set $(\Delta \cap \mathbb{Z}^2) \setminus V(S)$ is empty. Therefore, such subdivision S does NOT exist by the fact that a non-parallel quadrangle must satisfy either “the number of interior lattice points is non-zero” or “the polytope has an edge of length ≥ 2 ” in Lemma 4.3. Case (4) can be proved by the same argument. \square

REMARK 4.12. As mentioned in the introduction, this research aims to construct the tropical version of enumerative geometry of the 1-tacnodal curves. Therefore, we would like to lift the 1-tacnodal curve from a given degenerate 1-tacnodal curve by patchworking. It is known that there is no obstruction if the singular point is A_1 , and this is still true even if it is A_2 , which can be checked by a numerical criterion of the vanishing of the obstruction constructed by Shustin (See [7, Theorem 4.1], or [8, Lemma 5.4] for a tropical version). But, unfortunately, this criterion does not work if it is A_3 because of the following reason:

We recall a sufficient condition to apply patchworking [8, Lemma 5.5 (ii)], called *transversality*. Let S be the dual subdivision of a tropical curve T , $\Delta_1, \dots, \Delta_N$ be the 2-dimensional polytopes of S and (C_1, \dots, C_N) be a collection of complex curves such that the Newton polytope of the defining polynomial f_i of C_i is $\Delta_i \in S$ and, if $\sigma_{ij} := \Delta_i \cap \Delta_j \neq \emptyset$, $f_i^{\sigma_{ij}} = f_j^{\sigma_{ij}}$.

For an irreducible curve C_k for some $k \in \{1, \dots, N\}$, there is a union Δ_k^- of edges of Δ_k such that C_k satisfies the following inequality:

$$\sum' b(C_k, \xi) + \sum'' \tilde{b}(C_k, Q) + \sum''' ((C_k \cdot X(\sigma)) - \epsilon) < \sum_{\sigma \subset \partial \Delta} (C_k \cdot X(\sigma)),$$

where

- if C has a tacnode, then $b(C, \xi) = 1$ for both branches, if C is locally given by $\{x^{pr} + y^{qr} = 0\}$ for coprime integers p, q , then $\tilde{b}(C, \xi) = p + q - 1$ for each branch,
- \sum' ranges over all local branches ξ of C_k , centered at the points $z \in \text{Sing}(C_k) \cap (\mathbb{C}^*)^2$,
- \sum'' ranges over all local branches Q of C_k , centered at the points $z \in \text{Sing}(C_k) \cap X(\partial \Delta_k)$, and
- \sum''' ranges over all non-singular points z of C_k on $X(\partial \Delta_k)$ with $\epsilon = 0$ if $\sigma \subset \Delta_k^-$ and $\epsilon = 1$ otherwise,

then C_k is transversal.

Let $V \subset X(\Delta_{III})$ be a curve which is constructed in Lemma 3.6. We can easily check

$$\sum' b(V, \xi) = 0, \quad \sum'' \tilde{b}(V, Q) = 4, \quad \sum''' ((V \cdot X(\sigma)) - \epsilon) \geq 0 \quad \text{and} \quad \sum_{\sigma \subset \partial \Delta} (V \cdot X(\sigma)) = 4.$$

Therefore V does not satisfy the above inequality.

ACKNOWLEDGEMENTS. The author appreciates Professor Masaharu Ishikawa and Professor Takeo Nishinou for valuable discussions. He is also grateful to Professor Eugenii Shustin for giving comments on refinements of tropicalization. He is also grateful to the referee for careful reading of the manuscript and for giving useful comments.

References

[1] M. Gross: Tropical geometry and mirror symmetry. CBMS Regional Conference Series in Mathematics **114**, Published for the Conference Board of the Mathematical Sciences, Washington, DC; by the American Mathematical Society, Providence, RI, 2011.

- [2] J. Harris and D. Morrison: *Moduli of curves*, Graduate Texts in Mathematics **187**, Springer-Verlag, New York, 1998.
- [3] M.M. Kapranov: *Amoebas over non-Archimedean fields*, preprint (2000).
- [4] A.G. Kouchnirenko: *Polyédres de Newton et nombres de Milnor*, Invent. Math. **32** (1976), 1–31.
- [5] G. Mikhalkin: *Enumerative Tropical Algebraic Geometry in \mathbb{R}^2* , J. Amer. Math. Soc. **18** (2005), 313–377.
- [6] T. Nishinou and B. Siebert: *Toric degenerations of toric varieties and tropical curves*, Duke. Math. J. **135** (2006), 1–51.
- [7] E. Shustin: *Gluing of singular and critical points*, Topology **37** (1998), 195–217.
- [8] E. Shustin: *A tropical approach to enumerative geometry*, Algebra i Analiz **17** (2005), 170–214.

Mathematical Institute
Tohoku University
Aoba, Sendai, Miyagi, 980–8578
Japan
e-mail: ttaku32@gmail.com