



Title	On multiply transitive groups. VII
Author(s)	Oyama, Tuyosi
Citation	Osaka Journal of Mathematics. 1968, 5(2), p. 155-164
Version Type	VoR
URL	<a href="https://doi.org/10.18910/7692">https://doi.org/10.18910/7692</a>
rights	
Note	

*The University of Osaka Institutional Knowledge Archive : OUKA*

<https://ir.library.osaka-u.ac.jp/>

The University of Osaka

Oyama, T.  
Osaka J. Math.  
5 (1968), 155-164

## ON MULTIPLY TRANSITIVE GROUPS VII

TUYOSI OYAMA

(Received April 16, 1968)

### 1. Introduction

Let  $G$  be a 4-fold transitive group on  $\Omega = \{1, 2, \dots, n\}$ , and let  $P$  be a Sylow 2-subgroup of a stabilizer of four points in  $G$ . If  $P=1$ , then by a theorem of M. Hall [1. Theorem 5.8.1]  $G$  must be one of the following groups:  $S_4$ ,  $S_5$ ,  $A_6$ ,  $A_7$  or  $M_{11}$ . From a recent result of H. Nagao [7] it follows that, if  $P \neq 1$  is semi-regular and leaves exactly four or five points fixed, then  $G$  must be one of the following groups:  $S_6$ ,  $S_7$ ,  $A_8$ ,  $A_9$  or  $M_{12}$ .

The purpose of this paper is to extend the result of H. Nagao. Namely we shall prove the following

**Theorem.** *Let  $G$  be a 4-fold transitive group. If a Sylow 2-subgroup of a stabilizer of four points in  $G$  is semi-regular and not identity, then  $G$  must be  $S_6$ ,  $S_7$ ,  $A_8$ ,  $A_9$ ,  $M_{12}$  or  $M_{23}$ .*

**DEFINITION AND NOTATION.** A permutation  $x$  is called semi-regular if there is no point fixed by  $x$ . A permutation group  $H$  is called semi-regular if every nonidentity element of  $H$  is semi-regular on the points actually moved by  $H$ . For a permutation group  $G$  on  $\Omega$ , let  $G_{i,j,\dots}$  denote the stabilizer of the points  $i, j, \dots, r$  in  $G$ . For a subset  $S$  of  $G$  we denote the normalizer (or centralizer) of  $S$  in  $G$  by  $N_G(S)$  (or  $C_G(S)$ ). Let  $\alpha_i(x)$  denote the number of  $i$ -cycles of a permutation  $x$ . The totality of points left fixed by a set  $X$  of permutations is denoted by  $I(X)$ , and if a subset  $\Delta$  of  $\Omega$  is a fixed block of  $X$ , then the restriction of  $X$  on  $\Delta$  is denoted by  $X^\Delta$ .

### 2. Proof of the theorem

To prove the theorem we may assume that a stabilizer of four points in  $G$  fixes exactly four points (See [6]). In the proof of the theorem, we shall also make use of the fact [1. p. 80] that a 4-fold transitive group of degree less than 35 is one of the known groups.

Let  $P$  be a Sylow 2-subgroup of  $G_{1234}$ . Then  $|I(P)|$  is four, five, six, seven or eleven, and  $N_G(P)^{(\mathcal{P})}$  is  $S_4$ ,  $S_5$ ,  $A_6$ ,  $A_7$  or  $M_{11}$  (cf. [5]. Lemma 1). By the theorem of H. Nagao, we may treat only the last three cases.

Case I.  $|I(P)| = 11$ ,  $N_G(P)^{I(P)} = M_{11}$ .

Let  $a$  be a central involution of  $P$ , and suppose that  $P$  has an involution  $b$  different from  $a$ . Then  $a$  and  $b$  fix the same eleven points and generate a four group. Therefore we may assume that

$$\begin{aligned} a &= (1)(2)\cdots(11)(ij)(kl)\cdots, \\ b &= (1)(2)\cdots(11)(ik)(jl)\cdots. \end{aligned}$$

Then  $\langle a, b \rangle$  normalizes  $G_{ijkl}$  and hence it normalizes some Sylow 2-subgroup  $P'$  of  $G_{ijkl}$ . Since  $a^{I(P')}$  is an involution of  $M_{11}$ , it fixes three points. Now  $I(a) = I(b) = \{1, 2, \dots, 11\}$ . Hence  $b^{I(P')}$  fixes these three points and the four group  $\langle a, b \rangle^{I(P')}$  is contained in a stabilizer of three points of  $M_{11}$ . But this is impossible, because a stabilizer of three points of  $M_{11}$  is a quaternion group.

Thus  $P$  has only one involution, and hence  $P$  must be a cyclic group or a generalized quaternion group [1. Theorem 12.5.2.]. By Theorem 1 in [8]  $P$  is not cyclic, and by the following unpublished result of H. Nagao we have a contradiction.

**Lemma 1.** (H. Nagao) *Let  $G$  be a 4-fold transitive group, and  $P \neq 1$  a Sylow 2-subgroup of  $G_{1234}$ . If  $P$  fixes eleven points, then  $P$  is not a generalized quaternion group.*

Thus we have no group in this case.

Case II.  $|I(P)| = 6$  or  $7$ ,  $N_G(P)^{I(P)} = A_6$  or  $A_7$ .

In the proofs of the following series from i) to v) we assume that  $N_G(P)^{I(P)} = A_6$ , and we need the following result [8. Theorem 2] that any involution of  $G$  fixes exactly six points. The proofs in the case  $N_G(P)^{I(P)} = A_7$  are similar.

- i) *If an element  $a$  of  $G$  has a 4-cycle, then its order is an odd multiply of 4 or 8.*
- 1) *If  $a$  is of order 4, then  $\alpha_2(a)=2$  and  $\alpha_1(a)=2$ . (When  $N_G(P)^{I(P)} = A_7$ ,  $\alpha_2(a)=2$  and  $\alpha_1(a)=3$ ).*
- 2) *If  $a$  is of order  $4t$  with  $t$  odd, then  $\alpha_2(a)=2$  and  $\alpha_1(a)=2$ . (When  $N_G(P)^{I(P)} = A_7$ ,  $\alpha_2(a)=2$  and  $\alpha_1(a)=0$  or 3).*
- 3) *If  $a$  is of order  $8t$  with  $t$  odd, then  $\alpha_4(a)=1$  and  $\alpha_2(a)=1$ .*

Proof. 1) Let  $a$  be an element of order 4. Then we may assume that

$$a = (1\ 2\ 3\ 4)\cdots.$$

Since  $a$  normalizes  $G_{1234}$ ,  $a$  normalizes some Sylow 2-subgroup of  $G_{1234}$ . We may assume that  $a$  normalizes  $P$ . Since  $N_G(P)^{I(P)} = A_6$ ,  $a$  must be of the following form

$$a = (1 \ 2 \ 3 \ 4) (5 \ 6) \cdots,$$

where  $I(P) = \{1, 2, 3, 4, 5, 6\}$ . Since  $a^2$  is an involution and  $\alpha_1(a^2) = 6, \alpha_1(a) = 0, 2$  or  $4$ . If  $\alpha_1(a) = 0$ , then  $\alpha_2(a) = 3$  and  $a$  is of the form

$$a = (1 \ 2 \ 3 \ 4) (5 \ 6) (i_1 j_1) (i_2 j_2) \cdots.$$

From this form  $a$  normalizes some Sylow 2-subgroup  $P'$  of  $G_{56i_1j_1}$  and  $a^{I(P')} = (5 \ 6) (i_1 j_1) (i_2 j_2)$ , which is contrary to  $N_G(P')^{I(P')} = A_6$ . Therefore  $\alpha_1(a) \neq 0$ . Since  $P$  is semi-regular,  $P$  is elementary abelian by Lemma 2 in [8]. Therefore  $\alpha_1(a) \neq 4$ . Hence we have that  $\alpha_1(a) = 2$  and consequently  $\alpha_2(a) = 2$ .

2) Let  $a$  be of order  $4t$  with  $t$  odd, then  $\alpha_2(a^t) = 2$  and  $\alpha_1(a^t) = 2$ . Therefore  $\alpha_2(a) = 2$  and  $\alpha_1(a) = 2$ .

3) Let  $a$  be of order 8. Then from 1)  $\alpha_2(a^2) = 2$ . Hence  $\alpha_4(a) = 1$ . Thus we may assume that  $a$  is of the following form

$$a = (1 \ 2 \ 3 \ 4) \cdots.$$

Then  $a$  normalizes some Sylow 2-subgroup of  $G_{1234}$ . We may assume that  $a$  normalizes  $P$ . Since  $N_G(P)^{I(P)} = A_6$ ,  $a$  must be of the following form

$$a = (1 \ 2 \ 3 \ 4) (5 \ 6) \cdots,$$

where  $I(P) = \{1, 2, 3, 4, 5, 6\}$ . Since  $\alpha_1(a^2) = 2, \alpha_2(a) = 1$ . This is also true for an element of order  $8t$  with  $t$  odd.

Since an element of order 8 has only one 4-cycle,  $G$  has no element of order 16.

ii)  $P$  is an elementary abelian group of order 16.

Proof. By Lemma 2 in [8]  $P$  is elementary abelian. Therefore it suffices to prove that the order of  $P$  is 16. Let  $a = (1 \ 2) (3 \ 4) \cdots$  be an involution of  $G$ . Then  $a$  normalizes a Sylow 2-subgroup of  $G_{1234}$ . We may assume that  $a$  normalizes  $P$ . Let  $I(P) = \{1, 2, 3, 4, 5, 6\}$ . Since  $a^{I(P)}$  must be an even permutation,  $a$  is of the following form

$$a = (1 \ 2) (3 \ 4) (5) (6) \cdots.$$

Let  $a$  fixes the point 7 and let  $\Delta$  be the  $P$ -orbit containing 7. Then  $a$  fixes at most four points of  $\Delta$  and  $P$  is regular on  $\Delta$ . Therefore by Lemma in [4] we have that the order of  $P$  is at most 16.

Now let  $K$  be the kernel of the natural homomorphism  $N_G(P) \rightarrow N_G(P)^{I(P)}$ . Then  $KC_G(P)/C_G(P) \cong K/K \cap C_G(P)$ , and  $N_G(P) \trianglelefteq K \cdot C_G(P) \trianglelefteq K$ . Since  $N_G(P)/K = N_G(P)^{I(P)}$ ,  $N_G(P)/K$  is a simple group. Therefore  $N_G(P) = K \cdot C_G(P)$  or  $K \cdot C_G(P) = K$ . Since  $G_{1234} \geq K \geq K \cap C_G(P) \geq P$  and  $P$  is a Sylow 2-subgroup of  $G_{1234}$ ,  $K/K \cap C_G(P)$  is of odd order. If  $N_G(P) = K \cdot C_G(P)$ , then

from  $KC_G(P)/C_G(P) \cong K/K \cap C_G(P)$ ,  $N_G(P)/C_G(P)$  is of odd order. Hence any 2-element of  $N_G(P)$  belongs to  $C_G(P)$ . On the other hand there is an element  $x$  of order four, which is of the following form

$$x = (1\ 2\ 3\ 4)\cdots.$$

Then  $x$  normalizes  $G_{1234}$ , and hence we may assume that  $x$  normalizes  $P$ . From  $N_G(P)^{I(P)} = A_6$ ,  $x$  must be of the form

$$x = (1\ 2\ 3\ 4)(5\ 6)\cdots,$$

where  $I(P) = \{1, 2, 3, 4, 5, 6\}$ . By 1) of i)  $x$  fixes two points of  $\Omega - I(P)$ . Since  $P$  is semi-regular on  $\Omega - I(P)$ ,  $x$  commutes with exactly two elements of  $P$ . But by Theorem 1 in [8]  $|P| > 2$ . Therefore  $x \notin C_G(P)$ , which is a contradiction. Thus  $K \cdot C_G(P) = K$ .

Now  $N_G(P)/C_G(P)$  is a subgroup of the automorphism group of  $P$ . From  $N_G(P)/K \cong A_6$  and  $K \geq C_G(P)$ , the order of the automorphism group of  $P$  is not smaller than the order of  $A_6$ . Since  $P$  is elementary abelian and its order is at most 16, the order of  $P$  must be 16.

Next we also need a theorem of G. Frobenius (See [3]. Proposition 14.5), which will be stated here as Lemma 2.

**Lemma 2.** (G. Frobenius) *Let  $G \leq S_n$ , then*

$$\sum_{x \in G} \binom{\alpha_1(x)}{\kappa} \binom{\alpha_2(x)}{\lambda} \cdots = \frac{m \cdot |G|}{1^\kappa \cdot \kappa! \cdot 2^\lambda \cdot \lambda! \cdots}.$$

Here  $m$  is an integer obtained in the following way. Let  $\Omega^{(t)} = \{(i_1, \dots, i_\kappa, j_1, j_1', \dots, j_\lambda, j_\lambda', \dots)\}$  be a family of ordered sets consisting of  $t$  ( $= \kappa + 2\lambda + \dots$ ) points of  $\Omega$  such that there is at least one element  $x$  of  $G$  of the form

$$x = (i_1) \cdots (i_\kappa)(j_1 j_1') \cdots (j_\lambda j_\lambda') \cdots.$$

When  $G$  is regarded as a permutation group on  $\Omega^{(t)}$  by setting

$$(a_1, \dots, a_t)^x = (a_1^x, \dots, a_t^x)$$

for  $x \in G$  and  $(a_1, \dots, a_t) \in \Omega^{(t)}$ ,  $m$  is the number of  $G$ -orbits in  $\Omega^{(t)}$ .

iii) *Let  $x$  be an involution of  $N_G(P) - P$ . Then any fixed point of an element ( $\neq 1$ ) of  $\langle x, P \rangle^{\Omega - I(P)}$  is contained in exactly one orbit of  $P$ . The number of  $P$ -orbits in  $\Omega - I(P)$  is odd.*

Proof. Since the order of  $P$  is 16, by Lemma in [4]  $x$  commutes with four elements of  $P$ . Since  $P$  is semi-regular and  $x$  fixes four points of  $\Omega - I(P)$ , these points must be contained in the same  $P$ -orbit, say  $\Delta$ . Put  $Q = \langle x, P \rangle$ . The

order of  $Q$  is 32 and  $Q$  fixes  $\Delta$ . For an element  $a$  of  $P$  if  $x$  commutes with  $a$ , then  $xa$  is of order 2, and if not, then  $xa$  is of order 4, since  $(xax)a$  belongs to  $P$  and it is not the identity. Let  $xa$  be of order 4, then by 1) of i)  $\alpha_1(xa)=2$  and  $xa$  has no fixed point on  $\Omega-I(P)$ . Therefore

$$\begin{aligned}\sum_{y \in Q} \alpha_1(y^\Delta) &= \alpha_1(1^\Delta) + \sum'_{y'} \alpha_1(y'^\Delta) \\ &= 16 + \sum'_{y'} \alpha_1(y'^\Delta),\end{aligned}$$

where  $y'$  ranges over all involutions of  $Q-P$ . On the other hand from Lemma 2

$$\sum_{y \in Q} \alpha_1(y^\Delta) = |Q^\Delta| = 32.$$

Hence  $\sum'_{y'} \alpha_1(y'^\Delta)=16$ . Since  $Q-P$  has four involutions and these involutions have four fixed points in  $\Omega-I(P)$  respectively, these 16 points are all contained in  $\Delta$ . Hence  $Q$  is semi-regular on  $\Omega-I(P) \cup \Delta$ , in which any  $Q$ -orbit contains exactly two  $P$ -orbits. Thus the number of  $P$ -orbits in  $\Omega-I(P)$  is odd.

iv)  $G$  has an element of order 8.

Proof. Let  $a$  be an element of order four and of the following form

$$a = (1\ 2\ 3\ 4)\cdots.$$

Then  $a$  normalizes a Sylow 2-subgroup of  $G_{1234}$ . We may assume that  $a$  normalizes  $P$ . From  $N_G(P)^{I(P)}=A_6$ ,  $a$  must be of the following form

$$a = (1\ 2\ 3\ 4)(5\ 6)\cdots,$$

where  $I(P)=\{1, 2, 3, 4, 5, 6\}$ . By 1) of i)  $a$  fixes two points of  $\Omega-I(P)$ , and these points are contained in a  $P$ -orbit, say  $\Delta$ . Put  $Q=\langle P, a \rangle$ . Then the order of  $Q$  is  $4 \cdot 16$  and  $\Delta$  is a  $Q$ -orbit. Suppose that  $Q$  has no element of order 8. From iii) any fixed point of an element ( $\neq 1$ ) of  $\langle P, a^2 \rangle^{Q-I(P)}$  is contained in  $\Delta$ . Let  $a'$  be any element of  $Pa$  or  $Pa^{-1}$ . Then  $a'$  is of the following form

$$(1\ 2\ 3\ 4)(5\ 6)\cdots \text{ or } (1\ 4\ 3\ 2)(5\ 6)\cdots.$$

We assumed that  $Q$  has no element of order 8. Hence  $a'$  is of order 4, and  $a'$  has exactly two fixed points. Since  $Q/P$  is a cyclic group of order 4,  $a'^2$  belongs to  $\langle P, a^2 \rangle$ . Therefore  $a'$  fixes two points of  $\Delta$ .

From  $Q=P+Pa+Pa^2+Pa^{-1}$

$$\begin{aligned}\sum_{x \in Q} \alpha_1(x^\Delta) &\geq \sum_{x \in P} \alpha_1(x^\Delta) + \sum_{x \in Pa} \alpha_1(a^\Delta) + \sum_{x \in Pa^{-1}} \alpha_1(x^\Delta) \\ &= 16 + 2 \cdot 16 + 2 \cdot 16 = 5 \cdot 16.\end{aligned}$$

On the other hand by Lemma 2

$$\sum_{x \in Q} \alpha_1(x^\Delta) = |Q| = 4 \cdot 16 ,$$

which is a contradiction. Thus  $Q$  has an element of order 8.

Since  $G$  is 4-fold transitive, by Lemma 2

$$\sum_{x \in G} \alpha_4(x) = \frac{1}{4}g ,$$

and

$$\sum_{x \in G} \alpha_2(x) \cdot \alpha_4(x) = \frac{m \cdot g}{2 \cdot 4} ,$$

where  $g = |G|$ . From i) if  $\alpha_4(x) \neq 0$ , then  $\alpha_4(x) \cdot \alpha_2(x) = 2 \cdot \alpha_4(x)$  or  $\alpha_4(x)$ . Since there is an element of order 8, from i) we have

$$\sum_{x \in G} \alpha_4(x) < \sum_{x \in G} \alpha_2(x) \cdot \alpha_4(x) < 2 \cdot \sum_{x \in G} \alpha_4(x)$$

and hence  $1 < \frac{m}{2} < 2$ . Thus  $m = 3$ , and

$$\sum_{x \in G} \alpha_2(x) \cdot \alpha_4(x) = \frac{3}{8}g .$$

From two equations above, we obtain

$$\sum_y' \alpha_4(y) + \sum_{y'}' \alpha_4(y') = \frac{1}{4}g ,$$

$$\sum_y' 2 \cdot \alpha_4(y) + \sum_{y'}' \alpha_4(y') = \frac{3}{8}g ,$$

where  $y$  and  $y'$  range over all elements of order  $4t$  and  $8t$  ( $t$ : odd) respectively. Hence

$$\sum_y' \alpha_4(y) = \frac{1}{8}g .$$

On the other hand

$$\sum_{x \in G} \binom{\alpha_2(x)}{2} \cdot \alpha_4(x) = \frac{m' \cdot g}{2^2 \cdot 2 \cdot 4} .$$

Since an element of order  $8t$  with  $t$  odd has only one 2-cycle, and an element of order  $4t$  with  $t$  odd has two 2-cycles,

$$\sum_{x \in G} \binom{\alpha_2(x)}{2} \cdot \alpha_4(x) = \sum_y' \alpha_4(y) = \frac{1}{8}g ,$$

Therefore

$$\frac{m' \cdot g}{2^2 \cdot 2 \cdot 4} = \frac{1}{8} g,$$

hence

$$m' = 4.$$

From the remark of Lemma 2,  $G$  has four orbits on  $\Omega^{(8)} = \{(i_1, i_2, j_1, j_2, k_1, k_2, k_3, k_4) \mid x = (i_1 i_2)(j_1 j_2)(k_1 k_2 k_3 k_4) \dots \in G, x \text{ is of order 4}\}$ . Since  $G$  is 4-fold transitive on  $\Omega$ ,  $G_{1234} = H$  has four orbits on  $\Omega^{(4)} = \{(k_1, k_2, k_3, k_4) \mid a = (1 2)(3 4)(k_1 k_2 k_3 k_4) \dots \in G, a \text{ is of order 4}\}$ .

When  $H$  is regarded as a permutation group on  $\Omega^{(4)}$ , we denote it by  $H^*$ .

If  $(k_1, k_2, k_3, k_4) \in \Omega^{(4)}$ , then there is an element  $a$  of  $G$  of the form

$$a = (1 2)(3 4)(k_1 k_2 k_3 k_4) \dots.$$

Since  $a^{-1} = (1 2)(3 4)(k_1 k_4 k_3 k_2) \dots \in G$ , we have eight points  $(k_1, k_2, k_3, k_4)$ ,  $(k_2, k_3, k_4, k_1)$ ,  $(k_3, k_4, k_1, k_2)$ ,  $(k_4, k_1, k_2, k_3)$ ,  $(k_1, k_4, k_3, k_2)$ ,  $(k_4, k_3, k_2, k_1)$ ,  $(k_3, k_2, k_1, k_4)$  and  $(k_2, k_1, k_4, k_3)$  of  $\Omega^{(4)}$ .

v) Let  $(k_1, k_2, k_3, k_4) \in \Omega^{(4)}$ . Then  $(k_1, k_2, k_3, k_4)^{H^*}$  and  $(k_2, k_3, k_4, k_1)^{H^*}$  are the different  $H^*$ -orbits.

Proof. Since any 2-element of  $H$  is of order 2,  $H$  has no element as follows:

$$\begin{pmatrix} k_1 & k_2 & k_3 & k_4 & \dots \\ k_2 & k_3 & k_4 & k_1 & \dots \end{pmatrix} = (k_1 k_2 k_3 k_4) \dots.$$

Therefore  $(k_1, k_2, k_3, k_4)^{H^*} \neq (k_2, k_3, k_4, k_1)^{H^*}$ .

From now on we treat two cases  $N_G(P)^{I(P)} = A_6$  and  $A_7$  separately. For the proofs in these cases the result that the number of  $H^*$ -orbits is four is important.

A)  $N_G(P)^{I(P)} = A_6$ .

Let  $I(P) = \{1, 2, 3, 4, 5, 6\}$ . and  $G_{1234} = H$ . Then the points 5 and 6 are contained in  $H$ -orbits of odd length. Put  $5^H = \Delta_1$  and  $6^H = \Delta_2$ .

Suppose that  $\Delta_1 = \Delta_2$ . Since  $P$ -orbits in  $\Omega - I(P)$  are of even length, the length of  $\Delta_1$  is even, which is a contradiction. Therefore  $\Delta_1 \neq \Delta_2$ . Furthermore the other  $H$ -orbits in  $\Omega - I(H)$  are all of even lengths.

From  $N_G(P)^{I(P)} = A_6$  there is an element  $x$  of the following form

$$x = (1 2)(3)(4)(5 6) \dots.$$

Since  $x \in N_G(H)$ ,  $\Delta_1^x = \Delta_2$ . Hence  $|\Delta_1| = |\Delta_2|$ . Suppose that  $H$ -orbits in  $\Omega - I(H)$  are  $\Delta_1$  and  $\Delta_2$ . From iii) the number of  $P$ -orbits in  $\Omega - I(P)$  is odd,

and all  $P$ -orbits in  $\Omega - I(P)$  are of the same length. Hence  $|\Delta_1| \neq |\Delta_2|$ , which is a contradiction. Therefore  $H$  has at least three orbits in  $\Omega - I(H)$ .

If  $|\Delta_1| = |\Delta_2| = 1$ , then  $|I(H)| = 6$ , contradicting the assumption that  $|I(H)| = 4$ . Therefore  $|\Delta_1| = |\Delta_2| > 1$ .

Let  $(k_1, k_2, k_3, k_4)$  be a point of  $\Omega^{(4)}$ . Then there is an element

$$a = (1 2) (3 4) (k_1 k_2 k_3 k_4) \dots$$

in  $G$ . We may assume that  $a \in N_G(P)$  and  $a$  is of the form

$$a = (1 2) (3 4) (5) (6) (k_1 k_2 k_3 k_4) \dots .$$

By the assumption  $a$  is of order four, and from 1) of i) any point in  $\Omega - \{1, 2, \dots, 6\}$  appears in some 4-cycle of  $a$ . Since  $\Delta_1^a = \Delta_1$  and  $\Delta_2^a = \Delta_2$ , we may assume that  $\{i_1, i_2, i_3, i_4\} \subset \Delta_1$  and  $\{j_1, j_2, j_3, j_4\} \subset \Delta_2$ , where  $a = (i_1 i_2 i_3 i_4) (j_1 j_2 j_3 j_4) \dots$ . Then from v)  $(i_1, i_2, i_3, i_4)^{H^*}$ ,  $(i_2, i_3, i_4, i_1)^{H^*}$ ,  $(j_1, j_2, j_3, j_4)^{H^*}$  and  $(j_2, j_3, j_4, j_1)^{H^*}$  are all different  $H^*$ -orbits. Thus we have four  $H^*$ -orbits. But  $H$  has at least three orbits in  $\Omega - I(H)$ . Hence there is a 4-cycle  $(l_1 l_2 l_3 l_4)$  of  $a$  such that  $\{l_1, l_2, l_3, l_4\} \not\subset \Delta_1 \cup \Delta_2$ . Therefore  $(l_1, l_2, l_3, l_4)^{H^*}$  is the different  $H^*$ -orbit from these four  $H^*$ -orbits. Hence we have five  $H^*$ -orbits, which is a contradiction.

Thus we have no group in this case.

B)  $N_G(P)^{I(P)} = A_7$ .

Let  $I(P) = \{1, 2, 3, 4, 5, 6, 7\}$  and  $H = G_{1234}$ . Then from  $N_G(P)^{I(P)} = A_7$ , there is an element

$$x = (1) (2) (3) (4) (5 6 7) \dots .$$

Since  $x \in H$ , three points 5, 6 and 7 belong to the same  $H$ -orbit, say  $\Delta_1$ . Then  $\Delta_1$  is the only  $H$ -orbit in  $\Omega - I(H)$  of odd length. If  $H$  has only one orbit in  $\Omega - I(H)$ , namely,  $H$  is transitive on  $\Omega - I(H)$ , then a stabilizer of one point in  $G$  satisfies the assumption of Case II. A), which is a contradiction. Therefore  $H$  has at least two orbits, say  $\Delta_1$  and  $\Delta_2$ , in  $\Omega - I(H)$ .

Suppose that  $|\Delta_1| > 3$ . Let  $(k_1, k_2, k_3, k_4) \in \Omega^{(4)}$ . Then there is an element

$$a = (1 2) (3 4) (k_1 k_2 k_3 k_4) \dots$$

of  $G$ . By the assumption  $a$  is of order four, and from 1) of i) the cycles of  $a$  are all 4-cycles except two 2-cycles and three 1-cycles. Since  $a \in N_G(H)$ ,  $\Delta_2^a$  is an  $H$ -orbit. Assume that  $\Delta_2^a \neq \Delta_2$ . Since the length of  $\Delta_2$  is even,  $\Delta_2^a \neq \Delta_1$ . We may assume that  $k_1 \in \Delta_2$ , and hence  $k_2 \notin \Delta_1 \cup \Delta_2$ . Then we shall show that  $(k_1, k_2, k_3, k_4)^{H^*}$ ,  $(k_2, k_3, k_4, k_1)^{H^*}$ ,  $(k_1, k_4, k_3, k_2)^{H^*}$  and  $(k_4, k_3, k_2, k_1)^{H^*}$  are all different  $H^*$ -orbits. From v)  $(k_1, k_2, k_3, k_4)^{H^*} \neq (k_2, k_3, k_4, k_1)^{H^*}$  and  $(k_1, k_4, k_3, k_2)^{H^*} \neq (k_4, k_3, k_2, k_1)^{H^*}$ .

If  $(k_1, k_2, k_3, k_4)^{H^*} = (k_1, k_4, k_3, k_2)^{H^*}$ , then  $H$  has a following element

$$x = \begin{pmatrix} k_1 & k_2 & k_3 & k_4 & \cdots \\ k_1 & k_4 & k_3 & k_2 & \cdots \end{pmatrix} = (k_1)(k_3)(k_2 k_4) \cdots .$$

Since the order of  $x$  is even, there is a Sylow 2-subgroup of  $H$  fixing the point  $k_1$ . By the conjugacy of Sylow 2-subgroups of  $H$ ,  $k_1$  must be contained in  $\Delta_1$ , which is a contradiction. Therefore  $(k_1, k_2, k_3, k_4)^{H^*} \neq (k_1, k_4, k_3, k_2)^{H^*}$ . In the same way  $(k_2, k_3, k_4, k_1)^{H^*} \neq (k_4, k_3, k_2, k_1)^{H^*}$ .

If  $(k_2, k_3, k_4, k_1)^{H^*} = (k_1, k_4, k_3, k_2)^{H^*}$ , then  $H$  has a following element

$$\begin{pmatrix} k_2 & k_3 & k_4 & k_1 & \cdots \\ k_1 & k_4 & k_3 & k_2 & \cdots \end{pmatrix} = (k_1 k_2)(k_3 k_4) \cdots$$

But this is impossible, for  $k_1 \in \Delta_2$  and  $k_2 \notin \Delta_2$ . Therefore  $(k_2, k_3, k_4, k_1)^{H^*} \neq (k_1, k_4, k_3, k_2)^{H^*}$ .

Since  $a^2 = (1)(2)(3)(4)(k_1 k_3)(k_2 k_4) \cdots$  belongs to  $H$  and  $(k_4, k_3, k_2, k_1)^{a^2} = (k_2, k_1, k_4, k_3), (k_4, k_3, k_2, k_1)^{H^*} = (k_2, k_1, k_4, k_3)^{H^*}$ . Therefore in the same way  $(k_1, k_2, k_3, k_4)^{H^*} \neq (k_2, k_1, k_4, k_3)^{H^*} = (k_4, k_3, k_2, k_1)^{H^*}$ .

Thus we have four  $H^*$ -orbits. On the other hand since  $|\Delta_1| > 3$  and  $\Delta_1^a = \Delta_1$ , there is a 4-cycle of  $a$ , say  $(j_1 j_2 j_3 j_4)$ , such that  $\{j_1, j_2, j_3, j_4\} \subset \Delta_1$ . Then  $(j_1, j_2, j_3, j_4)^{H^*}$  is different from these four  $H^*$ -orbits. Thus we have five  $H^*$ -orbits which is a contradiction. Therefore  $\Delta_2^a = \Delta_2$ .

If  $H$  has an orbit different from  $\Delta_1$  and  $\Delta_2$  in  $\Omega - I(H)$ , then as proved above  $a$  fixes these three orbits respectively. By v)  $H^*$  has at least six orbits, which is a contradiction. Therefore  $H$ -orbits in  $\Omega - I(H)$  are  $\Delta_1$  and  $\Delta_2$ .

Now we shall show that  $|\Delta_1| > 3$  leads to a contradiction. Since  $\Delta_1^a = \Delta_1$  and  $\Delta_2^a = \Delta_2$ , we may assume that  $\{i_1, i_2, i_3, i_4\} \subset \Delta_1$  and  $\{j_1, j_2, j_3, j_4\} \subset \Delta_2$ , where  $a = (1 2)(3 4)(i_1 i_2 i_3 i_4)(j_1 j_2 j_3 j_4) \cdots$ . Since  $\Delta_1 \neq \Delta_2$ , by v)  $(i_1, i_2, i_3, i_4)^{H^*}, (i_2, i_3, i_4, i_1)^{H^*}, (j_1, j_2, j_3, j_4)^{H^*}$  and  $(j_2, j_3, j_4, j_1)^{H^*}$  are all different. We shall show that  $(i_1, i_4, i_3, i_2)^{H^*}$  is different from these four  $H^*$ -orbits. From  $\Delta_1 \neq \Delta_2$ ,  $(i_1, i_4, i_3, i_2)^{H^*} \neq (j_1, j_2, j_3, j_4)^{H^*}$  and  $(j_2, j_3, j_4, j_1)^{H^*}$ . If  $(i_1, i_2, i_3, i_4)^{H^*} = (i_1, i_4, i_3, i_2)^{H^*}$ , then  $H$  has a following element

$$x = \begin{pmatrix} i_1 & i_2 & i_3 & i_4 & \cdots \\ i_1 & i_4 & i_3 & i_2 & \cdots \end{pmatrix} = (i_1)(i_3)(i_2 i_4) \cdots .$$

Since  $x$  is of order even, there is a Sylow 2-subgroup of  $H$  fixing  $1, 2, 3, 4$   $i_1$  and  $i_3$ . Thus any Sylow 2-subgroup of  $H_{i_1 i_3}$  is a Sylow 2-subgroup of  $H$ . On the other hand  $a^2 = (1)(2)(3)(4)(i_1 i_3)(i_2 i_4) \cdots$  normalizes  $H_{i_1 i_3}$ . Hence  $a^2$  normalizes a Sylow 2-subgroup  $P'$  of  $H_{i_1 i_2}$ , and  $a^{I(P')} = (1)(2)(3)(4)(i_1 i_3) \cdots$ , contrary to  $N_G(P')^{I(P')} = A_7$ . Thus  $(i_1, i_2, i_3, i_4)^{H^*} \neq (i_1, i_4, i_3, i_2)^{H^*}$ . If  $(i_2, i_3, i_4, i_1)^{H^*} = (i_1, i_4, i_3, i_2)^{H^*}$ , then  $H$  has a following element

$$h = \begin{pmatrix} i_2 & i_3 & i_4 & i_1 & \cdots \\ i_1 & i_4 & i_3 & i_2 & \cdots \end{pmatrix} = (i_1 i_2) (i_3 i_4) \cdots.$$

Then

$$ah = (1 2) (3 4) (i_1) (i_3) (i_2 i_4) \cdots.$$

Since  $\alpha_2(ah) \geq 3$ , from 1)  $ah$  is of order  $2t$  with  $t$  odd. Put  $b = (ah)^t$ . Then  $b$  normalizes a Sylow 2-subgroup  $P''$  of  $H$ . Since  $\Delta_2$  contains an odd number of  $P''$ -orbits, from iii) the four points of  $I(b)$  are contained in  $\Delta_2$  and the three points  $i_1, i_3$  and some one of  $I(b)$  are contained in  $\Delta_1$ . Since  $b \in N_G(P'')$ ,  $I(P'') \supset \{1, 2, 3, 4, i_1, i_3\}$ . Therefore  $a^2 = (1) (2) (3) (4) (i_1 i_3) \cdots$  normalizes a Sylow 2-subgroup of  $H_{i_1 i_3}$ , which is also a Sylow 2-subgroup  $P'''$  of  $H$ . Thus  $(a^2)^{I(P''')} = (1) (2) (3) (4) (i_1 i_3) \cdots$ , which contradicts the assumption  $N_G(P''')^{I(P''')} = A_7$ . Therefore  $(i_2, i_3, i_4, i_1)^{H^*} \neq (i_1, i_4, i_3, i_2)^{H^*}$ .

Thus  $H^*$  has at least five orbits, which is a contradiction. Therefore  $|\Delta_1| = 3$ .

In the proof of Case II of [5. Theorem 2] we needed only the following condition: The number of the fixed points of an involution is seven, and every Sylow 2-subgroup of  $H$  fixes the same points. Therefore in the same way we have that  $G$  is  $M_{23}$ .

YAMAGUCHI UNIVERSITY

### References

- [1] M. Hall: *The Theory of Groups*, Macmillan, New York, 1959.
- [2] H. Wielandt: *Finite Permutation Groups*, Academic Press, New York, 1964.
- [3] H. Nagao: *Multiply Transitive Groups*, California Institute of Technology, California, 1967.
- [4] H. Nagao: *On multiply transitive groups* I, Nagoya Math. J. **27** (1966), 15–19.
- [5] H. Nagao and T. Oyama: *On multiply transitive groups* II, Osaka J. Math. **2** (1965), 129–136.
- [6] H. Nagao: *On multiply transitive groups* IV, Osaka J. Math. **2** (1965), 327–341.
- [7] H. Nagao: *On multiply transitive groups* V, J. Algebra. **9** (1968), 240–248.
- [8] R. Noda and T. Oyama: *On multiply transitive groups* VI, to appear in J. Algebra.