



Title	The fixed subrings of a finite group of automorphisms of $\aleph_0$ -continuous regular rings
Author(s)	Kado, Jiro
Citation	Osaka Journal of Mathematics. 1984, 21(3), p. 683-686
Version Type	VoR
URL	<a href="https://doi.org/10.18910/7699">https://doi.org/10.18910/7699</a>
rights	
Note	

*The University of Osaka Institutional Knowledge Archive : OUKA*

<https://ir.library.osaka-u.ac.jp/>

The University of Osaka

Kado, J.  
Osaka J. Math.  
21 (1984), 683-686

## THE FIXED SUBRINGS OF A FINITE GROUP OF AUTOMORPHISMS OF $\aleph_0$ -CONTINUOUS REGULAR RINGS

JIRO KADO

(Received June 13, 1983)

Let  $R$  be an associative ring,  $G$  a finite group of automorphisms of  $R$ , and let  $R^G$  be the fixed subring of  $G$  on  $R$ . A. Page has proved that if  $R$  is a left self-injective regular ring and the order  $|G|$  of  $G$  is invertible in  $R$ , then  $R^G$  is also a left self-injective regular ring [8]. This theorem is very useful when we investigate some structure of a nonsingular ring and the fixed subring of a finite group of automorphisms.

Recently D. Handelman has discovered an  $\aleph_0$ -continuous regular ring which coordinates the lattice of projections of a finite Rickart  $C^*$ -algebra as a subring of the maximal quotient ring of its  $C^*$ -algebra [4]. We shall prove in this note the following generalization of Page's theorem: if  $R$  is a left  $\aleph_0$ -continuous, left  $\aleph_0$ -injective regular ring and  $|G|$  is invertible in  $R$ , then  $R^G$  is again a left  $\aleph_0$ -continuous,  $\aleph_0$ -injective regular ring. We shall show as a corollary that if  $R$  is a left  $\aleph_0$ -continuous regular ring with  $|G|^{-1} \in R$ ,  $R^G$  is a left  $\aleph_0$ -continuous regular ring and  $S^G$  is the maximal left  $\aleph_0$ -quotient ring of  $R^G$ , where  $S$  is the maximal left  $\aleph_0$ -quotient ring of  $R$ .

### 1. Skew group rings

DEFINITION [7]. Let  $R$  be a ring with identity element 1 and  $G$  a finite group of automorphisms of  $R$ . The skew group ring,  $R \ast G$ , is defined to be a free left  $R$ -module with basis  $\{g: g \in G\}$  and multiplication given as follows: if  $r, s \in R$  and  $g, h \in G$ , then  $(rg)(sh) = rs^{g^{-1}}gh$ .

DEFINITION [3]. A regular ring  $R$  is left  $\aleph_0$ -continuous if the lattice of principal left ideals of  $R$  is upper  $\aleph_0$ -continuous. A ring  $T$  is left  $\aleph_0$ -injective if every homomorphism from a countably generated left ideal of  $T$  into  $T$  is extendable to a  $T$ -module endomorphism of  $T$ . For modules  $A$  and  $B$ ,  $A \subset_e B$  implies that  $A$  is an essential submodule of  $B$ .

A regular ring  $R$  has a maximal left  $\aleph_0$ -quotient ring  $S$  which is a quotient ring defined by the filter-like set  $\mathfrak{X}$  consisting of all countably generated, essen-

tial left ideals of  $R$  [3, § 14]. An element  $x$  in the maximal left quotient ring of  $R$  is contained in  $S$  if and only if there exists some  $J \in \mathfrak{X}$  such that  $Jx \subset R$ . Let  $g \in G$ . Then  $J^g$  is also contained in  $\mathfrak{X}$  and we define  $x^g: J^g \rightarrow R$  by setting  $rx^g = (r^{g^{-1}}x)^g$  for any  $r \in J^g$ . Then  $x^g$  determines a left  $R$ -homomorphism from  $J^g$  to  $R$  and thus  $x^g$  is a uniquely determined element of  $S$ .

K.R. Goodearl has proved the following fundamental result.

**Lemma 1** [3, Th. 14.12]. *Let  $R$  be a left  $\aleph_0$ -continuous regular ring, and let  $S$  be the maximal left  $\aleph_0$ -quotient ring of  $R$ . Then  $S$  is a left  $\aleph_0$ -continuous, left  $\aleph_0$ -injective, regular ring and  $R$  contains all the idempotents of  $S$ .*

It is well-known that if  $R$  is a left self-injective regular ring and  $|G|$  is invertible in  $R$ , then  $R*G$  is a left self-injective regular ring. We shall show an analogous result for left  $\aleph_0$ -continuous, left  $\aleph_0$ -injective regular rings. Of course, left  $\aleph_0$ -continuous, left  $\aleph_0$ -injective regular rings are not necessarily self-injective (See for example, [3, p. 174]).

**Theorem 1.** *Let  $S$  be a left  $\aleph_0$ -continuous, left  $\aleph_0$ -injective regular ring and  $G$  a finite group of automorphisms of  $S$  with  $|G|^{-1} \in S$ . Then the skew group ring  $S*G$  is a left  $\aleph_0$ -continuous, left  $\aleph_0$ -injective regular ring.*

Proof. By [5, Th. 3.2],  $S*G$  is already a regular ring. First we shall show the  $\aleph_0$ -injectivity. Let  $I$  be any countably generated left ideal of  $S*G$  and  $\phi$  any homomorphism from  $I$  to  $S*G$ .  $I$  is countably generated as an  $S$ -module. Then there exists an  $S$ -endomorphism  $\psi$  of  $S*G$  such that  $\psi$  is equal to  $\phi$  on  $I$  by [3, Prop. 14.19]. Define  $\bar{\psi}(x) = |G|^{-1} \sum_g g\psi(g^{-1}x)$  for any  $x$  in  $S*G$ . One easily checks that  $\bar{\psi}$  is an  $S*G$ -homomorphism and it is an extension of  $\phi$ . Since  $S$  is left  $\aleph_0$ -continuous and left  $\aleph_0$ -injective, any matrix ring over  $S$  is also a left  $\aleph_0$ -continuous, left  $\aleph_0$ -injective regular ring by [3, Prop. 14.19]. Therefore the lattice consisting of all finitely generated  $S$ -submodules of  $S*G$  is upper  $\aleph_0$ -continuous. Now let  $J$  be any countably generated left ideal of  $S*G$ . Then we have finitely generated  $S$ -submodule  $A$  of  $S*G$  such that  $J \subset_e A$  as an  $S$ -module. Put  $B = \bigcap_g gA$ , then it is finitely generated as an  $S$ -module and a left ideal of  $S*G$ . As  $B$  is a direct summand as  $S$ -module,  $B$  is a direct summand of  $S*G$  as  $S*G$ -module by Maschke's Theorem (See for example, [7, Th. 0.1]). Since  $J \subset_e B$  as an  $S$ -module, we have  $J \subset_e B$  as an  $S*G$ -module. Now our assertion follows by [3, Cor. 14.4].

**Corollary.** *Let  $G$  be a finite group of automorphisms of a left  $\aleph_0$ -continuous, left  $\aleph_0$ -injective regular ring  $S$ . Assume that  $|G|$  is invertible in  $S$ . Then  $S^G$  is again a left  $\aleph_0$ -continuous, left  $\aleph_0$ -injective regular ring.*

Proof. As in [7], consider  $S$  as an  $S*G$ - $S^G$ -bimodule. As a left  $S*G$ -

module,  $S$  is projective and isomorphic to the principal left ideal  $(S*G)e$ , where  $e=|G|^{-1}\sum_g g$ . Since  $S*G$  is left  $\aleph_0$ -continuous, left  $\aleph_0$ -injective regular ring,  $\text{End}_{S*G}(S)$  is a left  $\aleph_0$ -continuous, left  $\aleph_0$ -injective regular ring by [3, Prop. 14.19]. On the other hand we have  $S^G \cong \text{End}_{S*G}(S)$  by [7, Prop. 0.3] and the proof is complete.

## 2. The fixed subring in an $\aleph_0$ -continuous regular ring

Let  $R$  be a left  $\aleph_0$ -continuous regular ring,  $Q$  the maximal left quotient ring of  $R$  and  $S$  the maximal left  $\aleph_0$ -quotient ring of  $R$ . A finite group  $G$  acting on  $R$  may be extended to automorphisms on  $Q$  and on  $S$  as well. We assume that  $|G|^{-1} \in R$ . Then  $Q^G$  is the maximal left quotient ring of  $R^G$  by [8, Th. 3.6]. Hence it is natural to ask whether  $S^G$  is the maximal left  $\aleph_0$ -quotient ring of  $R^G$ . This is true. We need next two lemmas for its proof.

**Lemma 2.** *Let  $I$  be an essential, countably generated left ideal of  $R^G$ . Then  $RI$  is an essential, countably generated left ideal of  $R$ .*

Proof. Since  $RI$  is a countably generated left ideal of  $R$ , there exists a principal left ideal  $J$  such that  $RI \subset_e J$  by [3, Cor. 14.4]. Put  $Ra = \bigcap_g J^g$ , where  $a=a^2$ , then  $RI \subset_e Ra$ . Since  $Ra$  is  $G$ -invariant,  $(1-a)R$  is also  $G$ -invariant. If  $a \neq 1$ , then  $(1-a)R \cap R^G \neq 0$  by Bergman-Isaak Theorem [1, Prop. 2.3]. Choose some  $y \neq 0 \in (1-a)R \cap R^G$ . We have  $ay=0$  and so  $Iy=0$ . Then  $y$  must be zero since  $I \subset_e R^G$ . This is a contradiction and the proof is complete.

**Lemma 3.** *For any countably generated, essential,  $G$ -invariant left ideal  $I$ , there exists a countably generated, essential, left ideal  $A$  of  $R^G$  such that  $A \subset I \cap R^G$ .*

Proof. Put  $M=I*G$ . Then  $M$  is a countably generated, essential left ideal of  $R*G$ . Let  $M_1 \subset \dots \subset M_n \subset \dots$  be an increasing sequence of finitely generated left ideals such that  $M = \bigcup_n M_n$ . Put  $T = e(R*G)e$ , where  $e=|G|^{-1}\sum_g g$ . Each  $M_n e$  is a direct summand of  $(R*G)e$ . Let  $\phi_n$  be a projection from  $(R*G)e$  onto  $M_n e$ . We have  $\phi_n(e) \in T$  for all  $n$ . We claim that  $\sum_n T\phi_n(e)$  is an essential left ideal of  $T$ . Let  $Ta$  be any non-zero principal left ideal of  $T$ , where  $a^2=a$ . Since  $Me \subset_e (R*G)e$ , we have a non-zero principal left ideal  $(R*G)y \subset Me \cap (R*G)a$ . Let  $\psi$  be a projection from  $(R*G)e$  onto  $(R*G)y$ . Then we have  $\psi(e)a=\psi(e)$ . Since  $(R*G)y \subset M_n e$  for some  $n$ , we have  $\psi(e)=\phi_n(\psi(e)e)=\psi(e)\phi_n(e)$ . Thus  $Ta \cap \sum_n T\phi_n(e)$  contains a non-zero element  $\psi(e)$ .

Next consider the well-known isomorphism  $\theta: e(R*G)e \rightarrow R^G$  given by  $\theta[e(\sum_g r_g g)e] = \sum_g t(r_g)$ , where  $t(a) = |G|^{-1} \sum_g a^g$  ([6, Lemma 0.1]). Put  $A = \theta(\sum_n T\phi_n(e))$ ,

then  $A$  is a countably generated, essential left ideal of  $R^G$ . We claim that  $A \subset I \cap R^G$ . In fact each  $\phi_n(e)$  is contained in  $eMe = e(I \ast G)e$ . Since  $I$  is  $G$ -invariant, we have  $t(r) \in I$  for all  $r \in I$  and thus  $\theta(e(I \ast G)e) \subset I \cap R^G$ . Consequently we have  $A \subset I \cap R^G$ .

Now we shall prove our main theorem.

**Theorem 2.** *Let  $R$  be a left  $\aleph_0$ -continuous regular ring,  $G$  a finite group of automorphisms of  $R$  and  $S$  the maximal left  $\aleph_0$ -quotient ring of  $R$ . Assume  $|G|$  is invertible in  $R$ . Then  $R^G$  is a left  $\aleph_0$ -continuous regular ring and  $S^G$  is the maximal left  $\aleph_0$ -quotient ring of  $R^G$ .*

**Proof.** All idempotents of  $S^G$  are contained in  $R^G$  by Lemma 1. Then the lattice of principal left ideals of  $R^G$  is isomorphic to that of  $S^G$ . Since  $S^G$  is left  $\aleph_0$ -continuous regular ring by the Corollary of § 1,  $R^G$  is a  $\aleph_0$ -continuous regular ring. Let  $s$  be any element in  $S^G$ . There exists a countably generated left ideal  $J \subset_e R$  such that  $Js \subset R$  by [3, Prop. 14.11]. Put  $I = \bigcap_g J^g$ , then  $I$  is again countably generated, essential left ideal of  $R$  by [3, Lemma 14.10] and is  $G$ -invariant. By Lemma 3, we find a countably generated, left ideal  $A \subset_e R^G$  such that  $A \subset I \cap R^G$ . Therefore we have  $As \subset (I \cap R^G)s \subset R^G$ . On the other hand, let  $x$  be any element in  $Q^G$  such that  $Ix \subset R^G$  for some countably generated, essential left ideal  $I$  of  $R^G$ . By Lemma 2,  $RI$  is countably generated and  $RI \subset_e R$ . Since  $RIx \subset R$ ,  $x$  is contained in  $S^G$ . We complete the proof.

---

### References

- [1] G.M. Bergman and I.M. Isaak: *Rings with fixed-point free group actions*, Proc. London Math. Soc. **27** (1973), 69–87.
- [2] J.M. Goursaud, J. Osterburg, J.L. Pascaud and J. Valette: *Points fixes des anneaux réguliers auto-injectifs à gauche*, Comm. Algebra **9** (1981), 1343–1394.
- [3] K.R. Goodearl: *Von Neumann regular rings*, Pitman, 1979.
- [4] D. Handelman: *Finite Rickart  $C^*$ -algebras and their properties*, Advances in Math. **4** (1979), 171–196.
- [5] D. Handelman, J. Lawrence and W. Schelter: *Skew group rings*, Houston J. Math. **4** (1978), 175–190.
- [6] S. Montgomery: *Prime ideals in fixed rings*, Comm. Algebra **9** (1981), 423–449.
- [7] S. Montgomery: *Fixed rings of finite automorphism groups of associative rings*, Lecture Note in Math. 818. Springer-Verlag.
- [8] A. Page: *Actions de groupes*: Séminarie d'algèbre Paul Dubreil 1977–78, Lecture Note in Math. 740, Springer-Verlag.