

Title	NUMERICAL INVARIANTS AND MODULI SPACES FOR LINE ARRANGEMENTS
Author(s)	Dimca, Alexandru; Ibadula, Denis; Măcinic, Daniela Anca
Citation	Osaka Journal of Mathematics. 2020, 57(4), p. 847-870
Version Type	VoR
URL	https://doi.org/10.18910/77234
rights	
Note	

Osaka University Knowledge Archive : OUKA

<https://ir.library.osaka-u.ac.jp/>

Osaka University

NUMERICAL INVARIANTS AND MODULI SPACES FOR LINE ARRANGEMENTS

ALEXANDRU DIMCA, DENIS IBADULA and DANIELA ANCA MĂCINIC

(Received March 6, 2019, revised May 17, 2019)

Abstract

Using several numerical invariants, we study a partition of the space of line arrangements in the complex projective plane, given by the intersection lattice types. We offer also a new characterization of the free plane curves using the Castelnuovo-Mumford regularity of the associated Milnor/Jacobian algebra.

1. Introduction

Line arrangements in the complex projective plane \mathbb{P}^2 look like being simple objects, but a lot of questions related to them are still unanswered, e.g. Terao's conjecture saying that the freeness of such an arrangement is determined by the combinatorics, see Conjecture 6.1 below for a statement, [9, Chapter 8] for more information, and [36] for a survey. Or the conjecture that the monodromy of the associated Milnor fiber is determined by the combinatorics, see [28] for a recent result and [9] for more information.

In order to treat such questions, the study of parameter spaces (a.k.a moduli spaces) of line arrangements has been developed, centered especially on the irreducibility/connectivity questions, see [3], [4], [5], [26], [35].

In this paper, the new idea is to look at the way in which the parameter spaces $A(L)$ and $X(L)$ of line arrangements with a given intersection lattice L behave when the lattice L changes. In section 2 we describe two parameter spaces for the line arrangements $\mathcal{A} : f = 0$ in \mathbb{P}^2 having d lines, namely $A(d)$ and $X(d)$, which are both smooth irreducible varieties, see Corollary 2.3. To partition these two varieties $A(d)$ and $X(d)$ into finer strata, keeping track of the properties of the line arrangements, we use several numerical invariants and study their semi-continuity properties in Proposition 2.4. We consider in this section both line arrangements and arbitrary reduced curves in \mathbb{P}^2 , in order to point out that the numerical invariants associated to line arrangements enjoy special properties, see Corollary 2.6.

In section 3 we recall the definition and main properties of free and nearly free plane curves. Then we prove that a classical invariant in Commutative Algebra, namely the Castelnuovo-Mumford regularity, coincides, when applied to the Milnor/Jacobian algebra $M(f)$ of a reduced plane curve $C : f = 0$, to a naive invariant $st(f)$, coming from the Hilbert function of the graded algebra $M(f)$, exactly when the curve C is free, see Theorem 3.3 and Corollaries 3.4 and 3.5. Corollary 3.5 depends on a key result due to H. Schenck, see

2010 Mathematics Subject Classification. Primary 32S22; Secondary 14H50, 14B05, 13D02.

This work has been supported by the French government, through the UCA^{JEDI} Investments in the Future project managed by the National Research Agency (ANR) with the reference number ANR-15-IDEX-01.

[29]. We end this section by noting that our partitions of the spaces $A(d)$ and $X(d)$ are G -equivariant, where $G = \text{Aut}(\mathbb{P}^2)$ is acting in the obvious way on these parameter spaces. In Proposition 3.11 we describe the dimension of a line arrangement under this G -action.

In section 4 we fix an integer $d \geq 4$ and denote by $\mathcal{L}(d)$ the set of all possible intersection lattices of arrangements with d lines, modulo lattice isomorphisms. For each (isomorphism class of) lattice $L \in \mathcal{L}(d)$, we denote by $X(L)$ the subset of $X(d)$ consisting of line arrangements having an intersection lattice isomorphic to L . Hence the subsets $X(L)$ for $L \in \mathcal{L}(d)$ form the strata of a partition of the smooth irreducible variety $X(d)$. The first properties of these strata are given in Proposition 4.1. Then we discuss several examples of simple lattices L and of corresponding strata $X(L)$, e.g. the lattice L_{gen} corresponding to the generic line arrangement is discussed in Example 4.6 and an obvious generalization, the lattice $L(d, m)$, is considered in Proposition 4.7. Other lattices occurs in Proposition 4.10, Example 4.11, Example 4.15, the last two describing free (resp. nearly free) line arrangements. Note that our results on a stratum $X(L)$ (e.g. dimension, connectivity) easily translate into properties of the quotient $X(L)/G$. We also reprove in a new way a result due to Tohăneanu [34], giving the classification of line arrangements with a Jacobian syzygy of minimal degree 2, see Theorem 4.12.

In section 5 we point out the complexity of the stratification $(X(L))_{L \in \mathcal{L}(d)}$ of the space $X(d)$. First we describe all the strata when $d = 4, 5, 6$, and pay particular attention to the strata formed by (nearly) free arrangements. We explain just after Proposition 5.4 that these stratification do not satisfy the frontier condition in general, in particular they are not Whitney regular stratifications. In Remark 5.5 and in the answer to Question 5.6 we show that some nice features of this stratification noticed when $d \leq 6$ do not extend to higher degrees d .

In the final section we discuss Terao's conjecture in the case of line arrangements, recall the known results and give a new proof for Theorem 6.3. Finally, in Proposition 6.5 we give a generalization of the result saying that the generic arrangement is not free when $d \geq 4$.

The authors would like to thank the Oberwolfach Research Institute for Mathematics, where the major part on the work on this project was done during a RIP program.

We also thank Torsten Hoge for his useful remarks on the previous version of this paper and the referee for the careful reading of the manuscript and for his useful suggestions.

2. General facts on plane curves and line arrangements

2.1. Two parameter spaces for line arrangements: $A(d)$ and $X(d)$. Let $S = \mathbb{C}[x, y, z]$ be the graded polynomial ring in the variables x, y, z with complex coefficients, and S_m be the vector space of degree m homogeneous polynomials in S . Fix an integer $d \geq 1$ and regard the projective space $C(d) = \mathbb{P}(S_d)$ as the parameter space of degree d curves in \mathbb{P}^2 .

DEFINITION 2.1. We denote by $C(d)_0$ the subset in $C(d)$ corresponding to curves having only isolated singularities, and by $A(d)$ the subset in $C(d)_0$ corresponding to line arrangements consisting of d distinct lines.

Proposition 2.2. *The set $C(d)_0$ is Zariski open and dense in $C(d)$. The set $A(d)$ is a Zariski closed subset in $C(d)_0$.*

Proof. For the first claim, note that the complement $C(d) \setminus C(d)_0$ is the union of the finite family of Zariski closed subsets given by the images of the obvious mappings

$$\phi_m : \mathbb{P}(S_m) \times \mathbb{P}(S_{d-2m}) \rightarrow \mathbb{P}(S_d),$$

$(A, B) \mapsto A^2B$, for $m = 1, 2, \dots, [d/2]$. For the second claim, we consider the map

$$(2.1) \quad \psi : \mathbb{P}(S_1)^d \rightarrow \mathbb{P}(S_d),$$

given by $(\ell_1, \dots, \ell_d) \mapsto \ell_1 \cdot \ell_2 \cdot \dots \cdot \ell_d$. Then $A(d)$ is just the intersection of the set $C(d)_0$ with the image of the mapping ψ . □

Note that

$$(2.2) \quad X(d) := \psi^{-1}(A(d))$$

is exactly the set of linear forms $(\ell_1, \dots, \ell_d) \in \mathbb{P}(S_1)^d$ such that $\ell_i \neq \ell_j$ for $i \neq j$, and the restriction $\psi : X(d) \rightarrow A(d)$ is a Galois covering with structure group the symmetric group σ_d on d elements.

Corollary 2.3. *The parameter spaces $A(d)$ and $X(d)$ are smooth, irreducible algebraic varieties of dimension $2d$. The space $X(d)$ is simply connected and the fundamental group of the space $A(d)$ is given by*

$$\pi_1(A(d)) = \sigma_d.$$

Proof. The only claim that needs some explanation is the fact that $X(d)$ is simply connected. This follows from the fact that $X(d)$ is obtained from the simply connected variety $\mathbb{P}(S_1)^d$ by removing the codimension 2 linear subvarieties $\Delta_{ij} : \ell_i = \ell_j$ for all $1 \leq i < j \leq d$. □

2.2. Some numerical invariants for plane curves and line arrangements. For a polynomial $f \in S_d$, we denote by J_f the ideal in S generated by the partial derivatives f_x, f_y, f_z , and call J_f the Jacobian ideal of f . The graded ring $M(f) = S/J_f$ is called the Milnor or Jacobian algebra of f . We define

$$(2.3) \quad m_k(f) = \dim M(f)_k.$$

Note that one has $m_k(f) = \tau(f)$ for $k > 3(d - 2)$ and $f \in C(d)_0$, where $\tau(f)$ is the total Tjurina number of the reduced plane curve $C(f) : f = 0$, see [6]. We also denote by $\tau(C(f))$ this number, and note that it is nothing else but the degree of the Jacobian ideal J_f .

The minimal degree of a Jacobian syzygy for f is the integer $mdr(f)$ defined to be the smallest integer $r \geq 0$ such that there is a nontrivial relation

$$(2.4) \quad af_x + bf_y + cf_z = 0$$

among the partial derivatives f_x, f_y and f_z of f with coefficients a, b, c in S_r . We denote by $AR(f)$ the graded S -module consisting of all the triples $(a, b, c) \in S^3$ satisfying (2.4). In fact $AR(f)$ depends only on the class of $f \in S_d$ in $C(d) = \mathbb{P}(S_d)$.

Proposition 2.4. (1) *The subset $\{f \in S_d : m_k(f) \leq m\} \subset C(d)$ is Zariski open and dense in $C(d)$ for any $k \geq 0$ and any $m \geq 0$. In particular, the following two sets $\{f \in C(d)_0 : \tau(f) \leq m\} \subset C(d)_0$ and $\{f \in A(d) : \tau(f) \leq m\} \subset A(d)$, are Zariski*

open and dense in $C(d)_0$ (resp. in $A(d)$) for any $k \geq 0$ and any $m \geq 0$.

- (2) The subset $\{f \in S_d : mdr(f) \leq m\} \subset C(d)$ is Zariski closed in $C(d)$ for any $m \geq 0$. In particular, the following two sets $\{f \in C(d)_0 : mdr(f) \leq m\} \subset C(d)_0$ and $\{f \in A(d) : mdr(f) \leq m\} \subset A(d)$, are Zariski closed in $C(d)_0$ (resp. in $A(d)$) for any $m \geq 0$.

Proof. The first claim is clear by the semicontinuity properties of the rank of a matrix, whose rows are obtained by taking all the coefficients of the polynomials μg , where μ runs through the set of monomials of degree $k - d + 1$ in S and $g \in \{f_x, f_y, f_z\}$.

To prove the second claim, consider the closed subvariety Y_m in $\mathbb{P}(S_m^3) \times \mathbb{P}(S_d)$ given by

$$Y_m = \{((a, b, c), f) : af_x + bf_y + cf_z = 0\}.$$

Note that a polynomial $f \in S_d$ satisfies $mdr(f) \leq m$ if and only if $[f] \in \mathbb{P}(S_d)$ is in the image of Y_m under the second projection. □

DEFINITION 2.5. For a polynomial $f \in C(d)_0$, we recall the following invariants.

- (i) The coincidence threshold

$$ct(f) = \max\{q : \dim M(f)_k = \dim M(f_s)_k \text{ for all } k \leq q\},$$

with f_s a homogeneous polynomial in S of the same degree d as f and such that $C_s : f_s = 0$ is a smooth curve in \mathbb{P}^2 .

- (ii) The stability threshold $st(f) = \min\{q : \dim M(f)_k = \tau(f) \text{ for all } k \geq q\}$.
 (iii) The regularity $reg(f)$ is the Castelnuovo-Mumford regularity of the Milnor algebra $M(f)$, regarded as a graded S -module, see [18, Chapter 4].

The exact sequences

$$(2.5) \quad 0 \rightarrow AR(f)(-(d-1)) \rightarrow S(-(d-1))^3 \xrightarrow{(f_x, f_y, f_z)} S \rightarrow M(f) \rightarrow 0$$

and

$$(2.6) \quad 0 \rightarrow AR(f_s)(-(d-1)) \rightarrow S(-(d-1))^3 \xrightarrow{(f_{s,x}, f_{s,y}, f_{s,z})} S \rightarrow M(f_s) \rightarrow 0$$

and the fact that $AR(f_s)_k = 0$ for $k < d - 1$ imply that

$$(2.7) \quad ct(f) \geq mdr(f) + d - 2,$$

with equality for $mdr(f) < d - 1$. To have equality always, it is convenient to introduce the invariant $mdr_e(f)$, the minimal degree of an essential Jacobian relation for f , which is by definition the minimal degree of a relation (2.4), where the triple (a, b, c) does not belong to the S -submodule of $AR(f)$ generated by the Koszul relations $(f_y, -f_x, 0)$, $(f_z, 0, -f_x)$ and $(0, f_z, -f_y)$. With this definition we always have

$$(2.8) \quad ct(f) = mdr_e(f) + d - 2,$$

see [13].

Corollary 2.6. *With the above notation, the following hold.*

(1) Let $C : f = 0$ be a singular, reduced plane curve of degree $d \geq 3$ in \mathbb{P}^2 . Then

$$mdr(f) \leq d - 1, \quad mdr_e(f) \leq 2(d - 2) \text{ and } ct(f) \leq 3(d - 2).$$

Moreover, if $\tau(C) = 1$, all these inequalities are equalities.

(2) Let $\mathcal{A} : f = 0$ be an arrangement having $d \geq 2$ lines. Then

$$mdr(f) = mdr_e(f) \leq d - 2 \text{ and } ct(f) \leq 2(d - 2).$$

Moreover, both of these inequalities are equalities for a generic arrangement.

Proof. To prove (1), note that $mdr(f) \leq d - 1$ follows from the existence of Koszul relations. The inequality $mdr_e(f) \leq 2(d - 2)$ follows using (2.8) and the obvious fact that $ct(f) \leq T = 3(d - 2)$. When $\tau(C) = 1$, it follows from [13, Example 4.3] that $ct(f) = T$. Moreover, such a curve is nodal and irreducible, and hence $mdr(f) \geq d - 1$, by [13, Theorem 4.1]. This remark completes the proof of the first claim.

In view of Proposition 2.4 (2), to prove (2) it is enough to show that $mdr(f) = d - 2$ when \mathcal{A} is a generic arrangement. But this follows from [13, Theorem 4.1]. \square

The invariants $st(f)$ and $reg(f)$ are closely related, as the next section shows. However, they do not seem to satisfy semicontinuity properties similar to those in Proposition 2.4, see Remark 5.3.

3. Free and nearly free plane curves

3.1. Free plane curves. For the equivalence of the properties in the next definition, we refer to [33]. See also [9, Chapter 8].

DEFINITION 3.1. The curve $C : f = 0$ is a free divisor if the following equivalent conditions hold.

- (1) The Milnor algebra $M(f)$ is a Cohen-Macaulay S -module.
- (2) The minimal graded resolution of the Milnor algebra $M(f)$ as an S -module has the following form

$$0 \rightarrow S(-d_1 - d + 1) \oplus S(-d_2 - d + 1) \rightarrow S^3(-d + 1) \xrightarrow{(f_x, f_y, f_z)} S$$

for some positive integers d_1, d_2 .

- (3) The graded S -module $AR(f)$ is free of rank 2, i.e. there is an isomorphism

$$AR(f) = S(-d_1) \oplus S(-d_2)$$

for some positive integers d_1, d_2 .

When C is a free divisor, the integers $d_1 \leq d_2$ are called the exponents of C . They satisfy the relations

$$(3.1) \quad d_1 + d_2 = d - 1 \text{ and } \tau(C) = (d - 1)^2 - d_1 d_2,$$

where $\tau(C)$ is the total Tjurina number of C , see for instance [12]. For a free curve, one has $mdr(f) = d_1$, $ct(f) = d_1 + d - 2$ in view of (2.7), and $st(f) = d_2 + d - 3$, see for instance [10].

DEFINITION 3.2. The curve $C : f = 0$ is a nearly free divisor if the Milnor algebra $M(f)$ has a minimal graded resolution of the form

$$0 \rightarrow S(-d-d_2) \rightarrow S(-d-d_1+1) \oplus S^2(-d-d_2+1) \rightarrow S^3(-d+1) \xrightarrow{(f_x, f_y, f_z)} S$$

for some integers $1 \leq d_1 \leq d_2$, called the exponents of C .

For a nearly free curve, the exponents satisfy $d_1 + d_2 = d$, and one has $mdr(f) = d_1$, $ct(f) = d_1 + d - 2$ in view of (2.7), and $st(f) = d_2 + d - 2$ by the results in [10].

Theorem 3.3. *Let $C : f = 0$ be a reduced plane curve. Then*

$$st(f) - 1 \leq reg(f) \leq st(f),$$

and the equality $reg(f) = st(f)$ holds if and only if $C : f = 0$ is a free curve.

Proof. Let $H_{M(f)}$ (resp. $P_{M(f)}$) be the Hilbert function (resp. the Hilbert polynomial) of the graded S -module $M(f)$. Then [18, Theorem 4.2] implies that

$$H_{M(f)}(k) = P_{M(f)}(k)$$

for any $k \geq reg(f) + 1$. Since for a reduced plane curve the Hilbert polynomial $P_{M(f)}$ is just the constant $\tau(C)$, it follows from the definition of the stability threshold $st(f)$ that $st(f) \leq reg(f) + 1$, and hence $st(f) - 1 \leq reg(f)$.

To prove the other inequality, let J_f^{sat} be the saturation of the Jacobian ideal J_f , as discussed in a more general setting in [21]. Consider the exact sequence of graded S -modules

$$0 \rightarrow J_f^{sat}/J_f \rightarrow M(f) \rightarrow S/J_f^{sat} \rightarrow 0.$$

Then [17, Corollary 20.19] implies that

$$reg(f) = regM(f) \leq \max\{reg(J_f^{sat}/J_f), reg(S/J_f^{sat})\}.$$

Note the module J_f^{sat}/J_f has finite length, so [18, Corollary 4.4] implies that

$$reg(J_f^{sat}/J_f) = \max\{k : (J_f^{sat}/J_f)_k \neq 0\} = sat(J_f) - 1,$$

in the notation from [8]. Moreover, [8, Corollary 2] says that

$$sat(J_f) \leq \max\{T - ct(f), st(f)\},$$

where $T = 3(d-2)$. On the other hand the quotient S/J_f^{sat} is a Cohen-Macaulay module satisfying $\text{depth } S/J_f^{sat} = \dim S/J_f^{sat} = 1$, and [18, Corollary 4.8] tells us that $s = reg(S/J_f^{sat})$, where s is the smallest integer such that $k \geq s$ implies

$$H_{S/J_f^{sat}}(k) = P_{S/J_f^{sat}}(k) = \tau(C).$$

This integer s is determined in [8, Proposition 2], where it is shown that one has $s = T - ct(f)$. It follows that

$$reg(f) \leq \max\{T - ct(f), st(f) - 1\}.$$

When $C : f = 0$ is free, we have $ct(f) + st(f) = T$ by [10], and hence we get $reg(f) \leq st(f)$. A direct computation using the definition of the Castelnuovo-Mumford regularity in terms of a resolution (see [17], p.505) yields $reg(f) = d_2 + d - 3$ when C is free. The equality

$st(f) = reg(f)$ follows using the above formulas for $st(f)$. Note that in the free case, the equality $reg(f) = st(f)$ is also a consequence of Theorem 4.2 in [18], since $M(f)$ is Cohen-Macaulay in this case.

When $C : f = 0$ is not free, then it is shown in [10, Corollary 1.7] that $ct(f) + st(f) \geq T + 2$, which implies that $\max\{T - ct(f), st(f) - 1\} = st(f) - 1$, and this completes the proof. \square

Corollary 3.4. *Let $C : f = 0$ be a reduced plane curve of degree $d \geq 4$. Then the following hold.*

- (1) *C is free if and only if $reg(f) = 2(d - 2) - mdr_e(f)$.*
- (2) *C is nearly free if and only if $reg(f) = 2(d - 2) - mdr_e(f) + 1$.*
- (3) *If C is neither free nor nearly free, then $reg(f) \geq 2(d - 2) - mdr_e(f) + 2$.*

Proof. The claims (1) and (2) then follow from the equalities $ct(f) + st(f) = T$ (resp. $ct(f) + st(f) = T + 2$) which are shown in [10, Corollary 1.7] to characterize the free (resp. nearly free) curves. The claim in (3) follows from the equality (2.8) and the inequality $ct(f) + st(f) \geq T + 3$ which holds in this case by [10, Corollary 1.7]. \square

Note that in the cases (1) and (2) above one has $mdr(f) = mdr_e(f)$, while in the case (3) both $mdr(f) = mdr_e(f)$ and $mdr(f) = d - 1 < mdr_e(f)$ may occur.

Corollary 3.5. *If $\mathcal{A} : f = 0$ is any arrangement of $d \geq 4$ lines, then*

$$reg(f) \leq 2d - 5 \text{ and } st(f) \leq 2d - 4.$$

When $\mathcal{A} : f = 0$ is a generic arrangement of $d \geq 4$ lines, both of the above inequalities become equalities.

Proof. The inequality $reg(f) \leq 2d - 5$, with equality for a generic arrangement, follows from [29, Corollary 3.5]. The reader must notice that the regularity there is the Castelnuovo-Mumford regularity of the graded S -module $AR(f)$, and not as in our paper that of the Milnor algebra $M(f)$. The exact sequence (2.5) allows us to pass from one regularity to the other, namely one has

$$reg(f) = reg(M(f)) = reg(AR(f)) + d - 3.$$

The formula for $st(f)$ in the case of a generic arrangement follows from [13, Corollary 1.3]. In fact, using Theorem 3.3, we need only one of these two invariants, since it is known that such a line arrangement is not free, see for instance [12] or Proposition 6.5 below.

In the general case, if the arrangement \mathcal{A} is not free, then $st(f) = reg(f) + 1 \leq 2d - 4$. And for a free arrangement $\mathcal{A} : f = 0$, one has $st(f) = d + d_2 - 3 \leq 2d - 4$ since clearly $d_2 \leq d - 1$. \square

DEFINITION 3.6. We denote by $\mathcal{C}(d, \tau)$ (resp. $\mathcal{A}(d, \tau)$) the set of curves in $\mathcal{C}(d)_0$ (resp. the set of line arrangements in $\mathcal{A}(d)$) with a fixed global Tjurina number τ . We denote by $\mathcal{F}(d, \tau)$ (resp. $\mathcal{FA}(d, \tau)$) the set of free curves in $\mathcal{C}(d, \tau)$ (resp. the set of free line arrangements in $\mathcal{A}(d, \tau)$).

For the following result we refer to [11]. For the case of hyperplane arrangements in \mathbb{P}^n see [37].

Theorem 3.7. *The set $F(d, \tau)$ is a Zariski open subset in $C(d, \tau)$. The set $FA(d, \tau)$ is a Zariski open subset in $A(d, \tau)$.*

Using the Galois covering $\psi : X(d) \rightarrow A(d)$, we introduce the following notation.

$$(3.2) \quad X(d, m_k \leq m) = \psi^{-1}(\{f \in A(d) : m_k(f) \leq m\})$$

and $X(d, m_k = m) = X(d, m_k \leq m) \setminus X(d, m_k \leq m - 1)$.

$$(3.3) \quad X(d, \tau \leq m) = \psi^{-1}(\{f \in A(d) : \tau(f) \leq m\}) \text{ and } X(d, \tau) = \psi^{-1}(A(d, \tau)).$$

$$(3.4) \quad X(d, mdr \leq m) = \psi^{-1}(\{f \in A(d) : mdr(f) \leq m\})$$

and $X(d, mdr = m) = X(d, mdr \leq m) \setminus X(d, mdr \leq m - 1)$.

$$(3.5) \quad FX(d, \tau) = \psi^{-1}(FA(d, \tau)) \text{ and } FX(d) = \cup_{\tau} FX(d, \tau).$$

Then we have the following obvious consequence of Proposition 2.4 and Theorem 3.7.

Corollary 3.8. *The sets $X(d, m_k \leq m)$ and $X(d, \tau \leq m)$ are Zariski open in $X(d)$ for any positive integers k and m . The set $X(d; mdr \leq m)$ is Zariski closed in $X(d)$ for any positive integer m . Moreover, the set $FX(d, \tau)$ is Zariski open in the variety $X(d, \tau)$.*

It is known that $FA(d, \tau) \neq \emptyset$ implies that there is an integer $r \in [0, (d - 1)/2]$ such that

$$(3.6) \quad FX(d, \tau) \subset X(d, mdr = r), \text{ where } \tau = \tau(d, r) = (d - 1)^2 - r(d - 1 - r),$$

see [10], [14]. We set

$$(3.7) \quad \tau(d)_{\min} = \frac{3}{4}(d - 1)^2 \text{ for } d \text{ odd and } \tau(d)_{\min} = \lfloor \frac{3}{4}(d - 1)^2 \rfloor + 1 \text{ for } d \text{ even.}$$

With this notation we have the following result.

Corollary 3.9. *If $\mathcal{A} : f = 0$ is a free line arrangement in $A(d, \tau)$, then*

$$\tau = \tau(f) \geq \tau(d)_{\min}.$$

Proof. The inequalities follow from the formula for $\tau(d, r)$ given above. Here $\lfloor x \rfloor$ denotes the integral part of the real number x . \square

REMARK 3.10. It is shown in [10] that a line arrangement $\mathcal{A} : f = 0$ with $d = |\mathcal{A}|$ is nearly free with exponents $d_1 \leq d_2 = d - d_1$ if and only if

$$\tau(\mathcal{A}) = \tau(d, d_1) - 1.$$

3.2. Three group actions on parameter spaces. Consider the connected algebraic group $G = \text{Aut}(\mathbb{P}^2) = \text{PGL}(3, \mathbb{C})$ of dimension 8. This group acts naturally on the variety $C(d)$ and all the subsets $C(d)_0$, $A(d)$, $F(d, \tau)$, $FA(d, \tau)$ are in fact G -invariant, hence they inherit a natural G -action, and are unions of G -orbits $G \cdot f$, for some $f \in C(d)$. Moreover, G acts also on the variety $X(d)$ in a diagonal way, and such that the map $\psi : X(d) \rightarrow A(d)$ is G -equivariant. It follows that all the subsets $X(d, m_k \leq m)$, $X(d, \tau \leq m)$, $X(d, \tau)$, $X(d; mdr \leq m)$, $FX(d, \tau)$ are also G -invariant, so they consists of unions of G -orbits, denoted by $G \cdot (\ell_1, \dots, \ell_d)$,

for some $(\ell_1, \dots, \ell_d) \in X(d)$.

Proposition 3.11. *Let $(\ell_1, \dots, \ell_d) \in X(d)$ and denote $f = \psi(\ell_1, \dots, \ell_d) \in A(d)$. Then one has the following.*

- (1) $\dim G \cdot (\ell_1, \dots, \ell_d) = \dim G \cdot f$.
- (2) $\dim G \cdot f = 2$ if $d = 1$, $\dim G \cdot f = 4$ if $d = 2$ and for $d = 3$, one has $\dim G \cdot f = 5$ if $\mathcal{A} : f = 0$ consists of 3 concurrent lines, and $\dim G \cdot f = 6$ if $\mathcal{A} : f = 0$ consists of a triangle.
- (3) For $d \geq 4$, $\dim G \cdot f = 5$ if $\text{mdr}(f) = 0$, $\dim G \cdot f = 7$ if $\text{mdr}(f) = 1$ and $\dim G \cdot f = 8$ if $\text{mdr}(f) > 1$.

Proof. The first claim follows since the map ψ has finite fibers. To prove (2) and (3), note that one has

$$\dim G \cdot f = \dim G - \dim \text{Fix}(f),$$

where $\text{Fix}(f)$ is the stabilizer subgroup of f . The Lie algebra of $\text{Fix}(f)$ is exactly $AR(f)_1$, i.e. the linear Jacobian syzygies, see [16, Proposition 1.1]. When $d = 1$, we can take $f = x$ and it follows that $\dim AR(f)_1 = 6$, since in the notation from (2.4) one takes $a = 0$ and $b, c \in S_1$ arbitrary. When $d = 2$, we can take $f = xy$, and it follows that $\dim AR(f)_1 = 4$, since $a = \lambda x, b = -\lambda y$ and $c \in S_1$ arbitrary. When $d = 3$ there are two possibilities. The first one is $f = x^3 + y^3$, when $\dim AR(f)_1 = 3$, as $a = b = 0$ and $c \in S_1$. The second case is $f = xyz$ and then $\dim AR(f)_1 = 2$, since $AR(f)_1$ is spanned in this case by $(x, -y, 0)$ and $(x, 0, -z)$.

Assume now that $d \geq 4$. Then, if $r = \text{mdr}(f) = 0$, this means that $\dim AR(f)_0 = 1$, which implies $\dim AR(f)_1 = 3$. If $r = 1$, then it follows from [16, Proposition 2.2] that $\dim AR(f)_1 = 1$. When $r > 1$, one has $AR(f)_1 = 0$, so the claims in (3) are now proved. \square

Let \mathcal{G} be the Galois group of \mathbb{C} over \mathbb{Q} . Then \mathcal{G} acts on the parameter spaces $A(d)$ and $X(d)$ by acting on the coefficients of the defining equations. It follows that all the subsets $X(d, m_k \leq m), X(d, \tau \leq m), X(d, \tau), X(d; \text{mdr} \leq m), FX(d, \tau)$, as well as $A(d, m_k \leq m), A(d, \tau \leq m), A(d, \tau), A(d; \text{mdr} \leq m), FA(d, \tau)$ are also \mathcal{G} -invariant.

The symmetric group σ_d also acts on $X(d)$ by permuting the linear factors of the defining equation $f = 0$ of a line arrangement, and this is the reason why some strata in $X(d)$ are not irreducible, while their images in $A(d)$ have this property, see for instance Proposition 4.7 (1).

3.3. On rigid plane curves and line arrangements. We say that a plane curve $C : f = 0$ is algebraically rigid if $(J_f^{\text{sat}}/J_f)_d = 0$, where J_f^{sat} denotes as above the saturation of the Jacobian ideal J_f . Indeed, the vector space $(J_f^{\text{sat}}/J_f)_d$ is naturally identified to the space of first order locally trivial deformation of C in \mathbb{P}^2 , modulo the above G -action, see [20, 31, 32]. These deformations preserve the analytic isomorphism type of each singular point of C .

EXAMPLE 3.12. It is known that a curve $C : f = 0$ is free if and only if $J_f^{\text{sat}} = J_f$, see [33]. In particular, any free curve is algebraically rigid. A generic line arrangement of 4 lines in \mathbb{P}^2 is not free, but it is algebraically rigid by Proposition 3.11 (3) since $\text{mdr}(f) > 1$ in this case. In fact, in this case one has $\dim(J_f^{\text{sat}}/J_f)_3 = 1$ and $(J_f^{\text{sat}}/J_f)_k = 0$ for $k \neq 3$.

We say that a reduced plane curve $C : f = 0$ is topologically rigid if any deformation of C preserving the number of irreducible components of C , their degrees and the topological type of each singularity of C is trivial modulo the above G -action. For more on this type of rigidity see [25].

EXAMPLE 3.13. A line arrangement \mathcal{A} consisting in $d \geq 4$ lines passing through one point satisfies $\text{mdr}(f) = 0$, and it is free. Hence \mathcal{A} is an algebraically rigid curve. On the other hand, we can modify the cross-ratio of a subset of 4 lines in \mathcal{A} by moving one line, without changing the topology of the singularity, and hence such a family will not be contained in one G -orbit. Hence \mathcal{A} is not topologically rigid.

REMARK 3.14. If a reduced plane curve has only simple singularities of type A_k , D_k and E_6 , E_7 , E_8 , then C is algebraically rigid if and only if C is topologically rigid. Indeed, for a simple singularity, a topologically constant deformation is the same as an analytically constant deformation. In particular, for a line arrangement \mathcal{A} having only double and triple points, the two rigidity notions coincide. In such a case we will simply say that \mathcal{A} is rigid. For examples of this situation, see Remark 4.2 and the stratum $A(L(\Delta))$ in Proposition 5.4 below.

For more on the interest of rigidity in the study of line arrangements, see [2].

4. A partition of the parameter space $X(d)$

From now on in this paper we assume that $d \geq 4$. For a fixed integer $d \geq 4$, we denote by $\mathcal{L}(d)$ the set of all possible intersection lattices $L(\mathcal{A})$, for line arrangements in \mathbb{P}^2 consisting of d distinct lines. For a lattice $L \in \mathcal{L}(d)$, we denote by $X(L)$ the set of all elements $(\ell_1, \dots, \ell_d) \in X(d)$ such that the line arrangement

$$(4.1) \quad \mathcal{A} : \ell_1 = \dots = \ell_d = 0$$

has an intersection lattice $L(\mathcal{A})$ isomorphic to L , see [27] for more on intersection lattices. We also set $A(L) = \psi(X(L))$. Such a lattice gives in particular information on the multiple points p in the arrangement \mathcal{A} , and about their multiplicities, denoted by $m_p \geq 2$. In particular, we define

$$(4.2) \quad \tau(L) = \sum_p (m_p - 1)^2 = \tau(\mathcal{A}).$$

By definition, we have the following partitions

$$(4.3) \quad X(d) = \cup_{L \in \mathcal{L}(d)} X(L) \text{ and } X(d, \tau) = \cup_{L \in \mathcal{L}(d), \tau(L)=\tau} X(L)$$

and similarly

$$(4.4) \quad A(d) = \cup_{L \in \mathcal{L}(d)} A(L) \text{ and } A(d, \tau) = \cup_{L \in \mathcal{L}(d), \tau(L)=\tau} A(L).$$

One has the following.

Proposition 4.1. *For any lattice $L \in \mathcal{L}(d)$, the following hold.*

- (1) *The sets $X(L)$ and $A(L)$ are constructible; they are also G -invariant and \mathcal{G} -invariant.*
- (2) *$X(L) \subset X(d, \tau(L))$ and $A(L) \subset A(d, \tau(L))$.*

- (3) The function $mdr : A(L) \rightarrow \mathbb{N}$ attains its minimal value on a Zariski closed subset F of $A(L)$, and in general $F \neq A(L)$.
- (4) The function $m_k : A(L) \rightarrow \mathbb{N}$ attains its minimal value on a Zariski open subset U_k of $A(L)$, and in general $U_k \neq A(L)$.

Proof. The claim about the constructibility in (1) can be settled as follows. A point p of multiplicity $k \geq 3$ will give rise to a set $\mathcal{E}(L)_p$ of $k - 2$ equations to be satisfied by the set of coefficients $(a_i, b_i, c_i) \in \mathbb{P}(S_1)$, where $\ell_i = a_i x + b_i y + c_i z$. Indeed, if the lines passing through p are for instance $L_i : \ell_i = 0$ for $i = 1, 2, \dots, k$, then the fact that all these lines pass through p is expressed by the vanishing of $k - 2$ determinants $D(1, 2, j)$ of 3×3 matrices $A(1, 2, j)$, constructed using the coefficients of ℓ_1, ℓ_2 and ℓ_j to define the corresponding three rows, where $j = 3, 4, \dots, k$. Note that such determinants really define hypersurfaces in the product $\mathbb{P}(S_1)^d$. Moreover, when three lines L_u, L_v and L_w are not concurrent, we should add the condition that the corresponding determinant $D(u, v, w)$ is not zero. More details on this construction can be found in [26], see however Remark 4.4 below. The G -invariance and the \mathcal{G} -invariance of $X(L)$ is obvious.

The claim (2) is clear. For the first part in claim (3), use Proposition 2.4, (2). For the second part of claim (3), one may consider the example of two line arrangements

$$\begin{aligned} \mathcal{A} : f &= xy(x - y - z)(x - y + z)(2x + y - 2z)(x + 3y - 3z)(3x + 2y + 3z) \\ &\quad (x + 5y + 5z)(7x - 4y - z) = 0 \end{aligned}$$

and

$$\begin{aligned} \mathcal{A}' : f' &= xy(x + y - z)(5x + 2y - 10z)(3x + 2y - 6z)(x - 3y + 15z) \\ &\quad (2x - y + 10z)(6x + 5y + 30z)(3x - 4y - 24z) = 0, \end{aligned}$$

having isomorphic intersection lattices and constructed by Ziegler in [38]. A picture of these arrangements can be found in [9, Chapter 8]. They consists both of nine lines, and have only double and triple points. More precisely, they have $n_2 = 18$ double points and $n_3 = 6$ triple points, and hence $\tau(\mathcal{A}) = \tau(\mathcal{A}') = 42$. In the case of \mathcal{A} , the six triple points are on a conic, and a direct computation shows that $mdr(f) = 5$. For \mathcal{A}' , the six triple points are not on a conic, i.e. the arrangement \mathcal{A}' is a small deformation of the arrangement \mathcal{A} , and a direct computation shows that $mdr(f) = 6$. See also [30, Example 13]. The above example settles also the claim (4) by taking $k = 13$, since

$$m_{13}(f') = \tau(f') = 42 < m_{13}(f).$$

□

REMARK 4.2. In fact, it is clear that there is a topologically constant 1-parameter family of line arrangements \mathcal{A}_t such that $\mathcal{A}_0 = \mathcal{A}$ and \mathcal{A}_t for $t \neq 0$ has the same numerical invariants as \mathcal{A}' . This family is obtained by moving the sixth triple point till it gets onto the conic determined by the first 5 triple points. It follows that \mathcal{A} is not rigid, and one can check that $\dim(I/J_f)_9 = 4$. A direct computation shows that for \mathcal{A}' one has $\dim(I/J_{f'})_9 = 4$ as well, i.e. \mathcal{A}' is not rigid either.

REMARK 4.3. The above result says that the invariants mdr and m_k are not determined by the combinatorics in general. However, if \mathcal{A} is a free arrangement, both $mdr(f)$ and $m_k(f)$ are determined by the lattice $L(\mathcal{A})$. The claim for $mdr(f)$ follows from the formula (3.6). The claim for $m_k(f)$ follows from the fact that the exponents $d_1 = mdr(f)$ and $d_2 = d - 1 - mdr(f)$ determine the Hilbert function $H_{M(f)}$ via the resolution given in Definition 3.1 (2).

REMARK 4.4. The set of equations $\mathcal{E}(L) = \cup_p \mathcal{E}(L)_p$, with $\mathcal{E}(L)_p$ defined above in Proposition 4.1, is smaller than the set of equations constructed in [26], and which we call $\mathcal{E}'(L)$ here. Indeed, any point p of multiplicity $k \geq 3$ contributes $k - 2$ equations to our set $\mathcal{E}(L)$, and $\binom{k}{3}$ equations to the set $\mathcal{E}'(L)$. The two ideals $I(\mathcal{E}(L))$ and $I(\mathcal{E}'(L))$ are distinct. Indeed, the equations in $\mathcal{E}'(L)$ are linearly independent degree 3-forms, as each of them involves monomials in distinct set of variables. For instance the monomial $a_1 b_2 c_3$ occurs only in the equation associated to the triple of lines (L_1, L_2, L_3) , supposed to pass through a multiple point p .

On the other hand, it is clear that the two ideals $I(\mathcal{E}(L))$ and $I(\mathcal{E}'(L))$ both have $Y(L) = \overline{X(L)}$ as zero set, and hence one has in particular

$$(4.5) \quad \text{codim } X(L) = \text{codim } Y(L) \leq \sum_p (m_p - 2),$$

where $\text{codim } X(L)$ means the codimension of $X(L)$ in the corresponding $X(d)$. For lattices L coming from line arrangements with few lines, or of a reduced complexity, the above inequality is an equality, see for an example Proposition 4.7 (1) below. However, the monomial arrangement

$$\mathcal{A}(m, m, 3) : f = (x^m - y^m)(x^m - z^m)(y^m - z^m) = 0,$$

has $d = 3m$, 3 points of multiplicity m and m^2 points of multiplicity 3. It follows that

$$\sum_p (m_p - 2) = 3(m - 2) + m^2 > 6m = \dim X(3m)$$

for $m \geq 5$. Hence for these values of m , the inequality (4.5) is strict.

REMARK 4.5. The variety $X(L)$ corresponds exactly to the variety of all ordered complex realizations $\Sigma^{ord}(C)$ of the ordered combinatorics C^{ord} considered in [5], where C^{ord} is the ordered combinatorial type associated to the lattice L with a fixed numbering of the lines. The quotient $X(L)/G$ is the ordered moduli space $\mathcal{M}^{ord}(C)$ considered in [5]. The variety $A(L)$ corresponds exactly to the variety of all complex realizations $\Sigma(C)$ of the combinatorics C as considered in [5], while $A(L)/G$ is the moduli space $\mathcal{M}(C)$ of the combinatorics C . If L is the lattice corresponding to the MacLane line arrangement, it follows from [5, Example 1.7] that $X(L)$ is the union of two G -orbits and in particular is not connected, while $A(L)$ is just one G -orbit, and hence it is irreducible.

EXAMPLE 4.6. For any d , we denote by L_{gen} the lattice of the generic line arrangement of d lines. Then by the above description $X(L_{gen})$ is a Zariski open subset of $X(d)$, and hence $\dim X(L_{gen}) = \dim X(d) = 2d$. Moreover $\tau(L_{gen}) = \binom{d}{2}$ and in fact one has $X(d, \tau(L_{gen})) = X(L_{gen})$, i.e. any lattice $L \in \mathcal{L}(d)$ with $\tau(L) = \tau(L_{gen})$ is in fact isomorphic to the lattice L_{gen} .

To prove this claim, recall the formula

$$(4.6) \quad \sum_p \binom{m_p}{2} = \binom{d}{2},$$

valid for any line arrangement, see for instance [22]. Since

$$\binom{m_p}{2} \leq (m_p - 1)^2$$

for any $m_p \geq 2$, and the equality holds if and only if $m_p = 2$, the claim follows using the formula (4.2). This argument implies also that $\tau(L) > \tau(L_{gen})$ for any lattice $L \in \mathcal{L}(d)$, $L \neq L_{gen}$.

Moreover, in this case it follows that the function $mdr \circ \psi$ is constant on $X(L_{gen})$, and it takes the value $d - 2$, see [13, Theorem 4.1], as well as all the functions m_k , since one has $ct(f) = st(f) = 2d - 4$ in this case, recall Corollary 2.6 (2) and Corollary 3.5.

This example can be generalized as follows. For any m satisfying $2 \leq m \leq d$, let $L(d, m)$ denote the intersection lattice of a line arrangement in $A(d)$ having one point of multiplicity m and only double points in rest. Note that $L(d, 2) = L_{gen}$ for any integer d .

Proposition 4.7. *Assume that $d \geq 4$. Then the following hold.*

- (1) *The sets $X(L(d, m))$ and $A(L(d, m))$ are smooth of dimension $(2d - m + 2)$. Moreover $A(L(d, m))$ is irreducible.*
- (2) *$\tau(L(d, m)) = (m - 1)^2 + m + (m + 1) + \dots + (d - 1) = \binom{d}{2} + \binom{m-1}{2}$.*
- (3) *For $2m > d$, the function mdr is constant on $A(L(d, m))$, and it takes the values $d - m$.*
- (4) *For $m = d$ and $m = d - 1$, any arrangement in $A(L(d, m))$ is free. Any arrangement in $A(L(d, d - 2))$ is nearly free. For $2 \leq m \leq d - 3$, any arrangement in $A(L(d, m))$ is neither free, nor nearly free.*
- (5) *Any line arrangement $\mathcal{A} : f = 0$ with $mdr(f) = 0$ satisfies $L(\mathcal{A}) = L(d, d)$. Any line arrangement $\mathcal{A} : f = 0$ with $mdr(f) = 1$ satisfies $L(\mathcal{A}) = L(d, d - 1)$.*
- (6) *$A(L(d, d)) = A(d, \tau(L(d, d)))$ and $A(L(d, d - 1)) = A(d, \tau(L(d, d - 1)))$.*

Proof. To get an arrangement in $A(L(d, m))$, we have first to fix a point $p \in \mathbb{P}^2$, and then m distinct lines passing through p . These choices are parametrized by $B = \mathbb{P}^2 \times U$, where U is an open subset in $(\mathbb{P}^1)^m$. Note that B is smooth of dimension $m + 2$ and irreducible. The remaining $d - m$ lines are to be chosen in a Zariski open set $F \subset (\mathbb{P}(S_1))^{d-m}$, which is smooth of dimension $2d - 2m$ and irreducible. In this way we have constructed a fibration $F \rightarrow A(L(d, m)) \rightarrow B$, proving the first claim (1).

Note that $X(L(d, m))$ is not connected in general. Indeed, for $d = 4$ and $m = 3$, we cannot continuously deform within $X(L(4, 3))$ an element $(\ell_1, \ell_2, \ell_3, \ell_4)$ where the lines $L_j : \ell_j = 0$ are concurrent for $j = 1, 2, 3$ to an element $(\ell'_1, \ell'_2, \ell'_3, \ell'_4)$ where the lines $L'_j : \ell'_j = 0$ are concurrent for $j = 2, 3, 4$.

The second claim follows from the formula (4.2). The third claim follows from [11, Theorem 1.2]. The claim (4) follows from the formula for $\tau(L(d, m))$ given in (2). Indeed, if a line arrangement $\mathcal{A} : f = 0$ in $A(d)$ is free, then one has $\tau(\mathcal{A}) = \tau(d, r)$ where $r = mdr(f)$, as explained in (3.6).

Suppose first that $2m > d$, and hence $r = d - m$. The formula for $\tau(L(d, m))$ given in (2)

shows that

$$(4.7) \quad \delta = \tau(d, r) - \tau(\mathcal{A}) = \frac{(d-m)(d-m-1)}{2}.$$

Hence we have the equality $\delta = 0$ only for $m = d$ or for $m = d - 1$. Assume now that $r \leq d - m - 1$. Then [11, Theorem 1.2] implies that either $r = m - 1$ or $m \leq r$. In the first case the arrangement is free with exponents $d_1 = m - 1$ and $d_2 = d - m$, and the equation (4.7) shows that this is possible only if $m = d - 1$ or $m = d$, which is impossible. In the second case, [11, Theorem 1.1] shows that $\tau(\mathcal{A}) \leq \tau(d, m)$ and equality holds exactly when $r = m$ and \mathcal{A} is free. A direct computation shows that

$$(4.8) \quad \delta' = \tau(d, m) - \tau(\mathcal{A}) = \frac{(d-m-2)(d-m-1)}{2} + (m-1) > 0$$

for $2 \leq m \leq d - 2$. For claim involving the nearly free arrangements, use the above and Remark 3.10.

The first part in claim (5) is clear, since $mdr(f) = 0$ if and only if f does not depend on the variable z after a coordinate change. If $mdr(f) = 1$, then let $m \geq 2$ be the maximal multiplicity of an intersection point in \mathcal{A} . Using [11, Theorem 1.2], we deduce that 3 cases are possible.

(a) The case $mdr(f) = d - m = 1$, which clearly settles our claim.

(b) The case $mdr(f) = m - 1$, impossible, since this would imply that \mathcal{A} is a generic arrangement, for which $mdr(f) = d - 2 \geq 2$.

(c) The case $m \leq mdr(f)$, which is clearly impossible.

For claim (6), let $\mathcal{B} : g = 0$ be a line arrangement in $A(d, \tau(L(d, d)))$ (resp. in $A(d, \tau(L(d, d-1)))$). Then $mdr(g) \geq 0$ (resp. $mdr(g) \geq 1$) and the claim follows using [11, Theorem 1.1], which, though not stated there, holds for $mdr(g) \geq 0$ as well. \square

Corollary 4.8. *For $d \geq 4$ and $m \in \{2, d - 1, d\}$, one has*

$$A(L(d, m)) = A(d, \tau(L(d, m))) = A(d, mdr = d - m)$$

and

$$X(L(d, m)) = X(d, \tau(L(d, m))) = X(d, mdr = d - m).$$

In the following definition we introduce two simple combinatorics for line arrangements.

DEFINITION 4.9. Let $\tilde{L}(m_1, m_2)$ be the lattice of a projective line arrangement \mathcal{A} obtained by the generic intersection of two pencils of m_1 , respectively m_2 lines, with $m_2 \geq m_1 \geq 2$. Let $\hat{L}(m_1, m_2)$ be the lattice of a projective line arrangement \mathcal{A} having exactly one line containing one point of multiplicity m_1 and one point of multiplicity m_2 , $m_2 \geq m_1 \geq 3$, only double points apart from that, and $d = |\mathcal{A}| = m_1 + m_2 - 1$.

An arrangement with intersection lattice $\tilde{L}(m_1, m_2)$ has $d = (m_1 + m_2)$ lines, $m_1 m_2$ double points, one point of multiplicity m_1 and one point of multiplicity m_2 . We prove next that such an arrangement is never free.

Proposition 4.10. *With this notation, one has the following.*

- (1) *The set $A(\tilde{L}(m_1, m_2))$ is smooth, irreducible of dimension $d + 4$.*

- (2) $\tau(\tilde{L}(m_1, m_2)) = (d - 1)^2 - m_1 m_2 + 1$.
- (3) *The function mdr is constant on $A(\tilde{L}(m_1, m_2))$ and takes the value m_1 .*
- (4) *The intersection $FA(d) \cap A(\tilde{L}(m_1, m_2))$ is empty.*

Proof. The first claim can be proved by a similar argument as that used in the proof of Proposition 4.7 (1). The second claim is obvious using the formula

$$\tau(\tilde{L}(m_1, m_2)) = (m_1 - 1)^2 + (m_2 - 1)^2 + m_1 m_2.$$

By [11, Theorem 1.2] (applied for $m = m_2$), either $mdr(f) = m_1$, or $mdr(f) \leq m_1 - 1$. In the second case, one has one of the following two possibilities.

- (i) The arrangement $\mathcal{A} : f = 0$ is free, $mdr(f) = m_2 - 1$ and $2m_2 < m_1 + m_2 + 1$, which implies $m_1 = m_2 = m$. Then the exponents of \mathcal{A} are $(m - 1, m)$ and hence the formula (3.1) implies

$$\tau(\tilde{L}(m_1, m_2)) = (d - 1)^2 - m(m - 1) = (d - 1)^2 - m^2 + m > (d - 1)^2 - m^2 + 1,$$

a contradiction with (2).

- (ii) $m_2 \leq mdr(f) \leq m_1 - 1$, contradiction with $m_1 \leq m_2$.

In conclusion, $mdr(f) = m_1$ and this proves (3). To prove (4), note that the formula (3) for $\tau(\tilde{L}(m_1, m_2))$ can be rewritten in the form

$$\tau(\tilde{L}(m_1, m_2)) = (d - 1)^2 - m_1(m_2 - 1) + 1 - m_1.$$

Then, by (3.1), the arrangement \mathcal{A} is not free, since $m_2 - 1 = d - m_1 - 1$ and $m_1 > 1$. □

EXAMPLE 4.11. We consider now a line arrangement $\mathcal{A} : f = 0$, having as intersection lattice the lattice $\hat{L}(m_1, m_2)$ introduced in Definition 4.9 above. By [11, Theorem 1.2] (applied for $m = m_2$), either $mdr(f) = m_1 - 1$, or $mdr(f) \leq m_1 - 2$. In the second case, it can only happen that $mdr(f) \leq m_1 - 2 \leq m_2 - 1$ and $2m_2 < m_1 + m_2$, contradiction with the assumption $m_1 \leq m_2$. So, $mdr(f) = m_1 - 1$ and $\tau(\hat{L}(m_1, m_2)) = (m_1 - 1)^2 + (m_2 - 1)^2 + (m_1 - 1)(m_2 - 1)$. We already know that such an arrangement \mathcal{A} is free, since it is supersolvable (see [27, Prop 5.114] and [23, Theorem 4.2]). See also [15] for this family of line arrangements.

In general one has $X(\hat{L}(m_1, m_2)) \neq X(m_1 + m_2, \tau(\hat{L}(m_1, m_2)))$. Indeed, by [15] any exponents $2 \leq d_1 \leq d_2$ of a free line arrangement can be obtained by such an arrangement. But there are free arrangements \mathcal{B} which are not of this type, e.g. the monomial arrangements $\mathcal{A}(m, m, 3)$ for $m \geq 2$ considered in Remark 4.4.

The classification of the line arrangements $\mathcal{A} : f = 0$ with $mdr(f) = 2$ is given by the following theorem, which is one of the main results in [34]. We give a proof of this classification from a new viewpoint.

Theorem 4.12. *Let $\mathcal{A} : f = 0$ be a line arrangement in \mathbb{P}^2 , with $mdr(f) = 2$. Then $d = |\mathcal{A}| \geq 4$ and \mathcal{A} is one of the following type of line arrangements, described by their intersection lattices.*

- (1) $\mathcal{A} \in A(L(d, d - 2))$, or
- (2) $\mathcal{A} \in A(\hat{L}(3, d - 2))$ with $d \geq 5$, or
- (3) \mathcal{A} is linear equivalent to the monomial arrangement $\mathcal{A}(2, 2, 3)$.

Proof. Let m be the maximal multiplicity of an intersection point in \mathcal{A} . If we denote $d = |\mathcal{A}| \geq 2$ and we assume $mdr(f) = 2$, then [11, Theorem 1.2] implies that only the following cases are possible.

CASE 1. $mdr(f) = d - m$, in other words $m = d - 2$. This case covers the two cases (i) and (ii) in [34, Theorem 2]. Indeed, the case (i) corresponds to the case when \mathcal{A} has a triple point except the point of multiplicity $m = d - 2$ (e.g. for $d = 5$ we have two triple points in \mathcal{A}), while the case (ii) corresponds to the case when \mathcal{A} has only double points except the point of multiplicity $m = d - 2$. In fact, the line arrangements of type (ii) are exactly the line arrangement in $A(L(d, d - 2))$ considered in Proposition 4.7 (4) above, in particular they are all nearly free. When $d \geq 5$, the arrangements of type (i) are exactly the line arrangements with the intersection lattice of type $\hat{L}(3, d - 2)$ which is discussed in Example 4.11 above, in particular they are all free.

CASE 2. $mdr(f) = m - 1$ and \mathcal{A} is free. In particular, this implies that the exponents of \mathcal{A} are $d_1 = 2 \leq d_2$, and hence $d \geq 5$. Moreover $m = 3$, and hence \mathcal{A} has n_2 double points, n_3 triple points and no points of multiplicity > 3 . Using the formulas (3.1) and (4.6), we get the equations

$$n_2 + 4n_3 = d^2 - 4d + 7 \text{ and } 2n_2 + 6n_3 = d^2 - d.$$

They imply that $n_2 = 10d - d^2 - 21 \geq 0$ which yields $d \leq 7$. It is easy to classify the free line arrangements with $5 \leq d \leq 7$ and only double and triple points and we get in this way the lattice $\hat{L}(3, 3)$ already seen above, and the case (3) in [34, Theorem 2], which is essentially the monomial arrangement $\mathcal{A}(2, 2, 3)$.

CASE 3. $m = mdr(f)$, in other words \mathcal{A} has only double points, as in Example 4.6 above. But one knows that in this case $mdr(f) = d - 2$, see [13, Theorem 4.1], and hence we get again the case (ii) from [34, Theorem 2] for $d = 4$. □

The above Theorem and Proposition 4.7 imply the following.

Corollary 4.13. *Any line arrangement $\mathcal{A} : f = 0$ with $mdr(f) \leq 2$ is either free or nearly free. Moreover, when $mdr(f) \leq 2$, the lattice $L(\mathcal{A})$ determines the values of $mdr(f)$ and whether \mathcal{A} is free or nearly free. In fact one has the following, where $d = |\mathcal{A}|$.*

- (1) $mdr(f) = 0$ if and only if $L(\mathcal{A}) = L(d, d)$;
- (2) $mdr(f) = 1$ if and only if $L(\mathcal{A}) = L(d, d - 1)$;
- (3) $mdr(f) = 2$ if and only if $L(\mathcal{A})$ is one of the lattices $L(d, d - 2)$, $\hat{L}(3, d - 2)$ or $L(\mathcal{A}(2, 2, 3))$.

REMARK 4.14. The line arrangements $\mathcal{A} : f = 0$ with $mdr(f) = 3$ can be classified using the same approach, but the number of possibilities is much higher. Moreover, there are line arrangements with $mdr(f) = 3$ which are neither free, nor nearly free, for instance the generic arrangement of 5 lines.

EXAMPLE 4.15. We introduce a final lattice type. For two integers $i \leq j$ we define a homogeneous polynomial in $\mathbb{C}[u, v]$ of degree $j - i + 1$ by the formula

$$(4.9) \quad g_{i,j}(u, v) = (u - iv)(u - (i + 1)v) \cdots (u - jv).$$

Consider the line arrangement $\mathcal{A} : f = 0$ of $d = m_1 + m_2 \geq 4$ lines in \mathbb{P}^2 given by

$$f(x, y, z) = x(y - z)g_{1,m_1-1}(x, y)g_{2,m_2}(x, z) = 0$$

for $2 \leq m_1 \leq m_2$. Denote by $L'(m_1, m_2)$ the corresponding intersection lattice $L(\mathcal{A})$. One can show that the following hold, see for instance [15].

- (1) The line arrangement \mathcal{A} has one point of multiplicity m_1 , one point of multiplicity m_2 , in addition to $(m_1 - 2)$ points of multiplicity 3 and $m_1(m_2 - 3) + 6$ nodes;
- (2) $mdr(f) = m_1$;
- (3) $\tau(f) = (d - 1)^2 - m_1(m_2 - 1) - 1$.

REMARK 4.16. We say that a lattice L is rigid if the corresponding constructible set $A(L)$ is the disjoint union of finitely many G -orbits. It is clear that if $\mathcal{A} : f = 0$ corresponds to a point in $A(L)$ with L rigid, any topologically constant deformation of \mathcal{A} is in fact a path in the connected component of $A(L)$ containing \mathcal{A} , which is by definition a G -orbit. It follows that any such line arrangement \mathcal{A} is topologically rigid. Notice that the lattice $L(d, m)$ is rigid for $d \geq 4$ if and only if either $(d, m) = (4, 2)$ or $(d, m) = (4, 3)$. This follows from Proposition 3.11 and Proposition 4.7. Other examples of rigid lattices L are given in the next section.

A case of special interest is when the Galois group \mathcal{G} acts transitively on the set of orbits in $A(L)$ for a rigid lattice L , see [2], [3], [5].

5. On the partition $A(d) = \cup_{L \in \mathcal{L}(d)} A(L)$

In this section we describe the partition $A(d) = \cup_{L \in \mathcal{L}(d)} A(L)$ for $4 \leq d \leq 6$, and show that the complexity of this partition increases rapidly with d .

5.1. The case $d = 4$. For $d = 4$, the list $\mathcal{L}(4)$ consists of 3 lattices, namely $L(4, 2)$, $L(4, 3)$ and $L(4, 4)$ in the notation from Proposition 4.7. Hence we have the following partition

$$A(4) = A(L(4, 2)) \cup A(L(4, 3)) \cup A(L(4, 4)),$$

where $\dim A(L(4, 2)) = 8$, $\dim A(L(4, 3)) = 7$ and $\dim A(L(4, 4)) = 6$. Moreover, the sets $A(L(4, 2))$ and $A(L(4, 3))$ are G -orbits, i.e. the corresponding arrangements are rigid, while $A(L(4, 4))$ is the union of a 1-parameter family of G -orbits, as can be seen using Proposition 3.11 and its proof. Recall also Example 3.13. Note that the closure of $A(L(4, 2))$ in $A(4)$ is the whole space $A(4)$, while the closure of $A(L(4, 3))$ in $A(4)$ is $A(L(4, 3)) \cup A(L(4, 4))$, which follows from Corollary 3.8. Moreover the set $A(L(4, 4))$ is closed in $A(4)$.

In this case, one has

$$6 = \tau(L(4, 2)) < 7 = \tau(L(4, 3)) < 9 = \tau(L(4, 4))$$

and hence the corresponding 3 strata are distinguished by their Tjurina numbers. Moreover, one has

$$FA(4) = A(L(4, 3)) \cup A(L(4, 4)).$$

Note also that even in this simple case, the set

$$A(4, st \leq 3) := \{f \in A(4) : st(f) \leq 3\} = A(L(4, 3))$$

is neither open nor closed. Hence the invariant st does not have nice semicontinuity properties as τ or mdr . By inspection of this list, we can state the following result.

Proposition 5.1. *With this notation, one has the following complete list of free and nearly free line arrangements for $d = 4$.*

- (1) *The set $A(4, 6)$ is open, 8-dimensional, and consists only of nearly free arrangements with exponents $d_1 = d_2 = 2$.*
- (2) *The set $A(4, 7)$ is irreducible, 7-dimensional, and consists only of free arrangements with exponents $d_1 = 1, d_2 = 2$.*
- (3) *The set $A(4, 9)$ is irreducible, 6-dimensional, and consists only of free arrangements with exponents $d_1 = 0, d_2 = 3$.*

5.2. The case $d = 5$. For $d = 5$, the list $\mathcal{L}(5)$ consists of $L(5, 2), L(5, 3), L(5, 4), L(5, 5)$ and an additional lattice $L = L(\mathcal{A})$ where $\mathcal{A} : xyz(x + y)(x + z) = 0$. Note that the lattice L is just the lattice $\hat{L}(3, 3)$ from Example 4.11. In this case one has

$$A(5, mdr = 2) = A(L(5, 3)) \cup A(L),$$

with $\dim A(L(5, 3)) = 9$ and $\dim A(L) = 8$. Hence Corollary 4.8 does not hold for $m = d - 2$ in this case. One also has

$$10 = \tau(L(5, 2)) < 11 = \tau(L(5, 3)) < 12 = \tau(L) < 13 = \tau(L(5, 4)) < 16 = \tau(L(5, 5)).$$

Hence again the corresponding 5 strata are distinguished by their Tjurina numbers. Moreover, one has

$$FA(5) = A(L) \cup A(L(5, 4)) \cup A(L(5, 5)).$$

Note that

$$\overline{A(L)} \cap A(L(5, 4)) = \emptyset.$$

Though this might be obvious for some readers, we prefer to give an argument which is likely to work in many similar situation. Note that, using the Curve Selection Lemma, if $\overline{A(L)} \cap A(L(5, 4)) \neq \emptyset$, then we get a deformation of an ordinary singular point $(Y_4, 0)$ of multiplicity 4 into two ordinary singular points $(Y_3, 0)$ of multiplicity 3. Such a deformation is impossible, since it would contradict the semicontinuity of the spectrum on the interval $I = (-1/3, 2/3]$, see for details [24, Theorem (8.9.8)]. Indeed, one has

$$1 = \deg_I \text{spec}(Y_4) < 2 \deg_I \text{spec}(Y_3) = 2.$$

It follows that

$$\overline{A(L)} \cap \overline{A(L(5, 4))} = A(L(5, 5)).$$

By inspection of the list of lattices in $\mathcal{L}(5)$, we can state the following result.

Proposition 5.2. *With this notation, one has the following complete list of free and nearly free line arrangements for $d = 5$.*

- (1) *The set $A(5, 11)$ is irreducible, 9-dimensional, and consists only of nearly free arrangements with exponents $d_1 = 2, d_2 = 3$. Any $f \in X(5, 11)$ satisfies $st(f) = 6$ and $reg(f) = 5$.*

- (2) The set $A(5, 12)$ is irreducible, 8-dimensional, and consists only of free arrangements with exponents $d_1 = d_2 = 2$. Any $f \in X(5, 12)$ satisfies $st(f) = reg(f) = 4$.
- (3) The set $A(5, 13)$ is irreducible, 8-dimensional, and consists only of free arrangements with exponents $d_1 = 1, d_2 = 3$. Any $f \in X(5, 13)$ satisfies $st(f) = reg(f) = 5$.
- (4) The set $A(5, 16)$ is irreducible, 7-dimensional, and consists only of free arrangements with exponents $d_1 = 0, d_2 = 4$. Any $f \in A(5, 16)$ satisfies $st(f) = reg(f) = 6$.

Note that among the above sets, only $A(5, 12)$ is a G -orbit and hence the corresponding arrangements are rigid. The arrangements in $A(5, 13)$ are algebraically rigid, but not topologically rigid, recall Proposition 3.11.

REMARK 5.3. It is clear that $A(5, 12)$ is contained in the closure of $A(5, 11)$, and that $A(5, 16)$ is contained in the closure of $A(5, 12)$. The values given above for $st(f)$ and $reg(f)$ show that these invariants do not enjoy simple semicontinuity properties as in Proposition 2.4.

5.3. The case $d = 6$. For $d = 6$, the list $\mathcal{L}(6)$ consists of 10 lattices. We list them in increasing order of their Tjurina numbers.

- For $\tau = 15$, we have only the lattice $L(6, 2)$ as predicted by the general theory, recall Example 4.6. Moreover $X(L(6, 2))$ is an open subset in the 12-dimensional smooth variety $X(6)$

- For $\tau = 16$, we have only the lattice $L(6, 3)$ and the corresponding set $X(L(6, 3))$ has codimension 1 in $X(6)$.

- For $\tau = 17$, we have two lattices, namely $\tilde{L}(3, 3)$ and a new lattice, say $\tilde{L}'(3, 3)$. These two lattices have each 2 triple points and 9 nodes, and the invariant mdr takes the value 3 in both cases. In the lattice $\tilde{L}(3, 3)$ the 2 triple points are not on a line of the corresponding arrangement, while in the lattice $\tilde{L}'(3, 3)$ the 2 triple points are on such a line. In conclusion the corresponding two sets $X(\tilde{L}(3, 3))$ and $X(\tilde{L}'(3, 3))$ are not distinguished by the numerical invariants considered in this paper. Indeed, since $ct(f) = 7$ and $st(f) = 8$ in both cases, the invariants m_k 's also coincide for any k .

Both sets $X(\tilde{L}(3, 3))$ and $X(\tilde{L}'(3, 3))$ have codimension 2 in $X(6)$

- For $\tau = 18$, we have again two lattices, namely the lattice $L(6, 4)$, having a point of multiplicity 4, 9 nodes and $mdr = 2$ and the lattice $L'(3, 3)$ introduced in Example 4.15, and having 3 triple points, 6 nodes and $mdr = 3$.

The set $X(L(6, 4))$ has codimension 2 in $X(6)$, while the set $X(L'(3, 3))$ has codimension 3 in $X(d)$.

Comparing the values of $\tau(L) \leq 18$ and the corresponding values of the invariant mdr , we conclude that there are no free arrangements in this range.

- For $\tau = 19$, we have again two lattices, namely the lattice $\hat{L}(3, 4)$, having one point of multiplicity 4, one triple point and 6 nodes, and the lattice $L(\Delta)$ corresponding to the arrangement

$$\mathcal{A} : f = (x^2 - y^2)(x^2 - z^2)(y^2 - z^2) = 0,$$

and hence having 4 triple points and 3 nodes. Both of the corresponding sets $X(L)$ contain only free arrangements with $mdr = 2$.

- For $\tau = 21$, we have only the lattice $L(6, 5)$.

• For $\tau = 25$, we have only the lattice $L(6, 6)$. The properties of the last two lattices are discussed in Proposition 4.7 (4), (5) and (6). In particular, the sets $X(d, \tau)$ for $\tau = 19, 21, 25$ consist only of free arrangements, i.e. the last claim in Corollary 3.8 holds in a stronger version.

As a conclusion, we can state the following result.

Proposition 5.4. *With this notation, one has the following complete list of free and nearly free line arrangements for $d = 6$.*

- (1) *The set $A(6, 18)$ has two irreducible components, namely $A(L(6, 4))$ of dimension 10, and $A(L'(3, 3))$ of dimension 9; they consist only of nearly free arrangements with exponents $d_1 = 2, d_2 = 4$, and respectively $d_1 = d_2 = 3$.*
- (2) *The set $A(6, 19)$ has two irreducible components, namely $A(\hat{L}(3, 4))$ of dimension 9, and $A(L(\Delta))$ of dimension 8; they consist only of free arrangements with exponents $d_1 = 2, d_2 = 3$.*
- (3) *The set $A(6, 21)$ is irreducible, 9-dimensional, and consists only of free arrangements with exponents $d_1 = 1, d_2 = 4$.*
- (4) *The set $A(6, 25)$ is irreducible, 8-dimensional, and consists only of free arrangements with exponents $d_1 = 0, d_2 = 5$.*

Note that $Y = A(L(\Delta))$ is the only G -orbit in the list above, and hence consists only of rigid arrangements. Moreover $Z = \bar{Y} \setminus Y$ is a closed G -invariant subset of dimension < 8 . Using the same type of argument as in the case $d = 5$ above, one can show that $Z \subset A(L(6, 6))$. Since $A(L(6, 6))$ is 8-dimensional by Proposition 4.7 (1), it follows that Z is not a union of strata in the partition. This shows in particular that this partition is not Whitney regular, see [7, Chapter 1] for basic facts on regular stratifications.

REMARK 5.5. It is possible to extend this discussion to $d = 7$. For $d < 7$ we have seen that the closed set $X(d, \tau \geq \tau(d)_{\min})$ contains only free arrangements. For $d = 7$, one has $\tau(d)_{\min} = 27$. The new aspect occurring in this case is that the set $X(7, \tau \geq 27)$ contain free arrangements and one nearly free arrangement type with exponents $d_1 = 2, d_2 = 5$.

The above leads us to ask the following.

QUESTION 5.6. Is it true that, for any $d \geq 4$, the closed set $X(d, \tau \geq \tau(d)_{\min})$ in $X(d)$ contains only nearly free and free arrangements with d lines?

Unfortunately the answer to this question is negative. To see this, it is enough to consider the arrangements in $X(L(d, d-3))$ for $d \geq 11$. Then using Proposition 4.7 it is easy to check that $\tau(L(d, d-3)) > \tau(d)_{\min}$. On the other hand, the formula (4.7) and the characterization of free arrangements (resp. nearly free arrangements) by the property $\delta = 0$ (resp. $\delta = 1$) given in [10] show that any arrangement in $X(L(d, d-3))$ for $d \geq 11$ is neither free nor nearly free.

6. On Terao's conjecture

With the above notation, this conjecture in the case of line arrangements can be stated as follows.

CONJECTURE 6.1. [Terao’s Conjecture for the line arrangement \mathcal{A}] Let \mathcal{A} be a free line arrangement with $d = |\mathcal{A}|$. Then

$$A(L(\mathcal{A})) \subset FA(d).$$

Equivalently, $X(L(\mathcal{A})) \subset FX(d)$.

Assume that \mathcal{A} is free with exponents $d_1 \leq d_2$. Then the following are known.

Theorem 6.2. *Terao’s conjecture holds for the line arrangement \mathcal{A} if one has either $d = |\mathcal{A}| \leq 12$, or $d_1 \leq 5$.*

For the proofs of this result, see [1], [2], [19]. The next result was proved in [11], but we give below a new, clearer proof. Another rapid proof can be obtained by combining Theorem 2.7 and Lemma 2.10 in [2]. Let $m(\mathcal{A})$ denote the maximal multiplicity of an intersection point in \mathcal{A} .

Theorem 6.3. *Terao’s conjecture holds for the line arrangement \mathcal{A} if $m(\mathcal{A}) \geq d_1$.*

Proof. We apply first [11, Theorem 1.2] to the free arrangement \mathcal{A} . It follows that

$$(6.1) \quad d_1 \in \{d - m, m - 1, m\},$$

where $m = m(\mathcal{A})$. It follows that $m \in \{d - d_1, d_1, d_1 + 1\}$. Let now $\mathcal{B} \in A(L(\mathcal{A}))$, given by $g = 0$, and note that $m(\mathcal{B}) = m$. Now we apply [11, Theorem 1.2] to the line arrangement \mathcal{B} .

If we are in the case $mdr(g) = d - m(\mathcal{B}) = d - m \in \{d_1, d - d_1, d - d_1 - 1\}$, it follows that $mdr(g) \geq mdr(f) = d_1$. Since $\tau(\mathcal{B}) = \tau(\mathcal{A})$, the result [11, Theorem 1.1] implies that \mathcal{B} is free with the same exponents $d_1 \leq d_2$.

If we are in the case $mdr(g) = m(\mathcal{B}) - 1$, then the arrangement \mathcal{B} is free, so there is nothing to prove. Finally, we have to consider the case $m = m(\mathcal{B}) \leq mdr(g) \leq d - m - 1$. For $m = d - d_1$ this implies $d - d_1 \leq d_1 - 1$, which is impossible. For $m = d_1$ or $m = d - d_1 - 1 = d_2$, we are again in the case $mdr(g) \geq mdr(f) = d_1$, and we conclude as above. \square

As explained in [11], this Theorem implies the following.

Corollary 6.4. *With the above notation, one has the following.*

- (1) *Terao’s conjecture holds for the line arrangement \mathcal{A} if $m(\mathcal{A}) \geq d/2$.*
- (2) *Terao’s conjecture holds for the line arrangement \mathcal{A} if $d_1 \leq \sqrt{2d + 1} - 1$.*

We end with a result saying that a free arrangement cannot have too many singularities.

Proposition 6.5. *The intersection $FX(d) \cap X(L)$ is empty if*

$$(6.2) \quad \tau(L) < \frac{3}{4}(d - 1)^2.$$

In particular, the inequality (6.2) holds if

$$\sum_p (m_p - 1) > \frac{(d + 3)(d - 1)}{4}$$

where p runs through the set of multiple points of the lattice L , and $m_p \geq 2$ denotes the multiplicity of p .

Proof. The first claim follows from Corollary (3.9).

Then the formula for $\tau(L)$ given in (4.2) and the equality (4.6) imply that

$$\tau(L) = 2\binom{d}{2} - \sum_p (m_p - 1) < 2\binom{d}{2} - \frac{(d+3)(d-1)}{4} = \frac{3}{4}(d-1)^2.$$

□

EXAMPLE 6.6. Assume that the line arrangement \mathcal{A} is not generic, but has a lot of nodes, namely it has $N > \frac{(d+3)(d-1)}{4} - 2$ nodes, besides some other multiple points. Then \mathcal{A} is not free by the above result, since there is at least one point p with $m_p \geq 3$. When $d = 7$, this says that an arrangement \mathcal{A} having at least 14 nodes satisfies

$$\tau(\mathcal{A}) < \tau(7)_{\min} = \frac{3}{4}(d-1)^2 = 27$$

and hence it is not free. A detailed classification of the line arrangements \mathcal{B} with $|\mathcal{B}| = 7$, shows that there is a nearly free arrangement \mathcal{B} having 11 nodes and one point of multiplicity 5 such that $\tau(\mathcal{B}) = 27$. Moreover, for all arrangements \mathcal{B}' having 12 nodes (and some other multiple points), one has $\tau(\mathcal{B}') < 27$. Hence our bound is two units apart from the optimal one in this case.

References

- [1] T. Abe: *Roots of characteristic polynomials and intersection points of line arrangements*, J. Singul. **8** (2014), 100–117.
- [2] T. Abe, M. Cuntz, H. Kawanoue and T. Nozawa: *Non-recursive freeness and non-rigidity of plane arrangements*, arXiv:1411.3351.
- [3] M. Amram, M. Cohen, M. Teicher and F. Ye: *Moduli spaces of ten-line arrangements with double and triple points*, arXiv:1306.6105.
- [4] M. Amram, M. Teicher and F. Ye: *Moduli spaces of arrangements of 10 projective lines with quadruple points*, Adv. in Appl. Math. **51** (2013), 392–418.
- [5] E. Artal Bartolo, J. Carmona Ruber, J.I. Cogolludo Agustin and M.A. Marco Buzunariz: *Invariants of combinatorial line arrangements and Rybnikov’s example*; in Singularity theory and its applications, Adv. Stud. Pure Math. **43**, Math. Soc. Japan, Tokyo, 2006, 1–34.
- [6] A.D.R. Choudary and A.Dimca: *Koszul complexes and hypersurface singularities*, Proc. Amer. Math. Soc. **121** (1994), 1009–1016.
- [7] A. Dimca: *Singularities and topology of hypersurfaces*, Universitext, Springer-Verlag, New York, 1992.
- [8] A. Dimca: *Syzygies of Jacobian ideals and defects of linear systems*, Bull. Math. Soc. Sci. Math. Roumanie **56** (104) (2013), 191–203.
- [9] A. Dimca: *Hyperplane Arrangements: An Introduction*, Universitext, Springer, 2017.
- [10] A. Dimca: *Freeness versus maximal global Tjurina number for plane curves*, Math. Proc. Cambridge Phil. Soc. **163** (2017), 161–172.
- [11] A. Dimca: *Curve arrangements, pencils, and Jacobian syzygies*, Michigan Math. J. **66** (2017), 347–365.
- [12] A. Dimca and E. Sernesi: *Syzygies and logarithmic vector fields along plane curves*, J. Éc. polytech. Mat. **1** (2014), 247–267.
- [13] A. Dimca and G. Sticlaru: *Koszul complexes and pole order filtrations*, Proc. Edinb. Math. Soc. **58** (2015), 333–354.
- [14] A. Dimca and G. Sticlaru: *Free divisors and rational cuspidal plane curves*, Math. Res. Lett. **24** (2017), 1023–1042.

- [15] A. Dimca and G. Sticlaru: *On the exponents of free and nearly free projective plane curves*, Rev. Mat. Complut. **30** (2017), 259–268.
- [16] A.A. du Plessis and C.T.C. Wall: *Curves in $P^2(\mathbb{C})$ with 1-dimensional symmetry*, Rev. Mat. Complut. **12** (1999), 117–132.
- [17] D. Eisenbud: *Commutative Algebra*, Graduate Texts in Mathematics **150**, Springer NY 1995.
- [18] D. Eisenbud: *The Geometry of Syzygies: A Second Course in Algebraic Geometry and Commutative Algebra*, Graduate Texts in Mathematics, **229**, Springer, 2005.
- [19] D. Faenzi and J. Vallès: *Logarithmic bundles and line arrangements, an approach via the standard construction*, J. Lond. Math. Soc. **90** (2014), 675–694.
- [20] G.-M. Greuel, C. Lossen and E. Shustin: *Singular Algebraic Curves*, with an Appendix by O. Viro, Springer, Cham, 2018.
- [21] S.H. Hassanzadeh and A. Simis: *Plane Cremona maps: saturation and regularity of the base ideal*, J. Algebra **371** (2012), 620–652.
- [22] F. Hirzebruch: *Arrangements of lines and algebraic surfaces*; in Arithmetic and Geometry, Vol. II, Progress in Mathematics **36**, Birkhäuser, Boston, 1983, 113–140.
- [23] M. Jambu and H. Terao: *Free arrangements of hyperplanes and supersolvable lattices*, Adv. in Math. **52** (1984), 248–258.
- [24] V. Kulikov: *Mixed Hodge Structures and Singularities*, Cambridge Tracts in Mathematics **132**, Cambridge University Press, Cambridge, 1998.
- [25] V. Kulikov and E. Shustin: *On rigid plane curves*, Eur. J. Math. **2** (2016), 208–226.
- [26] S. Nazir and M. Yoshinaga: *On the connectivity of the realization spaces of line arrangements*, Ann. Sc. Norm. Super. Pisa Cl Sci. (5) **11** (2012), 921–937.
- [27] P. Orlik and H. Terao: *Arrangements of Hyperplanes*, Springer-Verlag, Berlin, 1992.
- [28] S. Papadima and A.I. Suciu: *The Milnor fibration of a hyperplane arrangement: from modular resonance to algebraic monodromy*, Proc. Lond. Math. Soc. **114** (2017), 961–1004.
- [29] H.K. Schenck: *Elementary modifications and line configurations in P^2* , Comment. Math. Helv. **78** (2003), 447–462.
- [30] H. Schenck: *Hyperplane arrangements: computations and conjectures*; in Arrangements of hyperplanes–Sapporo 2009, Adv. Stud. Pure Math. **62**, Math. Soc. Japan, Tokyo, 2012, 323–358.
- [31] E. Sernesi: *Deformations of Algebraic Schemes*, Grundlehren der Mathematischen Wissenschaften **334**, Springer-Verlag, Berlin, 2006.
- [32] E. Sernesi: *The local cohomology of the Jacobian ring*, Doc. Math. **19** (2014), 541–565.
- [33] A. Simis and S.O. Tohăneanu: *Homology of homogeneous divisors*, Israel J. Math. **200** (2014), 449–487.
- [34] S.O. Tohăneanu: *Projective duality of arrangements with quadratic logarithmic vector fields*, Discrete Math. **339** (2016), 54–61.
- [35] F. Ye: *Classification of moduli spaces of arrangements of nine projective lines*, Pacific J. Math. **265** (2013), 243–256.
- [36] M. Yoshinaga: *Freeness of hyperplane arrangements and related topics*, Ann. Fac. Sci. Toulouse Math. (6) **23** (2014), 483–512.
- [37] S. Yuzvinsky: *Free and locally free arrangements with a given intersection lattice*, Proc. Amer. Math. Soc. **118** (1993), 745–752.
- [38] G. Ziegler: *Combinatorial construction of logarithmic differential forms*, Adv. Math. **76** (1989), 116–154.

Alexandru Dimca
Université Côte d'Azur
CNRS, LJAD
France
e-mail: dimca@unice.fr

Denis Ibadula
Ovidius University, Faculty of Mathematics and Informatics
124 Mamaia Blvd.
900527 Constanța
Romania
e-mail: denis.ibadula@univ-ovidius.ro

Daniela Anca Măcinic
Simion Stoilow Institute of Mathematics
P.O. Box 1-764
RO-014700 Bucharest
Romania
e-mail: Anca.Macinic@imar.ro