Title	Public-Access Defibrillation in Japan
Author(s)	Kitamura, Tetsuhisa; Kiyohara, Kosuke; Iwami, Taku
Citation	New England Journal of Medicine. 376(7) p.e12
Issue Date	2017-02-16
oaire:version	VoR
URL	https://hdl.handle.net/11094/78398
rights	© 2017 Massachusetts Medical Society. Reprinted with permission.
Note	

Osaka University Knowledge Archive : OUKA

<https://ir.library.osaka-u.ac.jp/>

Osaka University

CORRESPONDENCE

Public-Access Defibrillation in Japan

TO THE EDITOR: In their study of the effect of public-access defibrillation on outcomes after out-of-hospital cardiac arrest in Japan, Kitamura et al. (Oct. 27 issue)¹ found an improved survival rate when a publicly accessible automated external defibrillator (AED) was used in the field. However, the frequency of AED use by bystanders before the arrival of emergency medical service (EMS) personnel remained very poor (approximately 2% of witnessed arrests of cardiac origin).

As compared with patients who did not receive public-access defibrillation, those who did had nearly twice the rate of cardiopulmonary resuscitation (CPR) (99.4% vs. 51.3%, $P < 0.001$), and the bystanders required dispatcher instructions less frequently during CPR. These findings suggest that patients who received public-access defibrillation were treated by trained bystanders, who were aware of the importance of CPR and of how to perform it.

The higher survival rate observed in this group highlights the importance of education, which most probably improved the rates of both public-access AED use and bystander CPR. In addition to AED use, bystander education constitutes the other component of any public-access defibrillation program^{2,3} and must be a priority.

Nicole Karam, M.D., M.P.H.

Xavier Jouven, M.D., Ph.D.

Eloi Marijon, M.D., Ph.D.

European Georges Pompidou Hospital
Paris, France
eloi_marijon@yahoo.fr

No potential conflict of interest relevant to this letter was reported.

1. Kitamura T, Kiyohara K, Sakai T, et al. Public-access defibrillation and out-of-hospital cardiac arrest in Japan. *N Engl J Med* 2016;375:1649-59.

2. Nichol G, Hallstrom AP, Kerber R, et al. American Heart Association report on the second public access defibrillation conference, April 17-19, 1997. *Circulation* 1998;97:1309-14.

3. The Public Access Defibrillation Trial Investigators. Public-access defibrillation and survival after out-of-hospital cardiac arrest. *N Engl J Med* 2004;351:637-46.

DOI: 10.1056/NEJMc1700160

TO THE EDITOR: Missing from the report by Kitamura et al. on survival with public-access defibrillation in Japan was a consideration of the cost-effectiveness of the program. The presence of 428,821 AEDs was credited with producing 201 survivors of out-of-hospital cardiac arrest with a favorable neurologic outcome in 2013. The typical price of an AED is \$1,500 to \$2,000, with an average warranty of 7 years.¹ Ongoing costs include the battery, at \$400 with a life of 4 years, and pads, at \$50 with a life of 2 years.¹ Additional expenses include maintenance (periodic tests and inspections), administrative costs, and training costs for personnel. I estimate an annual cost per AED of at least \$500, yielding a cost per survivor in excess of \$1 million. At an average age of 65 years, survivors might gain an additional 10 years of life,² yielding a cost of \$100,000 per year of life gained. Adjusting for quality of life and increased health care costs in these older, fairly sick persons (most of whom would have to live with an implanted defibrillator) would raise the cost considerably above \$100,000. This amount of money seems to exceed the commonly accepted definitions of cost-effective treatment.

Mayer Bassan, M.D.

11 Windmill St.
Jerusalem, Israel
mayer_bassan@yahoo.co.uk

No potential conflict of interest relevant to this letter was reported.

1. Altra Medical. AED cost of ownership and cost comparison. 2016 (<http://www.altramedical.com/aed-cost-of-ownership/>).

2. Bunch TJ, White RD, Gersh BJ, et al. Long-term outcomes of out-of-hospital cardiac arrest after successful early defibrillation. *N Engl J Med* 2003;348:2626-33.

DOI: 10.1056/NEJMc1700160

TO THE EDITOR: We have two concerns with the study by Kitamura et al. First, the authors adjust for some cardiac arrest characteristics but not for the location of arrest. A good outcome is more likely in public settings than in private settings.^{1,2}

Although some of this association may be explained by factors such as the use of bystander CPR and the initial rhythm, for which the authors adjusted, other important factors such as coexisting conditions and the quality of bystander CPR, which may be related to the location, were not adjusted for in the study. Because the location of arrest is included as an Utstein variable,³ we urge the authors to adjust for this.

Second, the authors included patients with witnessed, ventricular-fibrillation cardiac arrest with a cardiac cause. Because AEDs are applied to all cardiac arrests without prior knowledge of the rhythm or cause of arrest, the study by Kitamura et al. gives a misleading picture of the potential benefit of AEDs. The use of AEDs in patients with a nonshockable rhythm may be harmful owing to delays and interruptions in CPR.⁴ We encourage the authors to analyze the entire cohort.

Lars W. Andersen, M.D., Ph.D.

Aarhus University Hospital
Aarhus, Denmark
lwandersen@clin.au.dk

Bo Løfgren, M.D., Ph.D.

Regional Hospital Randers
Randers, Denmark

Asger Granfeldt, M.D., D.M.Sc.

Aarhus University Hospital
Aarhus, Denmark

No potential conflict of interest relevant to this letter was reported.

1. Weisfeldt ML, Everson-Stewart S, Sitlani C, et al. Ventricular tachyarrhythmias after cardiac arrest in public versus at home. *N Engl J Med* 2011;364:313-21.

2. Murakami Y, Iwami T, Kitamura T, et al. Outcomes of out-of-hospital cardiac arrest by public location in the public-access defibrillation era. *J Am Heart Assoc* 2014;3(2):e000533.

3. Jacobs I, Nadkarni V, Bahr J, et al. Cardiac arrest and cardiopulmonary resuscitation outcome reports: update and simplification of the Utstein templates for resuscitation registries: a statement for healthcare professionals from a task force of the International Liaison Committee on Resuscitation (American Heart Association, European Resuscitation Council, Australian Resuscitation Council, New Zealand Resuscitation Council, Heart and Stroke Foundation of Canada, InterAmerican Heart Foundation, Resuscitation Council of Southern Africa). *Resuscitation* 2004;63:233-49.

4. Chan PS, Krumholz HM, Spertus JA, et al. Automated external defibrillators and survival after in-hospital cardiac arrest. *JAMA* 2010;304:2129-36.

DOI: 10.1056/NEJMc1700160

public-access defibrillation programs. Efforts to increase the number of trained citizens would increase the rate of both CPR and AED use by lay rescuers, leading to an increase in survival with a favorable neurologic outcome after out-of-hospital cardiac arrest.

The economics of public-access defibrillation is an important issue, as suggested by Bassan. The cost-effectiveness of public-access defibrillation programs is still controversial. The Public Access Defibrillation Trial estimated that the cost-effectiveness of public-access defibrillation was similar to that of other medical interventions,¹ whereas another report mentioned that a nationwide public-access defibrillation program including dissemination of public-access AEDs is less likely to be cost-effective.² Importantly, the cost-effectiveness of public-access defibrillation is influenced by not only the number of AEDs but also the rate of AED use and the quality of both CPR and AED use. Therefore, cost-effectiveness should be examined after making efforts to maximize the rate of AED use with bystander education or the effective use of social-media technologies.³

Andersen et al. raise concerns about the lack of information on arrest location and on overall AED use (i.e., AED-pad application). Unfortunately, unlike the Utstein Osaka Project registry,^{4,5} the All-Japan Utstein Registry of the Fire and Disaster Management Agency did not have data on location or on AED pad application. Therefore, our study used type of bystander (family member or other) as a surrogate for location in the multivariable model, and this variable was well-balanced between the groups in a propensity-score-matching analysis (Table 1 of our article). In addition, because we do not have information on AED pad application, we focused on patients with bystander-witnessed ventricular-fibrillation arrests as the target population. The use of AEDs in patients with nonshockable rhythms may have a negative effect, whereas AED use may be beneficial if it encourages lay rescuers to perform CPR. Analysis of the entire cohort, with data on AED pad application, would be needed to address this question, as suggested by Andersen et al.

Tetsuhisa Kitamura, M.D., D.P.H.

Osaka University
Suita, Japan

THE AUTHORS REPLY: We agree with Karam et al. about the importance of bystander education in

Kosuke Kiyohara, D.P.H.

Tokyo Women's Medical University
Tokyo, Japan

Taku Iwami, M.D., Ph.D.

Kyoto University
Kyoto, Japan
iwami.taku.8w@kyoto-u.ac.jp

Since publication of their article, the authors report no further potential conflict of interest.

1. Nichol G, Huszti E, Birnbaum A, et al. Cost-effectiveness of lay responder defibrillation for out-of-hospital cardiac arrest. *Ann Emerg Med* 2009;54:226-35.
2. Moran PS, Teljeur C, Masterson S, O'Neill M, Harrington P,

Ryan M. Cost-effectiveness of a national public access defibrillation programme. *Resuscitation* 2015;91:48-55.

3. Zijlstra JA, Stieglis R, Riedijk F, Smeeke M, van der Worp WE, Koster RW. Local lay rescuers with AEDs, alerted by text messages, contribute to early defibrillation in a Dutch out-of-hospital cardiac arrest dispatch system. *Resuscitation* 2014;85:1444-9.

4. Kiyohara K, Kitamura T, Sakai T, et al. Public-access AED pad application and outcomes for out-of-hospital cardiac arrests in Osaka, Japan. *Resuscitation* 2016;106:70-5.

5. Murakami Y, Iwami T, Kitamura T, et al. Outcomes of out-of-hospital cardiac arrest by public location in the public-access defibrillation era. *J Am Heart Assoc* 2014;3(2):e000533.

DOI: 10.1056/NEJMc1700160

Correspondence Copyright © 2017 Massachusetts Medical Society.