

Title	Cohomology of discrete subgroups of $Sp(p, q)$
Author(s)	Konno, Yasuko
Citation	Osaka Journal of Mathematics. 1988, 25(2), p. 299-318
Version Type	VoR
URL	<a href="https://doi.org/10.18910/7870">https://doi.org/10.18910/7870</a>
rights	
Note	

*Osaka University Knowledge Archive : OUKA*

<https://ir.library.osaka-u.ac.jp/>

Osaka University

## COHOMOLOGY OF DISCRETE SUBGROUPS OF $Sp(p, q)$

Dedicated to Professor Shingo Murakami on his sixtieth birthday

YASUKO KONNO

(Received January 19, 1987)

### Introduction

Let  $G$  be a connected semi-simple Lie group with finite center and no compact factors. Let  $\Gamma$  be a uniform discrete subgroup of  $G$  and  $(\rho, F)$  be a finite dimensional irreducible representation of  $G$ . We are interested in the cohomology space  $H^*(\Gamma, F)$ . The purpose of this paper is to prove a non-vanishing theorem for  $H^*(\Gamma, F)$  in the case of  $G=Sp(p, q)$  ( $p \geq q \geq 1$ ).

As it is well-known, we can describe  $H^*(\Gamma, F)$  in terms of the relative Lie algebra cohomology. Let  $\mathfrak{g}$  be the Lie algebra of  $G$  and  $K$  be a maximal compact subgroup of  $G$ . Denote by  $\hat{G}$  the unitary dual of  $G$ . For  $(U, H_U) \in \hat{G}$ , we denote by  $H_U^0$  the space of  $K$ -finite vectors in  $H_U$ . Then  $H_U^0$  is an irreducible  $(\mathfrak{g}, K)$ -module. Also  $m(U, \Gamma)$  denotes the multiplicity with which  $U$  occurs in  $L^2(\Gamma \backslash G)$ . Define the subset  $\hat{G}_\rho$  of  $\hat{G}$  as follows;

$$\hat{G}_\rho = \{U \in \hat{G} | \chi_U = \chi_{\rho^*}\}$$

where  $\rho^*$  is the contragredient representation of  $\rho$  and  $\chi_U$  (resp.  $\chi_{\rho^*}$ ) is the infinitesimal character of  $U$  (resp.  $\rho^*$ ). Then, from the formula of Matsu-shima-Murakami ([1], VII, Theorem 6.1), we have

$$(0.1) \quad H^*(\Gamma, F) = \sum_{U \in \hat{G}_\rho} m(U, \Gamma) H^*(\mathfrak{g}, K; H_U^0 \otimes F).$$

From now on, we assume that  $G$  is simple. Depending on Kumaresan's work, Vogan and Zuckerman obtained the following precise vanishing theorem for the  $(\mathfrak{g}, K)$ -cohomology ([5], Theorem 8.1); if  $U$  is non-trivial, we have

$$H^i(\mathfrak{g}, K; H_U^0 \otimes F) = \{0\} \quad (i < r_G)$$

where  $r_G$  is the positive integer determined by  $G$  and given by Table 8.2 in [5] for non-complex groups. From this result and (0.1), if  $F$  is non-trivial, we have

$$H^i(\Gamma, F) = \{0\} \quad (i < r_G).$$

Note that  $r_G$  depends only on  $G$  and, in general,  $r_G \geq \text{rank}_{\mathbf{R}} G$ . On the other hand, the vanishing of  $H^i(\Gamma, F)$  below the  $\mathbf{R}$ -rank has been obtained in some papers ([1], VII, Proposition 6.4). There are some simple groups such that  $r_G = \text{rank}_{\mathbf{R}} G$ . In the case of  $G = SU(p, q)$  ( $p \geq q \geq 1$ ), where  $r_G = q = \text{rank}_{\mathbf{R}} G$ , Borel and Wallach showed that this vanishing theorem is best possible ([1], VIII, Corollary 5.9).

We concentrate our attention on the case of  $G = Sp(p, q)$ . In this case,  $r_G = 2q$  and hence  $r_G > q = \text{rank}_{\mathbf{R}} G$ . Therefore it is interesting to ask if the above vanishing theorem is best possible for  $G = Sp(p, q)$ . In this paper, we show that, in the case of  $G = Sp(p, q)$ , the first possible non-zero cohomology  $H^*(\Gamma, F)$  appears indeed at the degree  $2q = r_G$ . Main results are Theorem 3.4 and Theorem 4.2. In the case that  $F$  is trivial and  $q = 1$ , Theorem 3.4 is contained in the results of [3], Theorem 3.2 (see Remark 3.5). Also Theorem 4.2 for trivial  $F$  improves a part of the results of [4], Theorem 4.1 (see Remark 4.4).

Our method is similar to that in [1], VIII and depends heavily on the results there.

**1. The imbedding of  $Sp(p, q)$  into  $Sp(2n, \mathbf{R})$**

**1.1.** Throughout this paper,  $G$  will denote the group  $Sp(p, q)$  ( $p \geq q \geq 1$ ). At first we give our realization of  $G$  and provide some notations.

We set  $n = p + q$ . Let  $K_{p,q}$  be the  $2n \times 2n$  matrix given by

$$K_{p,q} = \left( \begin{array}{c|c|c} I_p & 0 & \\ \hline 0 & -I_q & \\ \hline & & I_p & 0 \\ & 0 & \hline & & 0 & -I_q \end{array} \right)$$

where  $I_m$  is the  $m \times m$  identity matrix. The group  $G$  is given by

$$G = \{g \in Sp(n, \mathbf{C}) \mid {}^t g K_{p,q} \bar{g} = K_{p,q}\}.$$

As a maximal compact subgroup of  $G$ , we choose  $K = G \cap U(2n)$ . Let  $\mathfrak{g}$  (resp.  $\mathfrak{k}$ ) be the Lie algebra of  $G$  (resp.  $K$ ) and  $\mathfrak{g} = \mathfrak{k} + \mathfrak{p}$  be the Cartan decomposition of  $\mathfrak{g}$ . For a real Lie algebra  $\mathfrak{u}$ , denote by  $\mathfrak{u}_c$  the complexification of  $\mathfrak{u}$ .

Let  $E_{ij}$  be the square matrix with 1 in the  $(i, j)$ -position and 0 elsewhere. For  $1 \leq i \leq n$ , set

$$T_i = \left( \begin{array}{c|c} E_{ii} & 0 \\ \hline 0 & -E_{ii} \end{array} \right)$$

and define

$$\mathfrak{t} = \left\{ \sum_{j=1}^n \mu_j T_j \mid \mu_j \in \sqrt{-1}\mathbf{R} \right\}.$$

Then  $\mathfrak{t}$  is a Cartan subalgebra of  $\mathfrak{g}$  such that  $\mathfrak{t} \subset \mathfrak{k}$ . Also define  $\lambda_i \in \mathfrak{t}_c^*$  ( $1 \leq i \leq n$ ) by

$$\lambda_i \left( \sum_{j=1}^n \mu_j T_j \right) = \mu_i.$$

The root system  $\Delta$  (resp.  $\Delta_{\mathfrak{t}}$ ) of the pair  $(\mathfrak{g}_c, \mathfrak{t}_c)$  (resp.  $(\mathfrak{k}_c, \mathfrak{t}_c)$ ) is given by

$$\begin{aligned} \Delta &= \{ \pm \lambda_i \pm \lambda_j \mid 1 \leq i, j \leq n \} \\ \text{(resp. } \Delta_{\mathfrak{t}} &= \{ \pm \lambda_i \pm \lambda_j \mid 1 \leq i, j \leq p \text{ or } p+1 \leq i, j \leq p+q \} ). \end{aligned}$$

We choose an order of  $(\sqrt{-1}\mathfrak{t})^*$  so that the set of simple roots in  $\Delta$  is  $\{ \lambda_1 - \lambda_2, \lambda_2 - \lambda_3, \dots, \lambda_{n-1} - \lambda_n, 2\lambda_n \}$ . Denote by  $\Delta^+$  (resp.  $\Delta_{\mathfrak{t}}^+$ ) the set of positive roots in  $\Delta$  (resp.  $\Delta_{\mathfrak{t}}$ ). Throughout this paper we fix this order.

For later use, we choose root vectors of  $\mathfrak{g}_c$  as follows;

$$\begin{aligned} X_{\lambda_i + \lambda_j} &= \left( \begin{array}{c|c} 0 & F_{ij} \\ \hline 0 & 0 \end{array} \right) & (1 \leq i, j \leq n) \\ X_{-\lambda_i - \lambda_j} &= \left( \begin{array}{c|c} 0 & 0 \\ \hline F_{ij} & 0 \end{array} \right) & (1 \leq i, j \leq n) \\ X_{\lambda_i - \lambda_j} &= \left( \begin{array}{c|c} E_{ij} & 0 \\ \hline 0 & -E_{ji} \end{array} \right) & (1 \leq i, j \leq n, i \neq j) \end{aligned}$$

where  $F_{ij} = E_{ij} + E_{ji}$  if  $i \neq j$  and  $F_{ij} = E_{ii}$  if  $i = j$ . Then  $\{T_i \mid 1 \leq i \leq n\} \cup \{X_{\alpha} \mid \alpha \in \Delta\}$  is a basis of  $\mathfrak{g}_c$ .

**1.2.** Now we construct an imbedding of  $G$  into  $Sp(2n, \mathbf{R})$ . Our imbedding is obtained by composing an imbedding of  $G$  into  $SU(2p, 2q)$  and an imbedding of  $SU(2p, 2q)$  into  $Sp(2n, \mathbf{R})$ . From now on,  $G'$  denotes the group  $SU(2p, 2q)$ . As a maximal compact subgroup of  $G'$ , we choose  $K' = G' \cap U(2n)$ . Let  $\mathfrak{g}'$  be the Lie algebra of  $G'$ .

The group  $G$  is naturally imbedded into the unitary group of the hermitian form on  $\mathbf{C}^{2n}$  defined by  $K_{p,q}$ . We put

$$Z = \left( \begin{array}{c|cc} I_p & 0 & 0 \\ \hline 0 & 0 & I_q \\ \hline & I_p & 0 \\ \hline 0 & 0 & I_q \end{array} \right)$$

Then  ${}^t Z K_{p,q} Z$  gives the standard hermitian form with signature  $(2p, 2q)$ . So, if we define

$$\psi(g) = {}^t Z g Z \quad (g \in G),$$

we obtain an imbedding  $\psi; G \rightarrow G'$ . Clearly we have  $\psi(K) \subset K'$ .

Moreover we will imbed  $G'$  into  $Sp(2n, \mathbf{R})$ . Naturally we consider  $GL(2n, \mathbf{C})$ , and hence  $G'$ , as to be the subgroups of  $GL(4n, \mathbf{R})$ . Define the orthogonal matrix  $Z'$  by

$$Z' = \left( \begin{array}{c|cc} I_{2p} & 0 & 0 \\ \hline 0 & -I_{2q} & 0 \\ \hline 0 & 0 & I_{2n} \end{array} \right)$$

Then it is easily checked that, if we define

$$\psi'(g) = {}^t Z' g Z' \quad (g \in G'),$$

we obtain an imbedding  $\psi'; G' \rightarrow Sp(2n, \mathbf{R})$ . This is the same imbedding that is constructed in [1], VIII, § 2.

In this way we obtain the imbedding

$$\iota = \psi' \circ \psi; G \rightarrow Sp(2n, \mathbf{R}).$$

These imbeddings  $\psi$ ,  $\psi'$  and  $\iota$  induce the imbeddings of Lie algebras and we use the same letters for them;

$$\begin{aligned} \psi &; \mathfrak{g}_c \rightarrow \mathfrak{g}'_c \\ \psi' &; \mathfrak{g}'_c \rightarrow \mathfrak{sp}(2n, \mathbf{C}) \\ \iota &; \mathfrak{g}_c \rightarrow \mathfrak{sp}(2n, \mathbf{C}). \end{aligned}$$

**1.3.** Here we give the explicit form of the image of  $\iota$ . It will be used in § 2. For this, we choose a basis of  $\mathfrak{sp}(2n, \mathbf{C})$  as follows;

$$S_i = \sqrt{-1} \left( \begin{array}{c|c} 0 & E_{ii} \\ \hline -E_{ii} & 0 \end{array} \right) \quad (1 \leq i \leq 2n),$$

$$\begin{aligned}
 Y_{ij}^+ &= \frac{1}{2} \left( \begin{array}{c|c} F_{ij} & -\sqrt{-1}F_{ij} \\ \hline -\sqrt{-1}F_{ij} & -F_{ij} \end{array} \right) & (1 \leq i, j \leq 2n) \\
 Y_{ij}^- &= \frac{1}{2} \left( \begin{array}{c|c} F_{ij} & \sqrt{-1}F_{ij} \\ \hline \sqrt{-1}F_{ij} & -F_{ij} \end{array} \right) & (1 \leq i, j \leq 2n) \\
 Z_{ij}^+ &= \frac{1}{2} \left( \begin{array}{c|c} E_{ij}-E_{ji} & \sqrt{-1}F_{ij} \\ \hline -\sqrt{-1}F_{ij} & E_{ij}-E_{ji} \end{array} \right) & (1 \leq i < j \leq 2n) \\
 Z_{ij}^- &= \frac{1}{2} \left( \begin{array}{c|c} E_{ji}-E_{ij} & \sqrt{-1}F_{ji} \\ \hline -\sqrt{-1}F_{ji} & E_{ji}-E_{ij} \end{array} \right) & (1 \leq i < j \leq 2n)
 \end{aligned}$$

where  $F_{ij}=E_{ij}+E_{ji}$  if  $i \neq j$  and  $F_{ij}=E_{ii}$  if  $i=j$ . By straightforward computations we obtain the following explicit description for the image of  $\iota$ ; for  $1 \leq i < j \leq p$  and  $p+1 \leq k < l \leq p+q$ ,

$$(1.1) \quad \left\{ \begin{aligned}
 \iota(T_i) &= S_i - S_{p+i} \\
 \iota(T_k) &= -S_{p+k} + S_{p+q+k} \\
 \iota(X_{\pm(\lambda_i + \lambda_j)}) &= Z_{i, p+j}^{\pm} + Z_{j, p+i}^{\pm} \\
 \iota(X_{\pm(\lambda_i + \lambda_k)}) &= -Y_{i, p+q+k}^{\pm} + Y_{p+i, p+k}^{\mp} \\
 \iota(X_{\pm(\lambda_k + \lambda_l)}) &= -Z_{p+k, p+q+l}^{\mp} - Z_{p+l, p+q+k}^{\mp} \\
 \iota(X_{\pm 2\lambda_i}) &= Z_{i, p+i}^{\pm} \\
 \iota(X_{\pm 2\lambda_k}) &= -Z_{p+k, p+q+k}^{\mp} \\
 \iota(X_{\pm(\lambda_i - \lambda_j)}) &= Z_{i, j}^{\pm} - Z_{p+i, p+j}^{\mp} \\
 \iota(X_{\pm(\lambda_i - \lambda_k)}) &= -Y_{i, p+k}^{\pm} + Y_{p+i, p+q+k}^{\mp} \\
 \iota(X_{\pm(\lambda_k - \lambda_l)}) &= -Z_{p+k, p+l}^{\mp} + Z_{p+q+k, p+q+l}^{\pm} .
 \end{aligned} \right.$$

**2. The construction of unitary representations**

In this section, we construct a certain series of irreducible unitary representations of  $G$ . In [1] Borel and Wallach constructed some irreducible representations of  $G'$  by using the oscillator representation. Our representations are obtained from these representations through the imbedding  $\psi$ ;  $G \rightarrow G'$ . We will often use the results and notations in [1], VIII.

**2.1.** First we sketch briefly the results in [1], VIII, § 2. Let  $Mp(2n, \mathbf{R})$  be the Metaplectic group and  $(W, L^2(\mathbf{R}^{2n}))$  be the oscillator representation of

$Mp(2n, \mathbf{R})$ . The imbedding  $\psi'; G' \rightarrow Sp(2n, \mathbf{R})$  lifts to an injective homomorphism  $\tilde{\psi}'; G' \rightarrow Mp(2n, \mathbf{R})$  ([1], VIII, Lemma 2.9). Define the unitary representation  $(V, L^2(\mathbf{R}^{2n}))$  of  $G'$  by

$$V(g) = W(\tilde{\psi}'(g)) \quad (g \in G').$$

Then  $(V, L^2(\mathbf{R}^{2n}))$  decomposes into the direct sum of irreducible representations of  $G'$ . In fact, for  $r \in \mathbf{Z}$ , define the subspace  $H_r$  of  $L^2(\mathbf{R}^{2n})$  by

$$H_r = \{ \phi \in L^2(\mathbf{R}^{2n}) \mid W(\text{Exp } tJ_{2p,2q})(\phi) = \exp(-\sqrt{-1}(p-q+r)t)\phi \}$$

where Exp is the exponential mapping of  $\mathfrak{sp}(2n, \mathbf{R})$  into  $Mp(2n, \mathbf{R})$  and

$$J_{2p,2q} = \left( \begin{array}{cc|cc} & & -I_{2p} & 0 \\ & 0 & 0 & I_{2q} \\ \hline I_{2p} & 0 & & \\ \hline 0 & -I_{2q} & & 0 \end{array} \right) \in \mathfrak{sp}(2n, \mathbf{R}).$$

Then  $H_r$  is stable under  $G'$  and so we put

$$V_r(g) = V(g)|_{H_r} \quad (g \in G').$$

From [1], VIII, Lemma 2.8, for each  $r \in \mathbf{Z}$ ,  $(V_r, H_r)$  is an irreducible unitary representation of  $G'$  and we have

$$L^2(\mathbf{R}^{2n}) = \bigoplus_{r \in \mathbf{Z}} H_r.$$

In the remainder of this section, we fix  $r \in \mathbf{Z}$ . Denote by  $\mathcal{S}(\mathbf{R}^{2n})$  the Schwartz space on  $\mathbf{R}^{2n}$  with the Schwartz topology and set  $H_r^\infty = H_r \cap \mathcal{S}(\mathbf{R}^{2n})$ . Then  $H_r^\infty$  is the space of  $C^\infty$ -vectors for  $V_r$  in  $H_r$ , ([1], VIII, Lemma 1.11). Also, we denote by  $H_r^0$  the space of  $K'$ -finite vectors for  $V_r$  in  $H_r$ . The space  $H_r^0$  is an irreducible admissible  $(\mathfrak{g}', K')$ -module.

In order to choose an orthogonal basis of  $H_r^0$ , we need some notations. Let  $(x_1, \dots, x_{2n})$  be the coordinates of  $\mathbf{R}^{2n}$ . Following [1], VIII, 1.16, for  $1 \leq j \leq 2n$ , define the operator  $D_j$  and  $A_j^\sharp$  by

$$D_j = \frac{1}{2} \left( \frac{\partial^2}{\partial x_j^2} - x_j^2 \right), \quad A_j^\sharp = \frac{1}{2} \left( \frac{\partial}{\partial x_j} \pm x_j \right).$$

Denote by  $\mathbf{Z}_+$  the set of non-negative integers. For  $m = (m_1, \dots, m_{2n}) \in (\mathbf{Z}_+)^{2n}$ , we set

$$\phi_m = (A_1^-)^{m_1} (A_2^-)^{m_2} \cdots (A_{2n}^-)^{m_{2n}} \phi_0$$

where  $\phi_0$  is the  $C^\infty$ -function on  $\mathbf{R}^{2n}$  defined by

$$\phi_0(x) = (2\pi)^{-n} \exp\left(-\frac{1}{2} \sum_{i=1}^{2n} x_i^2\right) \quad (x \in \mathbf{R}^{2n}).$$

(Note that  $\phi_m$  is equal to  $\psi_m$  in [1]. VIII 1.16, up to the multiplication by a constant.) Then, by [1], VIII, Lemma 1.17,  $\{\phi_m | m \in (\mathbf{Z}_+)^{2n}\}$  are mutually orthogonal in  $L^2(\mathbf{R}^{2n})$  and we have

$$(2.1) \quad H_r^0 = \bigoplus_{m \in \Phi_r} \mathbf{C} \phi_m$$

where  $\Phi_r = \{m \in (\mathbf{Z}_+)^{2n} | \sum_{i=1}^{2p} m_i - \sum_{i=2p+1}^{2n} m_i = r\}$ .

2.2. Now we construct unitary representations of  $G$ . Using the imbedding  $\psi; G \rightarrow G'$ , we define

$$U_r(g) = V_r(\psi(g)) \quad (g \in G).$$

Then we obtain the unitary representation  $(U_r, H_r)$  of  $G$ . Clearly, the subspace  $H_r^0$  of  $H_r$  is included in the space of  $K$ -finite vectors for  $U_r$  in  $H_r$  and stable under  $\mathfrak{g}$  and  $K$ . Thus  $H_r^0$  is a  $(\mathfrak{g}, K)$ -module. The infinitesimal representation of  $\mathfrak{g}$  on  $H_r^0$  induced from  $U_r$  is denoted by the same letter  $U_r$ .

We will examine the  $(\mathfrak{g}, K)$ -module  $H_r^0$  in detail. First we consider the infinitesimal representation  $(W, \mathcal{S}(\mathbf{R}^{2n}))$  of  $\mathfrak{sp}(2n, \mathbf{C})$  induced from  $(W, L^2(\mathbf{R}^{2n}))$ . By [2], p. 232, Theorem 5.4, the action of  $\mathfrak{sp}(2n, \mathbf{C})$  on  $\mathcal{S}(\mathbf{R}^{2n})$  is explicitly given as follows;

$$(2.2) \quad \begin{cases} W(S_i) = D_i & (1 \leq i \leq 2n) \\ W(Y_{ij}^\dagger) = \pm 2A_i^\dagger A_j^\dagger & (1 \leq i, j \leq 2n, i \neq j) \\ W(Y_{ii}^\dagger) = \pm A_i^\dagger A_i^\dagger & (1 \leq i \leq 2n) \\ W(Z_{ij}^\dagger) = 2A_i^\dagger A_j^\dagger & (1 \leq i < j \leq 2n). \end{cases}$$

Using the relation formulas among  $D_j$  and  $A_j^\dagger$  in [1], VIII, 1.16, we obtain

$$(2.3) \quad \begin{cases} D_j(\phi_m) = -\frac{1}{2}(2m_j+1)\phi_{m_1, \dots, m_{2n}} \\ A_i^\dagger A_j^\dagger(\phi_m) = \frac{1}{4}m_i m_j \phi_{m_1, \dots, m_{i-1}, \dots, m_{j-1}, \dots, m_{2n}} \\ A_i^\dagger A_i^\dagger(\phi_m) = \frac{1}{4}m_i(m_i-1)\phi_{m_1, \dots, m_{i-2}, \dots, m_{2n}} \\ A_i^- A_j^-(\phi_m) = \phi_{m_1, \dots, m_{i+1}, \dots, m_{j+1}, \dots, m_{2n}} \\ A_i^- A_i^-(\phi_m) = \phi_{m_1, \dots, m_{i+2}, \dots, m_{2n}} \\ A_i^\dagger A_j^-(\phi_m) = -\frac{1}{2}m_i \phi_{m_1, \dots, m_{i-1}, \dots, m_{j+1}, \dots, m_{2n}} \end{cases}$$



where  $m \in (\mathbf{Z}_+)^{2n}$ ,  $1 \leq i < j \leq 2n$  and  $\phi_{k_1, \dots, k_{2n}}$  is considered to be 0 if  $k_i < 0$  for some  $i$ . Therefore, combining (1.1), (2.2) and (2.3), we have the following formulas; for  $1 \leq i, j \leq p$  and  $p+1 \leq k, l \leq p+q$ ,

$$(2.4) \quad \begin{cases} U_r(T_i)(\phi_m) = (m_{p+i} - m_i)\phi_m \\ U_r(T_k)(\phi_m) = (m_{p+k} - m_{p+q+k})\phi_m \end{cases}$$

$$(2.5) \quad \begin{cases} U_r(X_{\lambda_i + \lambda_j})(\phi_m) = -m_i \phi_{m_1, \dots, m_i-1, \dots, m_{p+j+1}, \dots, m_{2n}} \\ \quad \quad \quad -m_j \phi_{m_1, \dots, m_j-1, \dots, m_{p+i+1}, \dots, m_{2n}} \\ U_r(X_{2\lambda_i})(\phi_m) = -m_i \phi_{m_1, \dots, m_i-1, \dots, m_{p+i+1}, \dots, m_{2n}} \\ U_r(X_{\lambda_k + \lambda_l})(\phi_m) = m_{p+q+l} \phi_{m_1, \dots, m_{p+k+1}, \dots, m_{p+q+l-1}, \dots, m_{2n}} \\ \quad \quad \quad + m_{p+q+k} \phi_{m_1, \dots, m_{p+l+1}, \dots, m_{p+q+k-1}, \dots, m_{2n}} \\ U_r(X_{2\lambda_k})(\phi_m) = m_{p+q+k} \phi_{m_1, \dots, m_{p+k+1}, \dots, m_{p+q+k-1}, \dots, m_{2n}} \\ U_r(X_{\lambda_i - \lambda_j})(\phi_m) = -m_i \phi_{m_1, \dots, m_i-1, \dots, m_{j+1}, \dots, m_{2n}} \\ \quad \quad \quad + m_{p+j} \phi_{m_1, \dots, m_{p+i+1}, \dots, m_{p+j-1}, \dots, m_{2n}} \\ U_r(X_{\lambda_k - \lambda_l})(\phi_m) = m_{p+l} \phi_{m_1, \dots, m_{p+k+1}, \dots, m_{p+l-1}, \dots, m_{2n}} \\ \quad \quad \quad - m_{p+q+k} \phi_{m_1, \dots, m_{p+q+k-1}, \dots, m_{p+q+l+1}, \dots, m_{2n}} \end{cases}$$

$$(2.6) \quad \begin{cases} U_r(X_{-\lambda_i - \lambda_j})(\phi_m) = -m_{p+i} \phi_{m_1, \dots, m_{j+1}, \dots, m_{p+i-1}, \dots, m_{2n}} \\ \quad \quad \quad - m_{p+j} \phi_{m_1, \dots, m_{i+1}, \dots, m_{p+j-1}, \dots, m_{2n}} \\ U_r(X_{-2\lambda_i})(\phi_m) = -m_{p+i} \phi_{m_1, \dots, m_{i+1}, \dots, m_{p+i-1}, \dots, m_{2n}} \\ U_r(X_{-\lambda_k - \lambda_l})(\phi_m) = m_{p+l} \phi_{m_1, \dots, m_{p+l-1}, \dots, m_{p+q+k+1}, \dots, m_{2n}} \\ \quad \quad \quad + m_{p+k} \phi_{m_1, \dots, m_{p+k-1}, \dots, m_{p+q+l+1}, \dots, m_{2n}} \\ U_r(X_{-2\lambda_k})(\phi_m) = m_{p+k} \phi_{m_1, \dots, m_{p+k-1}, \dots, m_{p+q+k+1}, \dots, m_{2n}} \\ U_r(X_{-\lambda_i + \lambda_j})(\phi_m) = -m_j \phi_{m_1, \dots, m_{i+1}, \dots, m_{j-1}, \dots, m_{2n}} \\ \quad \quad \quad + m_{p+i} \phi_{m_1, \dots, m_{p+i-1}, \dots, m_{p+j+1}, \dots, m_{2n}} \\ U_r(X_{-\lambda_k + \lambda_l})(\phi_m) = m_{p+k} \phi_{m_1, \dots, m_{p+k-1}, \dots, m_{p+l+1}, \dots, m_{2n}} \\ \quad \quad \quad - m_{p+q+l} \phi_{m_1, \dots, m_{p+q+k+1}, \dots, m_{p+q+l-1}, \dots, m_{2n}} \end{cases}$$

$$(2.7) \quad \begin{cases} U_r(X_{\lambda_i + \lambda_k})(\phi_m) = -\frac{1}{2} m_i m_{p+q+k} \phi_{m_1, \dots, m_i-1, \dots, m_{p+q+k-1}, \dots, m_{2n}} \\ \quad \quad \quad + 2\phi_{m_1, \dots, m_{p+i+1}, \dots, m_{p+k+1}, \dots, m_{2n}} \\ U_r(X_{\lambda_i - \lambda_k})(\phi_m) = -\frac{1}{2} m_i m_{p+k} \phi_{m_1, \dots, m_i-1, \dots, m_{p+k-1}, \dots, m_{2n}} \\ \quad \quad \quad - 2\phi_{m_1, \dots, m_{p+i+1}, \dots, m_{p+q+k+1}, \dots, m_{2n}} \end{cases}$$

$$(2.8) \quad \begin{cases} U_r(X_{-\lambda_i - \lambda_k})(\phi_m) = -\frac{1}{2} m_{p+i} m_{p+k} \phi_{m_1, \dots, m_{p+i-1}, \dots, m_{p+k-1}, \dots, m_{2n}} \\ \quad \quad \quad + 2\phi_{m_1, \dots, m_{i+1}, \dots, m_{p+q+k+1}, \dots, m_{2n}} \end{cases}$$

$$\left\{ \begin{aligned} U_r(X_{-\lambda_i+\lambda_k})(\phi_m) &= \frac{1}{2} m_{p+i} m_{p+q+k} \phi_{m_1, \dots, m_{p+i-1}, \dots, m_{p+q+k-1}, \dots, m_{2n}} \\ &\quad + 2\phi_{m_1, \dots, m_{i+1}, \dots, m_{p+k+1}, \dots, m_{2n}} \end{aligned} \right.$$

Of course, in these formulas,  $\phi_{k_1, \dots, k_{2n}}$  should be considered as to be 0 if  $k_i < 0$  for some  $i$ .

Now we can determine the set of weights of the  $\mathfrak{g}_c$ -module  $H_r^0$ . Let  $\phi_m$  be in  $H_r^0$ . By (2.4) we have

$$U_r(\sum_{i=1}^n \mu_i T_i)(\phi_m) = \{ \sum_{i=1}^p (m_{p+i} - m_i) \mu_i + \sum_{k=p+1}^{p+q} (m_{p+k} - m_{p+q+k}) \mu_k \} \phi_m$$

From this, the following lemma immediately follows.

**Lemma 2.1.** *Let  $m=(m_1, \dots, m_{2n})$  be in  $\Phi_r$ . In the  $\mathfrak{g}_c$ -module  $H_r^0$ ,  $\phi_m$  is a weight vector corresponding to the weight*

$$\Lambda_m = \sum_{i=1}^p (m_{p+i} - m_i) \lambda_i + \sum_{k=p+1}^{p+q} (m_{p+k} - m_{p+q+k}) \lambda_k.$$

We remark that the multiplicity of  $\Lambda_m$  in  $H_r^0$  is not finite.

**2.3.** Here we determine the  $K$ -spectrum of  $H_r^0$ . Let  $\hat{K}$  be the set of all equivalence classes of irreducible representations of  $K$ . Define the subset  $D_K$  of  $\mathfrak{t}_c^*$  by

$$D_K = \left\{ \lambda = \sum_{i=1}^n a_i \lambda_i \left| \begin{array}{l} a_i \in \mathbf{Z} \\ a_1 \geq a_2 \geq \dots \geq a_p \geq 0 \\ a_{p+1} \geq a_{p+2} \geq \dots \geq a_n \geq 0 \end{array} \right. \right\}.$$

Then there is the bijective correspondence between  $\hat{K}$  and  $D_K$ . That is,  $\lambda \in D_K$  corresponds to the irreducible  $K$ -module with highest weight  $\lambda$ . We denote by  $E_\lambda$  this  $K$ -module.

Let  $s \in \mathbf{Z}_+$  and  $s \geq -r$ . We define the finite dimensional subspace  $H_{r,s}^0$  of  $H_r^0$  by

$$H_{r,s}^0 = \bigoplus_{m \in \Phi_{r,s}} \mathbf{C} \phi_m,$$

where the subset  $\Phi_{r,s}$  of  $\Phi_r$  is given by

$$\Phi_{r,s} = \{ m \in (\mathbf{Z}_+)^{2n} \mid \sum_{i=1}^{2p} m_i = r + s, \sum_{i=2p+1}^{2n} m_i = s \}.$$

From (2.1), we have

$$H_r^0 = \bigoplus_{s \in \mathbf{Z}_+, s \geq -r} H_{r,s}^0.$$

**Proposition 2.2.** *Let  $s \in \mathbb{Z}_+$  and  $s \geq -r$ . Then  $H_{r,s}^0$  is the irreducible  $K$ -submodule of  $H_r^0$  with highest weight  $(r+s)\lambda_1 + s\lambda_{p+1} \in D_K$ . Hence we have*

$$H_r^0 = \bigoplus_{s \in \mathbb{Z}_+, s \geq -r} E_{(r+s)\lambda_1 + s\lambda_{p+1}}$$

as  $K$ -modules.

Proof. Put  $E_s = E_{(r+s)\lambda_1 + s\lambda_{p+1}}$ . Let  $X$  be in  $\mathfrak{k}_e$ . By (2.4), (2.5) and (2.6),  $U_r(X)(\phi_m)$  is a linear combination of  $\phi_{m'} = \phi_{m'_1, \dots, m'_{2n}}$  such that

$$\sum_{i=1}^{2p} m'_i = \sum_{i=1}^{2p} m_i, \quad \sum_{i=2p+1}^{2n} m'_i = \sum_{i=2p+1}^{2n} m_i.$$

Therefore  $H_{r,s}^0$  is stable under  $\mathfrak{k}_e$ .

Now we put  $\phi = \phi_{0, \dots, 0, r+s, 0, \dots, 0, s, 0, \dots, 0}$ , where  $r+s$  (resp.  $s$ ) appears in the  $(p+1)$ -th (resp.  $(2p+1)$ -th) position. Then  $\phi \in H_{r,s}^0$  and, by Lemma 2.1,  $\phi$  is a weight vector corresponding to the weight  $(r+s)\lambda_1 + s\lambda_{p+1}$ . It is easy to see that this weight is the highest among all the weights for  $H_{r,s}^0$ . Hence  $E_s$  certainly occurs in  $H_{r,s}^0$ .

We compare the dimension of  $H_{r,s}^0$  with that of  $E_s$ . Since  $\{\phi_m \mid m \in \Phi_{r,s}\}$  is a basis of  $H_{r,s}^0$ , we have

$$\begin{aligned} \dim H_{r,s}^0 &= \#\Phi_{r,s} \\ &= \binom{2p+r+s-1}{r+s} \cdot \binom{2q+s-1}{s} \\ &= \frac{(2p+r+s-1)!(2q+s-1)!}{(2p-1)!(r+s)!(2q-1)!s!}. \end{aligned}$$

On the other hand, Weyl's dimension formula gives the dimension of  $E_s$ . Denote by  $(\ , \ )_{\mathfrak{k}}$  the inner product in  $(\sqrt{-1}\mathfrak{k})^*$  induced from the Killing form of  $\mathfrak{k}_e$ . Recall that

$$\begin{aligned} (\lambda_i, \lambda_j)_{\mathfrak{k}} &= 0 && \text{if } i \neq j, \\ (\lambda_i, \lambda_i)_{\mathfrak{k}} &= \begin{cases} (4p+4)^{-1} & \text{if } 1 \leq i \leq p, \\ (4q+4)^{-1} & \text{if } p+1 \leq i \leq p+q. \end{cases} \end{aligned}$$

Also put  $\delta_{\mathfrak{k}} = \frac{1}{2} \sum_{\alpha \in \Delta_{\mathfrak{k}}^+} \alpha$ . Then we have

$$\delta_{\mathfrak{k}} = \sum_{i=1}^p (p-i+1)\lambda_i + \sum_{k=p+1}^{p+q} (p+q-k+1)\lambda_k.$$

From these formulas, easy calculations yield

$$\begin{aligned} \dim E_s &= \frac{\prod_{\alpha \in \Delta_{\mathfrak{k}}^+} ((r+s)\lambda_1 + s\lambda_{p+1} + \delta_{\mathfrak{k}}, \alpha)_{\mathfrak{k}}}{\prod_{\alpha \in \Delta_{\mathfrak{k}}^+} (\delta_{\mathfrak{k}}, \alpha)_{\mathfrak{k}}} \\ &= \frac{(2p+r+s-1)!(2q+s-1)!}{(2p-1)!(r+s)!(2q-1)!s!} \\ &= \dim H_{r,s}^0. \end{aligned}$$

Hence  $H_{r,s}^0$  is equivalent to  $E_s$ .

**2.4.** In this stage, we must determine the space of  $K$ -finite vectors in  $H_r$  for  $U_r$ .

**Lemma 2.3.** *The space of  $K$ -finite vectors in  $H_r$  for  $U_r$  coincides with  $H_r^0$ .*

Proof. For  $\tau \in \hat{K}$ , let  $H_r(\tau)$  be the isotypic  $K$ -submodule of  $H_r$  of type  $\tau$ . Clearly  $H_r^0$  is stable under  $K$  and  $H_r^0 \subset \bigoplus_{\tau \in \hat{K}} H_r(\tau)$ . Hence we have  $H_r^0 = \bigoplus_{\tau \in \hat{K}} H_r^0 \cap H_r(\tau)$ . Since  $H_r^0$  is dense in  $H_r$ , by [7], Chapter 4, Proposition 4.4.3.4, the closure of  $H_r^0 \cap H_r(\tau)$  is  $H_r(\tau)$ . By Proposition 2.2,  $H_r^0 \cap H_r(\tau)$  is finite dimensional. Therefore we have  $H_r^0 \cap H_r(\tau) = H_r(\tau)$  and hence  $H_r^0 = \bigoplus_{\tau \in \hat{K}} H_r(\tau)$ . The lemma is proved.

Together with Proposition 2.2, this lemma shows that  $(U_r, H_r)$  is admissible. Moreover we have the following proposition.

**Proposition 2.4.** *For  $r \in \mathbf{Z}$ , the unitary representation  $(U_r, H_r)$  of  $G$  is irreducible.*

Proof. From [7], Chapter 4, Theorem 4.5.5.4, it is sufficient to prove that the  $\mathfrak{g}$ -module  $H_r^0$  is algebraically irreducible. Let  $H$  be a non-zero  $\mathfrak{g}$ -stable subspace of  $H_r^0$ . Since  $H$  is stable under  $\mathfrak{k}$ , by Proposition 2.2, we have

$$H = \bigoplus_{s \in S(H)} H_{r,s}^0,$$

where  $S(H)$  is a non-empty subset of  $\mathbf{Z}_+$ . Suppose  $s_0 \in S(H)$ , that is,  $H_{r,s_0}^0 \subset H$ . We take a particular element

$$\phi = \phi_{0, \dots, 0, r+s_0, 0, \dots, 0, s_0, 0, \dots, 0}$$

in  $H_{r,s_0}^0$ , where  $r+s_0$  (resp.  $s_0$ ) appears in the  $(p+1)$ -th (resp.  $(2p+1)$ -th) position. Then, by (2.7), we have

$$U_r(X_{\lambda_1 + \lambda_{p+1}})(\phi) = 2\phi_{0, \dots, 0, r+s_0+1, 0, \dots, 0, s_0+1, 0, \dots, 0}.$$

Here the left hand side belongs to  $H$  and the right hand side belongs to  $H_{r,s_0+1}^0$ .

This implies  $H \cap H_{r,s_0+1}^0 \neq \{0\}$ . Therefore we have  $H_{r,s_0+1}^0 \subset H$ , that is,  $s_0+1 \in S(H)$ .

Similarly, if  $s_0 > \max\{0, -r\}$ , we have

$$U_r(X_{-\lambda_1-\lambda_{p+1}})(\phi) = -\frac{1}{2}(r+s_0)s_0\phi_{0,\dots,0,r+s_0-1,0,\dots,0,s_0-1,0,\dots,0} + 2\phi_{1,0,\dots,0,r+s_0,0,\dots,0,s_0,0,\dots,0,1,0,\dots,0}$$

where 1 appears in the first and  $(2p+q+1)$ -th position. In this formula, the first term of the right hand side belongs to  $H_{r,s_0-1}^0$  and the second term belongs to  $H_{r,s_0+1}^0$ . Since  $H_{r,s_0+1}^0 \subset H$ , we have  $H \cap H_{r,s_0-1}^0 \neq \{0\}$  and hence  $s_0-1 \in S(H)$ .

By the induction, we have  $S(H) = \{s \in \mathbb{Z}_+ \mid s \geq -r\}$ , that is,  $H = H_r^0$ . This proves the proposition.

After all we obtain a series of irreducible unitary representations of  $G$ ;  $\{(U_r, H_r) \mid r \in \mathbb{Z}\}$ .

### 3. The $(\mathfrak{g}, K)$ -cohomology

In this section, we study the  $(\mathfrak{g}, K)$ -cohomology space of the  $(\mathfrak{g}, K)$ -module  $H_r^0$  ( $r \in \mathbb{Z}$ ).

**3.1.** First of all we recall a known result which is our starting point. Let  $(U, H_U)$  be in  $\hat{G}$  and  $(\rho, F)$  be a finite dimensional irreducible representation of  $G$ . Denote by  $\mathfrak{U}(\mathfrak{g})$  the universal enveloping algebra of  $\mathfrak{g}_\mathbb{C}$ . The representation of  $\mathfrak{U}(\mathfrak{g})$  induced by  $U$  (resp.  $\rho$ ) is denoted by the same letter  $U$  (resp.  $\rho$ ). Let  $C$  be the Casimir element of  $\mathfrak{g}_\mathbb{C}$ . Then both the operators  $U(C)$  and  $\rho(C)$  are the scalar operators. Put  $U(C) = c_U \cdot \text{Id}$  and  $\rho(C) = c_\rho \cdot \text{Id}$ , where  $c_U, c_\rho \in \mathbb{C}$  and  $\text{Id}$  denotes the identity operator. If we note that  $K$  is connected, we have the following lemma.

**Lemma 3.1.** ([1], II, Proposition 3.1)

- (1). If  $c_U \neq c_\rho$ , then  $H^j(\mathfrak{g}, K; H_U^0 \otimes F) = \{0\}$  for all  $j \in \mathbb{Z}_+$ .
- (2). If  $c_U = c_\rho$ , then  $H^j(\mathfrak{g}, K; H_U^0 \otimes F) = \text{Hom}_K(\wedge^j \mathfrak{p}, H_U^0 \otimes F)$  for all  $j \in \mathbb{Z}_+$ .

**3.2.** For  $(U_r, H_r) \in \hat{G}$ , we will calculate the operator  $U_r(C)$ .

**Proposition 3.2.** For  $r \in \mathbb{Z}$ , we have

$$U_r(C) = (4n+4)^{-1}(r+2p)(r-2q) \cdot \text{Id}.$$

*Proof.* We use a concrete realization of  $C$  and calculate explicitly the action of  $U_r(C)$  on a particular element in  $H_r^0$ .

Recall that the Killing form of  $\mathfrak{g}_\mathbb{C}$  is given by

$$(X, Y) = 2(n+1) \text{Tr } XY \quad (X, Y \in \mathfrak{g}_\mathbb{C}).$$

Using the basis of  $\mathfrak{g}_e$  in 1.1., we have

$$\begin{aligned}
 4(n+1)C &= \sum_{i=1}^n T_i T_i + \sum_{1 \leq i < j \leq n} (X_{\lambda_i + \lambda_j} X_{-\lambda_i - \lambda_j} + X_{-\lambda_i - \lambda_j} X_{\lambda_i + \lambda_j}) \\
 &+ 2 \sum_{i=1}^n (X_{2\lambda_i} X_{-2\lambda_i} + X_{-2\lambda_i} X_{2\lambda_i}) \\
 &+ \sum_{1 \leq i < j \leq n} (X_{\lambda_i - \lambda_j} X_{\lambda_j - \lambda_i} + X_{\lambda_j - \lambda_i} X_{\lambda_i - \lambda_j}).
 \end{aligned}$$

First we consider the case that  $r \geq 0$ . Take a particular element  $\phi = \phi_{r, 0, \dots, 0} \in H_r^0$ . Using (2.4), ..., (2.8), we calculate straightforwardly  $4(n+1)U_r(C)(\phi)$ . Some terms turn out to vanish and the other terms are given as follows;

$$\begin{aligned}
 \sum_{i=1}^n U_r(T_i T_i)(\phi) &= r^2 \phi \\
 U_r(X_{\lambda_i \pm \lambda_k} X_{-\lambda_i \mp \lambda_k})(\phi) &= \begin{cases} -(r+1)\phi \pm 4\phi' & \text{if } i=1 \\ -\phi \pm 4\phi'' & \text{if } i \neq 1 \end{cases} \\
 U_r(X_{-\lambda_i \mp \lambda_j} X_{\lambda_i \pm \lambda_j})(\phi) &= \begin{cases} r\phi & \text{if } i=1 \\ 0 & \text{if } i \neq 1 \end{cases} \\
 U_r(X_{-\lambda_i \mp \lambda_k} X_{\lambda_i \pm \lambda_k})(\phi) &= \begin{cases} -\phi \pm 4\phi' & \text{if } i=1 \\ -\phi \pm 4\phi'' & \text{if } i \neq 1 \end{cases} \\
 2U_r(X_{-2\lambda_i} X_{2\lambda_i})(\phi) &= \begin{cases} 2r\phi & \text{if } i=1 \\ 0 & \text{if } i \neq 1 \end{cases}
 \end{aligned}$$

where  $1 \leq i < j \leq p$ ,  $p+1 \leq k \leq p+q$  and  $\phi'$ ,  $\phi''$  are certain elements in  $H_r^0$  determined by  $\phi$ . From these formulas, we can easily show that

$$4(n+1)U_r(C)(\phi) = (r+2p)(r-2q)\phi.$$

In the case that  $r < 0$ , if we take  $\phi = \phi_{0, \dots, 0, -r} \in H_r^0$ , similar calculations yield the above formula. Thus the proposition is proved.

**3.3.** Now we will show the non-vanishing of the  $(\mathfrak{g}, K)$ -cohomology of  $H_r^0$ . For this, we need the following lemma.

**Lemma 3.3.** For  $2q\lambda_1 \in D_K$ , we have

$$\dim \text{Hom}_K(\wedge^{2q}\mathfrak{p}, E_{2q\lambda_1}) = 1.$$

Proof. Any weight of  $\wedge^{2q}\mathfrak{p}_e$  is the sum of  $2q$  distinct non-compact roots of  $\mathfrak{g}_e$ . Since we have

$$2q\lambda_1 = \sum_{k=p+1}^{p+q} \{(\lambda_1 + \lambda_k) + (\lambda_1 - \lambda_k)\},$$

$2q\lambda_1$  is a weight of  $\wedge^{2q}\mathfrak{p}_e$  with multiplicity 1. It is easy to see that  $2q\lambda_1$  is the

highest among all the weights of  $\wedge^{2q}\mathfrak{p}_e$ . The lemma is proved.

For  $l \in \mathbf{Z}_+$ ,  $l\lambda_1$  is a dominant integral form for  $(\mathfrak{g}_e, \mathfrak{t}_e)$ . Denote by  $(\rho_l, F_l)$  the irreducible finite dimensional representation of  $G$  with highest weight  $l\lambda_1$ ; that is,  $(\rho_l, F_l)$  is the  $l$ -th symmetric tensor product of the standard representation of  $G$  on  $\mathbf{C}^{2n}$ . Let  $(\rho_l^*, F_l^*)$  be the contragredient representation of  $(\rho_l, F_l)$ .

**Theorem 3.4.** *If  $r \geq 2q$ , then we have*

$$H^{2q}(\mathfrak{g}, K; H_r^0 \otimes F_{r-2q}^*) \neq \{0\}.$$

Proof. As it is well-known, the operator  $\rho_{r-2q}^*(C)$  is given by

$$\rho_{r-2q}^*(C) = \{((r-2q)\lambda_1 + \delta, (r-2q)\lambda_1 + \delta) - (\delta, \delta)\} \cdot \text{Id},$$

where  $(\ , \ )$  is the inner product in  $(\sqrt{-1}\mathfrak{t})^*$  induced from the Killing form of  $\mathfrak{g}_e$  and  $\delta = \frac{1}{2} \sum_{\alpha \in \Delta^+} \alpha$ . Note that

$$\begin{aligned} \delta &= \sum_{i=1}^n (n-i+1)\lambda_i, \\ (\lambda_i, \lambda_j) &= (4n+4)^{-1} \delta_{ij} \quad (1 \leq i, j \leq n). \end{aligned}$$

By easy computations, we have

$$\rho_{r-2q}^*(C) = (4n+4)^{-1}(r+2p)(r-2q) \cdot \text{Id}.$$

From this and Proposition 3.2,  $U_r(C)$  and  $\rho_{r-2q}^*(C)$  act as the multiplication by the same scalar. Hence Lemma 3.1 implies that

$$\begin{aligned} \dim H^{2q}(\mathfrak{g}, K; H_r^0 \otimes F_{r-2q}^*) &= \dim \text{Hom}_K(\wedge^{2q}\mathfrak{p}, H_r^0 \otimes F_{r-2q}^*) \\ &= \dim \text{Hom}_K(\wedge^{2q}\mathfrak{p} \otimes F_{r-2q}, H_r^0). \end{aligned}$$

On the other hand, by Proposition 2.2, we have

$$(3.1) \quad \dim \text{Hom}_K(E_{r\lambda_1}, H_r^0) = 1.$$

Also, since  $r\lambda_1 = 2q\lambda_1 + (r-2q)\lambda_1$ , Lemma 3.3 implies that

$$(3.2) \quad \dim \text{Hom}_K(E_{r\lambda_1}, \wedge^{2q}\mathfrak{p} \otimes F_{r-2q}) \neq 0.$$

Therefore, combining (3.1) and (3.2), we have

$$\dim \text{Hom}_K(\wedge^{2q}\mathfrak{p} \otimes F_{r-2q}, H_r^0) \neq 0.$$

This proves the theorem.

REMARK 3.5. By Theorem 1.4 in [5], there is at most one irreducible unitary representation  $(U, H_U)$  such that  $U(C)$  acts by the same scalar as

$\rho_{r-2q}^*(C)$  and  $E_{r, \lambda_1}$  occurs in  $H_U^0$ . Our representation  $(U_r, H_r)$  is this very representation. Therefore we can determine the position of  $U_r$  in the Langlands' classification. In the case of  $q=1$ ,  $(U_2, H_2)$  is equivalent to the Langlands' representation  $J_{1,2}$  in [3], Theorem 3.2.

**4. The imbedding of  $U_r$  into  $L^2(\Gamma \backslash G)$**

In this section, we fix  $r \in \mathbf{Z}$ . We will construct a certain uniform discrete subgroup  $\Gamma$  of  $G$  such that  $m(U_r, \Gamma) \neq 0$ . Together with Theorem 3.4 and (0.1), this will prove the non-vanishing of the cohomology of  $\Gamma$ . The results in this section depend heavily on the results in [1], VIII, § 5.

**4.1.** Our discrete subgroup will be constructed arithmetically. First we realize  $G$  and  $G'$  as subgroups of linear algebraic groups.

Let  $k$  be a totally real finite extension of  $\mathbf{Q}$  and  $d$  be the degree of  $k$  over  $\mathbf{Q}$ . Assume that  $d \geq 2$ . Let  $\Sigma = \{\sigma_1, \dots, \sigma_d\}$  be the set of isomorphisms of  $k$  into  $\mathbf{R}$ . We regard  $k$  as a subfield of  $\mathbf{R}$  so that  $\sigma_1$  is the identity mapping. Put  $k' = k(\sqrt{-1})$ . We extend  $\sigma \in \Sigma$  to the imbedding of  $k'$  into  $\mathbf{C}$  which leaves  $\sqrt{-1}$  fixed. If  $H$  is a linear algebraic group in  $GL(l, \mathbf{C})$  defined over  $k$  or  $\mathbf{Q}$  and  $\mathbf{B}$  is a subfield of  $\mathbf{C}$ , we put  $H(\mathbf{B}) = H \cap GL(l, \mathbf{B})$ .

Denote by  $E_{k'}$  the vector space  $(k')^{2n}$ . We can choose  $a \in k$  so that  $a$  is positive and the conjugates  ${}^\sigma a$  by  $\sigma \in \Sigma$  ( $\sigma \neq \sigma_1$ ) are all negative. Fix such  $a$ . Let  $h$  (resp.  $b$ ) be a non-degenerate hermitian form (resp. a non-degenerate skew-symmetric bilinear form) on  $E_{k'}$  defined by the matrix

$$\left( \begin{array}{c|c|c} I_p & 0 & \\ \hline 0 & -aI_q & \\ \hline & & 0 \end{array} \right) \quad (\text{resp.} \quad \left( \begin{array}{c|c|c} & I_p & 0 \\ \hline & 0 & aI_q \\ \hline -I_p & 0 & \\ \hline 0 & -aI_q & \\ & & 0 \end{array} \right)).$$

Then  $h$  is an indefinite hermitian form with signature  $(2p, 2q)$  but the conjugates  ${}^\sigma h$  by  $\sigma$  ( $\sigma \neq \sigma_1$ ) are positive definite.

Using  $h$  and  $b$ , we can construct the linear algebraic group  $\mathbf{G}$  defined over  $k$  such that

$$\mathbf{G}(k) = \left\{ g \in SL(2n, k') \mid \begin{array}{l} h(gz, gw) = h(z, w) \\ b(gz, gw) = b(z, w) \end{array} \quad (z, w \in E_{k'}) \right\}.$$

Then  $\mathbf{G}(\mathbf{R})$  is isomorphic to  $G$  over  $\mathbf{R}$ . Similarly, using only  $h$ , we obtain the linear algebraic group  $\mathbf{G}'$  defined over  $k$  such that



$$G'(k) = \{g \in SL(2n, k) \mid h(gz, gw) = h(z, w) \quad (z, w \in E_{k'})\}.$$

Also,  $G'(\mathbf{R})$  is isomorphic to  $G'$  over  $\mathbf{R}$ .

Naturally, we have the rational imbedding of  $G$  into  $G'$  defined over  $k$ . We denote by  $\psi; G \rightarrow G'$  this imbedding. It should be noted that, up to conjugation over  $\mathbf{R}$ ,  $\psi|_{G(\mathbf{R})}; G(\mathbf{R}) \rightarrow G'(\mathbf{R})$  coincides with the imbedding  $\psi; G \rightarrow G'$  in 1.2.

**4.2.** Now we denote by  $\text{Res}_{k/\mathbf{Q}}$  the functor of the restriction of scalars from  $k$  to  $\mathbf{Q}$ . Let  $\mathcal{G} = \text{Res}_{k/\mathbf{Q}} G$  and  $\mathcal{G}' = \text{Res}_{k/\mathbf{Q}} G'$ . Then we have the canonical imbedding  $\text{Res}_{k/\mathbf{Q}} \psi; \mathcal{G} \rightarrow \mathcal{G}'$  defined over  $\mathbf{Q}$ . Put  $\Psi = \text{Res}_{k/\mathbf{Q}} \psi$ .

Over  $\mathbf{R}$ , we have the following isomorphisms ([2], 7.16);

$$\begin{aligned} \mathcal{G} &\cong {}^{\sigma_1}G \times {}^{\sigma_2}G \times \cdots \times {}^{\sigma_d}G \\ \mathcal{G}' &\cong {}^{\sigma_1}G' \times {}^{\sigma_2}G' \times \cdots \times {}^{\sigma_d}G', \end{aligned}$$

where, for  $\sigma \in \Sigma$ ,  ${}^{\sigma}G$  (resp.  ${}^{\sigma}G'$ ) denotes the conjugate of  $G$  (resp.  $G'$ ) by  $\sigma$ . So we have

$$(4.1) \quad \mathcal{G}(\mathbf{R}) \cong G \times Sp(n) \times \cdots \times Sp(n)$$

$$(4.2) \quad \mathcal{G}'(\mathbf{R}) \cong G' \times SU(2n) \times \cdots \times SU(2n).$$

Under these isomorphisms, the imbedding  $\Psi$  is the product of the conjugations  ${}^{\sigma_i}\psi; {}^{\sigma_i}G \rightarrow {}^{\sigma_i}G'$  ( $1 \leq i \leq d$ ) of  $\psi$ .

As in [1], VIII, 5.3,  $\mathcal{G}'$  is naturally imbedded into  $Sp_N$  over  $\mathbf{Q}$  where  $N = 2nd$ . In fact, consider  $E_{k'}$  as to be a  $4n$ -dimensional vector space over  $k$  and write  $E_k$  instead of  $E_{k'}$ . We define the skew-symmetric  $k$ -bilinear form  $\beta$  on  $E_k$  by

$$h(z, w) = \mu(z, w) + \sqrt{-1}\beta(z, w) \quad (z, w \in E_k).$$

Then  $G'$  is imbedded into the symplectic group  $Sp_{2n}$  defined by  $\beta$  over  $k$ . Further, if we consider  $E_{\mathbf{Q}} = \text{Res}_{k/\mathbf{Q}} E_k$  and  $\beta_{\mathbf{Q}} = \text{Res}_{k/\mathbf{Q}} \beta$ ,  $\mathcal{G}'$  is naturally imbedded into the group  $Sp_N$  defined by  $\beta_{\mathbf{Q}}$  over  $\mathbf{Q}$ . Denote by  $\Psi'; \mathcal{G}' \rightarrow Sp_N$  this imbedding.

Thus we obtain the imbedding  $\Psi' \circ \Psi; \mathcal{G} \rightarrow Sp_N$  defined over  $\mathbf{Q}$ . We choose a basis of  $E_{\mathbf{Q}}$  so that  $\beta_{\mathbf{Q}}$  is of standard form. With respect to this basis, we consider  $Sp_N$  as to be the subgroup of  $GL(2N, \mathbf{C})$ . Define

$$\begin{aligned} \mathcal{G}(\mathbf{Z}) &= \{g \in \mathcal{G}(\mathbf{Q}) \mid (\Psi' \circ \Psi)(g) \in Sp(N, \mathbf{Z})\} \\ \mathcal{G}'(\mathbf{Z}) &= \{g \in \mathcal{G}'(\mathbf{Q}) \mid \Psi'(g) \in Sp(N, \mathbf{Z})\}. \end{aligned}$$

Then  $\mathcal{G}(\mathbf{Z})$  (resp.  $\mathcal{G}'(\mathbf{Z})$ ) is an arithmetic subgroup of  $\mathcal{G}(\mathbf{R})$  (resp.  $\mathcal{G}'(\mathbf{R})$ ) ([2], 7.11, 7.12). By a standard argument about arithmetic subgroups,  $\mathcal{G}(\mathbf{Z})$  (resp.  $\mathcal{G}'(\mathbf{Z})$ ) turns out to be a uniform discrete subgroup of  $\mathcal{G}(\mathbf{R})$  (resp.  $\mathcal{G}'(\mathbf{R})$ )

([1], VIII, 5.4). In the direct product (4.1) (resp. (4.2)), denote by  $p_1; \mathcal{G}(\mathbf{R}) \rightarrow G$  (resp.  $p'_1; \mathcal{G}'(\mathbf{R}) \rightarrow G'$ ) the projection to the first component. Define

$$\Gamma_0 = p_1(\mathcal{G}(\mathbf{Z})), \quad \Gamma'_0 = p'_1(\mathcal{G}'(\mathbf{Z})).$$

Then  $\Gamma_0$  (resp.  $\Gamma'_0$ ) is a uniform discrete subgroup of  $G$  (resp.  $G'$ ) ([1], VIII, 5.5). Clearly we have

$$\psi(\Gamma_0) \subset \Gamma'_0.$$

As for the group  $G'$  and its representation  $(V_r, H_r)$ , Borel and Wallach obtained the following theorem.

**Theorem 4.1** ([1], VIII, Corollary 5.8). *There is a subgroup  $\Gamma'$  of finite index in  $\Gamma'_0$  such that  $m(V_r, \Gamma') \neq 0$ , where  $m(V_r, \Gamma')$  is the multiplicity of  $V_r$  in  $L^2(\Gamma' \backslash G')$ .*

As the proof of this theorem in [1] shows,  $\Gamma'$  is indeed a congruence subgroup of  $\Gamma'_0$ ; that is,  $\Gamma'$  is given by

$$\Gamma' = p'_1(\Omega')$$

where  $\Omega'$  is a congruence subgroup of  $\mathcal{G}'(\mathbf{Z})$ . Using this subgroup  $\Gamma'$ , we can construct our desired subgroup of  $G$ .

**Theorem 4.2.** *There is a subgroup  $\Gamma$  of finite index in  $\Gamma_0$  such that  $m(U_r, \Gamma) \neq 0$ .*

*Proof.* Let  $\Gamma'$  and  $\Omega'$  be as above. There is a congruence subgroup  $\Omega$  of  $\mathcal{G}(\mathbf{Z})$  such that  $\Psi(\Omega) \subset \Omega'$  ([2], 7.12). Put  $\Gamma = p_1(\Omega)$ . Then  $\Gamma$  is a subgroup of finite index in  $\Gamma_0$  and we have

$$(4.3) \quad \psi(\Gamma) \subset \Gamma'.$$

In the following, we will prove that  $m(U_r, \Gamma) \neq 0$ . As in 2.1, let  $H_r^\infty$  be the space of  $C^\infty$ -vectors in  $H_r$  for the representation  $(V_r, H_r)$  of  $G'$ . Since  $m(V_r, \Gamma') \neq 0$ , by [1], VIII, Theorem 4.3, there is a non-trivial continuous linear functional  $\lambda$  of  $H_r^\infty$  such that

$$\lambda \circ V_r(\gamma) = \lambda$$

for all  $\gamma \in \Gamma'$ . Using  $\lambda$ , we want to construct a non-trivial intertwining operator of  $H_r$  into  $L^2(\Gamma \backslash G)$ . For  $\phi \in H_r^\infty$ , define a function  $A'(\phi); G' \rightarrow \mathbf{C}$  by

$$A'(\phi)(g) = \lambda(V_r(g)\phi) \quad (g \in G').$$

Then  $A'(\phi)$  is a  $C^\infty$ -function on  $G'$  and left  $\Gamma'$ -invariant. Since  $G$  is imbedded into  $G'$  by  $\psi$  as a Lie subgroup,  $A'(\phi) \circ \psi; G \rightarrow \mathbf{C}$  is a  $C^\infty$ -function on  $G$ . Also,

by (4.3),  $A'(\phi) \circ \psi$  is left  $\Gamma$ -invariant. So we can define a linear mapping  $A; H_r^\infty \rightarrow C^\infty(\Gamma \backslash G)$  by

$$\begin{aligned} A(\phi)(\Gamma g) &= A'(\phi)(\psi(g)) \\ &= \lambda(U_r(g)\phi) \quad (\phi \in H_r^\infty, g \in G). \end{aligned}$$

Clearly we have

$$A(U_r(g)\phi) = U_\Gamma(g)A(\phi) \quad (\phi \in H_r^\infty, g \in G)$$

where  $U_\Gamma$  is the right regular representation of  $G$  on  $L^2(\Gamma \backslash G)$ . Moreover, from the continuity of  $\lambda$ , we have

$$(4.4) \quad A(U_r(X)\phi) = U_\Gamma(X)A(\phi) \quad (X \in \mathfrak{g}, \phi \in H_r^\infty).$$

Let  $\langle , \rangle$  (resp.  $\langle , \rangle_\Gamma$ ) be the inner product on  $H_r$  (resp.  $L^2(\Gamma \backslash G)$ ). For  $K$ -finite vectors  $\phi_1, \phi_2 \in H_r^0$ , set

$$(4.5) \quad (\phi_1, \phi_2) = \langle A(\phi_1), A(\phi_2) \rangle_\Gamma.$$

Then  $( , )$  defines a  $\mathfrak{g}$ -invariant hermitian form on the  $(\mathfrak{g}, K)$ -module  $H_r^0$ . Here, by Proposition 2.2,  $H_r^0$  decomposes into the sum of the isotypic  $K$ -submodules;

$$H_r^0 = \bigoplus_{s \in \mathbf{Z}_+, s \geq -r} H_{r,s}^0.$$

Since  $A|_{H_r^0}; H_r^0 \rightarrow L^2(\Gamma \backslash G)$  is  $K$ -equivariant, this decomposition is the orthogonal direct sum with respect to  $( , )$ , too. Also, each  $H_{r,s}^0$  is finite dimensional. From these facts, it is easy to see that there is a linear mapping  $B; H_r^0 \rightarrow H_r^0$  such that

$$(4.6) \quad (\phi_1, \phi_2) = \langle B\phi_1, \phi_2 \rangle \quad (\phi_1, \phi_2 \in H_r^0).$$

Then, by (4.4), we have

$$B(U_r(X)\phi) = U_r(X)(B(\phi)) \quad (X \in \mathfrak{g}, \phi \in H_r^0).$$

Since  $H_r^0$  is an irreducible  $(\mathfrak{g}, K)$ -module,  $B$  is a scalar operator  $\nu \cdot \text{Id}$  where  $\nu \in \mathbf{R}$  and  $\nu \geq 0$ . Combining (4.5) and (4.6), we have

$$\langle A(\phi_1), A(\phi_2) \rangle_\Gamma = \nu \langle \phi_1, \phi_2 \rangle \quad (\phi_1, \phi_2 \in H_r^0).$$

This implies that  $A|_{H_r^0}$  is continuous with respect to the topology of  $H_r^0$  in  $H_r$ . Hence the operator  $A|_{H_r^0}$  extends to a bounded operator

$$\bar{A}; H_r \rightarrow L^2(\Gamma \backslash G).$$

Note that  $H_r^0$  consists of analytic vectors for  $U_r$  and  $G$  is connected. Then (4.4) implies that

$$\bar{A}(U_r(g)\phi) = U_\Gamma(g)\bar{A}(\phi) \quad (\phi \in H_r, g \in G)$$

and hence  $\bar{A}$  is an intertwining operator of  $(U_r, H_r)$  into  $(U_\Gamma, L^2(\Gamma \backslash G))$ .

On the other hand,  $\lambda$  is non-trivial. From the density of  $H_r^0$  in  $H_r^\infty$ ,  $\lambda|_{H_r^0}$  is non-trivial. Hence  $\bar{A}$  is non-trivial. The theorem is proved.

**Corollary 4.3.** *For  $l \in \mathbf{Z}_+$ , there is a uniform discrete subgroup  $\Gamma$  of  $G$  such that*

$$H^{2q}(\Gamma, F_l^*) \neq \{0\}.$$

Proof. Theorem 3.4 implies that

$$H^{2q}(\mathfrak{g}, K; H_{l+2q}^0 \otimes F_l^*) \neq \{0\}.$$

Then, by [1], I, Theorem 5.3, the infinitesimal character of  $H_{l+2q}$  is equal to that of  $F_l$ . Applying Theorem 4.2 to  $U_{l+2q}$ , we obtain a uniform discrete subgroup  $\Gamma$  such that  $m(U_{l+2q}, \Gamma) \neq 0$ . Then, by (0.1), we have

$$H^{2q}(\Gamma, F_l^*) \neq \{0\}.$$

REMARK 4.4. In the above corollary, we consider the case  $l=0$ . Then we have

$$H^{2q}(\Gamma, \mathbf{C}) \neq \{0\}.$$

More precisely,  $H^{2q}(\Gamma, \mathbf{C})$  contains a cohomology class which corresponds to a non-trivial automorphic representation. This improves the result in [4]. In [4], Millson and Raghunathan showed that, for some  $\Gamma$ ,  $H^i(\Gamma, \mathbf{C})$  contains such a class for any  $i$  strictly between 0 and  $4pq$  and divisible by either  $4p$  or  $4q$  ([4], Theorem 4.1).

---

#### References

- [1] A. Borel, N. Wallach: Continuous cohomology, discrete subgroups and representations of reductive groups, Ann. Math. Studies, No. 94, Princeton University Press, 1980.
- [2] A. Borel: Introduction aux groupes arithmétiques, Hermann, 1969.
- [3] D.H. Collingwood: *A note on continuous cohomology for semi-simple Lie groups*, Math. Z. **189** (1985), 65–70.
- [4] J.J. Millson, M.S. Raghunathan: *Geometric construction of homology for arithmetic groups I*, Geometry and analysis, Indian Academy of Sciences and Tata Institute of Fundamental Research, Springer, 1981, 103–123.
- [5] D.A. Vogan, Jr, G.J. Zuckerman: *Unitary representations with non-zero cohomology*, Compositio Math. **53** (1984), 51–90.
- [6] N. Wallach: Symplectic geometry and Fourier analysis, Math. Sci. Press, Boston, 1977.

- [7] G. Warner: Harmonic analysis on semi-simple Lie groups I, *Grund. Math. Wiss.* 188, Springer, 1972.

Department of Integrated Arts and Sciences  
University of Osaka Prefecture  
Sakai, Osaka 591  
Japan