Title	Topological higgs inflation: Origin of standard model criticality
Author(s)	Hamada, Yuta; Oda, Kinya; Takahashi, Fuminobu
Citation	Physical Review D. 90(9) p.097301
Issue Date	2014-11-21
oaire:version	VoR
URL	https://hdl.handle.net/11094/78742
rights	© 2014 American Physical Society
Note	

Osaka University Knowledge Archive : OUKA

<https://ir.library.osaka-u.ac.jp/>

Osaka University

Topological Higgs inflation: Origin of Standard Model criticalityYuta Hamada,¹ Kin-ya Oda,² and Fuminobu Takahashi^{3,4}¹*Department of Physics, Kyoto University Kyoto 606-8502, Japan*²*Department of Physics, Osaka University, Osaka 560-0043, Japan*³*Department of Physics, Tohoku University, Sendai 980-8578, Japan*⁴*Kavli Institute for the Physics and Mathematics of the Universe (WPI), TODIAS, University of Tokyo, Kashiwa 277-8583, Japan*

(Received 24 August 2014; published 21 November 2014)

The measured values of the Higgs and top masses and of the strong gauge coupling constant point to the near-criticality of the Standard Model, where two vacua at the electroweak and Planck scales are quasidegenerate. We argue that the criticality is required by the occurrence of an eternal topological inflation induced by the Higgs potential. The role of this inflation is to continuously create a sufficiently flat and homogeneous Universe, providing the necessary initial condition for the subsequent slow-roll inflation that generates the density perturbations of the right magnitude. While the condition for the topological Higgs inflation is only marginally satisfied in the Standard Model, it can be readily satisfied if one introduces the right-handed neutrinos and/or the nonminimal coupling to gravity; currently unknown quantum gravity corrections to the potential may also help. We also discuss the $B - L$ Higgs inflation as a possible origin of the observed density perturbations. Its necessary initial condition, the restored $B - L$ symmetry, can be naturally realized by the preceding topological Higgs inflation.

DOI: [10.1103/PhysRevD.90.097301](https://doi.org/10.1103/PhysRevD.90.097301)

PACS numbers: 98.80.Cq, 11.10.Hi, 14.60.St, 14.80.Bn

I. INTRODUCTION

The discovery of the Standard Model (SM) Higgs boson only so far at the LHC [1,2] already provided us with various implications for theory beyond the SM as well as the Universe we live in. In particular, the experiments seem to suggest a special structure of the vacuum.

The measured Higgs boson mass about 125 GeV [3,4] implies that the SM could be valid all the way up to the Planck scale. If so, the Higgs potential may have another minimum around the Planck scale, depending on the top quark mass; see e.g. Refs. [5–17] for the latest analyses. If the extra minimum is the global one, our electroweak vacuum is unstable and decays through quantum tunneling processes with a finite lifetime. In contrast, nature may realize a critical situation where the two minima are degenerate in energy. Froggatt and Nielsen focused on this special case, the so-called Higgs criticality; they provided a theoretical argument to support this case, the multiple point criticality principle [18,19].¹ The near-critical behavior of the Higgs potential is a puzzle and has led to much excitement in the context of Higgs inflation [21–28].

In this paper we point out that the near-criticality can be understood if the Universe experiences eternal topological inflation [29–31] induced by the Higgs field at a very early stage. We study the condition for the topological Higgs inflation to occur in the SM and what kind of extensions can relax the condition. As we shall see shortly, while the

condition is only marginally satisfied in the SM, it can be readily satisfied if one extends the SM by introducing heavy right-handed neutrinos and/or a nonminimal coupling to gravity.

The topological Higgs inflation may be thought of as one of the variants of the Higgs inflation, but it is different in the following aspects. First, the topological inflation is free of the initial condition problem. If the Universe begins in a chaotic state at an energy close to the Planck scale, the Higgs field may take various field values randomly up to the Planck scale or higher [32]. As the Universe expands, the energy density decreases and the Higgs field finds itself either larger or smaller than the critical field value corresponding to the local maximum, and gets trapped in one of the two degenerate vacua with a more or less equal probability. This leads to formation of domain walls separating the two vacua. Interestingly, then, eternal inflation could take place inside the domain walls, if the thickness of the domain walls is greater than the Hubble radius [29,31]. In this sense, no special fine-tuning of the initial position of the inflaton is necessary for the inflation to take place. Specifically, the topological inflation occurs if the two minima are separated by more than the Planck scale, which was also confirmed by numerical calculations [33]. Secondly, the magnitude of density perturbations generated by topological Higgs inflation tends to be too large to explain the observed CMB temperature fluctuations. We need therefore another inflation after the end of the topological Higgs inflation, and we will return to this issue later in this paper. Thus, the role of the topological Higgs inflation is to continuously create a sufficiently flat Universe, solving the so-called longevity problem of

¹See e.g. Sec. 5.A.1 of Ref. [20] for a list of other possibilities to realize the (near) criticality.

inflation with a Hubble parameter much smaller than the Planck scale [32,34]; the Universe must be sufficiently flat and therefore long lived so that the subsequent slow-roll inflation with a much smaller Hubble parameter can take place.

II. TOPOLOGICAL HIGGS INFLATION IN SM

For a Higgs field value much larger than the electroweak scale, the effective potential is approximately given by²

$$V = \frac{1}{4} \lambda_{\text{eff}}(\varphi) \varphi^4, \quad (1)$$

where the effective coupling is expanded around its minimum as

$$\lambda_{\text{eff}} \simeq \lambda_{\text{min}} + \frac{\beta_2}{(16\pi^2)^2} \left(\ln \frac{\varphi}{\mu_{\text{min}}} \right)^2, \quad (2)$$

with $\beta_2 \simeq 0.5$; see e.g. Refs. [20,27].

Following the multipoint criticality principle, we assume $\lambda_{\text{min}} = 0$. The derivatives of the potential read

$$V_\varphi = \frac{\beta_2}{(16\pi^2)^2} \ln \frac{\varphi}{\mu_{\text{min}}} \left(\ln \frac{\varphi}{\mu_{\text{min}}} + \frac{1}{2} \right) \varphi^3, \quad (3)$$

$$V_{\varphi\varphi} = \frac{\beta_2}{2(16\pi^2)^2} \left(1 + \ln \frac{\varphi}{\mu_{\text{min}}} \right) \left(1 + 6 \ln \frac{\varphi}{\mu_{\text{min}}} \right) \varphi^2. \quad (4)$$

Note that $\varphi = \mu_{\text{min}}$ also gives the minimum of the potential when the criticality condition $\lambda_{\text{min}} = 0$ is met. The local maximum of the potential is at

$$\varphi_t = e^{-1/2} \mu_{\text{min}}. \quad (5)$$

Let us take a domain wall separating the two degenerate minima at $\varphi = \mu_{\text{min}}$ and $\varphi \approx 0$.³ The typical thickness of the domain wall is given by [33]

$$\delta \simeq |V_{\varphi\varphi}(\varphi_t)|^{-1/2} = \frac{16\pi^2}{\mu_{\text{min}}} \sqrt{\frac{2e}{\beta_2}}. \quad (6)$$

If the thickness is greater than the Hubble radius, the domain wall will expand, and topological inflation takes place [29,31]. The Hubble parameter at the maximum (i.e. around the center of the domain wall) is

²Here we can safely treat the Higgs inflation as a single-field model. See footnote 5 of Ref. [27] and Refs. [35,36]. Equation (1) can be obtained from the one-loop effective potential, including the loops from the NG bosons.

³Here and in what follows we neglect the electroweak scale vacuum expectation value compared to the Planck scale.

$$H^2 = \frac{V(\varphi_t)}{3M_P^2} = \frac{\beta_2}{(16\pi^2)^2} \frac{\mu_{\text{min}}^4}{48e^2 M_P^2} = (2.4 \times 10^{-4} M_P)^2 \left(\frac{\beta_2}{0.5} \right) \left(\frac{\mu_{\text{min}}}{M_P} \right)^4, \quad (7)$$

where $M_P \simeq 2.4 \times 10^{18}$ GeV is the (reduced) Planck scale. The condition for the domain wall to expand is [33]

$$H\delta = \sqrt{\frac{V(\varphi_t)}{3M_P^2 |V_{\varphi\varphi}(\varphi_t)|}} = \frac{1}{\sqrt{3|\eta(\varphi_t)|}} \gtrsim 0.48, \quad (8)$$

namely,

$$|\eta(\varphi_t)| \lesssim 1.4, \quad (9)$$

where $\eta \equiv M_P^2 V_{\varphi\varphi}/V$ is one of the slow-roll parameters. For the SM, this condition reads

$$\mu_{\text{min}} \gtrsim 3.9 M_P. \quad (10)$$

Note that the prefactor of this condition may contain an $\mathcal{O}(1)$ uncertainty, because it is derived for a Z_2 symmetric potential, and the precise condition depends on the detailed shape of the inflaton potential. In Fig. 1, we plot the scale μ_{min} at which the effective coupling λ_{eff} takes its minimum value, under the criticality condition $\lambda_{\text{min}} := \lambda_{\text{eff}}(\mu_{\text{min}}) = 0$; in the computation, we have used the two-loop renormalization group equations and the one-loop effective potential in the Landau gauge. We see that μ_{min} is indeed of the order of the Planck scale, given the measured mass of the Higgs boson. The condition (10) may be satisfied taking account of the uncertainty.

FIG. 1 (color online). The scale μ_{min} that gives the minimum of the effective coupling λ_{eff} , under the criticality condition $\lambda_{\text{min}} := \lambda_{\text{eff}}(\mu_{\text{min}}) = 0$, as a function of M_H . The band width corresponds to the 2σ deviation of α_s , with 1σ being given by $\alpha_s = 0.1185 \pm 0.0006$ [37]. The dotted lines show the 2σ band for M_H , with 1σ being given by $M_H = 125.9 \pm 0.4$ GeV [37].

FIG. 2 (color online). Allowed region in the prescription I (left) and II (right) in ξ vs $\mu_{\min}$ plane, where the dark (light) region satisfies $|\eta(\varphi_I)| < 1$ (1.4) and $|\eta(\varphi_{II})| < 1$ (1.4), respectively. In prescription I (left), $\mu_{\min} < M_P/\sqrt{\xi}$ is also imposed.

If one wanted to use this potential to generate the observed density fluctuation, the potential height must satisfy [38]

$$V(\varphi_t) = \frac{\beta_2}{(16\pi^2)^2} \frac{\mu_{\min}^4}{16e^2} < (1.94 \times 10^{16} \text{ GeV})^4 \frac{r_*}{0.12}, \quad (11)$$

namely,

$$\mu_{\min} < 0.4M_P \left(\frac{r_*}{0.12} \right)^{1/4} \left(\frac{0.5}{\beta_2} \right)^{1/4}, \quad (12)$$

which is inconsistent with the condition (10). Therefore, one cannot use this topological inflation directly to generate the observed density fluctuation, and we need another inflation after the topological Higgs inflation. The minimum scale $\mu_{\min}$ can be larger if one includes the right-handed neutrino [39,40]. If one embeds the SM in string theory, the string states may also change $\mu_{\min}$; see also Ref. [41] that considers the near-criticality in string theory context.

III. NONMINIMAL COUPLING TO GRAVITY

We may add the nonminimal coupling $\xi|H|^2 R$.⁴ The Einstein frame potential then becomes

$$U = \frac{\lambda_{\text{eff}}(\mu)}{4} \frac{\varphi^4}{(1 + \xi\varphi^2/M_P^2)^2}. \quad (13)$$

There are two prescriptions for the choice of the scale μ :

$$\mu = \frac{\varphi}{\sqrt{1 + \xi\varphi^2/M_P^2}} \quad (\text{prescription I}), \quad (14)$$

⁴The topological inflation in the Starobinsky model was considered in Ref. [42].

$$\mu = \varphi \quad (\text{prescription II}). \quad (15)$$

In the Einstein frame, the canonically normalized field χ is related to φ as

$$\frac{d\chi}{d\varphi} = \frac{\sqrt{1 + \xi \frac{\varphi^2}{M_P^2} + 6\xi^2 \frac{\varphi^2}{M_P^2}}}{1 + \xi \frac{\varphi^2}{M_P^2}}. \quad (16)$$

In the following, let us check whether the condition (9) can be satisfied in each prescription by estimating the slow-roll parameter defined by

$$\eta = M_P^2 \frac{U_{\chi\chi}}{U}. \quad (17)$$

A. Prescription I

The position of the local maximum and minimum of U , under the criticality condition $\lambda_{\min} = 0$, are respectively

$$\varphi_1 = \mu_{\min} \frac{1}{\sqrt{e - \xi \frac{\mu_{\min}^2}{M_P^2}}}, \quad \varphi_{\min} = \mu_{\min} \frac{1}{\sqrt{1 - \xi \frac{\mu_{\min}^2}{M_P^2}}}, \quad (18)$$

where $\mu_{\min}$ is defined by Eq. (2) even in presence of ξ . We see that $\mu_{\min} < M_P/\sqrt{\xi}$ is required in order to allow the local minimum $\varphi_{\min}$. At φ_1 , the slow-roll parameter becomes

$$\eta = -8 \frac{M_P^2}{\mu_{\min}^2} \frac{(e - \xi \frac{\mu_{\min}^2}{M_P^2})^2}{e + 6\xi^2 \frac{\mu_{\min}^2}{M_P^2}}. \quad (19)$$

In the left panel of Fig. 2, we plot the allowed region where the condition (9) with Eq. (19) is satisfied. We can see from

the figure that there is an allowed region with $\xi \sim 0.1$ and $\mu_{\min} \sim 3M_P$, which lies in the reasonable range of $\mu_{\min} \simeq (1-3)M_P$ in the SM, shown in Fig. 1.

B. Prescription II

In prescription II, the local maximum of U , under the criticality condition $\lambda_{\min} = 0$, is located at

$$\varphi_{\text{II}} = \frac{M_P}{\sqrt{\xi}} \sqrt{W\left(\frac{\xi \mu_{\min}^2}{e M_P^2}\right)}, \quad (20)$$

where W is the Lambert function defined by $z = W(z)e^{W(z)}$. The slow-roll parameter at this maximum is

$$\eta(\varphi_{\text{II}}) = -\frac{8M_P^2}{\varphi_{\text{II}}^2} \frac{1 + \xi \frac{\varphi_{\text{II}}^2}{M_P^2}}{1 + \xi(1 + 6\xi) \frac{\varphi_{\text{II}}^2}{M_P^2}}. \quad (21)$$

In the right panel of Fig. 2, we plot the allowed region. We can see that there is a large allowed region for $\xi \gtrsim \mathcal{O}(1-10)$ for $\mu_{\min} \simeq (1-3)M_P$. We have checked that the typical value of $|\eta|$ is greater than 0.1 for $\xi \lesssim 10^5$, and therefore, it is difficult to account for the observed density perturbations by this inflation alone.

IV. DISCUSSION

So far we have focused on the critical case where the two minima are degenerate. Successful topological Higgs inflation will be still possible even if the potential energy of the Planck-scale minimum is slightly higher or lower than the electroweak minimum. If the degeneracy is largely broken, however, the trapping probability of the Higgs field will be significantly biased to the minimum with a lower energy. This would reduce the number of domain walls, especially those with an infinite length. Also the domain walls become unstable and some of them will collapse before the topological inflation takes place. Thus, we expect that the topological Higgs inflation will become less likely as the Higgs potential goes away from the criticality. We argue, therefore, that the topological Higgs inflation which occurred at the very beginning of the Universe could be the reason why the measured Higgs boson mass points to the near-criticality of the SM vacuum. In order to estimate quantitatively to what extent the degeneracy can be broken, one must resort to numerical simulation, which is left for future work.

As we have mentioned earlier, while the topological Higgs inflation cannot account for the observed CMB temperature anisotropy, it creates a sufficiently flat and homogeneous Universe, and sets the required initial condition for the subsequent slow-roll inflation with a

much smaller Hubble parameter [32,34].⁵ There is a variety of possibilities, but one of the interesting candidates is the inflation based on the $B-L$ Higgs inflation [43–47], where the Coleman-Weinberg potential gives a dominant contribution to the tilt of the inflaton potential.⁶ The well-known problem of the density perturbations in the Coleman-Weinberg inflation model can be avoided if the $U(1)_{B-L}$ gauge coupling is small [48–50] or if there is a cancellation between the gauge and neutrino Yukawa contributions [43,51]. There are a couple of interesting implications. First, the inclusion of the right-handed neutrinos makes it easier to satisfy the condition for the topological Higgs inflation, since the $\mu_{\min}$ becomes larger for the fixed $\lambda_{\min} = 0$.⁷ With three right-handed neutrinos, adding the $U(1)_{B-L}$ is one of the plausible extensions of the SM as it is anomaly free. Secondly, the Higgs field will induce preheating as it oscillates about the origin after the topological inflation [52,53]. In order to estimate the reheating temperature precisely, one needs a detailed analysis of dissipation processes [54], but the reheating temperature is expected to be rather high. If so, the $U(1)_{B-L}$ symmetry is likely restored, setting the initial condition for the $B-L$ Higgs inflation.⁸ Thus, the topological Higgs inflation provides the required initial condition for the $B-L$ Higgs inflation. Thirdly, the $B-L$ Higgs field will decay into right-handed neutrinos, whose CP -violating decay can generate a right amount of the baryon asymmetry via leptogenesis [55].

ACKNOWLEDGMENTS

We are grateful to the Yukawa Institute for Theoretical Physics for the hospitality during the workshop PPP2014 (YITP-W-14-05), where the present work started, and to its participants for fruitful discussions. We thank Hikaru Kawai for useful comments. This work was in part supported by the Grant-in-Aid from the JSPS Fellows

⁵It is interesting to study various extensions of the SM to see if the topological Higgs inflation can generate the density perturbations of the right magnitude.

⁶This extension possibly changes the running of Higgs self-coupling λ due to the quartic coupling between the $B-L$ and SM Higgs bosons. However, such a coupling does not affect the running of λ if it is smaller than of order 10^{-2} . Unless this coupling is extremely small, it helps to keep the $B-L$ Higgs field at its origin during the first inflation, setting the initial condition for the subsequent $B-L$ Higgs inflation. Detailed study of this issue will be presented elsewhere.

⁷The neutrino Dirac-Yukawa couplings do not affect the Higgs potential significantly if they are smaller than of order 0.1; see Ref. [10].

⁸During the topological Higgs inflation, the $B-L$ Higgs field may be at the origin or may develop a nonzero value. This affects the contribution of the neutrino Yukawa coupling to the SM Higgs potential, whose quantitative estimate will be left for future work.

No. 25.1107 (Y.H.), the Grant-in-Aid for Scientific Research Nos. 23104009, 20244028, and 23740192 (K.O.), the Grant-in-Aid for Scientific Research on Innovative Areas [No. 23104008 (F.T.)], JSPS Grant-in-Aid for Young Scientists (B) [No. 24740135 (F.T.)],

Scientific Research (A) [No. 26247042 (F.T.)], Scientific Research (B) [No. 26287039 (F.T.)], and Inoue Foundation for Science (F.T.). This work was also supported by World Premier International Center Initiative (WPI Program), MEXT, Japan (F.T.).

-
- [1] G. Aad *et al.* (ATLAS Collaboration), *Phys. Lett. B* **716**, 1 (2012).
 - [2] S. Chatrchyan *et al.* (CMS Collaboration), *Phys. Lett. B* **716**, 30 (2012).
 - [3] G. Aad *et al.* (ATLAS Collaboration), *Phys. Rev. D* **90**, 052004 (2014).
 - [4] CMS Collaboration, Report No. CMS-PAS-HIG-14-009.
 - [5] M. Holthausen, K. S. Lim, and M. Lindner, *J. High Energy Phys.* **02** (2012) 037.
 - [6] F. Bezrukov, M. Y. Kalmykov, B. A. Kniehl, and M. Shaposhnikov, *J. High Energy Phys.* **10** (2012) 140.
 - [7] G. Degrandi, S. Di Vita, J. Elias-Miro, J. R. Espinosa, G. F. Giudice, and A. Strumia, *J. High Energy Phys.* **08** (2012) 098.
 - [8] S. Alekhin, A. Djouadi, and S. Moch, *Phys. Lett. B* **716**, 214 (2012).
 - [9] I. Masina, *Phys. Rev. D* **87**, 053001 (2013).
 - [10] Y. Hamada, H. Kawai, and K.-y. Oda, *Phys. Rev. D* **87**, 053009 (2013).
 - [11] F. Jegerlehner, [arXiv:1304.7813](#).
 - [12] D. Buttazzo, G. Degrandi, P. P. Giardino, G. F. Giudice, F. Sala, A. Salvio, and A. Strumia, *J. High Energy Phys.* **12** (2013) 089.
 - [13] V. Branchina and E. Messina, *Phys. Rev. Lett.* **111**, 241801 (2013).
 - [14] A. Kobakhidze and A. Spencer-Smith, [arXiv:1404.4709](#).
 - [15] A. Spencer-Smith, [arXiv:1405.1975](#).
 - [16] V. Branchina, E. Messina, and A. Platania, [arXiv:1407.4112](#).
 - [17] V. Branchina, E. Messina, and M. Sher, [arXiv:1408.5302](#).
 - [18] C. Froggatt and H. B. Nielsen, *Phys. Lett. B* **368**, 96 (1996).
 - [19] H. B. Nielsen, *Bled Workshops in Physics* **13**, 94 (2012).
 - [20] Y. Hamada, H. Kawai, K.-y. Oda, and S. C. Park, [arXiv:1408.4864](#).
 - [21] F. L. Bezrukov and M. Shaposhnikov, *Phys. Lett. B* **659**, 703 (2008).
 - [22] I. Masina and A. Notari, *Phys. Rev. D* **85**, 123506 (2012).
 - [23] I. Masina and A. Notari, *Phys. Rev. Lett.* **108**, 191302 (2012).
 - [24] K. Allison, *J. High Energy Phys.* **02** (2014) 040.
 - [25] Y. Hamada, H. Kawai, and K.-y. Oda, *Prog. Theor. Exp. Phys.* **2014**, 023B02 (2014).
 - [26] J. L. Cook, L. M. Krauss, A. J. Long, and S. Sabharwal, *Phys. Rev. D* **89**, 103525 (2014).
 - [27] Y. Hamada, H. Kawai, K.-y. Oda, and S. C. Park, *Phys. Rev. Lett.* **112**, 241301 (2014).
 - [28] F. Bezrukov and M. Shaposhnikov, [arXiv:1403.6078](#).
 - [29] A. D. Linde, *Phys. Lett. B* **327**, 208 (1994).
 - [30] A. D. Linde and D. A. Linde, *Phys. Rev. D* **50**, 2456 (1994).
 - [31] A. Vilenkin, *Phys. Rev. Lett.* **72**, 3137 (1994).
 - [32] A. D. Linde, *Phys. Lett.* **129B**, 177 (1983).
 - [33] N. Sakai, H.-A. Shinkai, T. Tachizawa, and K.-i. Maeda, *Phys. Rev. D* **53**, 655 (1996).
 - [34] K. Izawa, M. Kawasaki, and T. Yanagida, *Phys. Lett. B* **411**, 249 (1997).
 - [35] R. N. Greenwood, D. I. Kaiser, and E. I. Sfakianakis, *Phys. Rev. D* **87**, 064021 (2013).
 - [36] D. I. Kaiser and E. I. Sfakianakis, *Phys. Rev. Lett.* **112**, 011302 (2014).
 - [37] J. Beringer *et al.* (Particle Data Group), *Phys. Rev. D* **86**, 010001 (2012), and 2013 partial update for the 2014 edition.
 - [38] P. Ade *et al.* (Planck Collaboration), [arXiv:1303.5082](#).
 - [39] N. Haba and R. Takahashi, *Phys. Rev. D* **89**, 115009 (2014).
 - [40] Y. Hamada, H. Kawai, and K.-y. Oda, *J. High Energy Phys.* **07** (2014) 026.
 - [41] A. Hebecker, A. K. Knochel, and T. Weigand, *Nucl. Phys.* **B874**, 1 (2013).
 - [42] K. Kamada and J. Yokoyama, [arXiv:1405.6732](#) [*Phys. Rev. D* (to be published)].
 - [43] K. Nakayama and F. Takahashi, *J. Cosmol. Astropart. Phys.* **10** (2011) 033.
 - [44] K. Nakayama and F. Takahashi, *Phys. Lett. B* **707**, 142 (2012).
 - [45] K. Nakayama and F. Takahashi, *J. Cosmol. Astropart. Phys.* **05** (2012) 035.
 - [46] G. Barenboim, E. J. Chun, and H. M. Lee, *Phys. Lett. B* **730**, 81 (2014).
 - [47] S. Iso, K. Kohri, and K. Shimada, [arXiv:1408.2339](#).
 - [48] S. Hawking, *Phys. Lett.* **115B**, 295 (1982).
 - [49] A. A. Starobinsky, *Phys. Lett.* **117B**, 175 (1982).
 - [50] A. H. Guth and S. Pi, *Phys. Rev. Lett.* **49**, 1110 (1982).
 - [51] J. R. Ellis, D. V. Nanopoulos, K. A. Olive, and K. Tamvakis, *Phys. Lett.* **118B**, 335 (1982).
 - [52] F. Bezrukov, D. Gorbunov, and M. Shaposhnikov, *J. Cosmol. Astropart. Phys.* **06** (2009) 029.
 - [53] J. Garcia-Bellido, D. G. Figueroa, and J. Rubio, *Phys. Rev. D* **79**, 063531 (2009).
 - [54] K. Mukaida and K. Nakayama, *J. Cosmol. Astropart. Phys.* **03** (2013) 002.
 - [55] M. Fukugita and T. Yanagida, *Phys. Lett. B* **174**, 45 (1986).