

| | |
|--------------|---|
| Title | On the degrees of irreducible representations of finite groups |
| Author(s) | Watanabe, Atumi |
| Citation | Osaka Journal of Mathematics. 18(2) P.403-P.406 |
| Issue Date | 1981 |
| Text Version | publisher |
| URL | https://doi.org/10.18910/7939 |
| DOI | 10.18910/7939 |
| rights | |
| Note | |

Osaka University Knowledge Archive : OUKA

<https://ir.library.osaka-u.ac.jp/>

Osaka University

ON THE DEGREES OF IRREDUCIBLE REPRESENTATIONS OF FINITE GROUPS

ATUMI WATANABE

(Received November 29, 1979)

1. Introduction

Let G be a finite group of order $|G|$ and F be an algebraically closed field of characteristic 0. Let T be an irreducible representation of G over F and d_T be the degree of T . As is well known, d_T divides $|G|$. Furthermore there exists a sharper result due to Ito [2], namely, d_T divides the index in G of every abelian normal subgroup of G . Let s_T be the order of $\det T$, that is, s_T is the smallest natural number such that $|T(x)|^{s_T}=1$ for all $x \in G$, where $|T(x)|$ is the determinant of $T(x)$. In Lemma of [4] we showed the first part of the following

Theorem 1. *Let T be an irreducible representation of G over F . Then we have*

- (i) $d_T s_T \mid 2|G|$,
- (ii) if d_T or s_T is odd then $d_T s_T \mid |G|$.

The second part follows from (i) by considering the 2-part of $d_T s_T$, since both d_T and s_T divide $|G|$.

The purpose of the present paper is to prove the following theorems.

Theorem 2. *If G has an irreducible representation T over F with $d_T s_T \nmid |G|$, then the following holds.*

- (i) *A 2-Sylow subgroup P of G is cyclic and $P \neq 1$. Hence G has the normal 2-complement K .*
- (ii) $C_P(K)=1$.
- (iii) *T is induced from a representation of K .*

The converse of Theorem 2 is also true:

Theorem 3. *If G satisfies (i) and (ii) in Theorem 2, then G has an irreducible representation T with $d_T s_T \nmid |G|$.*

We also have the following

Theorem 4. *Let T be an irreducible representation of G over F . Then we have*

$$d_T s_T \leq |G|.$$

If $d_T s_T = G$, then G is cyclic.

I express my thanks to Professor H. Nagao for his valuable advice.

2. Proofs of the theorems

To prove our theorems we need the following Lemma.

Lemma. *Let T be an irreducible representation of G over F , H a normal subgroup of G of index n and T_0 be an irreducible component of T_H , T_H the restriction of T to H . Then we have the following.*

- (i) *If $T_H = T_0$, then $d_T = d_{T_0}$ and $s_T \mid n s_{T_0}$.*
- (ii) *If $T = T_0^G$, then $d_T = n d_{T_0}$ and $s_T \mid 2 s_{T_0}$. Furthermore if $2 \mid d_{T_0} s_{T_0}$ or s_T is odd, then $s_T \mid s_{T_0}$.*

Proof. (i) is clear. We prove (ii). Clearly $d_T = n d_{T_0}$. We set $s_T = s$, $d_{T_0} = d_0$ and $s_{T_0} = s_0$. Let x_1, \dots, x_n be a complete set of coset representatives of H in G . We extend T_0 to all elements of G by setting $T_0(x) = 0$ for all $x \notin H$. We may assume that $T(x)$ is a $n \times n$ matrix of blocks whose (i, j) -th entry is the $d_0 \times d_0$ matrix $T_0(x_i^{-1} x x_j)$:

$$T(x) = \begin{pmatrix} T_0(x_1^{-1} x x_1) & \dots & T_0(x_1^{-1} x x_n) \\ \dots & \dots & \dots \\ T_0(x_n^{-1} x x_1) & \dots & T_0(x_n^{-1} x x_n) \end{pmatrix} \quad (x \in G).$$

Hence for each $x \in G$, we have $|T(x)| = (-1)^{d_0 m} |T_0(y_1)| \dots |T_0(y_n)|$, where $y_i \in H$, $i = 1, \dots, n$, and m is an integer. Therefore, for each $x \in G$, $|T(x)|^{2s_0} = 1$ and hence $s \mid 2s_0$. If s is odd then $s \mid s_0$, and if d_0 or s_0 is even then $|T(x)|^{s_0} = 1$ for each $x \in G$ and hence $s \mid s_0$.

Proof of Theorem 2. We prove (i) by induction on $|G|$. Put $d_T = d$ and $s_T = s$. Since $ds \nmid |G|$, by Theorem 1, (ii) $2 \mid d$ and $2 \mid s$. In particular $P \neq 1$ and $2 \mid |G: G'|$, where G' is the commutator subgroup of G . Let H be a normal subgroup of G of index 2 and T_0 be an irreducible component of T_H . By Clifford's theorem, $T_H = T_0$ or $T = T_0^G$. Put $d_{T_0} = d_0$ and $s_{T_0} = s_0$.

(a) Suppose $T_H = T_0$. Since $ds \nmid |G|$, by Lemma, (i) $d_0 s_0 \nmid |H|$ and hence both d_0 and s_0 are even. Therefore by the induction hypothesis, $P \cap H$ is cyclic. Suppose P is not cyclic. For each $x \in P$, $\langle x^2 \rangle \neq P \cap H$ and $|T(x)|^2 = |T_0(x^2)|$. Hence $|T_0(x^2)|^{s_0/2} = 1$ and $|T(x)|^{s_0} = 1$. On the other hand, for each 2-regular element $x \in G$, $|T(x)|^{s_0} = 1$, because $s \mid 2s_0$. Therefore for each $x \in G$, $|T(x)|^{s_0} = 1$ and hence $s \mid s_0$. Thus $ds \mid d_0 s_0$ and $d_0 s_0 \mid 2|H| = |G|$, which is a contradiction. Therefore P is cyclic.

(b) Suppose $T = T_0^G$. We may assume $4 \mid |G|$. Suppose $d_0 s_0$ is odd.

Then since $d=2d_0$, s is even and $s \mid 2s_0$ we have $ds=4r$ with r odd. By Theorem 1, (i) $r \mid |G|$ and hence $ds \mid |G|$, which is a contradiction. Thus $2 \mid d_0s_0$ and by Lemma, (ii) $s \mid s_0$. Then $ds \mid 2d_0s_0$ and hence $d_0s_0 \nmid |H|$. By the induction hypothesis, $P \cap H$ is cyclic. Suppose P is not cyclic. For each $x \in P \cap H$, $|T(x)| = |T_0(x)| |T_0(y^{-1}xy)| = |T_0(xy^{-1}xy)|$, where y is an element of P which does not belong to $P \cap H$. Since $P \cap H$ is a cyclic 2-group and x and $y^{-1}xy$ are of the same order, $xy^{-1}xy$ does not generate $P \cap H$. Hence $|T_0(xy^{-1}xy)|^{s_0/2} = 1$ and $|T(x)|^{s_0/2} = 1$. For each $x \in P$ which does not belong to $P \cap H$, $|T(x)| = |T_0(x^2)|$, because d_0 is even. Since P is not cyclic, $|T_0(x^2)|^{s_0/2} = 1$, hence $|T(x)|^{s_0/2} = 1$. Therefore for each $x \in P$, $|T_0(x)|^{s_0/2} = 1$. On the other hand, for each 2-regular element $x \in G$, $|T_0(x)|^{s_0/2} = 1$ because $s \mid s_0$. Hence $s \mid s_0/2$ and $ds \mid 2d_0 \cdot s_0/2 = d_0s_0$. By Theorem 1, (i) we have $ds \mid |G|$. This is a contradiction. Therefore P is cyclic. By Burnside's theorem G has the normal 2-complement K . Thus (i) is proved.

Now we show (iii). Let T_1 be an irreducible component of T_K , \tilde{K} the inertial group of T_1 in G and let \tilde{T} be an irreducible representation of \tilde{K} such that $T = \tilde{T}^G$ and that T_1 is an irreducible component of \tilde{T}_K . Put $d_{T_1} = d_1$, $s_{T_1} = s_1$, $d_{\tilde{T}} = \tilde{d}$ and $s_{\tilde{T}} = \tilde{s}$. Since \tilde{K}/K is cyclic, $\tilde{T}_K = T_1$ and $\tilde{d} = d_1$ (see the proof of [1, (9.12)]). As \tilde{d} is odd, $\tilde{d}\tilde{s} \mid |\tilde{K}|$ by Theorem 1, (ii). If $2 \mid \tilde{d}\tilde{s}$, then $s \mid \tilde{s}$ by Lemma, (ii). Hence $ds \mid |G: \tilde{K}| \tilde{d}\tilde{s} \mid |G: \tilde{K}| |\tilde{K}| = |G|$. This yields a contradiction. Hence $2 \nmid \tilde{d}\tilde{s}$. Since $|\tilde{K}: K|$ is a power of 2, by Theorem 1, (i) $\tilde{d}\tilde{s} \mid |K|$. Therefore $ds \mid 2|G: \tilde{K}| \tilde{d}\tilde{s} \mid 2|G: \tilde{K}| |K|$. Thus we see $\tilde{K} = K$. This completes the proof of (iii).

Finally we prove (ii). From (iii), $|P| \mid d$. From (i), $C_P(K)$ is a central subgroup of G . Hence $d \mid |G: C_P(K)|$. Therefore $C_P(K) = 1$. This completes the proof of the theorem.

Proof of Theorem 3. We set $|P| = 2^a$, $P = \langle x \rangle$ and $y = x^{2^a-1}$. Since $C_P(K) = 1$, y induces a non-identity automorphism of K . By [3, Satz 108], there is a conjugate class of K which is not fixed by y . Hence y does not fix some irreducible representation of K over F , say T_0 . Since $\langle yK \rangle$ is the unique minimal subgroup of G/K , K is the inertial group of T_0 in G . Hence T_0^G is an irreducible representation of G . We set $T = T_0^G$. Then $2^a \mid d_T$ and we see $|T(x)| = -1$. Hence $d_T s_T \nmid |G|$.

Proof of Theorem 4. We prove by induction on $|G|$. If G is abelian, then the theorem is trivial. We assume that the theorem is true for any proper subgroup of G . First we prove $ds \leq |G|$. Suppose $ds > |G|$. By Theorem 1, (i) $ds = 2|G|$. Since $d \mid |G|$, s is even and $2 \mid |G: G'|$. Let H be a normal subgroup of G of index 2 and T_0 be an irreducible component of T_H . By the induction hypothesis and Lemma, $T = T_0^G$ and $d_{T_0} s_{T_0} = |H|$. By the induction hypothesis, H is cyclic, $d = 2$ and $s = |G|$. Hence G is abelian, which con-

tradicts $d=2$. Thus we have proved $ds \leq |G|$. Next we prove the remaining part of the theorem. Suppose $ds = |G|$. We may assume $G \neq 1$. Since $d < |G|$, $s \neq 1$ and hence $G \neq G'$. Let L be a normal subgroup of G of prime index p and T_1 be an irreducible component of T_L . We prove L is cyclic. By the induction hypothesis and Lemma, if $T_L = T_1$ or if $T = T_1^G$ and $d_{T_1} s_{T_1}$ is even, we see easily L is cyclic. Put $d_{T_1} = d_1$ and $s_{T_1} = s_1$. If $T = T_1^G$ and $d_1 s_1$ is odd, then we see $|L| = d_1 s_1$ or $|L| = 2d_1 s_1$. By the induction hypothesis, $|L| = d_1 s_1$ implies L is cyclic. In the case $|L| = 2d_1 s_1$, let U be the normal 2-complement of L . As $d_1 s_1$ is odd, by Clifford's theorem and Lemma, (ii) we see that $(T_1)_U$ is irreducible and s_1 is the order of $\det (T_1)_U$ and that $d_1 s_1 = |U|$. Hence by the induction hypothesis, U is cyclic. Hence $d_1 = 1$ and $s_1 = |U|$ and hence $|L'| \mid 2$. On the other hand $L' \subset U$. Therefore $L' = 1$ and L is cyclic. Thus we have proved that L is cyclic. If $T_L = T_1$, then $d = 1$ and $s = |G|$, hence G is cyclic. Suppose $T = T_1^G$, then $d = p$, $|G'| = p$ and $s = |G : G'|$. Let M be any normal subgroup of G of prime index. By the argument applied to L and by $d = p$, $|G : M| = p$. Hence G/G' is a p -group and hence G is a p -group. Since $|G'| = p$, G' is a central subgroup of G . By $s = |G : G'|$, G/G' is cyclic. Therefore G is abelian. This contradicts $d = p$, and this completes the proof.

References

- [1] W. Feit: Characters of finite groups, Benjamin, New York, 1967.
- [2] N. Ito: *On the degrees of irreducible representations of a finite group*, Nagoya Math. J. **3** (1951), 5-6.
- [3] A. Speiser: Die Theorie der Gruppen von endlicher Ordnung, 3rd., Berlin.
- [4] A. Watanabe: *On Fong's reductions*, Kumamoto J. Sci. (Math.) **13** (1979), 48-54.

Department of Mathematics
Faculty of Science
Kumamoto University
(860) Kurokami 2-39-1,
Kumamoto, Japan