

Title	On a theorem of Zariski-van Kampen type and its applications
Author(s)	Matsuno, Takanori
Citation	Osaka Journal of Mathematics. 32(3) P.645-P.658
Issue Date	1995
Text Version	publisher
URL	https://doi.org/10.18910/7960
DOI	10.18910/7960
rights	
Note	

Osaka University Knowledge Archive : OUKA

<https://ir.library.osaka-u.ac.jp/>

Osaka University

ON A THEOREM OF ZARISKI - VAN KAMPEN TYPE AND ITS APPLICATIONS

TAKANORI MATSUNO

(Received May 11, 1994)

1. Introduction. Zariski constructed a method to calculate $\pi_1(\mathbf{P}^2 - C)$, where \mathbf{P}^2 is the complex projective plane and C is a curve on it. In this paper, following the ideas of Zariski [5] and Van Kampen [4], we give a method to calculate $\pi_1(E - S)$, where E is a holomorphic line bundle over a complex manifold M and S is a hypersurface of E under certain conditions. Applying our method and the Reidemeister-Schreier method (see Rolfsen [3]), we can calculate the fundamental groups of regular loci of certain normal complex spaces. We give a few concrete examples in the final section.

This paper is a revised version of the author's master thesis [1]. The author would like to express his thanks to Professor M. Namba for his useful suggestions and encouragements and to Professor M. Sakuma whose suggestions about Lemma 1 (see section 2) was a great help to prove Main Theorem. He also expresses his thanks to the referee for useful comments.

2. Statement of Main Theorem. Let M be a connected n -dimensional complex manifold and $\mu: E \rightarrow M$ be a holomorphic line bundle over M and S be a hypersurface of E . We assume that E and S satisfy the following conditions:

(1) $\mu: S \rightarrow M$ is a finite proper holomorphic map, where μ' is the restriction of μ to $S(\mu' = \mu|_S)$.

(2) There is a hypersurface B of M such that $\mu'|_{S - \mu^{-1}(B)}: S - \mu^{-1}(B) \rightarrow M - B$ is an unbranched covering of degree d .

(3) $(d\mu')_p: T(S - \mu^{-1}(B))_p \rightarrow T(M - B)_{\mu'(p)}$ is isomorphic for every point $p \in S - \mu^{-1}(B)$.

Then we have a following lemma whose proof is given in section 4.

Lemma 1. $\mu|_{E - S - \mu^{-1}(B)}: E - S - \mu^{-1}(B)$ is a continuous fiber bundle.

We denote a standard fiber of $\mu: E \rightarrow M$ by \widehat{F} and that of $\mu|_{E - S - \mu^{-1}(B)}: E - S - \mu^{-1}(B) \rightarrow M - B$ by F . We assume that there is a continuous section $\xi: M \rightarrow E$ of $\mu: E \rightarrow M$ such that $\xi(M) \cap S = \emptyset$ (see Figure 1).




Figure 1

REMARK. Such a continuous section does not always exist. For example, if E is a negative line bundle and S is the image of the zero section, then there exists no such a continuous section.

In order to describe Main Theorem, we must prepare some more symbols. We choose $F \cap \xi(*)$ as a base point b_0 and we omit the base point hereafter. Since F can be identified with $C - \{n \text{ points}\}$, $\pi_1(F)$ is isomorphic to the n -th free group $F_n = \langle \gamma_1, \dots, \gamma_n \rangle$ (see Figure 2).




Figure 2

Let \widehat{Q} be the kernel of the surjective homomorphism

$$j_* : \pi_1(M - B) \rightarrow \pi_1(M),$$

induced from the injection $i : M - B \hookrightarrow M$. We assume that \widehat{Q} has a finite presentation as follows :

$$\widehat{Q} = \langle \beta_1, \dots, \beta_i \mid \square = 1, \dots, \square = 1 \text{ (some relations)} \rangle.$$

Let $\theta : \pi_1(M - B) \rightarrow B_n$ be the braid monodromy representation of the continuous fiber bundle $\mu|_{E - S - \mu^{-1}(B)}$ in Lemma 1, where B_n is the n -th braid group :

$$B_n = \langle \sigma_1, \dots, \sigma_{n-1} \mid [\sigma_i, \sigma_j] = 1 (|i - j| \geq 2), \sigma_i \sigma_{i+1} \sigma_i = \sigma_{i+1} \sigma_i \sigma_{i+1} (i = 1, \dots, n - 2) \rangle.$$

We define a homomorphism $\varphi : B_n \rightarrow \text{Aut}(\pi_1(F))$ as follows :

$$\begin{cases} \varphi(\sigma_j)(\gamma_j) &= \gamma_j^{-1} \gamma_{j+1} \gamma_j \\ \varphi(\sigma_j)(\gamma_{j+1}) &= \gamma_j \\ \varphi(\sigma_j)(\gamma_k) &= \gamma_k \text{ (if } k \neq j, j + 1). \end{cases}$$

Then we have the following theorem of Zariski-Van Kampen type :

Main Theorem. *If there is a continuous section ξ of $\mu : E \rightarrow M$ such that $\xi(M) \cap S = \phi$, then*

$$\pi_1(E - S) \cong \langle \gamma_1, \dots, \gamma_n \mid \gamma_j = \varphi(\theta(\beta_k))(\gamma_j) (1 \leq j \leq n, 1 \leq k \leq t) \rangle \rtimes \pi_1(M).$$

(a semi-direct product)

Here β_1, \dots, β_k generate the kernel of the homomorphism $j_* : \pi_1(M - B) \rightarrow \pi_1(M)$ and $\gamma_1, \dots, \gamma_n$ generate the image of the homomorphism $i_* : \pi_1(F) \rightarrow \pi_1(E - S - \mu^{-1}(B))$, where i_* is induced from injection $i : F \hookrightarrow E - Q - \mu^{-1}(B)$.

REMARK. In Main Theorem, the relations $\gamma_j = \varphi(\theta(\beta_k))(\gamma_j)$ are same as the usual monodromy relations, so it is not essential to factor the homomorphism

$$\pi_1(M - B) \longrightarrow \text{Aut}(\pi_1(F))$$

through the braid group.

Corollary. *Under the same assumptions in Main Theorem, assume moreover that M is simply connected (i.e. $\pi_1(M) = \{1\}$), then*

$$\pi_1(E - S) \cong \langle \gamma_1, \dots, \gamma_n \mid \gamma_j = \varphi(\theta(\beta_k))(\gamma_j) (1 \leq j \leq n, 1 \leq k \leq t) \rangle.$$

3. Proof of Main Theorem. Since $\mu : E - S - \mu^{-1}(B) \rightarrow M - B$ is a continuous fiber bundle, there is the following exact sequence :

$$\begin{aligned} \cdots \rightarrow \pi_2(F) &\xrightarrow{i_*} \pi_2(E - S - \mu^{-1}(B)) \xrightarrow{\mu_*} \pi_2(M - B) \xrightarrow{\Delta} \\ &\rightarrow \pi_1(F) \xrightarrow{i_*} \pi_1(E - S - \mu^{-1}(B)) \xrightarrow{\xi_*} \pi_1(M - B) \rightarrow \\ &\rightarrow \pi_0(F) \rightarrow \cdots \text{ (exact),} \end{aligned}$$

where μ_* and ξ_* are the homomorphisms induced by μ and ξ respectively.

$\mu_* \circ \xi_* = id|_{\pi_2(M - B)}$, since $\mu \circ \xi = id|_{M - B}$. Therefore we have $\Delta = \Delta \circ \mu_* \circ \xi_* = 0$. On the other hand $\pi_0(F) = \{1\}$, since F is connected. Hence we have the following exact sequence :

$$(1) \quad 1 \rightarrow \pi_1(F) \xrightarrow{i_*} \pi_1(E - S - \mu^{-1}(B)) \xrightarrow[\xi_*]{\mu_*} \pi_1(M - B) \rightarrow 1.$$

We denote $i_*\pi_1(F)$ by K , $\xi_*\pi_1(M - B)$ by H and $\pi_1(E - S - \mu^{-1}(B))$ by G . The short exact sequence means that :

$$G \cong K \rtimes H \text{ (a semi-direct product).}$$

Now let $B = B_1 \cup \dots \cup B_l$ be the irreducible decomposition of B and α_j be the meridian of B_j (see Figure 3).




Figure 3

REMARK. Here we assume that B has a finite irreducible decomposition for simplicity. But even if B has an infinite irreducible decomposition the following argument is the same.

From a theorem of Van Kampen [4] (see also Namba [2] Cor.1.2.8), we have the following exact sequence :

$$1 \rightarrow \langle\langle \alpha_1, \dots, \alpha_l \rangle\rangle^{\pi_1(M-B)} \rightarrow \pi_1(M - B) \rightarrow \pi_1(M) \rightarrow 1$$

where $\hat{Q} = \langle\langle \alpha_1, \dots, \alpha_l \rangle\rangle^{\pi_1(M-B)}$ is the smallest normal subgroup of $\pi_1(M - B)$ which contains $\alpha_1, \dots, \alpha_l$.

$\mu^{-1}(B)$ is a hypersurface of E , which has the irreducible decomposition

$$\mu^{-1}(B) = \mu^{-1}(B_1) \cup \dots \cup \mu^{-1}(B_l).$$

$\xi_*(\alpha_j)$ is a meridian of $\mu^{-1}(B_j)$, for $\mu : E \rightarrow M$ is a line bundle and so $d\mu : T_pM \rightarrow T_{\mu(p)}M$ is surjective. Then, from the theorem of Van Kampen again, we have the following exact sequence :

$$1 \rightarrow \langle\langle \xi_*(\alpha_1), \dots, \xi_*(\alpha_l) \rangle\rangle^G \rightarrow G \rightarrow \pi_1(E - S) \rightarrow 1,$$

where $\langle\langle \xi_*(\alpha_1), \dots, \xi_*(\alpha_l) \rangle\rangle^G$ is the smallest normal subgroup of G which contains $\xi_*(\alpha_1), \dots, \xi_*(\alpha_l)$.

We denote $\langle\langle \xi_*(\alpha_1), \dots, \xi_*(\alpha_l) \rangle\rangle^G$ by N , $\xi_*(\langle\langle \alpha_1, \dots, \alpha_l \rangle\rangle^{\pi_1(M-B)})$ by Q and KN by R .

Then we can easily check that

$$(2) \quad N \cap H = Q \text{ and } R \cap NH = N.$$

Consider the natural exact sequence

$$1 \rightarrow R/N \rightarrow G/N \rightarrow G/R \rightarrow 1.$$

Note that, by (1) and (2),

$$G/R = KH/R = (KN)(NH)/R = R(NH)/R \cong (NH)/(R \cap (NH)) = (NH)/N.$$

Hence, we have the exact sequence

$$(3) \quad 1 \rightarrow R/N \rightarrow G/N \xrightarrow{f} (NH)/N \rightarrow 1.$$

The homomorphism $g : (NH)/N \rightarrow G/N$ defined by

$$g : nh(\text{mod } N) \mapsto h(\text{mod } N) \quad (n \in N, h \in H)$$

is well-defined and satisfies $f \circ g =$ the identity. Hence the exact sequence (3) splits, so

$$G/N \cong (R/N) \rtimes (NH/N) \quad (\text{a semi-direct product}).$$

We can easily check that

$$K \cap N = \langle\langle a^{-1}qaq^{-1} \mid a \in K, q \in Q \rangle\rangle^K,$$

where $\langle\langle a^{-1}qaq^{-1} \mid a \in K, q \in Q \rangle\rangle^K$ is the smallest normal subgroup of K which contains $\{a^{-1}qaq^{-1} \mid a \in K, q \in Q\}$. Furthermore, note that if K and Q are respectively generated by $\{a_1, \dots, a_n\}$ and $\{q_1, \dots, q_t\}$, then

$$K \cap N = \langle\langle a_j^{-1}q_k a_j q_k^{-1} \mid 1 \leq j \leq n, 1 \leq k \leq t \rangle\rangle^K.$$

We assume that $\langle\langle \alpha_1, \dots, \alpha_l \rangle\rangle^{\pi_1(M-B)}$ has a finite presentation as follows :

$$\langle\langle \alpha_1, \dots, \alpha_l \rangle\rangle^{\pi_1(M-B)} = \langle \beta_1, \dots, \beta_t \mid \square = 1, \dots, \square = 1 \text{ (some relations)} \rangle.$$

Since $K = i_* \pi_1(F)$ is isomorphic to the n -th free group $\langle \gamma_1, \dots, \gamma_n \rangle$, we have :

$$K \cap N \cong \langle\langle \gamma_j^{-1} \xi_*(\beta_k) \gamma_j \xi_*(\beta_k)^{-1} \mid 1 \leq j \leq n, 1 \leq k \leq t \rangle\rangle^K.$$

Thus,

$$K/(K \cap N) \cong \langle \gamma_1, \dots, \gamma_n \mid \gamma_j^{-1} \xi_*(\beta_k) \gamma_j \xi_*(\beta_k)^{-1} = 1 \mid 1 \leq j \leq n, 1 \leq k \leq t \rangle.$$

Since G/N is isomorphic to $\pi_1(E-S)$, R/N is isomorphic to $K/(K \cap N)$ and $NH/N \cong H/(N \cap H) = H/Q$ is isomorphic to $\pi_1(M)$, we have :

$$\pi_1(E - S) \cong (K / (K \cap N)) \rtimes \pi_1(M) \text{ (a semi-direct product).}$$

Now using φ and θ (defined in section 2), we have :

$$\xi_*(\beta_k) \gamma_j \xi_*(\beta_k)^{-1} = \varphi(\theta(\beta_k))(\gamma_j) \text{ (see Figure 4).}$$

Hence,

$$K / (K \cap N) \cong \langle \gamma_1, \dots, \gamma_n | \gamma_j = \varphi(\theta(\beta_k))(\gamma_j) (1 \leq j \leq n, 1 \leq k \leq t) \rangle.$$

This completes the proof of Main Theorem.




Figure 4

4. Proof of Lemma 1. (Due to M.Sakuma): For a given point $q \in M - B$, we can take a neighborhood U of q such that

- (i) $\mu^{-1}(U) \underset{\text{homeomorphic}}{\approx} \coprod_{i=1}^n \tilde{U}_i (\mu : \tilde{U}_i \xrightarrow{\sim} U)$ We write $\mu^{-1}(q) \cap \tilde{U}_i = \{\tilde{q}_i\}$.
- (ii) The following diagram is commutative.

$$\begin{array}{ccc} \mu^{-1}(U) & \xrightarrow{\sim} & U \times C \\ \mu \searrow \cap \nearrow P_1 & & \text{(where } P_1 : (p, z) \mapsto p \text{)} \\ U & & \end{array}$$

Here we define a map $h_i : \tilde{U}_i \rightarrow C$ as follows :

$$h_i : \tilde{U}_i \rightarrow \mu^{-1}(U) \cong U \times C \xrightarrow{\text{projection}} C.$$

Then we can write \tilde{U}_i as follows :

$$\tilde{U}_i = \{(x, h_i(x)) \in U \times C | x \in U\}.$$

We write $z_j = h_i(\tilde{q}_i)$, then there exists a positive number $\varepsilon > 0$ such that

- (1) $\text{Im } h_i \subset \text{Int}(D_\varepsilon(z_i))$, where $D_\varepsilon(z_i)$ is an ε -disk whose center is z_i and $\text{Int}(D_\varepsilon(z_i))$ is the interior of $D_\varepsilon(z_i)$.
- (2) $D_\varepsilon(z_1), \dots, D_\varepsilon(z_n)$ are disjoint each other.

From Lemma 2 bellow, there exists a fiber preserving homeomorphism Φ such

that $\Phi(\tilde{U}_i) = U \times \{x\}$.

$$\begin{array}{c} \mu^{-1}(U) \cong U \times \mathbf{C} \xrightarrow{\phi} U \times \mathbf{C} \\ \mu \searrow \cap \downarrow P_1 \cap \nearrow P_1 \\ U \end{array}$$

So we can take local coordinates of $\mu : E - S - \mu^{-1}(B) \rightarrow M - B$. This shows Lemma 1.

Lemma 2. *Let D be an ε -disk of \mathbf{C} whose center is the origin. Let U be the neighborhood of q as above. Let $h : U \rightarrow \text{Int}(D)$ be a continuous map such that $h(q) = 0$, where $\text{Int}(D)$ is the interior of D . Put $\tilde{U} = \{(x, h(x)) \in U \times \text{Int}(D) \mid x \in U\} \subset U \times \text{Int}(D)$. Then there exists a homeomorphism $\Psi : U \times D \xrightarrow{\sim} U \times D$ such that*

- (i) $\Psi(\tilde{U}) = U \times \{0\}$.
- (ii) Ψ is fiber preserving. (i.e. the following diagram is commutative.)

$$\begin{array}{ccc} & \Psi & \\ U \times D & \xrightarrow{\quad} & U \times D \\ & \searrow \cap \nearrow & \\ & U & \end{array}$$

- (iii) $\Psi|_{U \times \partial D} : U \times \partial D \rightarrow U \times \partial D$ is the identity map.

Proof of Lemma 2. First we define a homeomorphism $H_x : D \rightarrow D$ for each point $x \in U$ as follows:

- (i) $H_x(h(x)) = 0$.
- (ii) $H_x|_{\partial D} = id|_{\partial D}$.
- (iii) H_x is extended to D with radial extension (see Figure 5).




Figure 5

Second we define a homeomorphism $\Psi : U \times D \xrightarrow{\sim} U \times D$ as follows :

$$\Psi(x, z) = \Psi(x, H_x(z)).$$

Ψ satisfies the above conditions. (q.e.d.)

5. Case of Trivial Line Bundle. In Main Theorem, we assumed the existence of a continuous section ξ such that $\xi(M) \cap S = \phi$. In the case of the trivial line bundle we can prove the following proposition :

Proposition 1. *Let M be a connected complex manifold and $\mu : E \rightarrow M$ be a trivial line bundle on M ($i, e, E = M \times C$ and $\mu(p, z) = p$ for every point $(p, z) \in M \times C$). Let f_1, \dots, f_n be holomorphic functions on M and S be the hypersurface of E defined by*

$$S = \{(p, z) \in E \mid z^n + f_1(p)z^{n-1} + \dots + f_n(p) = 0\}.$$

Then there is a continuous section ξ of $\mu : E \rightarrow M$ such that $\xi(M) \cap S = \phi$.

Proof. We define a continuous function $h : M \rightarrow C$ by

$$h(p) = |f_1(p)| + \dots + |f_n(p)| + 1.$$

We define a section $\xi : M \rightarrow E$ by

$$\xi(p) = (p, h(p)).$$

One can easily see that this section ξ of μ satisfies $\xi(M) \cap S = \phi$. In fact, if there is a point $p \in M$ such that $\xi(p) \in S$, then

$$\{h(p)\}^n + f_1(p)\{h(p)\}^{n-1} + \dots + f_n(p) = 0.$$

Since $h(p) \geq 1$

$$1 = \frac{f_1(p)}{h(p)} - \frac{f_2(p)}{\{h(p)\}^2} - \dots - \frac{f_n(p)}{\{h(p)\}^n}.$$

Hence

$$1 \leq \frac{|f_1(p)|}{h(p)} + \frac{|f_2(p)|}{\{h(p)\}^2} + \dots + \frac{|f_n(p)|}{\{h(p)\}^n}.$$

Since $\{h(p)\}^k \geq h(p)$ ($k = 1, 2, \dots$),

$$\begin{aligned} 1 &\leq \frac{|f_1(p)|}{h(p)} + \frac{|f_2(p)|}{h(p)} + \dots + \frac{|f_n(p)|}{h(p)} \\ &= \frac{|f_1(p)| + \dots + |f_n(p)|}{|f_1(p)| + \dots + |f_n(p)| + 1} < 1. \end{aligned}$$

A contradiction.

(q.e.d.)

Let $\mu : C^{m+1} \rightarrow C^m$ be the trivial line bundle on C^m defined by

$$\mu : (z_1, \dots, z_m, z_{m+1}) \rightarrow (z_1, \dots, z_m).$$

Let S be the hypersurface of C^{m+1} defined by

$$S = \{(z_1, \dots, z_m, z_{m+1}) \in \mathbb{C}^{m+1} \mid z_{m+1}^n + f_1(z)z_{m+1}^{n-1} + \dots + f_n(z) = 0\} \dots (1),$$

where $z = (z_1, \dots, z_m)$ and $f_1(z), \dots, f_n(z)$ are polynomials.

By Corollary to Main Theorem and Proposition 1, we have

Theorem 1. *Let S be the hypersurface of \mathbb{C}^{m+1} defined by (1). Then,*

$$\pi_1(\mathbb{C}^{m+1} - S) \cong \langle \gamma_1, \dots, \gamma_n \mid \gamma_j = \varphi(\theta(\beta_k))(\gamma_j), (1 \leq j \leq n, 1 \leq k \leq t) \rangle.$$

Furthermore, let $(X_0 : X_1 : \dots : X_{m+1})$ be homogeneous coordinates of \mathbb{P}^{m+1} such that $(X_1/X_0, \dots, X_{m+1}/X_0) = (z_1, \dots, z_{m+1}) \in \mathbb{C}^{m+1}$ and \bar{S} be the closure of S in \mathbb{P}^{m+1} . Then we have the following theorem of Zariski :

Theorem 2(Zariski [5]).

Suppose that $p_\infty = (0 : \dots : 0 : 1)$ is not contained in \bar{S} . Then

$$\begin{aligned} & \pi_1(\mathbb{P}^{m+1} - \bar{S}) \\ & \cong \langle \gamma_1, \dots, \gamma_n \mid \gamma_n \gamma_{n-1} \dots \gamma_1 = 1, \gamma_j = \varphi(\theta(\beta_k))(\gamma_j), (1 \leq j \leq n, 1 \leq k \leq t) \rangle. \end{aligned}$$

Proof. Let H_∞ be the hypersurface of \mathbb{P}^{m+1} defined by $H_\infty = \{X_0 = 0\}$, (i.e. hyperplane at infinity) and α be a meridian of H_∞ in $\mathbb{P}^{m+1} - \bar{S} - H_\infty$ (see Figure 6).




Figure 6

From the theorem of Van Kampen [4], we have the following exact sequence :

$$1 \rightarrow \langle \alpha \rangle \cong \pi_1(\mathbb{C}^{m+1} - S) \rightarrow \pi_1(\mathbb{C}^{m+1} - S) \rightarrow \pi_1(\mathbb{P}^{m+1} - \bar{S}) \rightarrow 1 \text{ (exact).}$$

We can take α as $(\gamma_n \gamma_{n-1} \dots \gamma_1)^{-1}$ in $\mathbb{C}^{m+1} - S$ (see Figure 7).

Thus,

$$\pi_1(\mathbb{P}^{m+1} - \bar{S}) \cong \pi_1(\mathbb{C}^{m+1} - S) / \langle \gamma_n \gamma_{n-1} \dots \gamma_1 \rangle \cong \pi_1(\mathbb{C}^{m+1} - S).$$

This shows Theorem 2.




Figure 7

REMARK. A similar theorem to Theorem 1 holds for $\mu: \mathbf{B}^m(\varepsilon) \times \mathbf{B}^1(\varepsilon') \rightarrow \mathbf{B}^m(\varepsilon)$, where $\mathbf{B}^m(\varepsilon)$ is a m -dimensional complex ball; $\mathbf{B}^m(\varepsilon) = \{(z_1, \dots, z_m) \in \mathbf{C}^m \mid |z_1|^2 + \dots + |z_m|^2 < \varepsilon^2\}$. In this case, the existence of continuous section with a similar conditions to Theorem 1 is obvious.

6. Calculations of Fundamental Groups of Finite Branched Coverings

EXAMPLE 1.

Let X be the surface in \mathbf{C}^3 defined by

$$X = \{(\lambda, x, y) \in \mathbf{C}^3 \mid y^2 = x(x-1)(x-\lambda)\}.$$

X has two isolated singular points at $(0,0,0)$ and $(1,1,0)$. Hence X is normal. Let $\pi: X \rightarrow \mathbf{C}^2$ be the projection map defined by

$$\pi(\lambda, x, y) = (\lambda, x).$$

Then π is a double branched covering of \mathbf{C}^2 . The branch locus S of π is a curve in \mathbf{C}^2 and is written as:

$$S = \{(\lambda, x) \in \mathbf{C}^2 \mid x(x-1)(x-\lambda) = 0\}.$$

According to Theorem 1, we can calculate $\pi_1(\mathbf{C}^2 - S)$. Let $\mu: \mathbf{C}^2 \rightarrow \mathbf{C}$ be the trivial line bundle on \mathbf{C} defined by

$$\mu(\lambda, x) = \lambda.$$

The branch locus B of μ is $\{0, 1\} \subset \mathbf{C}$ and $\pi_1(\mathbf{C} - B)$ is isomorphic to the free group $\langle \beta_1, \beta_2 \rangle$, where β_1 and β_2 are its generators and can be considered as the




Figure 8

meridians of $\{0\}$ and $\{1\}$, respectively. We may take $q_0 = \frac{1}{2}$ as a reference point of $\pi_1(C-B)$. In this case the standard fiber F of $\mu|_{C-S-\mu^{-1}(s)}$ is $C - \{3\text{-points}\}$. We define γ_1, γ_2 and γ_3 as the meridians of $(\frac{1}{2}, 1), (\frac{1}{2}, \frac{1}{2})$ and $(\frac{1}{2}, 0)$, respectively. The image of β_1 and β_2 by $\theta : \pi_1(C-B) \rightarrow B_3$ are σ_1^2 and σ_2^2 , respectively. Then we have

$$\begin{aligned} \pi_1(C^2 - S) &\cong \langle \gamma_1, \gamma_2, \gamma_3 \mid \gamma_j = \varphi(\theta(\beta_k))(\gamma_j), j=1, 2, 3, k=1, 2 \rangle \\ &\cong \langle \gamma_1, \gamma_2, \gamma_3 \mid \gamma_2\gamma_3 = \gamma_3\gamma_2, \gamma_1\gamma_2 = \gamma_2\gamma_1 \rangle. \end{aligned}$$

By using the Reidemeister-Schreier method (c.f. Rolfsen [3] P.315-P.316), we can calculate $\pi_1(\text{Reg}X)$, where $\text{Reg}X$ is the set of regular points of X . Since $\pi_1(C^2 - B)$ is generated by three elements and since π is a double branched covering, we take the 3-th free group F_3 and the 5-th free group F_5 . As in Figure 9, we take their generators $\{\gamma_1, \gamma_2, \gamma_3\}$ and $\{b_1, b_2, b_3, b_4, b_5\}$, respectively, where $\pi^{-1}(\gamma_1) = \{x_1, x_2\}$, $\pi^{-1}(\gamma_2) = \{y_1, y_2\}$, $\pi^{-1}(\gamma_3) = \{z_1, z_2\}$, and

$$\begin{aligned} b_1 &= y_1x_1^{-1} \\ b_2 &= x_1y_2x_2^{-1}x_1^{-1} \\ b_3 &= z_1x_1^{-1} \\ b_4 &= x_1z_2x_2^{-1}x_1^{-1} \\ b_5 &= x_1x_2. \end{aligned}$$

Then we transfer the relation of $\pi_1(C^2 - S)\gamma_2\gamma_3\gamma_2^{-1}\gamma_3^{-1} = 1$ in the words of F_5 :

$$y_1z_2y_2^{-1}z_1^{-1} = b_1b_4b_5b_5^{-1}b_2^{-1}b_3^{-1} = 1$$




Figure 9

$$x_1 y_2 z_1 y_1^{-1} z_2^{-1} x_1^{-1} = b_2 b_5 b_3 b_1^{-1} b_5^{-1} b_4^{-1} = 1.$$

In a similar way, we transfer the relation $\gamma_1 \gamma_2 \gamma_1^{-1} \gamma_2^{-1} = 1$ in the words of F_5 :

$$\begin{aligned} x_1 y_2 x_2^{-1} y_1^{-1} &= b_2 b_5 b_5^{-1} b_1^{-1} = 1 \\ x_1 x_2 y_1 x_1^{-1} y_2^{-1} x_1^{-1} &= b_5 b_1 b_5^{-1} b_2^{-1} = 1. \end{aligned}$$

We also transfer the relations $\gamma_1^2 = 1, \gamma_2^2 = 1, \gamma_3^2 = 1$ since the ramification index of each irreducible component of S is equal to 2 :

$$\begin{aligned} x_1 x_2 &= b_5 = 1 \\ y_1 y_2 &= b_1 b_2 b_5 = 1 \\ z_1 z_2 &= b_3 b_4 b_5 = 1. \end{aligned}$$

Putting $\alpha_1 = b_1$ and $\alpha_2 = b_4$, we have

$$\begin{aligned} \pi_1(\text{Reg } X) &\cong \langle \alpha_1, \alpha_2 \mid \alpha_1^2 = 1, (\alpha_1 \alpha_2)^2 = 1 \rangle \\ &\cong (\mathbb{Z}/2\mathbb{Z}) * (\mathbb{Z}/2\mathbb{Z}) \text{ (free product)}. \end{aligned}$$

EXAMPLE 2.

Let X be the hypersurfaces of C^4 defined by

$$X = \{(x, y, z, w) \in C^4 \mid w^n = z^2 - xy^2\} (n \geq 2).$$

The singular locus of X is the line $\{(x, y, z, w) \in X \mid y = z = w = 0\}$. Hence X is normal. Let $\pi : X \rightarrow C^3$ be the projection map defined by :

$$\pi(x, y, z, w) = (x, y, z).$$

Then π is a cyclic branched covering of C^3 . The branch locus S of π is a surface in C^3 and is written as :

$$S = \{(x, y, z) \in \mathbb{C}^3 \mid z^2 - xy^2 = 0\} \text{ (the Cartan umbrella).}$$

According to Theorem 1, we can calculate $\pi_1(\mathbb{C}^3 - S)$. The result is

$$\pi_1(\mathbb{C}^3 - S) \cong \langle \gamma \rangle \text{ (the free group).}$$

From the Reidemeister-Schreier method again, we have :

$$\pi_1(\text{Reg}X) \cong \{i.e. \text{ Reg}X \text{ is simply connected}\}.$$

EXAMPLE 3.

Let X be the hypersurface of \mathbb{C}^{m+2} defined by

$$X = \{(z_1, \dots, z_{m+2}) \in \mathbb{C}^{m+2} \mid z_{m+2}^2 + z_{m+1}^2 + g(z_1, \dots, z_m) = 0\},$$

where g is a polynomial which is not constant. The singular locus of X is at most $(m-1)$ -dimensional. Hence X is normal. Let $\pi : X \rightarrow \mathbb{C}^{m+1}$ be the projection map defined by :

$$\pi(z_1, \dots, z_{m+1}, z_{m+2}) = (z_1, \dots, z_{m+1}).$$

Then π is a branched covering of \mathbb{C}^{m+1} . The branch locus S of π is a hypersurface in \mathbb{C}^{m+1} and is written as :

$$S = \{z_{m+1}^2 + g(z_1, \dots, z_m) = 0\}.$$

By Theorem 1, $\pi_1(\mathbb{C}^{m+1} - S)$ can be written as :

$$\pi_1(\mathbb{C}^{m+1} - S) \cong \langle \gamma_1, \gamma_2 \mid \square = 1, \dots, \square = 1 \rangle.$$

From the Reidemeister-Schreier method again, we have :

$$\pi_1(\text{Reg}X) \cong \begin{cases} \{1\} \text{ or} \\ \mathbb{Z}/q\mathbb{Z} (\exists q \in \mathbb{Z}) \text{ or} \\ \mathbb{Z} \end{cases}$$

(i.e. $\pi_1(\text{Reg}X)$ is isomorphic to a cyclic group).

References

- [1] T. Matsuno : *Normal singularities and fundamental groups*, Osaka Univ. master theses (in Japanese).
- [2] M. Namba : *Branched coverings and algebraic functions*, Pitman Res. Notes Math. Ser., Vol.161, Longman Sci. Tech., Harlow, 1987.
- [3] D. Rolfsen : *Knots and links*, Publish or Perish, Berkeley, 1976.
- [4] van Kampen : *On the fundamental group of an algebraic curves*, Amer.J.Math. **55** (1933), 255-260.
- [5] O. Zariski : *Collected Papers* Vol 3, MIT Press, Cambridge, 1973.

Department of Mathematics
Faculty of Science
Osaka University
Toyonaka, Osaka 560, Japan