

Title	On the boundary behavior of harmonic morphisms at the boundary of compactifications of Martin type
Author(s)	Ikegami, Teruo
Citation	Osaka Journal of Mathematics. 25(4) P.729-P.736
Issue Date	1988
Text Version	publisher
URL	<a href="https://doi.org/10.18910/8031">https://doi.org/10.18910/8031</a>
DOI	10.18910/8031
rights	
Note	

*Osaka University Knowledge Archive : OUKA*

<https://ir.library.osaka-u.ac.jp/>

Osaka University

Ikegami, T.  
Osaka J. Math.  
25 (1988), 729-736

## ON THE BOUNDARY BEHAVIOR OF HARMONIC MORPHISMS AT THE BOUNDARY OF COMPACTIFICATIONS OF MARTIN TYPE

TERUO IKEGAMI

(Received September 25, 1987)

### Introduction

The boundary behavior of analytic functions on the unit disc has been an interesting material of the complex analysis. Among various investigations on this problem some results such as the theorem of Fatou-Plessner and the theorem of Riesz-Frostman-Nevalinna were generalized in a far extensive context, that is, for analytic mappings of Riemann surfaces and their relevant compactifications. These were found in the book of Constantinescu-Cornea [1].

Later [2], they considered harmonic morphisms between BreLOT's harmonic spaces under Wiener's compactification and revealed that above mentioned theorems are based essentially on the potential theoretic character. By using relations between the Wiener and the Martin compactifications of BreLOT's harmonic spaces with additional assumptions, the author obtained informations on the boundary behavior of harmonic morphisms at the Martin boundary [5], [6].

Under the framework of Constantinescu-Cornea [3], K. Oja [10] generalized theorems of Fatou-Plessner and of Riesz. Recently [7], in order to discuss the Naïm theory [9] in the context of harmonic spaces of Constantinescu-Cornea the author defined the compactifications of Martin type. Here we give a supplementary remark on mappings of type  $B1$  and translate the informations on the Wiener boundary into those on the Martin type compactification following the spirit of [6]. It is still possible to discuss the problem without any compactification as in [11], [12].

Acknowledgment. The author wishes his herty thanks to Prof. F-Y. Maeda for the valuable comments.

### 1. Preliminaries.

Let  $X$  be a  $\mathcal{P}$ -harmonic space in the sense of Constantinescu-Cornea [3]. We assume that  $X$  has a countzble base and 1 is superharmonic on  $X$ . Further, we assume that  $X$  has a compactification of Martin type  $X^*$  [7], i.e., (1)  $X^*$  is

metrizable and resolutive, (2) there is a finite continuous function  $K(x, z)$  defined on  $X \times \Delta (\Delta = X^* \setminus X)$  such that  $k_z: x \rightarrow k(x, z)$  is non-negative and harmonic on  $X$  for every  $z \in \Delta$ , (3) there is a non-negative Borel measure  $\mu$  on  $\Delta$  and a set  $\Delta_1$  corresponding to a certain class of minimal harmonic functions  $k_z$  such that  $\mu(\Delta \setminus \Delta_1) = 0$  and  $\mu(T) = 0$  if  $T$  is negligible, (4) for every  $u \in \text{HB}(X)$  there is a resolutive function  $f$  such that  $u(x) = H_f(x) = \int k(x, z) f(z) d\mu(z)$  for every  $x \in X$ . Then,  $X$  has the Wiener compactification  $X^W$  which is resolutive [4] and  $X^*$  is a quotient of  $X^W$ , i.e., there is a continuous surjection  $\pi: X^W \rightarrow X^*$  satisfying  $\pi(x) = x$  for every  $x \in X$ .

Let  $X'$  be a second harmonic space such that  $X'$  has a countable base and 1 is superharmonic on  $X'$ . We suppose further that  $X'$  is *connected* and *elliptic* if  $X' \notin \mathcal{P}$ , i.e., if  $X'$  is not  $\mathcal{P}$ -harmonic. For a compactification  $X'^*$  of  $X'$  we make the following assumption:

Case (i)  $X'^* = X'$  if  $X'$  is compact,

Case (ii)  $X'^*$  is an arbitrary metrizable compactification if  $X'$  is non-compact and  $X' \notin \mathcal{P}$ ,

Case (iii)  $X'^*$  is an arbitrary metrizable and resolutive compactification if  $X' \in \mathcal{P}$ .

*Remarks.* (1) The case (iii) occurs since 1 is harmonizable [4], and  $X'$  is non-compact if  $X' \in \mathcal{P}$ .

(2) If  $X' \notin \mathcal{P}$ , then  $X'$  is completely degenerated, i.e., 0 is the only one potential on  $X'$  and all non-negative superharmonic functions are harmonic and they are proportional ([3], Exercise 3.1.10).

(3) If  $F' \subset X'$  is closed and non-polar then  $X' \setminus F' \in \mathcal{P}$  ([3], Exercise 6.2.5).

### Proposition 1.

Let  $f' \in \mathcal{C}(X'^*)$ ,  $F'$  be a compact non-polar subset of  $X'$  with non-empty interior and  $G' = X' \setminus F'$ . Then the restriction of  $f'$  to  $G'$  is harmonizable on  $G'$ .

*Proof.* Case (i):  $G'$  is relatively compact and is resolutive since, in virtue of the above Remark (3),  $G'$  is an open subset of a  $\mathcal{P}$ -harmonic space ([3], Th. 2.4.2). Hence by the same argument as in [8], 2.1.4,  $h_{f', G'}^{G', X'} = H_{f', X'}^{G', X'}$ . In the following we refer to [8] without mentioning any possible modification.

Case (ii): We show first that  $h_1^{G', X'} = 0$ . This function  $h_1^{G', X'}$  is associated with 0 on  $\partial G'$  ([8], 2.1.5). Let  $p'_0$  be an Evans function on  $G'$  of some potential  $p' > 0$  on  $G'$ . Then,  $\liminf [p'_0 - h_1^{G', X'}] \geq 0$  on  $\partial G'$  ([8] 1.2.14). The function

$$v' = \begin{cases} \inf (p'_0 - h_1^{G', X'}, 0) & \text{on } G' \\ 0 & \text{on } F' \end{cases}$$

is superharmonic on  $X'$ . Since, by Remark (2), the non-negative superharmonic function  $1+v'$  on  $X'$  is harmonic and is proportional to 1, we have  $v'=0$ , which implies  $h_1^{G',X'}=0$ , since  $p'_0-h_1^{G',X'}\geq 0$ . Considering  $f'^+$  and  $f'^-$  separately, we obtain  $h_{f'}^{G',X'}=0$ . Thus,  $\bar{h}_{f'}^{G',G'}=\bar{h}_{f'}^{G',X'}+H_{f'}^{G',X'}=H_{f'}^{G',X'}$  ([8], 2.1.4) and  $\underline{h}_{f'}^{G',G'}=-\bar{h}_{(-f')}^{G',G'}=-H_{(-f')}^{G',X'}=H_{f'}^{G',X'}$ .

Case (iii): Since  $f'$  is harmonizable on  $X'$  ([8], 3.2.9) it is harmonizable on  $G'$  ([8], 2.2.3), q.e.d..

Let  $\varphi: X\rightarrow X'$  be a harmonic morphism [10], i.e.,  $\varphi: X\rightarrow X'$  is continuous and for every open subset  $U'$  of  $X'$  with  $\varphi^{-1}(U')\neq\emptyset$  and for every hyperharmonic function  $u'$  on  $U'$ ,  $u'\circ\varphi$  is hyperharmonic on  $\varphi^{-1}(U')$ .

$\varphi$  is called *locally polarly non-constant* [10] if there is no open subset  $U$  of  $X$  such that  $\varphi(U)$  is a singleton and polar.

$\varphi$  is of *type Bl* at  $x'\in X'$  if there is an open neighborhood  $U'$  of  $x'$  such that  $\varphi^{-1}(U')=\emptyset$  or  $h_1^{\varphi^{-1}(U'),X}=0$ . A harmonic morphism which is of type *Bl* at each point of  $X'$  is called *of type Bl*.

## 2. The boundary behavior of harmonic morphisms at Wiener boundary

We consider the Wiener compactification  $X^W$  of  $X$ . For the definition and properties of  $X^W$ , one may refer to [4]. The following results are obtained, except for the theorem on mappings of type *Bl*, by K. Oja and we quote them with brief proof.

Let  $\varphi: X\rightarrow X'$  be a harmonic morphism. We define, for  $\tilde{x}\in\Delta^W=X^W\setminus X$ , the cluster set

$$\varphi^*(\tilde{x}) = \cap \overline{\{\varphi(U^* \cap X)\}}; U^* \text{ is a neighborhood of } \tilde{x} \text{ in } X^W,$$

where the closure is taken in the compact space  $X'^*$ .

### Theorem of Fatou-Plessner.

$\varphi^*(\tilde{x})$  is either  $X'^*$  or a singleton, and only the latter case occurs if  $X'\in\mathcal{P}$ .

For the case (i) and the case (ii), in view of Prop. 1, we may use a result of K. Oja, [10], Lemma 3.1, and in the case (iii)  $f'\circ\varphi$  is a Wiener function on  $X$  for every Wiener function  $f'$  on  $X'$  ([10], Lemma 4.7), and every Wiener function on  $X$  can be extended continuously to  $X^W$ .

Putting

$$\begin{aligned} \tilde{P} &= \{\tilde{x}\in\Delta^W; \varphi^*(\tilde{x}) = X'^*\}, \\ \tilde{F} &= \{\tilde{x}\in\Delta^W; \varphi^*(\tilde{x}) \text{ is a singleton}\} \end{aligned}$$

we can rewrite above theorem as

$$\Delta^W = \tilde{P} \cup \tilde{F} \text{ and } \Delta^W = \tilde{F} \text{ provided } X'\in\mathcal{P}.$$

Further, we have  $\Delta^w \setminus \Gamma^w \subset \tilde{F}$ , where  $\Gamma^w$  is the harmonic boundary of  $X^w$  ([10], Cor. 6.14).

In the same way as [2], Th. 6.3, it is shown that  $\tilde{P}$  is an open and closed subset of  $\Gamma^w$ .

We call a set  $A' \subset X'^*$  is *polar in  $X'^*$*  if  $A' \cap X'$  is polar and if for any open subset  $G' \subset X'$  with  $G' \in \mathcal{P}$  and for every  $a'_0 \in G'$  there is a non-negative hyperharmonic function  $u'$  on  $G'$  such that  $u'(a'_0) < +\infty$  and

$$\lim_{\substack{a' \rightarrow x' \\ a' \in G'}} u'(a') = +\infty \quad \text{for all } x' \in \overline{G'} \cap A' \cap \Delta'.$$

**Theorem of Riesz.**

Let  $\varphi$  be locally polarly non-constant, and  $\tilde{A}$  be a subset of  $\Delta^w$ . Then, if  $A' = \bigcup_{\tilde{x} \in \tilde{A}} \varphi^*(\tilde{x})$  is polar in  $X'^*$  then  $\bar{H}_{1\tilde{A}}^w = 0$ .

Proof. Let  $K'_1$  and  $K'_2$  be disjoint compact subsets of  $X'$  with non-empty interiors and  $G'_i = X' \setminus K'_i$  ( $i = 1, 2$ ). Then, there is a non-negative hyperharmonic functions  $u'_i$  on  $G'_i$  such that  $u'_i$  is finite on a dense subset of  $G'_i$  and  $\lim_{\substack{a' \rightarrow x' \\ a' \in G'}} u'_i(a') = +\infty$  for every  $x' \in \overline{G'_i} \cap A'$  ([10], Lemma 6.9, Th. 6.11 and [3],

Excercise 3.1.11). Putting  $u_i = u'_i \circ \varphi$ ,  $G_i = \varphi^{-1}(G'_i)$  and  $\tilde{A}_i = \{\tilde{x} \in \Delta^w \setminus (\overline{X \setminus G_i}) ; \lim_{\substack{a \rightarrow \tilde{x} \\ a \in G_i}} u_i(a) = +\infty\}$  we have  $\tilde{A} \subset \tilde{A}_1 \cup \tilde{A}_2$  and  $\bar{H}_{1\tilde{A}_i}^w = 0$ . The last assertion is

derived if we prove the following:

(1)  $\hat{R}_u^{X \setminus G_i} = u$ , where  $u = \bar{H}_{1\tilde{A}_i}^w$ ,

and

(2)  $u \leq \hat{R}_{\frac{X \setminus G_i}{H^w}} = q$  and  $q$  is a potential on  $X$ , where  $f = 1_{\Delta^w \setminus (\overline{X \setminus G_i})}$ .

Let  $v_1$  be a non-negative hyperharmonic function on  $X$  and  $v_1 \geq 1$  on  $X \setminus G_i$ . The function

$$v_2 = \begin{cases} 1 & \text{on } X \setminus G_i \\ \inf(1, v_1 + \varepsilon u_i) & \text{on } G_i \end{cases}$$

satisfies  $v_2 \geq u$  for every  $\varepsilon > 0$ . In particular,  $v_1 + \varepsilon u_i \geq u$  on  $G_i$ . Since  $P = \{x \in G_i : u_i(x) = \infty\} = \varphi^{-1}(\{x' \in G'_i : u'_i(x') = \infty\})$  is polar,  $v_1 \geq u$  on  $G_i$ . Hence  $v_1 \geq u$ , for  $v_1 \geq 1 \geq u$  on  $X \setminus G_i$ . Now, let  $v$  be a non-negative hyperharmonic function on  $X$  satisfying  $v \geq u$  on  $X \setminus G_i$ . Then  $v_1 + v - u$  is non-negative and hyperharmonic on  $X$  and  $v_1 + v - u \geq 1$  on  $X \setminus G_i$ , which means  $v_1 + v - u \geq R_1^{X \setminus G_i}$  and therefore  $R_1^{X \setminus G_i} + R_u^{X \setminus G_i} - u \geq R_1^{X \setminus G_i}$ . Thus we have  $R_u^{X \setminus G_i} \geq u$ , which completes the proof of (1).

To prove (2), let  $\vartheta$  be a non-negative hyperharmonic function on  $X$  satisfying  $\liminf \vartheta \geq \alpha > 1$  at every point of  $\Delta^w \cap (\overline{X \setminus G_i})$ . Then, there exists an open subset  $V^*$  of  $X^w$  such that  $V^* \supset \Delta^w \cap (\overline{X \setminus G_i})$  and  $\vartheta \geq 1$  on  $V = V^* \cap X$ . Letting

$K=(X\setminus G_i)\setminus V$ , we have  $\hat{R}_1^K + \vartheta \geq \hat{R}_1^{X/G_i} \geq \hat{R}_{H^W}^{X/G_i} = q \geq h_q^{X,X}$ , hence  $\hat{R}_1^K + \alpha H_g^W \geq h_q^{X,X}$ , where  $g = 1_{\Delta^W \cap (\overline{X \setminus G_i})}$ . Since  $K$  is compact and  $\alpha (>1)$  is arbitrary, we have  $h_q^{X,X} \leq H_g^W$ , therefore  $h_q^{X,X} \leq H_{\min(f,g)}^W = 0$ , i.e.,  $q$  is a potential.

The following results on mappings of type *Bl* are obtained quite in the same way as in [2].

**Proposition 2.**

$$\{\varphi^*(\tilde{x}); \tilde{x} \in \Gamma^W \setminus \tilde{P}\} \cap X' = \{x' \in X'; \varphi \text{ is not of type Bl at } x'\}.$$

Proof. Let  $\tilde{x} \in \Gamma^W \setminus \tilde{P}$  and  $x' = \varphi^*(\tilde{x}) \in X'$ . For any open neighborhood  $U'$  of  $x'$ , since  $U = \varphi^{-1}(U') \neq \emptyset$ , we have to show that  $\bar{h}_1^{U',X} \neq 0$ . Suppose for a moment that  $\bar{h}_1^{U',X} = 0$ . If  $(\bar{U} \setminus U) \cap X \neq \emptyset$ ,  $1 = \bar{h}_1^{U',X} + H_1^{U',X}$  implies  $1 = H_1^{U',X} = R_1^{X \setminus U}$  on  $U$ , and hence  $R_1^{X \setminus U} = 1$  in  $X$ . Select a neighborhood  $V^*$  of  $\tilde{x}$  such that  $V^* \cap \overline{X \setminus U} = \emptyset$ . This is possible since  $\tilde{x} \in \overline{X \setminus U}$ . Then  $p = \inf \{\hat{R}_1^{V^* \cap X}, \hat{R}_1^{X \setminus U}\}$  is a potential ([8], 3.2.23) and  $0 = \liminf_{\tilde{x}} p = \liminf_{\tilde{x}} \hat{R}_1^{V^* \cap X} = 1$  which is a contradiction. If  $(\bar{U} \setminus U) \cap X = \emptyset$ , we have  $1 = \bar{h}_1^{U',X}$  and this contradicts our assumption.

Next, we prove that  $\varphi$  is of type *Bl* at each point  $x' \in \varphi^*(\Gamma^W \setminus \tilde{P}) \cap X'$ . Since  $\varphi^*(\Gamma^W \setminus \tilde{P}) \cap X'$  is a closed subset of  $X'$ , we can find an open neighborhood  $U'$  of  $x'$  such that  $\bar{U}' \cap \varphi^*(\Gamma^W \setminus \tilde{P}) = \emptyset$ . We may suppose that  $U'$  is relatively compact and  $U' \neq X'$ . Then we have  $\Gamma^W \subset \overline{X \setminus \varphi^{-1}(U')}$  and hence  $R_1^{X \setminus \varphi^{-1}(U')} = 1$ , which shows as above that  $\bar{h}_1^{\varphi^{-1}(U'),X} = 0$ .

**Theorem (Characterization of mappings of type Bl)**

- (1) When  $X' \notin P$ ,  $\varphi$  is of type *Bl* if and only if  $\Gamma^W = \tilde{P}$ .
- (2) When  $X' \in P$ ,  $\varphi$  is of type *Bl* if and only if  $\varphi^*(\Gamma^W) \cap X' = \emptyset$ .

Proof. (1): If  $\varphi$  is of type *Bl*, by Prop. 2,  $\varphi^*(\Gamma^W \setminus \tilde{P}) \cap X' = \emptyset$ , which implies  $\Gamma^W = \tilde{P}$  whenever  $X'$  is compact. When  $X'$  is non-compact, we can deduce that  $\Gamma^W \setminus \tilde{P}$  is of harmonic measure zero and hence  $\Gamma^W \setminus \tilde{P} = \emptyset$  since  $\tilde{P}$  is open and closed in  $\Gamma^W$ . The converse is trivial.

(2): In this case,  $\tilde{P} = \emptyset$  and  $\varphi$  is extended continuously on  $X^W$ . Hence the assertion is an immediate consequence of Prop. 2.

REMARK. If  $X' \in \mathcal{P}$  and  $\varphi$  is of type *Bl* then we can show  $\varphi^*(\Gamma^W) = \Gamma'$  ([2] Cor. 6.2, [8] 3.1.7).

**3. The fine boundary behavior of harmonic morphisms**

Let  $\varphi$  be a harmonic morphism between  $X$  and  $X'$  and  $X^*$  be a compactification of Martin type. In [7], we have shown that each point  $z \in \Delta_1$  has the fine filter  $\mathcal{G}_z = \{E \subset X; R_{k_z}^{X,E} \neq k_z\}$  and the set

$$\Delta_2 = \{z \in \Delta_1; \mathcal{G}_z \text{ converges to } z\}$$

contains  $\mu$ -almost all boundary points, i.e.,  $\mu(\Delta \setminus \Delta_2) = 0$ . For  $z \in \Delta_2$  we define

$$\hat{\phi}(z) = \cap \{\overline{\varphi(E)}; E \in \mathcal{Q}_z\},$$

where the closure is taken in  $X'^*$ . Let

$$\begin{aligned} \hat{P} &= \{z \in \Delta_2; \hat{\phi}(z) = X'^*\}, \\ \hat{F} &= \{z \in \Delta_2; \hat{\phi}(z) \text{ is a singleton}\}. \end{aligned}$$

Then as in [5], [6] we can prove the following key theorem which reveals the relation between  $\hat{\phi}$  and  $\varphi^*$ :

**Key Theorem.**

*Let*

$$N = (\Delta \setminus \Delta_2) \cup [\Delta_2 \setminus (\hat{F} \cup \hat{P})]$$

*and*

$$\tilde{N} = (\Delta^w \setminus \Gamma^w) \cup [\pi^{-1}(\Delta \setminus \hat{P}) \cap \tilde{P}] \cup (\tilde{F} \setminus \tilde{F}_1),$$

where  $\tilde{F}_1 = \{\tilde{x} \in \tilde{F}; \pi(\tilde{x}) \in \Delta_2, \varphi^*(\tilde{x}) = \hat{\phi}(\pi(\tilde{x}))\}$ .

*Then,  $\mu(N) = 0$  and  $\tilde{N}$  is of harmonic measure zero and*

$$\hat{\phi}(z) = \varphi^*(\tilde{x}) \text{ if } z \in \Delta \setminus N \text{ and } \tilde{x} \in (\pi^{-1}(z) \cap \Gamma^w) \setminus \tilde{N}.$$

The above theorem makes it possible for us to translate informations about  $\varphi^*$  into those of  $\hat{\phi}$ .

**Theorem of Fatou-Plessner.**

$$\hat{\phi}(z) = X'^* \text{ or a singleton for } \mu\text{-almost all } z.$$

This is immediate from the definition of  $N$  in the key theorem.

**Theorem of Riesz.**

*Suppose  $\varphi$  is locally polarly non-constant and let  $A$  be a  $\mu$ -measurable subset of  $\Delta_2$ . If  $\cup_{z \in A} \hat{\phi}(z)$  is polar in  $X'^*$ , then  $A$  is polar.*

In fact, using notations in the key theorem and putting

$$\tilde{A} = [\pi^{-1}(A \setminus N) \cap \Gamma^w] \setminus \tilde{N}$$

we have

$$\cup_{\tilde{x} \in \tilde{A}} \varphi^*(\tilde{x}) \subset \cup_{z \in A} \hat{\phi}(z),$$

which implies that  $\tilde{A}$  is of harmonic measure zero, i.e.,  $\bar{H}_{1_{\tilde{A}}}^w = 0$ . Since  $\pi^{-1}(A) \cap \Gamma^w \subset \tilde{A} \cup \pi^{-1}(N) \cup \tilde{N}$ , we have  $H_{1_{\pi^{-1}(A)}}^w = 0$ , and hence  $H_{1_A} = H_{1_{\pi^{-1}(A)}}^w = 0$ . Thus  $A$  is polar ([10], Cor. 6.6).

**Theorem on the boundary characterization of mappings of type *Bl*.**

- (1) When  $X' \notin \mathcal{P}$ ,  $\varphi$  is of type *Bl* if and only if  $\mu(\tilde{F})=0$ .
- (2) When  $X' \in \mathcal{P}$ ,  $\varphi$  is of type *Bl* if and only if  $\phi(z) \cap X' = \emptyset$  for  $\mu$ -almost all  $z$ .

Proof. (1): If  $\varphi$  is of type *Bl*, then  $\tilde{P} = \Gamma^w$  so that  $\mu(\tilde{F})=0$ , by the key theorem. Conversely, if  $\varphi$  is not of type *Bl*, then  $\Gamma^w \cap \tilde{F} \neq \emptyset$ . Since  $\Gamma^w \cap \tilde{F}$  is an open subset of  $\Gamma^w$ , it follows that it has a positive harmonic measure at some point of  $X$ . Hence  $\mu(\pi(\Gamma^w \cap \tilde{F})) > 0$ , which implies  $\mu(\tilde{F}) > 0$  by the key theorem.

(2): Since  $\tilde{P} = \emptyset$  in this case  $\mu(\Delta \setminus \pi(\tilde{F}_1)) = 0$  by the key theorem. Since  $\phi(z) \in \varphi^*(\Gamma^w)$  for  $z \in \pi(\tilde{F}_1)$ , it follows that  $\varphi^*(\Gamma^w) \cap X' = \emptyset$  is equivalent to  $\phi(z) \cap X' = \emptyset$  for  $\mu$ -almost all  $z$  (cf. the proof of [6], Th. 5.3).

REMARK. It is readily seen that if  $X' \in \mathcal{P}$  and if  $\varphi$  is of type *Bl* then  $\Gamma \setminus \bigcup_{z \in \tilde{F}} \phi(z)$  is of harmonic measure zero.

**References**

- [1] C. Constantinescu-A. Cornea: Ideale Ränder Riemannscher Flächen, Ergebnisse der Math. ihrer Grenzgebiete, Neue Folge Bd. 32, Springer 1963.
- [2] C. Constantinescu-A. Cornea: Compactifications of harmonic spaces, Nagoya Math. J. **25** (1965) 1–57.
- [3] C. Constantinescu-A. Cornea: Potential theory of harmonic spaces, Berlin-Heidelberg-New York, Springer 1972.
- [4] J. Hyvönen: On resolute compactifications of harmonic spaces, Ann. Acad. Sci. Fenn. Dissertations 1976.
- [5] T. Ikegami: On the boundary behavior of harmonic maps, Osaka J. Math. **10** (1973) 641–653.
- [6] T. Ikegami: The boundary behavior of analytic mappings of Riemann surfaces, Complex Analysis Joensuu 1978, Lecture Notes in Math, 747, Springer 1979.
- [7] T. Ikegami: Compactifications of Martin type of harmonic spaces, Osaka J. Math. **23** (1986) 653–680.
- [8] C. Meghea: Compactification des espaces harmoniques, Lecture Notes in Mat. 222, Springer. 1971.
- [9] L. Naïm: Sur le rôle de la frontière de R.S. Martin dans la théorie du potentiel, Ann. Inst. Fourier, Grenoble **7** (1957) 183–281.
- [10] K. Oja: On cluster sets of harmonic morphisms between harmonic spaces, Ann. Acad. Sci. Fenn. Dissertations 1979.
- [11] K. Oja: Theorems of the Riesz type for co-fine cluster sets of harmonic morphisms, Ann. Acad. Sci. Fenn. **6** (1981) 77–88.
- [12] D. Sibony: Allure à la frontière minimale d’une classe de transformations. Théorème de Doob généralise, Ann. Inst. Fourier, Grenoble **18** (1968) 91–120.



Department of Mathematics  
Osaka City University  
Sugimoto 3-3-138  
Osaka 558, Japan