



Title	On the Borsuk-Ulam theorem for $Z_{p^a}$ actions on $S^{2n-1}$ and maps $S^{2n-1} \rightarrow R^m$
Author(s)	Munkholm, Hans J.
Citation	Osaka Journal of Mathematics. 1970, 7(2), p. 451-456
Version Type	VoR
URL	<a href="https://doi.org/10.18910/8171">https://doi.org/10.18910/8171</a>
rights	
Note	

*The University of Osaka Institutional Knowledge Archive : OUKA*

<https://ir.library.osaka-u.ac.jp/>

The University of Osaka

Munkholm, H. J.  
 Osaka, J. Math.  
 7 (1970), 451-456

## ON THE BORSUK-ULAM THEOREM FOR $Z_{p^a}$ ACTIONS ON $S^{2n-1}$ AND MAPS $S^{2n-1} \rightarrow R^m$

HANS J. MUNKHOLM

(Received April 13, 1970)

### 1. Introduction

In [3] we raised the following question:

Let  $G$  be a finite group acting properly (as a group of *homeomorphisms*) on the  $n$ -sphere. For any (topological)  $m$ -manifold  $M$  and any map  $f: S^n \rightarrow M$  let  $A(f) = \{x \in S^n \mid f(x) = f(xg), \text{ all } g \in G\}$ . What can be deduced about  $\dim A(f)$ ?

In case  $M = R^m$ , euclidean  $m$ -space,  $A(f)$  is the set of solutions of  $(|G| - 1)m + 1$  equations in  $n + 1$  unknowns so one might hope to get

$$(1.1) \quad \dim A(f) \geq n - (|G| - 1)m.$$

If  $G$  is cyclic of prime order then (1.1) actually holds even for maps  $f: S^n \rightarrow M^m$  provided  $M^m$  is compact (for  $G = Z_2$  and  $m = n$  assume also that  $f_* = 0: H_n(S^n; Z_2) \rightarrow H_n(M^m; Z_2)$ ), see [3]. In this note we consider  $G = Z_{p^a}$ , cyclic of *odd* prime power order, and we restrict attention to maps into  $R^m$ . Our results are expressed in two theorems:

**Mod  $p^a$  Borsuk-Ulam theorem:** *For any proper action of  $Z_{p^a}$  on  $S^{2n-1}$ ,  $p$  an odd prime, and any map  $f: S^{2n-1} \rightarrow R^m$  one has*

$$\dim A(f) \geq (2n - 1) - (p^a - 1)m - [m(a - 1)p^a - (ma + 2)p^{a-1} + m + 3].$$

**Mod  $p^a$  Borsuk-Ulam anti-theorem:** *Consider the standard linear action of  $Z_{p^a}$  on  $S^{2n-1}$ . Assume  $a > 1$  and  $p^a \neq 9$ . If  $2n - 1 \leq (p^a - 1)m + (2p - 3)m - 1$  then there exists a map  $f: S^{2n-1} \rightarrow R^m$  with  $A(f) = \emptyset$ .*

Notice that the anti-theorem says that (1.1) fails whenever  $a > 1$  and  $p^a \neq 9$ ; the theorem gives  $m(a - 1)p^a - (ma + 2)p^{a-1} + m + 3$  as an upper bound for this failure. For  $a = 1$  this upper bound is 1, so for  $G = Z_p$  we are 1 off our previous results [3].

**REMARKS.** 1.  $\dim$  means covering dimension.  
 2. For  $p^a = 9$  and  $m > 1$  there is a result similar to the anti-theorem. We leave that to the interested reader.

3. In private correspondence with M. Nakaoka I have recently learned that (1.1) holds for  $Z_p$ -actions on mod  $p$  homology spheres  $S^n$  and maps  $f: S^n \rightarrow M^m$  without the restriction of niceness of  $f$  imposed by me in [3].

## 2. Proof of theorem

Let  $\mu: S \times G \rightarrow S$  be a proper action of the cyclic group  $G$  of *odd* prime power order  $p^a = q = 2k+1$  on the  $(2n-1)$  sphere  $S$ . Denote by  $\eta$  the corresponding principal  $G$ -bundle  $S \rightarrow S/\mu$  over the orbit space  $S/\mu$ . For a complex  $G$ -module  $M$  let  $\eta[M]$  be the complex vector bundle  $S \times_G M \rightarrow S/\mu$ . The correspondence  $M \mapsto \eta[M]$  gives rise to a ring homomorphism  $\alpha: \mathcal{R}G \rightarrow K^0(S/\mu)$  where  $\mathcal{R}G$  is the complex representation ring for  $G$  while  $K^0$  denotes complex  $K$ -theory. Denote by  $L$  the standard 1-dimensional complex  $G$ -module, i.e.  $L = \mathbb{C}$ , the field of complex numbers, and, fixing a generator  $g_0$  for  $G$ ,  $g_0 c = \exp(2\pi q^{-1} \sqrt{-1})c$ . Then  $\mathcal{R}G = \mathbb{Z}[\rho]/(\rho^q - 1)$  where  $\rho$  is the class of  $L$ . Finally, put  $\lambda = \eta[L]$  and for any map  $f: S \rightarrow R^m$  let  $\lambda_f$  be the restriction of  $\lambda$  to  $A(f)/\mu \subseteq S/\mu$ .

Now the mod  $p^a$  Borsuk-Ulam theorem is essentially contained in

**Lemma 1.** *If  $d\lambda_f$  has a never vanishing section then  $d \geq n-1+p^{a-1} - \frac{1}{2}am(p^a-p^{a-1})$ .*

**Proof.** Assume that  $d\lambda_f$  has a never vanishing section. We first show

$$(2.1) \quad P(\rho) := (\rho-1)^d [(\rho-1)(\rho^2-1) \cdots (\rho^k-1)]^m \in (\rho-1)^n \cdot \mathbb{Z}[\rho]/(\rho^q-1).$$

Recall that the  $i^{\text{th}}$  Atiyah class  $a_i(\xi)$  of an  $n$ -dimensional complex vector bundle  $\xi$  is given by  $a_i(\xi) = \gamma^i(\xi - n)$  with  $\gamma^i$  as in [1]. Then we have the usual Whitney duality, namely  $a_i(\xi_1 \oplus \xi_2) = \sum_{j+k=i} a_j(\xi_1) a_k(\xi_2)$ , also for any line bundle  $\xi$ ,  $a_1(\xi) = \xi - 1$ . Therefore it is immediate that  $\alpha P(\rho) = a_{d+mk}(\Lambda)$  where  $\Lambda$  is the vector bundle  $d\lambda \oplus m[\lambda \oplus \lambda^2 \oplus \cdots \oplus \lambda^k]$ , and so (2.1) follows from

$$(2.2) \quad \text{Ker } (\alpha: \mathcal{R}G \rightarrow K^0(S/\mu)) \subseteq (\rho-1)^n \cdot \mathcal{R}G,$$

$$(2.3) \quad \Lambda \text{ admits a never vanishing section.}$$

To get (2.2) we compare  $\mu$  with the standard linear action  $\mu_0: S \times G \rightarrow S$  obtained by viewing  $S$  as the unit sphere in  $nL = L \oplus \cdots \oplus L$ .  $S^{2n-1}/\mu_0$  is a  $(2n-1)$ -dimensional cell complex and  $\eta: S \rightarrow S/\mu$  is  $(2n-1)$ -universal in the sense of [5]. Hence there is a bundle map

$$\begin{array}{ccc} S & \xrightarrow{b} & S \\ \downarrow \eta_0 & \bar{b} & \downarrow \eta \\ S/\mu_0 & \xrightarrow{\bar{b}} & S/\mu. \end{array}$$

Furthermore, it is obvious that

$$\begin{array}{ccc} \mathcal{R}G & \xrightarrow{\alpha} & K^0(S/\mu) \\ & \searrow \alpha_0 & \downarrow \bar{b}^* \\ & & K^0(S/\mu_0) \end{array}$$

commutes, so  $\text{Ker } \alpha \subseteq \text{Ker } \alpha_0$ . But  $\alpha_0$  fits into an exact sequence (see [1])

$$\mathcal{R}G \xrightarrow{\varphi} \mathcal{R}G \xrightarrow{\alpha_0} K^0(S/\mu_0)$$

where  $\varphi$  is multiplication by  $\lambda_{-1}(n\rho) = (1-\rho)^n$ , so  $\text{Ker } \alpha_0 \subseteq (\rho-1)^n \cdot \mathcal{R}G$ .

In [3] it is shown that  $f$  gives rise to a section  $S$  of  $m(\lambda \oplus \lambda^2 \oplus \dots \oplus \lambda^k)$  which vanishes precisely on  $A(f)/\mu$  (see especially Digression 1, p. 171–2 and Step 3, p. 180–1 of [3]).  $s$  and the given section  $s_0$  of  $d\lambda_f$  go together to prove (2.3). This completes the proof of (2.1).

Our next step is to show that (2.1) is actually equivalent to the inequality  $d \geq n-1+p^{a-1}-\frac{1}{2}am(p^a-p^{a-1})$ . The equivalence is obvious if  $n < d+mk$ , so assume  $n \geq d+mk$ . Lift (2.1) to the polynomial ring  $Z[x]$  to get the equivalent

$$(2.1.1) \quad \exists g, h \in Z[x]: P(x) = g(x)(x-1)^n + h(x)(x^q-1).$$

Now  $P(x) = (x-1)^{d+mk} \cdot \prod_{j=2}^k f_j(x)^{m[k/j]}; (x^q-1) = (x-1) \cdot \prod_{i=1}^a f_{p^i}(x)$  where  $f_j$  is the  $j^{\text{th}}$  cyclotomic polynomial and  $[k/j]$  is the integral part of  $k/j$ . Hence, if (2.1.1) holds then  $g$  is divisible by  $\prod_{i=1}^{a-1} f_{p^i}(x)$  and  $h$  is divisible by  $(x-1)^{d+km-1}$ . So, putting  $\varepsilon_j = 0$  if  $j \nmid p^a$ ,  $\varepsilon_j = 1$  if  $j \mid p^a$ , (2.1.1) implies (and is clearly implied by)

$$(2.1.2) \quad \exists \bar{g}, \bar{h} \in Z[x]: \prod_{j=2}^k f_j(x)^{m[k/j]-\varepsilon_j} = \bar{g}(x) \cdot (x-1)^{n-d-km} + \bar{h}(x) \cdot f_q(x).$$

Let  $\gamma$  be a primitive  $q^{\text{th}}$  root of unity and consider the projection  $Z[x] \rightarrow Z[\gamma] \subseteq C$ . Its kernel is the ideal generated by  $f_q(x)$  so (2.1.2) is equivalent to

$$(2.1.3) \quad (\gamma-1)^{n-d-km} \mid \prod_{j=2}^k f_j(\gamma)^{m[k/j]-\varepsilon_j} \quad \text{in } Z[\gamma].$$

Now  $Z[\gamma]$  is precisely the algebraic integers of the field  $Q(\gamma)$  and  $(\gamma-1)Z[\gamma]$  is the *unique* prime ideal in  $Z[\gamma]$  lying above  $pZ$ , see e.g. [6]. Let  $\mathcal{N}: Q(\gamma) \rightarrow Q$  be the norm map for the extension  $Q(\gamma)/Q$ . It is then an immediate consequence of classical ideal theory for Dedekind extensions that (2.1.3) is equivalent to

$$(2.1.4) \quad \mathcal{N}(\gamma-1)^{n-d-km} \mid \prod_{j=1}^k \mathcal{N}(f_j(\gamma))^{m[k/j]-\varepsilon_j} \quad \text{in } Z.$$

The norms involved here are not hard to compute, so rearranging (2.1.4) slightly it takes the desired form  $d \geq n-1+p^{a-1}-\frac{1}{2}am(p^a-p^{a-1})$ .

If  $A(f)/\mu$  happens to be a  $CW$  complex then of course we have  $(\dim A(f)/\mu < 2d) \Rightarrow (d\lambda_f \text{ has a never vanishing section})$ , and the above lemma can then be translated into a condition on  $\dim A(f)/\mu$ . Since also  $\dim A(f) = \dim A(f)/\mu$  (because  $A(f) \rightarrow A(f)/\mu$  is a finite covering and  $\dim$  has the monotonicity and sum-properties, see [4]) this completes the proof of the mod  $p^a$  Borsuk-Ulam theorem.  $A(f)/\mu$ , however, need not be a  $CW$  complex so we need to know the following

**Lemma 2.** *If  $\lambda$  is a complex line bundle over a compact metric space  $X$  of covering dimension  $< 2d$  then  $d\lambda$  admits a never vanishing section.*

I certainly do not believe that this lemma is unknown. However, nor do I know of any reference for it, so a proof of it is given as an appendix.

### 3. Proof of the anti-theorem

Consider the standard linear action  $\mu_0$  of  $G = Z_{p^a}$  on  $S^{2N-1}$ ,  $N$  big, i.e. view  $S^{2N-1}$  as the unit sphere in  $NL = L \oplus \dots \oplus L$ .  $S^{2N-1}/\mu_0$  is a  $CW$ -complex with  $S^{2N-1}/\mu_0$  as  $(2n-1)$ -skeleton. Let  $\xi$  be the vector bundle  $S^{2N-1} \times_G IG \rightarrow S^{2N-1}/\mu_0$  where  $IG$  is the augmentation ideal of the real group algebra  $RG$ . We notice that the anti-theorem is a consequence of

$$(3.1) \quad \begin{aligned} m\xi \text{ admits a never vanishing section over the} \\ [(p^a-1)m + (2p-3)m-1]\text{-skeleton.} \end{aligned}$$

Indeed, it is well known how a section  $s$  of  $m\xi$  over the  $(2n-1)$ -skeleton corresponds to an equivariant map  $F: S^{2n-1} \rightarrow m(IG) = IG \oplus IG \oplus \dots \oplus IG$ . If  $i: IG \rightarrow RG$  is the inclusion then equivariance of  $F$  means that  $(i \oplus \dots \oplus i)F$  has the form  $(i \oplus \dots \oplus i)F(x) = (\sum_g f_i(xg^{-1})g, \dots, \sum_g f_m(xg^{-1})g)$  for well defined continuous maps  $f_i: S^{2n-1} \rightarrow R$ . Put  $f = (f_1, \dots, f_m)$  and notice that  $A(f) = \phi$  is equivalent to  $s$  having no zeros.

If we have shown (3.1) for  $m=1$  then it follows for general  $m$  by noticing that  $m\xi \cong \Delta^*(\xi \times \dots \times \xi)$  for any skeletal approximation  $\Delta: S^{2N-1}/\mu_0 \rightarrow S^{2N-1}/\mu_0 \times \dots \times S^{2N-1}/\mu_0$  to the  $m$ -fold diagonal. Hence, assume  $m=1$ . In [3] we showed that the mod  $p$  Euler class of  $\xi$  vanishes whenever  $a > 1$ . If we further exclude the case  $p^a = 9$  then the same proof shows that the integral Euler class vanishes. Hence  $\xi$  does have a never vanishing section over the  $2k$  skeleton. The obstructions to extending this section over the successive skeletons lie in  $H^{2k+i}(S^{2N-1}/G; \pi_{2k-1+i}(S^{2k-1})) \cong H^{2k+i}(G; \pi_{2k-1+i}(S^{2k-1}))$ . For  $0 < i < 2p-3$  the homotopy group in question has vanishing  $p$ -primary component so the obstructions vanish and we do have our desired section over the  $(2k+2p-4)$ -skeleton.

REMARK. In the above we have made strong use of the fact that  $\xi$  admits

a complex structure so that  $\xi$  is orientable and hence no twisting of coefficients occur.

#### 4. Remarks on the case $G=Z_{25}$ , $m=1$ , linear action

For  $G=Z_{25}$  and  $m=1$  our results show that there exists a map  $f: S^{29} \rightarrow R$  with  $A(f)=\phi$  whereas every map  $f: S^{33} \rightarrow R$  has  $A(f)\neq\phi$ . In fact every map  $f: S^{31} \rightarrow R$  has  $A(f)\neq\phi$  as we now show. Suppose that  $A(f_0)=\phi$  for some  $f_0: S^{31} \rightarrow R$ . Then

$$s_0(xG) = (x, \pi(\Sigma_g f_0(xg^{-1})g))G$$

defines a cross-section  $s_0$  of  $\xi$  over the 31-skeleton. ( $\pi: RG \rightarrow IG$  is given by  $\pi(\Sigma r_g/g) = \Sigma r_g(g-1)$ ). The obstruction to extending  $s_0$  further lie in  $H^{32+i}(S^{2N-1}/Z_{25}; \pi_{31+i}(S^{23}))$ . Since the 5-primary component of  $\pi_{31+i}(S^{23})$  is zero for  $0 \leq i < 6$  we get a never vanishing section over the 37-skeleton. As in §3 this gives an  $f: S^{37} \rightarrow R$  with  $A(f)=\phi$ . But that contradicts the above result for maps  $S^{33} \rightarrow R$ .

Unfortunately for  $p^a > 25$  our positive and negative results are too far apart to close the gap between them by means as trivial as the above.

#### Appendix. Proof of lemma 2

Let  $\Delta$  be the abstract  $4d-1$  simplex and  $|\Delta|$  its standard realization in  $R^{4d}$ . By the general embedding theorem for compacta (see e.g. p. 139 of [2])  $X$  can be taken as a closed subspace of  $|\Delta|$ . Let  $K_n$  be the subcomplex of  $\Delta^{(n)}$  ( $=n^{\text{th}}$  barycentric subdivision of  $\Delta$ ) spanned by all  $4d-1$  simplices  $\tau$  for which  $|\tau| \cap X \neq \phi$ . Then  $K_n$  is a subcomplex of the barycentric subdivision of  $K_{n-1}$  so the inclusion  $i_n: |K_n| \rightarrow |K_{n-1}|$  admits a simplicial approximation  $\varphi_n: K_n \rightarrow K_{n-1}$ . Also  $\{|K_n|\}$  is cofinal in the (downward) directed set of all neighborhoods of  $X$  in  $|\Delta|$ , so for any abelian group  $A$  we have  $\check{H}^*(X; A) \cong \lim_{\rightarrow} H^*(|K_n|; A)$ , where as usual  $\check{H}^*$  is Čech cohomology, while  $H^*$  can be taken as any ordinary cohomology theory. Since line bundles are characterized by the first Chern class  $c_1 \in \check{H}^2(-; Z)$  it follows that  $\lambda$  admits an extension  $\lambda_N$  over  $|K_N|$  for  $N$  sufficiently large. Fix such an  $N$  and define (inductively, for  $n > N$ )  $\lambda_n = |\varphi_n|_* \lambda_{n-1}$ . Let  $\sigma_n$  be the sphere bundle associated with  $d\lambda_n$ . Since  $\lambda \cong \lambda_n|_X$ ,  $n \geq N$ , it is clearly sufficient to show that  $\sigma_n$  admits a cross-section when  $n$  is sufficiently large, in other words, if we let  $k$  be the maximal number such that for some  $n \geq N$   $\sigma_n$  admits a cross-section over the  $k$ -skeleton  $|K_n^k|$  of  $K_n$ , then we must show  $k \geq 4d-1$ . Suppose  $k < 4d-1$ . Choose  $n \geq N$  such that  $\sigma_n|_{|K_n^k|}$  has a cross-section,  $s$ , say. Consider the restriction  $s'$  of  $s$  to the  $(k-1)$ -skeleton and the obstruction  $c$  to extending  $s'$  over the  $(k+1)$ -skeleton (obstruction in the sense of [5]).  $c \in H^{k+1}(|K_n|; \pi)$  where  $\pi = \pi_k(S^{2d-1})$ , and —

by maximality of  $k$ — $c \neq 0$ . Since  $k$  is clearly  $\geq 2d-1$  our assumption on  $\dim X$  assures that  $H^{k+1}(X; \pi) = \lim_{\rightarrow_j} H^{k+1}(|K_j|; \pi)$  vanishes so there is an  $m > n$  such that  $c|K_m| = |\varphi|^*c = 0$ ; here  $\varphi$  is an abbreviation for  $\varphi_{n+1}\varphi_{n+2}\dots\varphi_m: K_m \rightarrow K_n$ . Now  $\sigma_m = |\varphi|^*\sigma_n$  and  $|\varphi|$  is skeleton preserving so  $s$  gives rise to a cross-section  $s_1$  of  $\sigma_m|K_m^k|$ . Moreover, if  $s_1'$  is the restriction of  $s_1$  to  $|K_m^{k-1}|$  then the obstruction to extending  $s_1'$  over  $|K_m^{k+1}|$  is precisely  $|\varphi|^*c$ . But  $|\varphi|^*c = 0$  so  $s_1'$  does extend over  $|K_m^{k+1}|$ , thus contradicting the maximality of  $k$ .

UNIVERSITY OF ILLINOIS AT CHICAGO CIRCLE AND AARHUS UNIVERSITY

---

### References

- [1] M.F. Atiyah: K-theory, Notes by D.W. Anderson, Harvard University, 1964.
- [2] W. Franz: Topologie I, Allgemeine Topologie, Walter de Gruyter and Co., Berlin, 1960.
- [3] H.J. Munkholm: *Borsuk-Ulam type theorems for proper  $Z_p$ -actions on (mod  $p$  homology)  $n$ -spheres*, Math. Scand. **24** (1969), 167–185.
- [4] J. Nagata: Modern Dimension Theory, North Holland Publ. Co., Amsterdam, 1965.
- [5] N.E. Steenrod: The Topology of Fibre Bundles, Princeton University Press, 1951.
- [6] H. Weyl: Algebraic Theory of Numbers, Princeton University Press, 1940.