

Title	Immunoadjuvant Activities of the Enzymatic Digests of Bacterial Cell Walls Lacking Immunoadjuvancy by Themselves
Author(s)	Kotani, Shozo; Watanabe, Yoshiro; Kinoshita, Fumio et al.
Citation	Biken journal : journal of Research Institute for Microbial Diseases. 1977, 20(2), p. 87-90
Version Type	VoR
URL	https://doi.org/10.18910/82596
rights	
Note	

Osaka University Knowledge Archive : OUKA

<https://ir.library.osaka-u.ac.jp/>

Osaka University

SHORT COMMUNICATION

IMMUNOADJUVANT ACTIVITIES OF THE ENZYMATIC DIGESTS OF BACTERIAL CELL WALLS LACKING IMMUNOADJUVANCY BY THEMSELVES

SHOZO KOTANI, YOSHIRO WATANABE, FUMIO KINOSHITA and KEIJIRO KATO

Department of Microbiology, Osaka University Dental School, Joan-cho, Kita-ku, Osaka

HAROLD R. PERKINS

Department of Microbiology, University of Liverpool, Life Science Building, Liverpool L69 3BX, U. K.

(Received December 28, 1976)

Previous works in this series (Kotani et al., 1975, 1977; Kotani, 1976) have shown that the cell walls from *Staphylococcus epidermidis* (ATCC 155), *Micrococcus lysodeikticus* (NCTC 2665) and *Arthrobacter* sp. (NCIB 9423) were exceptionally adjuvant-inactive, among the test walls from all 25 species (27 strains) of gram-positive bacteria whose peptidoglycans were the group A types (Schleifer and Kandler, 1972), in both induction of delayed-type hypersensitivity and stimulation of antibody production when injected with ovalbumin into the foot pads of guinea pigs as a water-in-oil emulsion. It has also been demonstrated (Kotani et al., 1977) that the cell walls of the group B peptidoglycan types obtained from five bacterial species such as *Corynebacterium poinsettiae* (NCPP 177) and other plant-pathogenic corynebacteria were devoid of the immunoadjuvancy.

To elucidate the reasons of inabilities as an immunoadjuvant of the above cell walls, the walls of *S. epidermidis* (ATCC 155), *M. lysodeikticus* (NCTC 2665) and *C. poinsettiae* (NCPP 177) were degraded into soluble "frag-

ments" by treatment with appropriate cell wall lytic enzymes of different attack points on peptidoglycans, and the immunoadjuvant activities of the enzymatic digests were assayed. Two kinds of the peptidoglycan-degrading enzymes were used for this purpose; one of them was an endo-*N*-acetylmuramidase which hydrolyzed the β -1,4 glycosidic linkage between the *N*-acetylmuramic acid and *N*-acetylglucosamine residues in a glycan chain and the other was endopeptidases which cleaved the cross-links between the neighbouring stem peptide subunits in the peptide portion (Kato, 1975).

The cell wall specimens prepared as described in previous papers (Kotani et al., 1975; Perkins, 1971) were submitted to digestion with both Mutanolysin (Yokogawa et al., 1974; 1975) and the L-11 enzyme (Hamada et al., 1971) or L-3 enzyme (Katayama et al., 1976) under the conditions as shown in Table 1. A solubilized material separated by centrifugation at $10,000 \times g$ for 30 min was, without fractionation, examined for the immunoadjuvant activities as previously described (Kotani

TABLE 1. *Digestion of cell walls with peptidoglycan-degrading enzymes*

Constituent	<i>S. epidermidis</i>		<i>M. lysodeikticus</i>		<i>C. poinsettiae</i>	
	Mutanolysin (mg or U)	L-11 55 U ^a	Mutanolysin 0.5 mg	L-11 18 U ^b	Mutanolysin 0.15 mg	L-3 0.25 mg
Cell walls (mg)	52.0 mg	32.2 mg	51.0 mg	41.2 mg	1.5 mg	1.5 mg
Buffer (pH, molar con- centration)	Na-acetate 6.5, 0.005	Na-phosphate 8.0, 0.01	Na-acetate 6.5, 0.005	Na-phosphate 8.0, 0.01	Na-phosphate 6.5, 0.01	Na-phosphate 7.8, 0.01
Total volume (ml)	3.0	2.6	3.0	2.6	2.4	2.4
Reaction time (hr)	48	48	24	48	90	90
Insoluble re- sidue (mg)	6.98	6.80	0.50	0.65	0.60	0.40

^a Lytic units against *S. epidermidis* cell walls.

^b Lytic units against *M. lysodeikticus* cell walls.

et al., 1975). The amount of solubilized cell walls was calculated by subtracting the amount of insoluble residue from that of a cell wall specimen used for digestion.

Table 2 indicates that the enzymatic digest, either by the muramidase or the endopeptidase, of *S. epidermidis* cell walls showed the strong adjuvant activities in terms of positive corneal and delayed-type skin responses and elevation of precipitating antibody level to a test protein antigen (ovalbumin) when administered to guinea pigs as a water-in-oil emulsion. It has also become apparent, unexpectedly, that the solubilized product by the L-11 endopeptidase, but not the Mutanolysin digest, of *M. lysodeikticus* walls exhibited the weak but definite immunopotentiating effects which had never been recognized with the cell walls themselves isolated from this organism (We found that the walls from two strains other than NCTC 2665 were also devoid of the adjuvant activities). The cell walls of *C. poinsettiae*, on the other hand, did not show the immunoadjuvant activities even after solubilization with either the muramidase or the L-3 endopeptidase.

The results reported in this paper are of interest in view of the following facts. All available data on the chemical structures of

bacterial cell wall peptidoglycans indicate that disaccharide tri- or tetra-peptide subunits are highly polymerized by extended glycan chain (e.g. the average chain length of a glycan part of *S. epidermidis* cell walls was reported to be around 13 hexosamine residues, Tipper, 1969), but they are not assembled in more than dimeric units (at most trimer) by cross-linking between the peptide subunits (Kato, 1975). Consequently, the endopeptidase digest of bacterial cell walls has been found to have on the whole a much bigger molecular size or a much more repetitive structure than the endo-*N*-acetylmuramidase digest. In this connection, we have found that the water-soluble adjuvant-active component(s) obtained from the endopeptidase digest of the cell walls from various gram-positive bacteria exhibited the immunoadjuvant activities to induce delayed-type hypersensitivity to ovalbumin and had definite arthritogenic activity in Lewis inbred rats, but the component(s) obtained by digestion with glycan-splitting enzymes of the same cell wall specimens failed to induce delayed-type anti-ovalbumin hypersensitivity and to produce arthritis in Lewis rats (Koga et al., 1976; Kohashi et al., 1976; Kotani et al., unpublished observation). The findings by Ciorbaru et al. (1976) would also be significant by indi-

TABLE 2. *Immunoadjuvant activities of the products solubilized by the action of endo-N-acetylmuramidase or endopeptidase on the adjuvant-inactive cell walls from S. epidermidis, M. lysodeikticus and C. poinsettiae*

Experimental group	Cell walls from	Enzymatic treatment with	Dose (μ g)	Corneal response (48 hr) ^a Mean (Range)	Skin response (48 hr) ^a Erythema (mm) ^f , Induration ^b		Antibody level (ratio) ^c	
					Mean \pm S.E.	Mean \pm S.E.	Mean	S.E.
6	<i>S. epidermidis</i> (ATCC 155)	None	200	0	7 \pm 0.49	ND ^g	1.1	\pm 0.17
45		Mutanolysin	200	2.5 (1.0-3.0)	18 \pm 3.2 ^{*,d}	2.4 \pm 0.34	4.4	\pm 1.1*
47		L-11	200	3.0	18 \pm 1.7*	2.9 \pm 0.14 ^{**}	4.4	\pm 0.90 ^{**}
3	<i>M. lysodeikticus</i> (NCTC 2665)	None	100	0	ND	ND	1.2	\pm 0.43
46		Mutanolysin	100	0	9 \pm 0.55	1.9 \pm 0.36	0.55	\pm 0.08
60		L-11	200	2.3 (0.5-3.0)	15 \pm 1.2 ^{**}	2.2 \pm 0.15 ^{**}	3.6	\pm 0.85*
43	<i>C. poinsettiae</i> (NCPP 177)	None	100	0.3 (0 -1.0)	6 \pm 1.7	1.5 \pm 0.17	0.65	\pm 0.13
60		Mutanolysin	100	0.5 (0 -1.5)	11 \pm 0.64	1.8 \pm 0.03	1.6	\pm 0.24
62		L-3	100	0	9 \pm 2.0	1.4 \pm 0.13	0.08	\pm 0.015 ^{**}
3	None (Freund incomplete type)	None	None	0	9 \pm 0.93	1.4 \pm 0.10	[62 \pm 20.3] ^e	
62				0.6 (0 -2.0)	11 \pm 1.2	1.8 \pm 0.13	[167 \pm 17.8]	

^a Corneal test and skin test were performed 3 and 4 weeks after the sensitization, respectively.

^b Ratio of double thickness of the skin injected with 100 μ g ovalbumin/0.1 ml saline to that of the skin of the opposite side.

^c Ratio of antibody nitrogen (μ gN/ml serum) in the test group to that in the respective FIA group.

^d The difference between the test and respective control group was significant at a level of 5% (*) or 1% (***) by the "Student" t-test.

^e μ g Antibody nitrogen/ml serum specimen.

^f Average diameter of redness (mm).

^g Not determined.

cating that the products solubilized by digestion with *Streptomyces albus* G (endo)peptidases of *Nocardia rubra* cell wall peptidoglycans were mitogenic, but the lysozyme digests were not. Further work is required to obtain the logical interpretation of the observations reported here.

REFERENCES

Ciorbaru, R., J.-F. Petit, E. Lederer, E. Zissman, C. Bona, and L. Chedid. 1976. Presence and subcellular localization of two distinct mitogenic fractions in the cells of *Nocardia rubra* and *Nocardia opaca*: Preparation of soluble mitogenic peptidoglycan fractions. *Infect. Immun.* 13: 1084-1090.

ACKNOWLEDGMENTS

This research was supported in part by Grants in Aid for Scientific Research from the Ministry of Education, Science and Culture (Nos. 148149 and 112118).

Hamada, S., T. Narita, S. Kotani, and K. Kato. 1971. Studies on cell walls of group A *Streptococcus pyogenes*, type 12. II. Pyrogenic and related biological activities of the higher molecular weight fraction of an enzymatic digest of the cell walls. *Biken J.* 14: 217-231.

- Katayama, T., T. Matsuda, K. Kato, and S. Kotani. 1976. Isolation and purification of D-alanyl-meso-2,6-diaminopimelic acid endopeptidase of *Sterptomyces* L-3 enzyme using soluble substrates of known chemical structure from *Lactobacillus plantarum* cell wall digests. *Biken J.* 19: 75-91.
- Kato, K. 1975. Degradation of Bacterial Cell Walls. p. 198-232. In S. Suzuki, T. Yamada, and I. Yamashina [ed.] *Methods Biochemistry Vol. 10: Metabolism of Saccharides*, Tokyo Kagaku Dōjin. [In Japanese]
- Koga, T., S. Kotani, T. Narita, and C. M. Pearson. 1976. Induction of adjuvant arthritis in the rat by various bacterial cell walls and their water-soluble components. *Int. Arch. Allergy Appl. Immunol.* 51: 206-213.
- Kohashi, O., C. M. Pearson, Y. Watanabe, S. Kotani, and T. Koga. 1976. Structural requirements for arthritogenicity of peptidoglycans from *Staphylococcus aureus* and *Lactobacillus plantarum* and analogous synthetic compounds. *J. Immunol.* 116: 1635-1639.
- Kotani, S., T. Narita, D. E. S. Stewart-Tull, T. Shimono, Y. Watanabe, K. Kato, and S. Iwata. 1975. Immunoadjuvant activities of cell walls and their water-soluble fractions prepared from various gram-positive bacteria. *Biken J.* 18: 77-92.
- Kotani, S. 1976. Biological activities of bacterial cell wall peptidoglycans and their subunits, with special reference to the immunoadjuvant actions. *Seikagaku* 48: 1081-1107. [In Japanese]
- Kotani, S., Y. Watanabe, F. Kinoshita, K. Kato, K. H. Schleifer, and H. R. Perkins. 1977. Inabilities as an immunoadjuvant of cell walls of the group B peptidoglycan types and those of arthro-bacters. *Biken J.* 20: 1-4.
- Perkins, H. R. 1971. Homoserine and diamino-butyric acid in the mucopeptide-precursor-nucleotides and cell walls of some plant-pathogenic corynebacteria. *Biochem. J.* 121: 417-423.
- Schleifer, K. H., and O. Kandler. 1972. Peptidoglycan types of bacterial cell walls and their taxonomic implications. *Bacteriol. Rev.* 36: 407-477.
- Tipper, D. J. 1969. Structures of the cell wall peptidoglycans of *Staphylococcus epidermidis* Texas 26 and *Staphylococcus aureus* Copenhagen. II. Structure of neutral and basic peptides from hydrolysis with the *Myxobacter* AL-1 peptidase. *Biochemistry* 8: 2192-2202.
- Yokogawa, K., S. Kawata, S. Nishimura, Y. Ikeda, and Y. Yoshimura. 1974. Mutanolysin, bacteriolytic agent for cariogenic streptococci: Partial purification and properties. *Antimicrob. Agents Chemother.* 6: 156-165.
- Yokogawa, K., S. Kawata, T. Takemura, and Y. Yoshimura. 1975. Purification and properties of lytic enzymes from *Streptomyces globisporus* 1829. *Agr. Biol. Chem.* 39: 1533-1543.