

Title	Immunological Unresponsiveness in Mice. I. Immunological Unresponsiveness Induced in Embryonic Mice by Maternofetal Transfer of Human γ -Globulin
Author(s)	Shinka, Sohei; Dohi, Yoshitane; Komatsu, Tosinori et al.
Citation	Biken journal : journal of Research Institute for Microbial Diseases. 1974, 17(2), p. 59-72
Version Type	VoR
URL	https://doi.org/10.18910/82674
rights	
Note	

Osaka University Knowledge Archive : OUKA

<https://ir.library.osaka-u.ac.jp/>

Osaka University

IMMUNOLOGICAL UNRESPONSIVENESS IN MICE.

I. IMMUNOLOGICAL UNRESPONSIVENESS INDUCED IN EMBRYONIC MICE BY MATERNOFETAL TRANSFER OF HUMAN γ -GLOBULINSOHEI SHINKA, YOSHITANE DOHI, TOSINORI KOMATSU, RAMAMURTHI NATARAJAN¹ and TSUNEHISA AMANO

Department of Immunology, Research Institute for Microbial Diseases, Osaka University, Yamada-kami, Suita, Osaka

(Received February 28, 1974)

SUMMARY Immunological unresponsiveness to human γ -globulin (H_7G) was induced in fetal mice by maternofetal transfer after a single injection of tolerogen (deaggregated H_7G) into the mothers. The degree and duration of this unresponsiveness depended upon the dose of tolerogen injected and the time of its administration during pregnancy.

Injection of 5 mg of tolerogen into pregnant mice on the 14-15th day of gestation or later caused stable unresponsiveness in their offspring which lasted for more than 56 weeks after challenge with H_7G in Freund's incomplete adjuvant. However, injection of tolerogen before the 14-15th day of gestation, resulted in transient unresponsiveness in the offspring, followed by antibody formation. This unresponsiveness was specific to the tolerogen and was not terminated by challenge with a cross-reacting antigen, such as rabbit γ -globulin. When the challenge was given in Freund's complete adjuvant, only injection on the 14th day resulted in lasting unresponsiveness. Thus the 14-15th day seems to be a critical stage for induction of the immunological unresponsiveness. The significance of these results is discussed in the relation of the ontogenic development of immunocompetent cells, that is, thymus-derived cells and bone marrow-derived cells.

Similar results were obtained in rats, but no relationship could be found between the time of induction of unresponsiveness and its specificity.

INTRODUCTION

There are several reports of attempts to induce immunological unresponsiveness (or tolerance)

of humoral antibody responses by prenatal exposure alone to soluble protein antigens (Dresser and Mitchison, 1968). Induction of unresponsiveness has mainly been demonstrated in chickens (Hirata and Schechtman,

¹ Present address: Department of Microbiology, Maulana Azad Medical College, New Delhi, India.

1960; Tempelis, Wolfe and Mueller, 1958) and not seen in mammals (Hanan and Oyama, 1954; Downe, 1955), though there have been a few experiments on rabbits (Trench, Gardner and Green, 1964). So far no studies on unresponsiveness in embryo mice have been reported.

As is well known, it is relatively simple to produce unresponsiveness by neonatal injection of a variety of soluble antigens into mice and it is generally accepted that such susceptibility to tolerance may be due to immaturity of the immune system in neonatal animals (Burnet, 1969). Therefore, more younger embryos could be tolerated more readily on contact with a certain amount of antigen.

In mice, the cellular events involved in the immune response have been well characterized. For antibody response to antigens, in general, the cooperation of at least two specific cell types is required. (Mitchell and Miller, 1968; Claman and Chaperon, 1969; Rajewsky et al., 1969). One type is bone marrow-derived lymphocytes (B cells) which are the direct precursors of antibody-forming cells and respond to the antigenic determinant on a given antigen. The other type is thymus-derived lymphocytes (T cells) which respond to the carrier portion of the antigen and have helper function for B cells (Mitchell and Miller, 1968). Moreover, the immunological unresponsiveness induced by an antigen (i.e. tolerogen) is understood to be a function of the unresponsiveness of T or B cells (Miller and Mitchell, 1969; Taylor, 1969; Playfair, 1969; Roelants and Goodman, 1970; Chiller, Habicht and Weigle, 1970).

Recently, much attention has been focused on the ontogenic development of the immune system in embryonic mice (Decker et al., 1972; Dwyer and Mackay, 1972; Chiscon and Golub, 1972; Nossal and Pike, 1972, 1973; Spear et al., 1973). The times of appearance and the kinetic pattern of T cells, B cells and specific antigen-binding cells in various organs have been studied by measuring the reactivities of test cells to antibodies specific to T or B

cells and radioactive antigens at different embryonic ages. The development of the functions of T or B cells have also been investigated by determination of the ability of test cells to cooperate with adult B or T cells. However, there are no reports on the onset of susceptibility to tolerance of T or B cells in embryonic mice.

In studies on this problem we plan to induce unresponsiveness to a given antigen as early during embryonic life as possible and to see when susceptibility to tolerance starts and whether the time is the same for different antigens. For this purpose, H₇G seemed a suitable antigen because this protein can be transferred across the hemochorial barrier (Gitlin and Koch, 1968; Gitlin and Morphis, 1969; Morphis and Gitlin, 1970) and because H₇G in its monomeric form (deaggregated H₇G) has been widely used as a tolerogen in experiments on immunological tolerance (Dresser 1962; Golub and Weigle, 1967a, 1967b; Chiller et al., 1970).

This paper reports that unresponsiveness to H₇G could be induced in embryonic mice by its maternofetal transfer between the 14th day of gestation and term. This unresponsiveness was lost on challenge with Freund's complete adjuvant. From these results it is suggested that the cellular development of unresponsiveness varies with the time of its induction.

MATERIALS AND METHODS

1. *Animals*

Mice of the C3H strain were obtained from the Shionogi Lab., Osaka, and the Donryu rats from Nihon Rat Co., Saitama. They were fed on stock diet MF obtained from the Oriental Kobo Co., Osaka. Breeding animals were kept as groups of 20 female and 5 male mice per cage and 5 female and 2 male rats per cage and females were examined daily for vaginal plugs. Females possessing the plug were removed, caged separately, and treated with the tolerogen at the required time of gestation. The offspring were kept with their mothers until 4 weeks of age and then separated. The offspring were not segregated according to sex.

2. Reagents

Cohn fraction II of human γ -globulin ($H\gamma G$) and rabbit γ -globulin ($R\gamma G$) were obtained from AB KABI, Stockholm, Sweden. Both γ -globulins were purified by two chromatographies on a column of DEAE-cellulose. Purified γ -globulins (IgG) from monkey (Mon), horse (Hor), bovine (Bov), sheep (Sh) and goat (Goa), were obtained by same procedure from their globulin fractions. These sera were obtained from Kanonji Institute, The Research Foundation for Microbial Diseases of Osaka University, Kanonji, Kagawa. Six times recrystallized hen egg-white lysozyme (HL) were purchased from the Seikagakukogyo Co. Ltd., Osaka.

Two fragments, Fab and Fc, of $H\gamma G$ were prepared by digestion of $H\gamma G$ with crystalline mercuripapain, following the procedure of Porter (1959). These fractions were purified by the methods of Edelman (1960).

3. Protein concentration

Protein concentration were determined from the absorption at $280\text{ m}\mu$, taking the $E_{1\text{cm}}^{1\%}$ values of $H\gamma G$, $R\gamma G$ and HL as 14, 15 and 26.4, respectively.

4. Induction of unresponsiveness

Pregnant 6 month-old mice or rats were given a single intravenous injection of deaggregated protein (d- $H\gamma G$ or d- $R\gamma G$, referred to as tolerogen) in 0.1 ml of phosphate buffered saline, pH 7.0 (PBS-7). The protein, 5% solution in PBS-7, was deaggregated by centrifugation at $100,000\times g$ for 120 min in the SW 50 rotor of a Beckman ultracentrifuge. The upper third of the preparation was diluted with PBS-7 and used as tolerogen. In mice, the day of gestation when tolerogen was injected was estimated, taking the day when the litter was found as day 21 of gestation. On the average the period of gestation was 20 days.

5. Immunization procedures

The offspring of mothers which had received the tolerogen during pregnancy were immunized at 4 or 8 weeks of age, by subcutaneous injection of 50 μg (for mice) or 500 μg (for rats) of the protein in 0.1 ml of Freund's incomplete (FICA) or complete adjuvant (FCA). Booster injections were given 4 and 26 weeks after the primary immunization (referred to as challenge) using the same antigen in FICA. In




FIGURE 1. Outline of schedule for induction of unresponsiveness, immunization and bleeding of mice or rats.

some experiments, heat-aggregated H₇G (a-H₇G) was used to immunized mice. Heat-aggregation was performed by Gamble's methods (Gamble, 1966). The method of immunization used was as described by Chiller (1971).

6. Determination of serum antibody level

At intervals after the challenge, blood was taken from the ophthalmic vein of mice and the tail of rats. The antibody titers of the sera were determined by the passive hemagglutination (PHA) test, using 0.5% sheep erythrocytes coated with the antigen by glutaraldehyde (Segre and Segre, 1968). Results are expressed as log₂ (PHA titer)⁻¹. When diluted serum was used, the dilution factor is shown. PHA titers of groups of mice or rats are expressed as geometric means with the standard deviations.

7. Hemolytic plaque assay

The method of Jerne and Nordin (1963) was used with the modifications of Cunningham and Szenberg (1968). H₇G was covalently coupled to indicator sheep erythrocytes with 1-ethyl-3-(3-dimethylamino-propyl) carbodiimide HCl (Golub et al., 1968), obtained from the K & K Lab. Inc., Plainview, N. Y.. Indirect plaque-forming cells (PFC) were developed using an optimal concentration of goat anti-mouse γ -globulin.

The schedule for induction of unresponsiveness, immunization and bleeding of mice and rats is outlined in Fig. 1.

RESULTS

1. Immunological unresponsiveness induced in fetal mice

1) Effect of the dose of tolerogen on induction of immunological unresponsiveness in fetal mice

Three groups of mice in late pregnancy were given single intravenous injections of 0.1 ml of solutions containing 5 mg, 0.5 mg and 0.05 mg of d-H₇G, respectively and their offspring were challenged by subcutaneous injection of 50 μ g of H₇G or HL in FICA at 8 weeks of age. Four weeks later, sera were taken and their PHA titers against H₇G were examined. As shown in Table 1, treatment of mothers in

TABLE 1. Effect of the dose of d-H₇G injected into pregnant mice on the induction of unresponsiveness to H₇G in fetal mice^a

H ₇ G treatment		Litter No.	Challenge antigen		
Dose (mg)	Day of gestation		H ₇ G HL		
		PHA ^b titer against			
		H ₇ G	R ₇ G	HL	
5	18	1	—	—	
			—	—	
			—	—	
	15	2	—	—	
			—	—	
			—	—	
		3	—	—	2
			—	—	1
			—	—	2
0.5	18	4	—	—	3
			—	—	2
			—	—	3
	16	5	—	—	
			—	—	
			—	—	
		6	—	—	
			—	—	
			—	—	
0.05	18	7	—	—	
			—	—	
			—	—	
	8	3	—		
		2	—		
		1	—		
9	4	1			
	5	—			
	—	—			
None	11	3	2	3	
		5	2	2	
		7	3	1	
	12	7	4	3	
		3	—		
		6	4		
7	7	—	2		
	7	—			
	7	1			
7	7	2			
	7	—			
	7	2			

^a Mice were challenged subcutaneously with 50 μ g of H₇G or HL in FICA.

^b Values are the PHA titers of 10-fold diluted sera taken from the mice 4 weeks after the challenge. (—), PHA titer below 1:10.

late pregnancy with 5 or 0.5 mg of d-H₇G induce complete unresponsiveness to H₇G in the offspring, while 0.05 mg of d-H₇G induced partial unresponsiveness.

This unresponsiveness was specific to H₇G, and the immune response to HL was not suppressed.

2) Effect of time of administration of tolerogen on persistence of the state of the unresponsiveness

Six groups of pregnant mice were given a single intravenous injections of 5 mg of d-H₇G on days (9, 13, 14, 15, 18 and 20) of gestation, respectively, and their offspring were challenged by subcutaneous injection of 50 μg of H₇G in FICA at 4 weeks of age and given booster injections 4 and 26 weeks after the challenge. The offspring of untreated mice were used as controls. The PHA titers to H₇G in the sera of these mice were followed until 56 weeks after the challenge. Fig. 2 shows the changes in the PHA titers in each group. Four weeks after the challenge, no experimental groups showed antibody titers against H₇G although the control group showed a significant antibody titer. After the booster injection, offspring of mice injected with tolerogen on the 9th day of gestation (9 day-group) began to produce antibodies and later some mice in the 13 day-group also began to show antibody titers. However, most mice in the 14, 15 and 18 day-groups remained unresponsive until at least 56 weeks after the challenge. Mice in the 20 day-group also showed prolonged unresponsiveness, but it was not so complete as in the above groups.

3) Effect of cross-reacting IgG on the state of unresponsiveness

Two groups of offspring from mice injected with 5 mg of d-H₇G on the 16th and 18th day of gestation respectively were subdivided into two sections, and these were challenged subcutaneously with 50 μg of H₇G and R₇G in FICA at 4 weeks of age, respectively. A further two groups of mice from mothers injected with 5 mg of d-R₇G on days 16 and 18 were treated similarly. Sera were taken 4 to

20 weeks after the challenge and antibody titers against both H₇G and R₇G were determined by PHA tests. The antibody titers 12 weeks after the challenge are given in Fig. 3.

The results show that the offspring of mice treated with d-H₇G on day 16 or 18 day of gestation and challenged with H₇G showed complete unresponsiveness to both H₇G and and cross-reacting R₇G. However, mice challenge with R₇G demonstrated slightly suppressed responses to R₇G compared with control mice but no antibody titers to H₇G. On the other hand, offspring of mice treated with d-R₇G on day 16 or 18 of gestation and challenged with R₇G demonstrated almost no antibody titers to either antigen, whereas those challenged with H₇G manifested a normal immune response to H₇G but did not produce any detectable antibody to R₇G. The degree of serological cross-reactivity between H₇G and R₇G in control groups was very low.

Thus, it did not seem possible to terminate the state of unresponsiveness by challenge with cross-reacting antigen, using any combination of antigens.

4) Effect of adjuvant on persistence of unresponsiveness

Three groups of offspring of mice which had received 5 mg of d-H₇G on the 14th, 16th and 18th day of gestation were each subdivided into two sections. One section from each group was challenged with H₇G in FICA and the other section with the same antigen in FCA at 4 weeks of age. The antibody titers against H₇G 4 to 20 weeks after the challenge are shown in Fig. 4.

Mice challenged with H₇G in FICA showed persistence of the state of complete unresponsiveness until at least 20 weeks after the challenge, whereas mice challenged with H₇G in FCA showed different patterns of unresponsiveness. Mice treated at the 16th or 18th day of embryonic life started to show antibody titers 8 weeks after the challenge and these titers increased progressively with time, but mice treated at the 14th day of embryonic life did not show any antibody titer for at least




FIGURE 2. Kinetics of serum antibody titers in mice made tolerant to H_7G by maternofetal transfer at various embryonic ages and challenged with H_7G in FICA. Groups of mice from mothers which had received a single injection of 5 mg of $d-H_7G$ at various days of gestation, were challenged subcutaneously with 50 μg of H_7G in FICA at 4 weeks of age and given booster injections 4 and 26 weeks after the challenge. Values are geometric mean PHA titers against H_7G of the groups after the challenge. Standard deviations are shown as vertical bars. Shaded columns represent the mean PHA titers of control groups. *a*: Numbers shown above each column are numbers of mice per group. *b*: Numbers in parentheses are numbers of litters in each group. *c*: All mice in this group died 40–56 weeks after the challenge.




FIGURE 3. Specificity of unresponsiveness to H_7G and R_7G induced in fetal mice. Groups of mice from mothers which had received a single injection of 5 mg of $d-H_7G$ or $d-R_7G$ on the 16th day or 18th day of gestation, respectively, were challenged with 50 μg of H_7G or R_7G in FICA at 4 weeks of age and serum antibody titers to both H_7G and R_7G were determined 12 weeks after the challenge. For details, see Fig. 2. *a*: The two columns on the right show mean PHA titers of control groups.

20 weeks after the challenge.

5) Absence of antibody-forming cells in mice treated during the embryonic period

Next these results were compared with results on adult mice treated with $d-H_7G$ and challenged with heat-aggregated H_7G ($a-H_7G$) in saline, and studies were made to see whether the state of unresponsiveness induced in fetal mice was due to deficiency of antibody-forming cells. For these purposes experiments were made using $a-H_7G$ without Freund's adjuvant as the immunogen and the state of unresponsiveness was examined by the PHA test and the enumeration of plaque-forming cells. Two groups of mice treated with $d-H_7G$ on the 14th and 18th days of embryonic life, respectively, were injected intravenously with 400 μg of $a-H_7G$ at 4 weeks old and ten days later were given a booster injection of 400 μg of $a-H_7G$ intraperitoneally. Their sera and spleens were examined 5 days after the booster injection. As can be seen in Table 2, neither serum antibody nor 19S and 7S antibody-forming cells, represented as direct and indirect PFC, respectively, were detected in the tolerant mice.

2. Immunological unresponsiveness induced in fetal rats

Similar results to those described above were obtained in preliminary experiments using Donryu rats. So an attempt was made to induce partial tolerance in fetal rats by maternofetal




FIGURE 4. Effect of mycobacteria in the adjuvant on persistence of unresponsiveness induced in fetal mice. Three groups of mice from mothers which had received 5 mg of d-H γ G at different days of gestation were challenged with H γ G in FICA or in FCA at 4 weeks of age and the serum antibody titers against H γ G were determined 4 to 20 weeks after the challenge. For details, see Fig. 2. a: The two columns on the right are mean PHA titers of control groups.

transfer of a decreased dose of the tolerogen per body weight and to analyze the specificity of the antibody produced as a result of this partial tolerance. As the rats showed strong antibody response against H γ G compared with C3H mice, it was possible to analyze the speci-

ficity of it by immunodiffusion.

Eight week old rats whose mothers had been treated with 0.5 mg or 5 mg of d-H γ G at various times during pregnancy were challenged with 500 μ g of H γ G in FICA and sera were taken 4, 8 and 16 weeks after the challenge and analyzed by the PHA test and by immunodiffusion using Fab or Fc of H γ G and various cross-reacting IgGs.

As seen in Fig. 5 and Table 3, rats treated with d-H γ G in during the embryonic period showed a state of partial unresponsiveness, that is a considerable decrease of the serum antibody titer in comparison with the control group for 16 weeks after the challenge. Precipitin tests were made on individual sera using whole molecules of H γ G (IgG) and its two fragments (Fab and Fc). It was found that the sera of some rats from groups treated on the 16th or 18th day of gestation did not show any precipitation line with IgG or with the two fragments. The sera of a few rats showed precipitation lines with IgG and its fragments. The sera of most rats showed a particular state of partial unresponsiveness, termed "split tolerance", namely, their sera gave precipitation lines with Fc but not with Fab. Similar analyses were carried out using various cross-reacting IgGs and it was also noted that the sera of most rats did not give precipitation lines with any cross-reacting IgGs tested except monkey IgG. However, a few exceptional sera reacted with several kinds of IgGs while some did not react with any IgG tested. However, no relation was found between the dose of the tolerogen or the time of induction of the unresponsiveness and the specificity of the reaction.

TABLE 2. Serum antibody titers and number of antibody-forming cells in mice rendered tolerant to H₇G by maternofetal transfer at various embryonic ages and challenged with heat-aggregated H₇G^a

H ₇ G treatment		No. of mice ^b	H ₇ G PFC/10 ⁶ spleen cells ^c		PHA titer ^d against H ₇ G
Dose (mg)	Day of gestation		direct PFC	indirect PFC	
5	14	7	4±1(N=4)	3±2 (N=5)	0
	18	7	2±1(N=4)	4±2 (N=5)	0
None		8	10±4(N=4)	1010±281(N=4)	9.3±0.7

^a Mice were challenged intravenously with 400 µg of a-H₇G at 4 weeks of age and 10 days later a booster injection of the same antigen was given intraperitoneally. Five days after the booster injection, serum antibody titers and the numbers of PFC were determined.

^b Each group consisted of 2 litters.

^c Geometric mean numbers of PFC counted in spleen cells pooled according to group ± standard deviation. N, number of slides counted.

^d Geometric mean PHA titer of group ± standard deviation.

DISCUSSION

An immunological unresponsive state to H₇G, as judged by failure to form humoral antibody, was induced across the maternofetal barrier in fetal mice by a single intravenous injection of d-H₇G into pregnant mice. This unresponsiveness is specific to the tolerogen, and its degree and duration depend upon the dose of the tolerogen and the time of its administration during pregnancy.

The dose of tolerogen affected the state of unresponsiveness, since injection of increasing amounts (0.05, 0.5 and 5 mg) of d-H₇G into mice in late pregnancy changed the unresponsiveness from partial to complete. It is not known how these differences in the amount of d-H₇G injected into the pregnant mother affected the concentration of d-H₇G transferred to the fetal mice. However, Gitlin and Koch (1968) found that on increasing the maternal serum concentration of H₇G up to a level of 2 mg/ml, there was a rapid increase in the amount transferred to the fetus on the 15th to 17th day of gestation in Swiss albino mice.

The effect of the time of administration of tolerogen on the state of the unresponsiveness was studied using a constant amount of 5 mg

of d-H₇G as tolerogen and H₇G in FICA as immunogen. The results indicated that between the 14th day of gestation and term, and particularly the 14–15th day, is a highly susceptible period to tolerance and the unresponsiveness induced during this period is stable, lasting for at least 56 weeks after challenge. However, the unresponsiveness induced before the 14th day is transient and gradually disappears. Therefore, about the 14–15th day of gestation seems to be a critical stage for induction of the unresponsiveness. One possible explanation of this is that before the 14th day of gestation, H₇G cannot be transferred through the maternofetal barrier efficiently, if at all. No information is available on the catabolism of deaggregated H₇G transferred to fetal mice by the maternofetal route. However, Morphis (1970) found that the extent of maternofetal transfer of H₇G between the 15th day of gestation and term was 100 times that on the 11th day in Swiss albino Webster mice. Moreover, the half-life of H₇G in adult mice was found to be only about 2 days by determining the rate of elimination of ¹³¹I and ¹²⁵I-labeled H₇G myeloma proteins from circulation (Spiegelberg and Grey, 1968). Thus, it is quite likely that the poor induction of stable unresponsive-




FIGURE 5. Kinetics of serum antibody titers in rats rendered partially tolerant to $H\gamma G$ by maternofetal transfer at various embryonic ages and challenged with $H\gamma G$ in FICA. Seven groups of rats from mothers which had received a single injection of $H\gamma G$ at different doses at various days of gestation were challenged subcutaneously with $500 \mu g$ of $H\gamma G$ in FICA at 8 weeks of age and the serum antibody titers against $H\gamma G$ were determined 4, 8 and 16 weeks after the challenge. For details, see Fig. 2. a: The column on the right shows the mean PHA titer of the control group.

ness before the 14th day of gestation is due to an insufficient concentration of transferred tolerogen, irrespective of whether the system of antigen recognition has developed in the embryo at this time, or not. An alternative interpretation is that the susceptibility develops from the 14th day of gestation may due to development of the immune system, in-

volving antigenic recognition to tolerance by contact with a specific antigen (or tolerogen). Recently, there have been many reports on the ontogenic development of the immune system in embryonic mice. Dwyer and Mackay showed that specific binding of antigen to lymphocytes was first detected on thymic lymphocytes, at the 14th day of gestation in CBA mice. Spear et al. also found that T cells, B cells and antigen-binding cells appeared in the spleen of Swiss-L mice on the 15–16th day of gestation. On the other hand, Chiscon and Golub (1972) first detected B cell function capable of cooperating with adult T cells, on the 16th day of gestation in the liver of CBA mice, but could not detect T cell function in fetal or neonate liver and lung. Although there are small discrepancies between these results, in general they indicate that the immune system in fetal mice begins to develop on the 14–16th day of gestation. This is compatible with the second interpretation for the reason for a critical stage at the 14–15th day of gestation, given above.

Mice which became tolerant to $H\gamma G$ by maternofetal transfer of the tolerogen remained fully responsive to an unrelated antigen, such as hen egg-white lysozyme. Thus, the unresponsiveness seems to be specific to the tolerogen. However, the immune response of these mice to the cross-reacting antigen, $R\gamma G$, was slightly suppressed. This suppression may

be attributed to the unresponsiveness of cross-reacting T cells or/and B cells which are a minority of the total cells which respond to $R\gamma G$. No antibody against $H\gamma G$ was detected in these mice, so the unresponsiveness to $H\gamma G$ was not terminated by challenge with cross-reacting antigen. These results do not agree with the report of Scales and Cruse

TABLE 3. *Specificity of antibody produced in rats rendered partially tolerant to H₇G by maternofetal transfer at various embryonic ages and challenged with H₇G in FICA^a*

H ₇ G treatment		Litter No.	PHA titer against H ₇ G ^b	Precipitation with various IgGs in gel ^c									
Dose (mg)	Day of gestation			Human			Rab	Mon	Hor	Bov	Sh	Goa	
				IgG	Fab	Fc							
5	10	1	11	+	•	+	•	+	•	•	•	•	
			12	+	•	+	•	+	•	•	•	•	
			10	+	•	+	•	+	•	•	•	•	
	16	2	12	+	•	+	•	+	•	•	•	•	
			8	±	•	•	•	•	•	•	•	•	
			8	•	•	•	•	•	•	•	•	•	
	18	3	12	+	•	+	•	+	•	•	•	•	
			9	•	•	•	•	•	•	•	•	•	
			13	+	•	+	•	+	•	•	•	•	
0.5	16	4	10	+	•	+	•	+	•	•	•	•	
			9	+	•	+	•	+	•	•	•	•	
			11	+	+	+	•	+	•	•	•	•	
	18	5	12	+	•	+	±	•	•	•	•	+	
			9	+	•	•	•	•	•	•	•	•	
			11	+	•	+	•	•	•	•	+	+	
	18	6	8	12	+	•	+	•	•	•	•	•	+
				8	•	•	•	•	•	•	•	•	•
				11	+	•	+	•	•	•	•	•	•
7		8	11	•	•	•	•	•	•	•	•	•	
			8	•	•	•	•	•	•	•	•	•	
			10	+	•	+	•	+	•	•	•	•	
None	8	8	6	+	•	•	•	•	•	•	•	•	
			13	+	+	+	+	+	+	+	+	+	
			13	+	+	+	+	+	+	+	+	+	
	9	8	13	13	+	+	+	+	+	+	+	+	
				14	+	+	+	+	+	+	+	+	
				14	+	+	+	+	+	+	+	•	+
	9	9	12	12	+	•	+	+	+	+	+	+	
				13	+	+	+	•	+	•	•	•	•
				13	+	+	+	•	+	•	+	•	+
9	9	14	14	+	+	+	+	+	•	+	+		
			12	+	•	+	+	+	+	+	+	+	
			13	+	+	+	•	+	•	+	•	+	

^a Sera of rats of the 7 groups described in Fig. 5 taken 16 weeks after the challenge were analyzed by the PHA test and by gel precipitation.

^b PHA titers of 10-fold diluted sera from each rat.

^c +, strong precipitation line, ±, faint line, •, no line.

(1970) that the unresponsiveness to H₇G in adult mice may be terminated by injection of bovine γ -globulin or its fragments, but the discrepancy may be due to differences in many factors, such as the phylogenic relationship among antigens, the manner of induction of unresponsiveness and the strain of mice used. Lack of termination of the unresponsiveness was probably due to unresponsiveness of B cells which respond to H₇G or of cross-reacting T cells, capable of cooperating with the B cells. Further work is required on this.

Gitlin and Morphis (1969) found that there is no difference in the facilities for maternofetal transfer of H₇G and R₇G in mice. Further studies showed that mice which were tolerant to R₇G showed almost complete unresponsiveness to homologous R₇G but remained normally responsive to cross-reacting H₇G. This may be because the population of cross-reacting T or B cells is only a small proportion of the total T or B cells which respond to H₇G. Moreover, the unresponsiveness was not terminated by challenge with H₇G. This failure may also be explained as described above.

When Freund's complete adjuvant was used for the challenge, mice which had become tolerant to H₇G on the 16–18th day of embryonic life developed an antibody response 8 weeks after the challenge, but mice which had become tolerant on the 14th day did not, and the unresponsiveness persisted for at least 20 weeks after the challenge. These results can be explained on the basis of the following two predictions; first, the unresponsiveness induced on the 16 or 18th day of embryonic life may be restricted to specific T cells, while that induced on the 14th day may involve both T cells and B cells. Second, mycobacteria (H₃₇RA strain) in FCA may have a helper function for T cells, as suggested by Lind (1968) to explain his finding that unresponsiveness to a salmonella flagellin in rats can be terminated by challenge with flagellin in FCA. Recently, Chiller et al. (1973) found that bacterial endotoxin had a similar capacity to terminate the unresponsiveness of mice to H₇G when ad-

ministered with challenging antigen.

Previously Chiller et al. (1971) found a difference between the kinetic patterns of unresponsiveness of T cells and B cells induced in adult mice: T cells became tolerant more easily and remained for longer than B cells. A similar difference seems to exist between the susceptibilities of T cells and B cells to tolerance in fetal mice; namely, the T cells can probably easily be rendered tolerant at any time from the 14 days of gestation to term, whereas B cells are less susceptible to tolerance and in order to become tolerant a suitable amount of the tolerogen must be administered during a short critical period. The fate of B cells may be decided by various factors, such as the quantitative relationship between the concentration of tolerogen and the number of B cells or the degree of differentiation of B cells when they are exposed to the tolerogen. These prediction require further studies, including studies on the reconstitution of T and B cells. The reason for the differences found between the responses of mice which became tolerant on the 16 and the 18th days of gestation is unknown.

Heat-aggregated H₇G is highly immunogenic, without added adjuvant, and has been widely employed as challenge immunogen in previous studies on unresponsiveness to H₇G induced in adult mice (Weigle, 1972). In the present work this form of antigen gave similar results to those obtained using Freund's adjuvant. Complete unresponsiveness was observed in mice for at least 8 weeks after birth at both the serum and cellular level. No antibody was detected in the serum or antibody-forming cells in the spleen. This shows that the unresponsiveness observed in this work is due to lack of antibody formation, that is, absence of antibody forming cells.

Studies on the induction of the unresponsiveness by maternofetal transfer of d-H₇G were also made in Donryu rats. Attempts were made to produce partial tolerance in fetal rats. This was done by inoculation of decreasing amounts of d-H₇G to one tenth of

the amount employed in mice. In these partially tolerant rats, the relationship between the specificity of the depleted antibody among the anti-H₇G antibodies produced and the dose of the tolerogen or time of induction of unresponsiveness was studied.

All the rats treated on the 10th day of gestation and half the rats treated on the 16th or 18th day showed the phenomenon termed "split tolerance" (Taussig, 1971). That is, gel precipitation showed that they produced detectable antibody to Fc but not to Fab. However, the analyses were not very sensitive or quantitative, so it is still uncertain whether

"split tolerance" in the strict sense was actually induced. No relationship could be found using cross-reacting IgGs between the specificity of the depleted antibody at the determinant level and the dose of time or inoculation of the tolerogen.

ACKNOWLEDGMENTS

We wish to thank Dr. Shizuo Tanabe, Department of Bacteriology, Medical School, Osaka University for supplying C3H mice and also to express our gratitude to Miss Misako Fujiwara of this Institute for her excellent technical assistance.

REFERENCES

- Burnet, M. 1969. Cellular Immunology Books 1 and 2, Melbourne University Press and Cambridge University Press, London.
- Chiller, J. M., G. S. Habicht, and W. O. Weigle. 1970. Cellular sites of immunologic unresponsiveness. Proc. Natl. Acad. Sci. U.S.A. 65: 551-556.
- Chiller, J. M., G. S. Habicht, and W. O. Weigle. 1971. Kinetic differences in unresponsiveness of thymus and bone marrow cells. Science 171: 813-815.
- Chiller, J. M., and W. O. Weigle. 1973. Termination of tolerance to human gamma globulin in mice by antigen and bacterial lipopolysaccharide (Endotoxin). J. Exp. Med. 137: 740-750.
- Chiscon, M. O., and E. S. Golub. 1972. Functional development of the interacting cells in the immune response. I. Development of T cell and B cell function. J. Immunol. 108: 1379-1386.
- Claman, H. N., and E. A. Chaperon. 1969. Immunologic complementation between thymus and marrow cells—A model for the two-cell theory of immunocompetence. Transplant. Rev. 1: 92-113.
- Cunningham, A. J., and A. Szanberg. 1968. Further improvements in the plaque technique for detecting single antibody-forming cells. Immunology 14: 599-600.
- Decker, J., J. Clarke, L. MacPherson, R. Weinstein, and E. E. Sercarz. 1972. Early appearance of antigen-binding cells to two different antigens during fetal lymphoid development. Adv. Exp. Med. Biol. 29: 269-275.
- Downe, A. E. R. 1955. Inhibition of the production of precipitating antibodies in young rabbits. Nature 176: 740-741.
- Dresser, D. W. 1962. Specific inhibition of antibody production. II. Paralysis induced in adult mice by small quantities of protein antigen. Immunology 5: 378-388.
- Dresser, D. W., and N. A. Mitchison. 1968. The mechanism of immunological paralysis. Adv. Immunol. 8: 129-181.
- Dwyer, J. M., and I. R. Mackay. 1972. The development of antigen-binding lymphocytes in foetal tissues. Immunology 23: 871-879.
- Edelman, G. M., J. F. Heremans, M. T. Heremans, and H. G. Kunkel. 1960. Immunological studies of human γ -globulin. Relation of the precipitin lines of whole γ -globulin to those of the fragments produced by papain. J. Exp. Med. 112: 203-223.
- Gamble, C. N. 1966. The role of soluble aggregates in the primary immune response of mice to human gamma-globulin. Int. Arch. Allerg. 30: 446-455.
- Gitlin, D., and C. Koch. 1968. On the mechanisms of maternofetal transfer of human albumin and γ G globulin in the mouse. J. Clin. Invest. 47: 1204-1209.
- Gitlin, D., and L. G. Morphis. 1969. Systems of materno-fetal transport of γ G immunoglobulin in the mouse. Nature 223: 195-196.
- Golub, E. S., R. I. Mishell, W. O. Weigle, and R. W. Dutton. 1968. A modification of the hemolytic plaque assay for use with protein antigens. J. Immunol. 100: 133-137.

- Golub, E. S., and W. O. Weigle. 1967a. Studies on the induction of immunologic unresponsiveness. I. Effects of endotoxin and phytohemagglutinin. *J. Immunol.* 98: 1241-1247.
- Golub, E. S., and W. O. Weigle. 1967b. Studies on the induction of immunologic unresponsiveness. II. Kinetics. *J. Immunol.* 99: 624-628.
- Hanan, R., and J. Oyama. 1954. Inhibition of antibody formation in mature rabbits by contact with the antigen at an early age. *J. Immunol.* 73: 49-53.
- Hirata, A. A., and A. M. Schechtman. 1960. Studies on immunologic depression in chickens. *J. Immunol.* 85: 230-239.
- Jerne, N. K., and A. A. Nordin. 1963. Plaque formation in agar by single antibody-producing cells. 140: 405.
- Lind, P. E. 1968. The immune responses of normal and tolerant rats to salmonella adelaide flagellin in Freund's complete adjuvant. *Aust. J. Exp. Biol. Med. Sci.* 46: 179-188.
- Miller, J. F. A. P., and G. F. Mitchell. 1969. Thymus and antigen-reactive cells. *Transplant. Rev.* 1: 3-42.
- Mitchell, G. F., and J. F. A. P. Miller. 1968. Cell to cell interaction in the immune response. II. The source of hemolysin-forming cells in irradiated mice given bone marrow and thymus or thoracic duct lymphocytes. *J. Exp. Med.* 128: 821-837.
- Morphis, L. G., and D. Gitlin. 1970. Maturation of the maternofetal transport system for human γ -globulin in the mouse. *Nature* 228: 573.
- Nossal, G. J. V., and B. L. Pike. 1972. Differentiation of B lymphocytes from stem cell precursors. *Adv. Exp. Med. Biol.* 29: 11-18.
- Nossal, G. J. V., and B. L. Pike. 1973. Studies on the differentiation of B lymphocytes in the mouse. *Immunology* 25: 33-45.
- Playfair, J. H. L. 1969. Specific tolerance to sheep erythrocytes in mouse bone marrow cells. *Nature* 222: 882-883.
- Porter, R. R. 1959. The hydrolysis of rabbit γ -globulin and antibodies with crystalline papain. *Biochem. J.* 73: 119-126.
- Rajewsky, K., V. Schirmacher, S. Nase, and N. K. Jerne. 1969. The requirement of more than one antigenic determinant for immunogenicity. *J. Exp. Med.* 129: 1131-1143.
- Roelants, G. E., and J. W. Goodman. 1970. Tolerance induction by an apparently non-immunogenic molecule. *Nature* 227: 175-176.
- Scales, R. W., and J. M. Cruse. 1970. Abrogation of immunologic unresponsiveness by whole molecules and fragments of cross-reacting immunoglobulin. *J. Immunol.* 105: 1072-1081.
- Segre, D., and M. Segre. 1968. Hemolytic plaque formation by mouse spleen cells producing antibodies to ovalbumin. *Immunochemistry* 5: 206-212.
- Spear, P. G., A. Wang, U. Rutishauser, and G. M. Edelman. 1973. Characterization of splenic lymphoid cells in fetal and newborn mice. *J. Exp. Med.* 138: 557-573.
- Spiegelberg, H. L., and H. M. Grey. 1968. Catabolism of human γ G immunoglobulins of different heavy chain subclasses. II. Catabolism of γ G myeloma proteins in heterologous species. *J. Immunol.* 101: 711-716.
- Taussig, M. J. 1971. Studies on the induction of immunological tolerance. The inhibition of tolerance-induction by antiserum: split tolerance and the time-course of tolerance induction. *Eur. J. Immunol.* 1: 367-371.
- Taylor, R. B. 1969. Cellular cooperation in the antibody response of mice to two serum albumin: specific function of thymus cells. *Transplant. Rev.* 1: 114-149.
- Tempelis, C. H., H. R. Wolfe, and A. Mueller. 1958. The production of immunological unresponsiveness by the intravenous injection of bovine serum albumin into the chick embryo. *Br. J. Exp. Pathol.* 39: 323-327.
- Trench, C. A. H., P. S. Gardner, and C. A. Green. 1964. Induction of immunological tolerance to human gamma-globulin in rabbits using the maternal route of inoculation. *Immunology* 7: 567-569.
- Weigle, W. O. 1972. Immunological unresponsiveness. *Adv. Immunol.* 16: 61-122.