



Title	Temperature Sensitive Growth of A Herpes Type Virus Isolated from A Chicken with Marek's Disease
Author(s)	Onoda, Tetsuo; Konobe, Takeo; Takaku, Keisuke et al.
Citation	Biken journal : journal of Research Institute for Microbial Diseases. 1971, 14(3), p. 357-360
Version Type	VoR
URL	https://doi.org/10.18910/82753
rights	
Note	

The University of Osaka Institutional Knowledge Archive : OUKA

<https://ir.library.osaka-u.ac.jp/>

The University of Osaka

SHORT COMMUNICATION

TEMPERATURE SENSITIVE GROWTH OF A HERPES TYPE VIRUS ISOLATED FROM A CHICKEN WITH MAREK'S DISEASE¹

TETSUO ONODA, TAKEO KONOBE and KEISUKE TAKAKU

Kanonji Institute, The Research Foundation for Microbial Diseases of Osaka University, Kanonji, Kagawa

MATAO NAITO, KOICHI ONO and SHIRO KATO

Department of Pathology, Research Institute for Microbial Diseases, Osaka University, Yamada-kami, Suita, Osaka

(Received May 4, 1971)

Temperature sensitive (*ts*) mutants isolated from animal viruses are useful in genetic and physiological analyses of animal viruses (Fenner, 1969). The present communication briefly describes the isolation of a *ts* virus of the Biken C2 strain of a herpes-type virus (HTV) isolated by Kato et al. (1970) from a chicken with Marek's disease (MD).

The Biken C2 strain of HTV was successfully propagated in quail embryo fibroblasts (QUEF) (Onoda et al., 1970). A characteristic of this virus is that its infectivity in cell culture is largely associated with cells. The virus was assayed by a plaque method utilizing 0.9% agar overlay. Cultures of QUEF were maintained in Eagle's minimum essential medium (MEM) with 5% calf serum and 10% trytose phosphate broth. Monolayers in dishes were inoculated with HTV-infected cells. After incubation at 36°C for 24 hr most cells had become infected. Then the growth medium was replaced by medium containing nitrosoguanidine (NG) as a mutagen. This drug has been used extensively as a mutagen for DNA

type viruses (Tegtmeyer et al., 1970). The infected cells were exposed to various concentrations of NG (5, 10, 20 and 40 µg/ml) at 36°C for 5 days. NG at a concentration of 40 µg/ml caused cell degeneration. Therefore it was used at a concentration of 20 µg/ml. For screening of potential viruses, the permissive and nonpermissive temperatures used were 36°C and 41°C, respectively. After treatment with NG HTV-infected cells with an infectivity of approximately 10 plaque forming units (PFU) per dish were plated onto QUEF and kept at the permissive temperature (36°C) for 4 days. Then the plaques were carefully picked out and were each seeded in dishes containing fresh, normal QUEF monolayers. They were transferred 2 to 3 times at intervals of 4 days at 36°C. Potential viruses were then screened in QUEF at permissive and nonpermissive temperatures, using 4-fold dilutions and 2 dishes for each dilution. Viruses which showed some reduction in growth at 41°C were purified twice by plaque transfer at 36°C. Viruses harvested from single plaques of this type were tested for their abilities to multiply at 36°C and 41°C. Of 120 plaques isolated, only one was strongly suppressed by the high

¹ This work was supported in part by a Scientific Research Grant from the Ministry of Education.




FIGURE 1. *Growth curve of parental herpes type virus (HTV).* QUEF monolayers infected with HTV (1.1×10^2 PFU) were incubated at 36 C or 41 C. Samples were removed at intervals. Volumes of 0.2 ml of serial 4-fold dilutions of a suspension of infected cells were inoculated into 5.0 ml of overlay medium on each culture. Plaque assays were done 6 days after infection.




FIGURE 3. *Temperature shift-up experiment on ts HTV.* Three sets of QUEF cultures were infected with ts HTV (1.5×10^2 PFU) and incubated at 36 C. Two sets of them were transferred to 41 C 1 and 2 days after infection respectively and incubated until 4 days after infection (---○---). Plaque assays were done by the same way as for Fig. 1.




FIGURE 2. *Growth curve and temperature shift-down experiment on ts HTV.* Conditions were the same as for Fig. 1. In the temperature shift-down experiment, 3 sets of QUEF cultures were infected with ts HTV (1.3×10^2 PFU). One set of cultures was incubated at 36 C and two sets of them were incubated at 41 C. One of the latter sets was shifted to 36 C 2 days after infection and incubated for 2 days (---○---). Plaque assays were done by the same way as for Fig. 1.

temperature when compared with the parent strain. Virus titration was carried out as follows. Monolayer cultures of QUEF in 60 mm Falcon plastic dishes were infected with the parental HTV and ts HTV, respectively. At various times after infection, pairs of cultures were removed from the incubator, treated with trypsin and pooled. Volumes of 0.2 ml of serial 4-fold dilutions of a suspension of the cultured cells were used for inoculation. Plaque counts are made on four plates. The parent HTV grew equally well at 36 C and 41 C (Fig. 1). Growth of ts HTV resembled that of the parent strain at 36 C, but was hardly observed at 41 C (Fig. 2). However, growth of ts HTV occurred when infected cultures were shifted from 41 C to 36 C 2 days after infection, as shown in Fig. 2. On the other hand, when cultures infected with ts HTV were transferred from 36 C to 40 C at intervals after infection, scarcely any growth occurred after the transfer (Fig. 3). QUEF cultures infected with ts HTV were also ex-




FIGURE 4. *Fluorescence photomicrograph of QUEF infected with ts HTV for 4 days at 41 C.*




FIGURE 5. *Fluorescence photomicrograph of QUEF infected with ts HTV for 4 days at 36 C.*

amined by the fluorescent antibody technique (FAT). Monolayers of QUEF on coverslips were inoculated with *ts* HTV and the parental HTV. The coverslips were removed at intervals, fixed in cold acetone and stained by the direct FAT method (Naito et al., 1969). When QUEF cultures infected with *ts* HTV were incubated at the nonpermissive temperature for 4 days, scarcely any fluorescent cells were detected in them (Fig. 4). However, when cells infected with *ts* HTV were incubated for 4 days at the permissive temperature, many plaques consisting of numerous fluorescent cells were observed in them (Fig. 5). On the other hand, when cultures infected with the parental HTV were incubated for 4 days at the permissive and nonpermissive temperatures, many plaques consisting of numerous fluorescent cells were observed in both cultures. When

cultures infected with *ts* HTV, were transferred from 41C to 36C or from 36C to 41C at intervals increase in the number of fluorescent cells was parallel with the growth at these permissive and nonpermissive temperatures. The morphological characteristics of plaques of *ts* HTV in duck embryo fibroblasts and QUEF were examined at 36C and found to be the same as those of the parental HTV (Onoda et al., 1970).

ts HTV seems to be a kind of mutant, but it is unknown whether its appearance is related with NG treatment, or it is a spontaneous phenomenon.

REFERENCES

Fenner, F. 1969. Conditional lethal mutants of animal viruses. *Current Topics in Microbiol. Immunol.* 48: 1-28.

Kato, S., K. Ono, M. Naito, T. Doi, Y. Mori, N. Iwa and T. Onoda. 1970. Isolation of herpes type virus from chickens with Marek's disease using duck embryo fibroblast cultures. *Biken J.* 13: 193-203.

Onoda, T., K. Ono, T. Konobe, M. Naito, Y. Mori and S. Kato. 1970. Propagation of herpes type virus isolated from chickens with Marek's disease in Japanese quail embryo fibroblasts. *Biken J.* 13: 219-228.

Onoda, T., K. Koyama, T. Konobe, K. Takaku, K. Ono and S. Kato. 1970. Studies on the characteristics of herpes-type virus isolated from a chicken with Marek's disease. *Biken J.* 14: 167-176.

Naito, M., K. Ono, S. Tanabe and S. Kato. 1969. Immunofluorescent studies using fractionated immunoglobulin of sera of fowls with Marek's disease. *Biken J.* 12: 257-261.

Tegtmeyer, P., C. Dohan and C. Reznikoff. 1970. Inactivating and mutagenic effects of nitrosoguanidine on simian virus 40. *Proc. Nat. Acad. Sci. U.S.A.* 66: 745-752.