



| | |
|--------------|---|
| Title | Immunological Properties of Chikungunya Virus and Its Components |
| Author(s) | Igarashi, Akira; Nithiuthai, Pannee; Rojanasuphot, Suntharee |
| Citation | Biken journal : journal of Research Institute for Microbial Diseases. 1970, 13(3), p. 229-231 |
| Version Type | VoR |
| URL | https://doi.org/10.18910/82802 |
| rights | |
| Note | |

The University of Osaka Institutional Knowledge Archive : OUKA

<https://ir.library.osaka-u.ac.jp/>

The University of Osaka

SHORT COMMUNICATION

IMMUNOLOGICAL PROPERTIES OF CHIKUNGUNYA VIRUS AND ITS COMPONENTS

AKIRA IGARASHI, PANNEE NITHIUTHAI¹ AND SUNTHAREE ROJANASUPHOT²

Department of Preventive Medicine, Research Institute for Microbial Diseases, Osaka University, Yamada-kami Suita, Osaka

(Received May 30, 1970)

When the protein components of purified Chikungunya virus (ChV) were analyzed, two major structural proteins were found on electrophoresis in polyacrylamide gel containing sodium dodecylsulfate. One protein was associated with hemagglutinin (HANin) and the other with the core of the ChV particle. When HANin was obtained from purified ChV by Tween 80-ether treatment followed by CsCl density gradient centrifugation, it was shown to contain only the HANin protein. While the intracellular nucleoprotein core, formerly named "X", contained only the core protein of ChV (Igarashi et al., 1970). The antibody responses to these structural components of ChV and purified virus were studied.

Purified virus, HANin and the "X"-component were prepared as reported before (Igarashi et al., 1970). The purified virus was obtained by sucrose gradient sedimentation, it was diluted with PBS (Dulbecco and Vogt, 1954) and concentrated by ultracentrifugation. The precipitate was resuspended in PBS at a protein concentration of 400 μ g protein/ml, determined by the method of Lowry et al. (1951). The protein contents of HANin and the "X" component determined were found to be 150 μ g/ml before dialysis against PBS (for HANin) and KTM³ (for "X") in a cold room overnight. Then the concentrated purified ChV and the two dialyzed components, were each mixed with an equal volume of 0.5 M glycine containing 0.02 % of gelatin. To samples of ChV and "X", formalin was added at a final concentration of 0.05 %, and the mixture was kept for 4 weeks in a refrigerator at 4 C before immunization. Before immunization, the residual infectivity of all antigens was checked by inoculation into BHK21 cell cultures. No infectivity was detected.

One ml of each antigen was emulsified with an equal volume of Freund's complete adjuvant and injected intramuscularly into 2 kg female rabbit.

A preimmune serum specimen was taken from each rabbit, and samples of blood were

1 Participant in the Biken Microbiology course in 1969 arranged by the Overseas Technical Cooperation Agency (OTCA), Japan; Present address: Department of Microbiology, Suandok Hospital, Chiang-Mai, Thailand.

2 Fellow of the OTCA, Japan, under the Colombo plan; Present address: Virus Research Institute, Department of Medical Sciences, Ministry of Public Health, Bangkok, Thailand.

3 KTM: 0.01M KCl, 0.01M Tris-HCl, 0.0001 M MgCl₂, pH 7.6.

4 Virus diluent: 5% bovine serum in YLE (0.5% lactalbumin hydrolysate, 0.1% yeast extract in Earle's balanced salt solution).

removed one month later. These paired serum specimens from each rabbit were tested for antibodies against “X”, HAnin and ChV by the hemagglutination-inhibition (HI), complement fixation (CF) and neutralization (N) tests. The HI and CF tests were carried out according to the methods of Clarke and Casals (1958) and Hammon and Work (1964), modified for use in a microtiter system (Sever, 1962). The N test by the 50% plaque reduction method was performed as follows: ten fold dilutions of the test sera in virus diluent⁴ were mixed with an equal volume of 200 PFU/0.2 ml of fresh ChV at 37 C for 90 min. The mixture was then inoculated onto monolayer cultures of

VERO cells to detect plaques (Igarashi and Tuchinda, 1967). The plaque reduction rate was plotted on a probit chart (Finley, 1952) to estimate the serum dilution that reduced the plaque count to 50% of the control.

The results of serological tests are shown in Tables 1 and 2. Preimmune sera had no detectable antibody in any rabbit (Table 1). The sera from rabbits immunized with the “X”-component showed no homologous antibody detectable by the HI, N and CF tests using HAnin as antigen, but they gave a positive CF reaction when antigen from infected suckling mouse brain or “X” was used.

The sera from rabbits immunized with HAnin showed a significant positive reaction by the HI, N and CF tests with HAnin or antigen from infected suckling mouse brain but their response against “X” was not so high in the CF test. Similar results were obtained when rabbits were immunized with formalin-treated purified ChV.

These results show that immunization with HAnin is sufficient to evoke an antibody response detectable by HI and N tests in the arbovirus system as in the influenza system (Webster and Laver, 1967).

Viral antigen prepared from infected suckling mouse brains reacted with the three types of immune sera, showing that the former con-

TABLE 1. Serological tests on serum specimens taken from test rabbits before immunization.

| Rabbit No. | HI | N | CF (SMB ^a) |
|------------|-----|-----|------------------------|
| 2 | <20 | <10 | <4 |
| 3 | <20 | <10 | <4 |
| 5 | <20 | <10 | <4 |
| 6 | <20 | <10 | <4 |
| 7 | <20 | <10 | <4 |
| 8 | <20 | <10 | <4 |
| 9 | <20 | <10 | <4 |

^a The antigen was prepared from infected suckling mouse brains by sucrose-aceton extraction method of Clarke and Casals (1958).

TABLE 2. Serological tests on serum specimens taken from test rabbits one month after immunization.

| Immunogen | Rabbit No. | HI | N | CF ^a | | |
|----------------------------------|------------|------|--------------------|-----------------|-------|-----|
| | | | | SMB | HAnin | “X” |
| “X”, formalin-treated | 2 | <20 | <10 | 64 | 8 | 32 |
| | 3 | <20 | 20 | 64 | 8 | 64 |
| HAnin | 5 | 640 | 10 ^{4.35} | 128 | 128 | 8 |
| | 6 | 1280 | 10 ^{4.01} | 256 | 256 | 16 |
| Purified virus, formalin-treated | 7 | 640 | 10 ^{3.46} | 256 | 128 | 8 |
| | 8 | 640 | 10 ^{4.36} | 256 | 128 | 8 |
| | 9 | 640 | 10 ^{4.24} | 256 | 64 | 32 |

^a Three kinds of antigens were used in the CF tests.

tained both HAnin and "X" antigens. This seems significant in connection with the cross reactivity of the arboviruses, because antigens from infected suckling mouse brains are routinely used in standard serological tests. Formalin-treated vaccines of group A arbovirus have been reported to be potentially dangerous because in practice it is rather difficult to inactivate the virus particles completely in this way (Sutton and Brooke, 1954; Smith et al., 1956). Thus, if a vaccine is developed using HAnin as antigen, especially a highly virulent

arbovirus, it may be more reliable and effective than the formalin-treated virus vaccine because the HAnin is free from the viral nucleoprotein core.

ACKNOWLEDGEMENTS

We thank Prof. K. Fukai of this Department and Dr. S. Tanabe of the Department of Microbiology, Osaka University, Dr. M. Watanabe and Mrs. S. Saito of the Osaka Public Health Institute for their kind advices in this work.

REFERENCES

- Clarke, D. H. and J. Casals. 1958. Techniques for hemagglutination and hemagglutination-inhibition with arthropod-borne viruses. *Amer. J. Trop. Med. Hyg.* 7: 561-573.
- Dulbecco, R. and M. Vogt. 1954. Plaque formation and isolation of pure lines with poliomyelitis virus. *J. Exp. Med.* 99: 167-182.
- Finley, D. J. 1952. Probit analysis. A statistical treatment of the sigmoid response curve. Cambridge University Press.
- Hammon, W. M. and T. H. Work. 1964. Arbovirus infection in man p. 268-311. In E. H. Lennette and N. J. Schmidt [ed.]. Diagnostic procedures for viral and rickettsial diseases. 3rd ed. American Public Health Association Inc. New York.
- Igarashi, A., T. Fukuoka, P. Nithiuthai, L.-C. Hsu and K. Fukai. 1970. Structural components of Chikungunya virus. *Biken J.* 13: 93-110.
- Igarashi, A. and P. Tuchinda. 1967. Studies on Chikungunya virus. I. Plaque titration on an established cell line. *Biken J.* 10: 37-39.
- Lowry, O. H., N. J. Rosebrough, A. L. Farr and R. J. Randall. 1951. Protein measurement with the Folin phenol reagent. *J. Biol. Chem.* 193: 265-276.
- Sever, J. L. 1962. Application of a microtechnique to viral serological investigations. *J. Immunol.* 88: 320-329.
- Smith, D. G., H. K. Mamay, R. G. Marshall and J. C. Wagner. 1956. Venezuelan equine encephalomyelitis. Laboratory aspects of fourteen human cases following vaccination and attempts to isolate the virus from the vaccine. *Amer. J. Hyg.* 63: 150-164.
- Sutton, L. S. and C. C. Brooke. 1954. Venezuelan equine encephalomyelitis due to vaccination. *J. Amer. Med. Assoc.* 155: 1473-1476.
- Webster, R. G. and W. G. Laver. 1967. Preparation and properties of antibody directed specifically against the neuraminidase of influenza virus. *J. Immunol.* 99: 49-55.