



Title	Isolation of Temperature-Sensitive Conditional Lethal Mutants of An Avian Adenovirus (CELO), and Localization of Viral Antigen in Cells Infected with Them
Author(s)	Ishibashi, Masahide
Citation	Biken journal : journal of Research Institute for Microbial Diseases. 1970, 13(1), p. 59-62
Version Type	VoR
URL	<a href="https://doi.org/10.18910/82818">https://doi.org/10.18910/82818</a>
rights	
Note	

*The University of Osaka Institutional Knowledge Archive : OUKA*

<https://ir.library.osaka-u.ac.jp/>

The University of Osaka

## PRELIMINARY REPORT

## ISOLATION OF TEMPERATURE-SENSITIVE CONDITIONAL LETHAL MUTANTS OF AN AVIAN ADENOVIRUS (CELO), AND LOCALIZATION OF VIRAL ANTIGEN IN CELLS INFECTED WITH THEM

MASAHIDE ISHIBASHI

Department of Experimental Chemotherapy, Research Institute for Microbial Diseases, Osaka University, Yamada-kami Suita, Osaka

(Received December 1, 1969)

Chicken embryo lethal orphan (CELO) virus was first described by Yates and Fry (1957). Several characters of this virus are quite similar to those of mammalian adenoviruses (Petek et al., 1963a; Petek et al., 1963b; Dutta and Pomeroy, 1963). From studies on adenovirus DNA (Green et al., 1967), the viral chromosome seems to consist of 30-50 genes. This virus is readily propagated in primary chicken kidney cell (CKC) cultures so I select it as a model system to analyze the replication and oncogenesis of adenovirus using temperature-sensitive conditional lethal (*ts*) mutants, which replicate well at the non-restrictive temperature of 31°C but scarcely at all at the restrictive temperature of 40°C. This paper reports the isolation of these *ts* mutants and the localization of viral antigen in infected cells under restrictive and non-restrictive conditions.

Virus propagation and plaque formation were done in CKC cultures as originally described by Kawamura et al. (1963). The Ote strain of CELO virus, isolated and identified by the cross neutralization test with Phelp's strain of Yates (Kawamura et al., 1963), was kindly supplied by Dr. H. Kawamura. A26, a clone of the Ote strain was obtained by five repeated single plaque isolations in CKC cultures, and was used as the original wild type strain in the

present study. Before treatment with mutagen A26 was propagated in CKC at 40°C to dilute out possible spontaneous *ts* mutants in the viral population. The A26 stock thus obtained (infectious titer,  $2 \times 10^9$  PFU/ml) was exposed to hydroxylamine, as described by Freese and Freese (1965). Then, the virus was seeded into CKC cultures after appropriate dilution, and incubated at 31°C for plaque formation. Clones were isolated from well separated plaques and their growth at 40°C and 31°C was examined. About 1,800 clones were examined and 49 *ts* mutants were picked up. Preliminary results indicated that they did not form plaques when  $10^3$ - $10^5$  PFU were inoculated onto *ca*  $10^6$  cells and incubated at 40°C (Table 1). The ratio of infectious

TABLE 1.

Efficiency of Plating at 40°C to at 31°C	
<i>ts</i> 1	$<4.3 \times 10^{-5}$
<i>ts</i> 2	$<1.3 \times 10^{-5}$
<i>ts</i> 3	$<1.3 \times 10^{-4}$
<i>ts</i> 4	$<3.8 \times 10^{-5}$
<i>ts</i> 5	$<2.0 \times 10^{-4}$
<i>tt</i> 8	$<1.1 \times 10^{-4}$
<i>ts</i> 10	$<8.2 \times 10^{-5}$
<i>ts</i> 11	$<2.5 \times 10^{-6}$
A 26	$7.0 \times 10^{-1}$

TABLE 2.

Infectious Virus Yields at 40 C and 31 C			
mutant	40 C	31 C	Ratio : 31 C/40 C
ts 1	$1.0 \times 10^4$	$4.9 \times 10^7$	$4.9 \times 10^3$
ts 2	$2.1 \times 10^3$	$7.0 \times 10^7$	$3.3 \times 10^4$
ts 3	$2.0 \times 10^4$	$4.2 \times 10^7$	$2.1 \times 10^3$
tt 4	$5.2 \times 10^3$	$7.1 \times 10^7$	$1.4 \times 10^4$
ts 5	$2.3 \times 10^3$	$4.7 \times 10^7$	$2.0 \times 10^4$
ts 8	$2.0 \times 10^4$	$5.2 \times 10^7$	$2.6 \times 10^3$
ts 10	$1.7 \times 10^3$	$1.0 \times 10^8$	$5.9 \times 10^4$
ts 11	$7.1 \times 10^2$	$1.6 \times 10^8$	$2.3 \times 10^5$
A26	$5.4 \times 10^3$	$3.4 \times 10^7$	$6.3 \times 10^{-2}$

CKC cultures were infected by adding 10–20 PFU/cell and incubated at 40 C for 2 hr for adsorption. Then, cultures were washed once with fresh medium and incubated at 40 C or 31 C. Cultures incubated at 40 C were harvested 34 hr after infection, and cultures incubated at 31 C were harvested when the cytopathic change of infected cells was nearly complete

virus yield after one growth cycle of *ts* mutants at 31 C to that at 40 C was  $10^3$ – $10^5$  (Table 2).

Immunofluorescent studies on the viral antigen were done as follows. Partially purified virus was obtained from a lysate of CKC infected with A26 by treatment with fluorocarbon and centrifugation at 70,000×g for 90 min. The virus was emulsified with complete Freund's adjuvant and antiserum was prepared by infecting it intramuscularly into a rabbit.  $\gamma$ -Globulin conjugated with fluorescein isothiocyanate was prepared from the

serum as described by Shimojo et al. (1967). Coverslip cultures of CKC infected with one of the *ts* mutants (input multiplicity 10–20 PFU/cell) were fixed with acetone at –20 C for 15 min and allowed to react with the conjugated  $\gamma$ -globulin for 1 hr. The wild type strain, A26, accumulated viral antigen




FIGURE 1. Nuclear accumulation of viral antigen in infected with the wild type strain, A26, after 18 hr incubation at 40 C.



FIGURE 2. Nuclear accumulation of viral antigen in cells infected with ts8 after 68 hr incubation at 31 C.

TABLE 3. Classification of *ts* mutants on the basis of the localization of viral antigen in infected cells at the restrictive temperature.

Type of mutant	Location of viral antigen	Strain number of <i>ts</i> mutant	Total
N type	nucleus	<i>ts</i> 3, 7, 9, 12, 16, 18, 24, 25, 27, 31, 32, 33, 34, 36, 38, 40, 41, 42, 43, 45.	20
C type	cytoplasm	<i>ts</i> 5, 8, 10, 13, 19, 28, 29, 30, 49.	9
FA <sup>–</sup> type	neither in cytoplasm nor nucleus	<i>ts</i> 4, 6, 11, 14, 15, 17, 22, 44.	8

in the nucleus at either 40°C or 31°C (Fig. 1). At the non-restrictive temperature of 31°C, all the *ts* mutants accumulated viral antigen in the nucleus (Fig. 2), although some *ts* mutants retained some antigen in the cytoplasm during an early stage of incubation. As summarized in Table 3, *ts* mutants can be classified into 3

3a) had inclusion-like structures in the nucleus (Fig. 3b). However, this staining procedure does not allow such clear elucidation of intranuclear inclusions as hematoxylin-eosin staining after Carnoy's fixation. The third group (FA<sup>-</sup> type) produced no, or only slight antigen even after prolonged incubation at




FIGURE 3. a) Cytoplasmic retention of viral antigen in cells infected with *ts*8 after 25 hr incubation at 40°C.  
b) The area shown in a) was observed after Giemsa's staining.

groups on the basis of the localization of viral antigen in infected cells at the restrictive temperature of 40°C. The first group of *ts* mutants (N type) accumulated the antigen in the nucleus like the wild type strain. The second group (C type) retained viral antigen in the cytoplasm but not in the nucleus even after prolonged incubation (Fig. 3 and 4). By staining immunofluorescent samples with Giemsa's solution it was found that with *ts* 8 cells in which the cytoplasm showed fluorescence (Fig.




FIGURE 4. Cytoplasmic retention of viral antigen in cells infected with *ts*10 after 31 hr incubation at 40°C.

40°C.

When cytosine arabinoside, an inhibitor of DNA synthesis (10 or 40 µg/ml) was added at the time of virus inoculation, neither cells infected with A26 nor those infected with *ts* 8 showed viral antigen detectable by antibody-staining. This suggests that the antigen observed with the present antibody represented viral structural proteins produced after synthesis of viral DNA.

The above immunofluorescent study demonstrated an unexpected new class of *ts* mutants designated as the C type. At the restrictive temperature they seem to be defective in transportation from the cytoplasm to the nucleus of viral structural protein(s), claimed to be synthesized on cytoplasmic polysomes in a human adenovirus-KB cell system (Thomas and Green, 1966; Velicer and Ginsberg, 1968).

A more detailed report of this work, including results on complementation between *ts* mutants and on temperature-shift will appear elsewhere.

## ACKNOWLEDGEMENT

The author is grateful to Dr. H. Kawamura for his generous supply of the Ote strain and for technical

advice, and to Dr. J. Kawamata and Dr. K. Mutai for helpful discussion.

## REFERENCES

Dutta, S. K. and B. S. Pomeroy. 1963. Electron-microscopic structure of chicken embryo lethal orphan virus. *Proc. Soc. Exp. Biol. Med.* 114: 539-541.

Freese, E. and E. B. Freese. 1965. The oxygen effect on deoxyribonucleic acid inactivation by hydroxylamines. *Biochemistry* 4: 2419-2433.

Green, M., M. Piña, R. Kimes, P. C. Wensink, L. A. MacHattie and C. A. Thomas, Jr. 1967. Adenovirus DNA, I. Molecular weight and conformation. *Proc. Nat. Acad. Sci. U.S.* 57: 1302-1309.

Kawamura, H., T. Sato, H. Tsubahara and S. Isogai 1963. Isolation of CELO virus from chicken trachea. *Jap. Nat. Inst. Anim. Health Quart.* 3: 1-10.

Petek, M., B. Felluga and R. Zoleto. 1963a. Biological properties of CELO virus: stability to various agents, and electron-microscopic study. *Avian Dis.* 7: 38-49.

Petek, M., B. Felluga, R. Zoleto and G. Bersani. 1963b. Further studies on CELO virus: its relationship to the adenovirus group. *Arch. ges. Virusforsch.* 14: 637-649.

Shimojo, H., H. Yamamoto and C. Abe. 1967. Differentiation of adenovirus 12 antigens in cultured cells with immunofluorescent analysis. *Virology* 31: 748-752.

Thomas, D. C. and M. Green. 1966. Biochemical studies on adenovirus multiplication XI. Evidence of a cytoplasmic site for the synthesis of viral-coded proteins. *Proc. Nat. Acad. Sci. U.S.* 56: 243-246.

Velicer, L. F. and H. S. Ginsberg. 1968. Cytoplasmic synthesis of type 5 adenovirus capsid proteins. *Proc. Nat. Acad. Sci. U.S.* 61: 1264-1271.

Yates, V. J. and D. E. Fry. 1957. Observations on a chicken embryo lethal orphan (CELO) virus. *Amer. J. Vet. Res.* 18: 657-660.