

Title	Changes in Susceptibility of Staphylococcus aureus and Corynebacterium diphtheriae Cell Walls to Egg White Lysozyme, the L ₃ -and L ₁₁ -Enzymes Caused by Trichloroacetic Acid Treatment
Author(s)	Kotani, Shozo; Kato, Keijiro; Matsubara, Toshiro et al.
Citation	Biken journal : journal of Research Institute for Microbial Diseases. 1963, 6(4), p. 317-320
Version Type	VoR
URL	https://doi.org/10.18910/82985
rights	
Note	

Osaka University Knowledge Archive : OUKA

<https://ir.library.osaka-u.ac.jp/>

Osaka University

Changes in Susceptibility of *Staphylococcus aureus* and *Corynebacterium diphtheriae* Cell Walls to Egg White Lysozyme, the L₃- and L₁₁-Enzymes Caused by Trichloroacetic Acid Treatment

In the course of studies of the cell walls of some lysozyme-insensitive, Gram-positive bacteria, two cell wall lytic enzymes have been isolated in this laboratory. One of them, the *Flavobacterium* L₁₁ enzyme, exerts its lytic activity towards the cell walls of *Staphylococcus aureus*, strain Newman 1, and the other, the *Streptomyces* L₃ enzyme, is active against the walls of *Corynebacterium diphtheriae*, strain Park-Williams No. 8 (Kotani *et al.*, 1959a; Kotani *et al.*, 1959b; Mori *et al.*, 1960; Mori and Kotani, 1962; Kato *et al.*, 1962). It has been shown recently (Kato *et al.*, 1962; unpublished observations) that from the susceptible cell walls both the L₃- and L₁₁-enzymes liberate into solution compounds having free NH₂-terminal groups, without any significant release of reducing and/or hexosamine-reactive substances. There is, however, a distinct difference in the lytic activity ranges of these two enzymes against various bacterial species, as shown in Table 1.

Recent studies, on the other hand, demonstrated that whereas the whole walls of *S. aureus* were not attacked by egg white lysozyme, the residue, exhaustively extracted with trichloroacetic acid to remove ribitol teichoic acid, was dissolved by this enzyme (Mandelstam and Strominger, 1961; Morse, 1962; Kato *et al.*, 1962) and that removal of the rhamnose-glucosamine polymer from the cell walls of group A streptococci rendered them sensitive to lysozyme (Krause and McCarty, 1961).

Table 1. Lytic Activity Range of the L₃- and L₁₁-Enzymes

Test organisms	Cell wall lytic enzymes		
	L ₃ enzyme	L ₁₁ enzyme	Egg white lysozyme
<i>Staphylococcus aureus</i> (Newman 1)	—	+	—
<i>Corynebacterium diphtheriae</i> (Park-Williams No. 8)	+	—	—
BCG (Takeo)	—	—	—(+)*
<i>Streptococcus pyogenes</i> Group A (089)	—	—	—
<i>Micrococcus lysodeikticus</i> (2665)	—	+	+
<i>Sarcina lutea</i> ** (3232)	—	+	+
<i>Bacillus megaterium</i> ** (KM)	+	+	+

Constructed from the data of Mori *et al.* (1960) and Kato *et al.* (1962).

*While intact cells are resistant, isolated cell walls are partially lysed.

**Only intact cells are used as a substrate for the assay

Experiments reported here have been undertaken to see if the susceptibility

of *S. aureus* and *C. diphtheriae* cell walls to egg white lysozyme, the L₃- and L₁₁-enzymes is influenced by removal of their 'special structures' (Strominger, 1962) by trichloroacetic acid treatment.

A specimen of the cell walls of *C. diphtheriae*, Park-Williams No. 8, which had been thoroughly extracted with ethanol-ether (1:1, v/v) and chloroform, was suspended in 5 per cent trichloroacetic acid solution at a rate of 50 mg/ml and extracted in the cold for 24 hours, with stirring. The extraction was repeated in all eight times. By this treatment, cell wall components which were mainly composed of galactose, mannose and arabinose and amounted to about 16 per cent of the original walls, were released into solution (unpublished observations). Removal of a teichoic acid fraction from a cell wall preparation of *S. aureus*, Newman 1, was effected by suspending the preparation in 10 per cent trichloroacetic acid at a rate of 24 mg/ml and heating this suspension at 60°C for 6 hours. About 80 per cent of the phosphorus in the cell walls became soluble during the extraction. The treated cell walls of both organisms were thoroughly washed with distilled water by centrifugation to remove trichloroacetic acid.

The treated and non-treated cell walls were suspended in 0.01 M Na-phosphate buffer containing none or one of the lytic enzymes and 0.1 per cent sodium azide as preservative. Buffer of pH 6.8 was used in the assay with lysozyme and the L₁₁ enzyme, and the pH of the buffer was adjusted to 8.0 in systems containing the L₃ enzyme. The final optical densities of the reaction mixtures were about 0.5. The tubes (18±0.5 mm external diameter) containing the reaction mixtures were incubated at 37°C for 24 hours and changes in the optical density were followed at intervals with a Hitachi photoelectric colorimeter, Type EPO-B, using a No. 55 filter.

The following points will be apparent from the results summarized in Fig. 1: (1) Whereas the whole walls of *S. aureus* are not attacked by either lysozyme or the L₃ enzyme, the cell walls treated with hot trichloroacetic acid are dissolved by both enzymes. The rate of optical density reduction under the action of the L₃ enzyme is definitely slower than that with the same enzyme units of L₁₁ enzyme, but there is no difference between the final optical density reductions with the L₁₁- and L₃-enzymes. The per cent optical density reduction by lysozyme treatment, on the other hand, is nearly 20 per cent less than by the L₁₁- and L₃-enzymes. (2) After removal of the cell wall components, consisting mainly of hexoses and pentose, the walls of *C. diphtheriae* exhibit a definite increase in susceptibility to egg white lysozyme, but not to the L₁₁ enzyme. The maximum per cent optical density reduction by lysozyme treatment, however, is only 50 per cent as compared with 70 per cent in the reaction mixture containing L₃ enzyme. In this connection, it is worth mentioning that treatment of *C. diphtheriae* cell walls with 10 per cent trichloroacetic acid at 60°C for 18 hours rather decreased the susceptibility to the L₃ enzyme and that the treated cell walls were not attacked by either lysozyme or the L₁₁ enzyme.




Fig. 1. Changes in Susceptibility of *S. aureus* and *C. diphtheriae* Cell Walls to Egg White Lysozyme, the L₃- and L₁₁-Enzymes Caused by Trichloroacetic Acid Treatment

*C: no enzyme, Ly: egg white lysozyme, L₃: L₃ enzyme, and L₁₁: L₁₁ enzyme. Final concentrations of the lytic enzymes:

(A) and (B): lysozyme 0.05 mg/ml, L₃ enzyme 1.25 unit/ml, and L₁₁ enzyme 2.5 unit/ml.

(C) and (D): lysozyme 0.05 mg/ml, L₃ enzyme 1.25 unit/ml, and L₁₁ enzyme 1.25 unit/ml.

The reason why *C. diphtheriae* cell walls, which became sensitive to lysozyme by removal of the 'special structure', do not exhibit any sensitivity to the L₁₁ enzyme seems to deserve further investigation, in view of the facts that there is a close similarity between the attack points towards the susceptible cell walls of the L₁₁- and L₃-enzymes and that *S. aureus* cell walls, originally resistant to the L₃ enzyme, become sensitive to this enzyme by removal of the teichoic acid.

REFERENCES

- Kato, K., Kotani, S., Matsubara, T., Kogami, J., Hashimoto, S., Chimori, K. and Kazekawa, I. (1962). Lysis of *Staphylococcus aureus* cell walls by a lytic enzyme purified from culture supernatants of *Flavobacterium* species. *Biken J.* **5**, 155-179.
- Kotani, S., Hirano, T., Kitaura, T., Kato, K. and Matsubara, T. (1959a). Studies on the isolation of bacteria capable of lysing the cell walls of various lysozyme-resistant, pathogenic bacteria. *Biken's J.* **2**, 143-150.
- Kotani, S., Kato, K., Matsubara, T., Hirano, T. and Higashigawa, T. (1959b). *Staphylolytic* activity of a culture filtrate of a *Flavobacterium* species, isolated from soil. *Biken's J.* **2**, 211-213.
- Krause, R. M. and McCarty, M. (1961). Studies on the chemical structure of the streptococcal cell wall. I. The identification of a mucopeptide in the cell walls of groups A and A-variant streptococci. *J. Exptl. Med.* **114**, 127-140.
- Mandelstam, M. H. and Strominger, J. L. (1961). On the structure of the cell walls of *Staphylococcus aureus* (Copenhagen). *Biochem. Biophys. Res. Commun.* **5**, 466-471.
- Mori, Y., Kato, K., Matsubara, T. and Kotani, S. (1960). Lysis of the cell walls of *Corynebacterium diphtheriae* by extracellular enzymes of *Streptomyces sp.* I. Production and concentration of the lytic factor. *Biken's J.* **3**, 139-149.
- Mori, Y. and Kotani, S. (1962). Lysis of the cell walls of *Corynebacterium diphtheriae* by extracellular enzymes of *Streptomyces sp.* 2. Production of "protoplasts" by treatment with a purified cell wall lytic enzyme. *Biken J.* **5**, 55-66.
- Morse, I. (1962). Studies on the chemistry and immunochemistry of cell walls of *Staphylococcus aureus*. *J. Exptl. Med.* **116**, 229-245.
- Strominger, J. L. (1962). Biosynthesis of bacterial cell walls. In *The Bacteria. A Treatise on Structure and Function* (edited by Gunsalus, I. C. and Stanier, R. Y.), Vol. III: Biosynthesis, pp. 413-470. Academic Press, New York.

SHOZO KOTANI
 KEIJIRO KATO
 TOSHIRO MATSUBARA
 YUJI MORI
 MASAKAZU CHIMORI
 HIDEO KISHIDA

Department of Bacteriology
Nara Medical College
Kashiwara, Nara
(Received on November 28, 1963)