

Title	Interchangeability of Components between C ₃ in Human and Guinea Pig Complement
Author(s)	Inai, Shinya; Kishimoto, Susumu; Hirao, Fumio et al.
Citation	Biken's journal : journal of the Research Institute for Microbial Diseases. 1961, 4(2), p. 131-132
Version Type	VoR
URL	https://doi.org/10.18910/83061
rights	
Note	

Osaka University Knowledge Archive : OUKA

<https://ir.library.osaka-u.ac.jp/>

Osaka University

Interchangeability of Components between C'₃ in Human and Guinea pig Complement

There has been much progress by many workers in the isolation and purification of C'₃ in guinea pig serum. Recently, Rapp *et al.*^{1,2)} reported that C'₃ in guinea pig serum could be separated into two components either by chromatography or by precipitation with methanol and acid. Further, C'₃ has been isolated from human serum, although the preparation obtained was incapable of reacting with EAC'_{1,4,2} according to Pillemer³⁾.

We became interested in the serological characteristics of C'₃ through studies on the properdin system in 1956. In the course of properdin assay by zymosan, we happened to find that a combination of appropriate amounts of human R₃ and guinea pig R₃ caused hemolysis of sensitized sheep erythrocytes suspension (EA). This interesting fact leads us to a comparative study of human and guinea pig C'₃ components. The separation of the components of C'₃ was studied with human and guinea pig R₃ by chromatography by the method of Rapp *et al.*²⁾.

It was found that human C'₃ could be separated into two components C'a' and C'b' like those of guinea pig C'₃ (C'a, C'b). Further, human R₃ in combination with either the C'b of guinea pig serum or the C'b' of human serum caused hemolysis of sensitized erythrocytes, whereas it did not in combination with either C'a (guinea pig) or C'a' (human). On the other hand, guinea pig R₃ in combination with either C'a (guinea pig) or C'a' (human) caused hemolysis, whereas it did not in combination with either C'b (guinea pig) or C'b' (human).

This suggested that whereas human R₃ has been depleted of its C'b' rather than its C'a' component, guinea pig R₃ has been depleted of its C'a rather than its C'b component. Thus guinea pig R₃ in combination with human R₃ causes hemolysis of EA, because the combination forms a complete complement system.

Work is in progress on the manner in which these incomplete systems (R₃ human and R₃ guinea pig) serve as complements to one another.

REFERENCES

- 1) Rapp, H. J. (1959). Mechanism of immune hemolysis: Recognition of two steps in the conversion of EAC'_{1,4,2} to E.* *Science* **127**, 234-236.
- 2) Rapp, H. J., Sims, M. R. and Borsos, T. (1959). Separation of components of guinea pig complement by chromatography. *Proc. Soc. Exptl. Biol. Med.* **100**, 730-733.
- 3) Arday, F. R., Pillemer, L. and Lepow, I. H. (1959). The properdin system and immune hemolysis: Studies on the purification and properties of the third component of human complement. *J. Immunol.* **82**, 458-466.

* Editor's Note: After this article had been received, we happened to know that A. B. Taylor and M. A. Leon had reported more advanced results on C'₃ components in the Federation Proceedings (Vol. 20 Part I, p. 19, 1961), which arrived at this institute in the end of April.

SHINYA INAI
SUSUMU KISHIMOTO
FUMIO HIRAO
KAZUYOSHI NAGAKI
YOSHIHIKO SAIKI

*The Third Department of Internal Medicine,
Osaka University Medical School,
Osaka
Received on April 15, 1961*