

Title	The fundamental group of the smooth part of a log Fano variety
Author(s)	Zhang, D.-Q.
Citation	Osaka Journal of Mathematics. 32(3) P.637-P.644
Issue Date	1995
Text Version	publisher
URL	https://doi.org/10.18910/8455
DOI	10.18910/8455
rights	
Note	

Osaka University Knowledge Archive : OUKA

<https://ir.library.osaka-u.ac.jp/>

Osaka University

THE FUNDAMENTAL GROUP OF THE SMOOTH PART OF A LOG FANO VARIETY

D.-Q. ZHANG

(Received March 22, 1994)

Introduction

Let X be a normal projective variety over the complex number field C . We call X a *Fano variety* if X is \mathbf{Q} -Gorenstein and the anti-canonical divisor $-K_X$ is ample. A Fano variety X is said to be a *log Fano variety* if X has only log terminal singularities (cf. [6]). A Fano variety X is called a *canonical Fano variety* if X has only canonical singularities (cf. [6]). The *Cartier index* $c(X)$ is the smallest positive integer such that $c(X)K_X$ is a Cartier divisor. The *Fano index*, denoted by $r(X)$, is the largest positive rational number such that $-K_X \sim_{\mathbf{Q}} r(X)H$ (\mathbf{Q} -linear equivalence) for a Cartier divisor H .

This note consists of two sections. In §1, we shall consider canonical Fano 3-folds and prove the following :

Theorem 1. *Let X be a canonical Fano 3-fold. Let $X^\circ := X - \text{Sing } X$ be the smooth part of X . Assume that X has only isolated singularities. Then we have :*

- (1) *Suppose the Fano index $r(X)$ is 1. Then $\pi_1(X^\circ) = \mathbf{Z}/c(X)\mathbf{Z}$ (cf. Remark 1.1 in §1).*
- (2) *Suppose that the canonical divisor K_X is a Cartier divisor. Then X° is simply connected.*

REMARK. (1) The assumption that X has only isolated singularities is used to prove Lemma 1.3 in §1.

(2) Using the same proof (see §1) one can show that $\pi_1(X^\circ) = \mathbf{Z}/c(X)\mathbf{Z}$ when X is a log Fano variety of Fano index one and with only isolated singularities because even in this case the $\mathbf{Z}/c(X)\mathbf{Z}$ -covering Y constructed in §1 has only isolated canonical singularities.

In [19], we shall give a universal bound for $c(X)$. Under the much stronger condition that X has only terminal and cyclic quotient singularities, T. Sano

proved that $c(X) \leq 2$ (cf. [15]).

In §2, we shall consider n -dimensional log Fano varieties and prove the following :

Theorem 2. *Let X be a log Fano variety of Fano index $r(X) > \dim X - 2$. Let $X^\circ := X - \text{Sing } X$ be the smooth part of X . Then we have :*

- (1) *The fundamental group $\pi_1(X^\circ)$ of X° is a finite group.*
- (2) *Suppose that X has only canonical singularities. Then $\pi_1(X^\circ)$ is an abelian group of order ≤ 9 . (See [9] for the unique canonical Fano surface X with $|\pi_1(X^\circ)| = 9$.)*
- (3) *Suppose that $r(X) \geq \dim X - 1$. Then $\pi_1(X^\circ)$ is a finite abelian group generated by two elements, and has order ≤ 9 .*
- (4) *Suppose $r(X) > \dim X - 1$. Then the smooth part X° of X is simply connected.*

To prove Theorem 2, we manage to reduce to the dimension two case. In the dimension two case, Theorem 2(1) was proved in our joint works [3, 4] (see [18], and also [2] for a differential geometric proof), and Theorem 2(2) proved in [8, 9]. In this reduction process, in order to apply the crucial theorem due to Alexeev, we need the hypothesis about the Fano index $r(X)$. We hope this hypothesis can be eventually dropped.

Note that in Theorem 2(2) the hypothesis “ X has only canonical singularities” is necessary. Indeed, in [17] we have a lot of examples of log Fano surfaces X with exactly one triple point and several double points but with a non-abelian $\pi_1(X^\circ)$.

We want to remove the condition about the Fano index $r(X)$ in the above theorems and raise the following question :

Let X be a Fano variety. Let $X^\circ := X - \text{Sing } X$ be the smooth part of X . Suppose that X has only log-terminal (or canonical, or terminal) singularities. Is $\pi_1(X^\circ)$ a finite group ?

In general, it is not true that the smooth part of a rational variety with only log-terminal singularities has finite fundamental group. Just consider the example $(\mathbf{P}^1 \times E)/\tau$ in [3, §1.15] where E is an elliptic curve and τ is an involution acting on both \mathbf{P}^1 and E non-trivially and diagonally. So, the ampleness of $-K_X$ is an essential condition.

This work was started during the author’s stay in Max-Planck-Institut in Bonn. The author thanks very much to the institute and Professor F. Hirzebruch for the hospitality and financial support. The author also thanks the referee for very careful reading.

1. Proof of Theorem 1

We shall first show that Theorem 1(1) follows from Theorem 1(2). Let X be as in Theorem 1(1). We have a linear equivalence $-cK_X \sim cH$ where $c=c(X)$ is the Cartier index and H is a primitive Cartier divisor. Here we say that a Cartier divisor H on X is primitive if $\mathcal{O}(H)$ is not divisible in the Picard group of X . Let

$$Y := \text{Spec } \bigoplus_{i=0}^{c-1} \mathcal{O}_X(i(K_X + H)).$$

Then the natural $\mathbb{Z}/c\mathbb{Z}$ -covering morphism $f: Y \rightarrow X$ is etale over the smooth part $X^\circ := X - \text{Sing } X$ of X . Moreover, $-K_Y = -f^*K_X \sim f^*H$. So, Y is a (Gorenstein) Fano variety with only canonical singularities. Thus, the hypotheses in Theorem 1(2) are satisfied by Y . Note that $\pi_1(Y^\circ) = \pi_1(f^{-1}(X^\circ))$ because $Y^\circ - f^{-1}(X^\circ)$ has codimension ≥ 2 in the smooth 3-fold $Y^\circ := Y - \text{Sing } Y$. So, Theorem 1(1) will follow from Theorem 1(2). Moreover, we have

REMARK 1.1. $f^{-1}(X^\circ)$ is the universal covering of X° .

Now we shall prove Theorem 1(2). Let X be as in Theorem 1(2). We can write $-K_X = rH$ where $r=r(X) \in \mathbb{Z}_{>0}$ is the Fano index and H an ample Cartier divisor. Let $S \in |-K_X|$ be a general member. By [12], we know that S is a K3-surface possibly with rational double singularities. Let $\sigma: T \rightarrow S$ be a minimal resolution of singularities. Then T is a K3-surface. The first assertion of the following Lemma 1.2 is from [13] or [10, Theorem 5] and the second is a consequence of the first.

Lemma 1.2. *Let T be a K3-surface defined over \mathbb{C} . Let L be a nonzero numerically effective divisor on T . Then we have :*

- (1) *$|L|$ has base points if and only if there exist irreducible curves E, Γ , and an integer $k \geq 2$ such that $L \sim kE + \Gamma$, $(E^2) = 0$, $(\Gamma^2) = -2$, $E \cdot \Gamma = 1$. In this case, every member of $|\Gamma|$ is of the form $E' + \Gamma$, where E' is a sum of k effective divisors E_1, \dots, E_k such that $E_i \sim E$ for all i ; in particular, there is an elliptic fibration $\varphi: T \rightarrow \mathbb{P}^1$ such that E is a fiber and Γ is a cross-section.*
- (2) *$|sL|$ is base point free for all $s \geq 2$.*

We need the following lemma which is proved in [16, Theorem 0.5].

Lemma 1.3. (1) *The singular locus $\text{Sing } S$ of a general member $S \in |-K_X|$ contains $S \cap \text{Sing } X$.*

(2) *If $r(X) > 1$ then $|-K_X|$ is base point free. Hence a general member $S \in |-K_X|$ is disjoint from $\text{Sing } X$.*

(3) *Let $\text{Bs}|-K_X|$ be the base locus. If $\dim \text{Bs}|-K_X| = 1$ then $\text{Bs}|-K_X|$ is*

disjoint from $\text{Sing } X$. Hence a general member $S \in |-K_X|$ is disjoint from $\text{Sing } X$.

(4) If $\dim \text{Bs}|-K_X|=0$ then $P := \text{Bs}|-K_X|$ is a single point and P is a rational double point of S of Dynkin type A_1 . So, $S \cap \text{Sing } X = \emptyset$ or $\{P\}$. (Indeed, $S \cap \text{Sing } X = \{P\}$ (cf. [16])).

Proof. (1) follows from the condition that the divisor S is a Cartier divisor on X .

(2) and (3) are proved in [16]. Moreover, we have $\text{Bs}|-K_X| = \text{Bs}|(-K_X)_{|S}|$.

(4) By Lemma 1.2 and by $\dim \text{Bs}|(-K_X)_{|S}| (= \dim \text{Bs}|-K_X|) = 0$, we see that $|\sigma^*((-K_X)_{|S})| = |kE| + \Gamma$. Here E is a fiber of an elliptic fibration $\varphi : T \rightarrow \mathbf{P}^1$, Γ is a cross-section of φ and $P := \sigma(\Gamma)$ is a singularity on S with $\sigma^{-1}(P) = \Gamma$. So, P is a rational double point of S of Dynkin type A_1 . We have also $\text{Bs}|-K_X| = \{P\}$. (4) is proved.

Since the Kodaira D -dimension $\kappa(X, S) \geq 2$, the following natural homomorphism is surjective (see Theorem 2.4 below and the arguments after Theorem 2.4)

$$\pi_1(S - S \cap \text{Sing } X) \rightarrow \pi_1(X^\circ).$$

To finish the proof of Theorem 1(2), we need to prove that $\pi_1(S - S \cap \text{Sing } X) = (1)$. If $S \cap \text{Sing } X = \emptyset$, then $\pi_1(S) = (1)$ and we are done because S is a K3-surface possibly with rational double singularities. Suppose $P := S \cap \text{Sing } X \neq \emptyset$, then by Lemma 1.3, P is a single point and a rational double point of S of Dynkin type A_1 . Using the notations in Lemmas 1.2 and 1.3, we have $S - S \cap \text{Sing } X = S - \{P\} = T' - \Gamma$. Here T' is the resolution of the singularity P on S . Without loss of generality, we may assume that $T' = T$ and it suffices to prove the following lemma in order to finish the proof of Theorem 1(2).

Lemma 1.4. *Let T be a K3-surface. Let $\varphi : T \rightarrow \mathbf{P}^1$ be an elliptic fibration. Let Γ be a (-2) -curve which is a cross-section of φ . Then $T - \Gamma$ is simply connected.*

Proof. Consider the long cohomology exact sequence :

$$H^2(T, \mathbf{Z}) \rightarrow H^2(\Gamma, \mathbf{Z}) \rightarrow H^3(T, \Gamma; \mathbf{Z}) \rightarrow H^3(T, \mathbf{Z}).$$

Since T is simply connected we have $H^3(T, \mathbf{Z}) = 0$. Let E be a fiber of φ . Then $E \cdot \Gamma = 1$. So the map $H^2(T, \mathbf{Z}) \rightarrow H^2(\Gamma, \mathbf{Z})$ takes E to the generator. Thus, by the duality we have $H_1(T - \Gamma, \mathbf{Z}) = (0)$.

To prove Lemma 1.4, we have only to show that $\pi_1(T - \Gamma)$ is abelian. Let E_1 be a singular fiber of φ . Since the blowing-up of smooth points in the open surface $T - \Gamma$ does not affect the fundamental group of this open surface, one may assume that E_1 is simple normal crossing and hence E_1 consists of \mathbf{P}^1 's. Clearly, $\kappa(T,$

$E_1 + \Gamma = 2$. So, we can apply [11, Cor. 2.3, see Theorem 2.4 below]. Let U_1, U_2 be (open) tubular neighborhoods of sufficiently small radii in T of the curves E_1, Γ , respectively. Then $U_1 \cup U_2$ is an open tubular neighborhood of $E_1 + \Gamma$. Hence the following natural homomorphism is surjective :

$$\pi_1((U_1 - \Gamma \cap U_1) \cup (U_2 - \Gamma)) \rightarrow \pi_1(T - \Gamma).$$

Thus, to finish the proof of Lemma 1.4, we have only to prove that $\pi_1((U_1 - \Gamma \cap U_1) \cup (U_2 - \Gamma))$ is abelian. Applying Van-Kampen Theorem, we see easily that $\pi_1((U_1 - \Gamma \cap U_1) \cup (U_2 - \Gamma)) = \pi_1(U_1 - \Gamma \cap U_1)$. Indeed, $\pi_1(U_2 - \Gamma)$ is generated by a loop σ around Γ , and we may assume that σ is taken from $(U_1 - \Gamma \cap U_1) \cap (U_2 - \Gamma)$. We may even assume that σ is a loop in $E_1 - \Gamma \cap E_1$ around the point $\Gamma \cap E_1$. So, σ is contractible in $U_1 - \Gamma \cap U_1$ because $E_1 - \Gamma \cap E_1$ is a union of one A^1 and several P^1 's.

Now the proof of Lemma 1.4 is reduced to the proof that $\pi_1(U_1 - \Gamma \cap U_1)$ is abelian. Note that $\pi_1(U_1 - \Gamma \cap U_1) = \pi_1(E_1 - \text{the smooth point } \Gamma \cap E_1 \text{ of } E_1)$ because E_1 is a strong deformation retract of U_1 , and $\pi_1(E_1 - \text{the smooth point } \Gamma \cap E_1 \text{ of } E_1) = \pi_1(E_1)$ because E_1 is a divisor with simple normal crossing whose irreducible components are all isomorphic to P^1 and E_1 meets Γ transversally in a single point. Note that $\pi_1(E_1)$ is equal to (1) when E_1 is a tree of P^1 's, and equal to Z when E_1 is a simple loop of P^1 's. So $\pi_1(U_1 - \Gamma \cap U_1)$ is abelian.

The proof of Lemma 1.4 is completed. This proof does not work when Γ is a multiple section.

2. Proof of Theorem 2

Let X be a log Fano variety of dimension $d (d \geq 2)$ satisfying the hypothesis in Theorem 2. Write $-K_X \sim_q rH$ where r is a positive rational number such that $r > d - 2$ and H is an ample Cartier divisor. This is possible by the hypothesis in Theorem 2. We need the following :

Theorem 2.1 (cf. Theorem 0.5 in [1]). *Let X be a log Fano variety. With the above notations and assumptions, then $|H|$ is non-empty and base component free, and a general member X_{d-1} in $|H|$ is a normal projective variety with only log terminal singularities.*

By the adjunction formula, we have $-K_{X_{d-1}} \sim_q (r-1)H|_H$. Hence when $d \geq 3$, X_{d-1} is a log Fano variety by the definition and its Fano index $r(X_{d-1}) \geq r - 1 > \dim X_{d-1} - 2$.

Applying Alexeev's Theorem $(d-1)$ -times, we get a ladder below such that the assertions in the following lemma hold true :

$$(X_d = X, H_d = H), (X_{d-1}, H_{d-1}), \dots, (X_1, H_1).$$

Lemma 2.2. (1) *We have $X_i \in |H_{i+1}|$, $H_i = H_{i+1}|_{X_i}$ and $-K_{X_i} \sim_q (r+i-d)H_i$ for all $i \geq 1$. Moreover, X_i 's ($i \geq 2$) are log Fano varieties of Fano index $r(X_i) > \dim X_i - 2$.*

(2) *If $r = d - 1$, then X_1 is a nonsingular elliptic curve.*

(3) *If $r > d - 1$, then X_1 is a nonsingular rational curve.*

The following lemma is, though easy, crucial in order to reduce to the dimension two case.

Lemma 2.3. *Let X be a log Fano variety of dimension d ($d \geq 2$). With the notations and assumptions at the beginning of the section, we have :*

(1) *$|H|$ has at most isolated base points and the base locus $\text{Bs}|H|$ is contained in the smooth part X° of X .*

(2) *The singular locus $\text{Sing } X_{d-1}$ of X_{d-1} contains $X_{d-1} \cap \text{Sing } X$.*

(3) *If X has only canonical singularity then so does X_{d-1} .*

Proof. Note that (2) follows from the condition that X_{d-1} is a Cartier divisor on X . The assertion (3) follows from (1) and the definition of canonical singularity.

Now we shall prove (1). Consider the following exact sequence :

$$0 \rightarrow \mathcal{O}_X \rightarrow \mathcal{O}_X(H) \rightarrow \mathcal{O}_{X_{d-1}}(H) \rightarrow 0.$$

By Kawamata's vanishing theorem (cf. [6, Theorem 1-2-5]), we have $H^1(X, \mathcal{O}_X) = 0$ and hence we have a surjection :

$$(*) \quad H^0(X, \mathcal{O}_X(H)) \rightarrow H^0(X_{d-1}, \mathcal{O}_{X_{d-1}}(H)).$$

By the result (*), we have $\text{Bs}|H| = \text{Bs}|H_{d-1}|$ where $H_{d-1} := H|_{X_{d-1}}$. So if $|H_{d-1}|$ has at most isolated base points then the same is true for $|H|$. If a point P in $\text{Bs}|H_{d-1}|$ is a smooth point on X_{d-1} then P is also a smooth point on X because X_{d-1} is a Cartier divisor on X . Thus, we are reduced to prove a statement similar to (1) for X_{d-1} (cf. Lemma 2.2). By the same argument, we can reduce to prove (1) for X_2 . So to prove (1), we may assume that $\dim X = 2$.

By Alexeev's Theorem, we may assume that H is normal. Hence H is nonsingular because $\dim H = 1$. This, together with the condition that H is a Cartier divisor on X , implies that H is contained in the smooth part of X . Hence follows the second part of (1). Thus (1) follows because $|H|$ is base component free by Alexeev's Theorem.

Now we shall apply the following :

Theorem 2.4 (cf. [11, Cor. 2.3]). *Let \tilde{X} be a nonsingular projective variety. Let \tilde{H} be a divisor on \tilde{X} such that the Kodaira D -dimension $\kappa(\tilde{X}, \tilde{H}) \geq 2$. Let Δ be a Zariski-closed proper subset. Let U be any open tubular neighborhood of*

\tilde{H} in \tilde{X} . Then the following natural homomorphism is a surjection :

$$\pi_1(U - \Delta \cap U) \rightarrow \pi_1(\tilde{X} - \Delta).$$

Let $f : \tilde{X} \rightarrow X$ be a resolution of singularities such that $f^*X_{d-1} + \Delta$ is a normal crossing divisor. Here Δ is the exceptional divisor of f . By Lemma 2.3(2), $X_{d-1}^\circ := X_{d-1} - \text{Sing } X_{d-1}$ is a Zariski-open subset of $X_{d-1} - \text{Sing } X = f^*X_{d-1} - \Delta$. So one has a surjective homomorphism

$$\pi_1(X_{d-1}^\circ) \rightarrow \pi_1(f^*X_{d-1} - \Delta).$$

Applying Nori's Theorem to $\tilde{H} : = f^*X_{d-1}$, one obtains a surjective homomorphism :

$$\pi_1(f^*X_{d-1} - \Delta) \rightarrow \pi_1(X^\circ).$$

Combining these two surjections, we have proved the following Lemma 2.5 when $i = d - 1$. Applying the same arguments several times, one can prove Lemma 2.5 for all $i \geq 1$.

Lemma 2.5. *The natural homomorphism $\pi_1(X_i^\circ) \rightarrow \pi_1(X_{i+1}^\circ)$ is surjective for all $i \geq 1$.*

Now Theorem 2(1) follows from Lemma 2.5 because it is true in the dimension two case by [3, 4]. Theorem 2(2) follows from Lemma 2.5 and Lemma 2.3(3) since it is true in the dimension two case by [8, 9].

Theorem 2(4) follows from Lemma 2.5 and Lemma 2.2(3) because now $X_1^\circ = X_1 \cong \mathbf{P}^1$ and $\pi_1(X_1^\circ) = (1)$.

In view of Lemma 2.5 and Theorem 2(4), it suffices to prove Theorem 2(3) in the case where $d = \dim X = 2$ and $r(X) = d - 1 = 1$. It is easy to see that $H_1(X^\circ, \mathbf{Z})$ is finite (cf. e.g. [18, Lemma 1.3]). By Lemma 2.5, $\pi_1(X^\circ)$ is a surjective image of $\pi_1(X_1^\circ) = \mathbf{Z} \times \mathbf{Z}$ because $X_1^\circ = X_1$ is a nonsingular elliptic curve by Lemma 2.2(2). Theorem 2(3) is proved via [9] for X_2 is Gorenstein now by Th. 2.1.

We have proved Theorem 2 stated in the Introduction. Actually, the proofs for (3) and (4) of Theorem 2 are self-contained. In other words, we obtained a simpler proof for the result in [3, 4] when $r(X) \geq \dim X - 1 = 1$.

References

- [1] V.A. Alexseev : *Theorems about good divisors on log Fano varieties (Case of index $r > n - 2$)*, Lecture Notes in Mathematics, Vol. **1479**, Springer-Verlag (1989), 1-9.
- [2] A. Fujiki, R. Kobayashi and S. Lu : *On the fundamental group of certain open normal surfaces*, Saitama Math. J. **11** (1993), 15-20.
- [3] R.V. Gurjar and D.-Q. Zhang : *π_1 of smooth points of a log del Pezzo surface is finite : I, J.*

- Math. Sci. Univ. Tokyo, **1** (1994), 137-180.
- [4] R.V. Gurjar and D-Q. Zhang : π_1 of smooth points of a log del Pezzo surface is finite : II, J. Math. Sci. Univ. Tokyo **2** (1995).
- [5] Y. Kawamata : *Boundedness of \mathbb{Q} -Fano threefolds*, Contemporary Math. **131** (1989), Part 3, 439-445.
- [6] Y. Kawamata, K. Matsuda and K. Matsuki : *Introduction to the Minimal Model Problem*, Advanced Studies in Pure Math., Vol. **10**, Kinokuniya, North-Holland (1987), 283-360.
- [7] J. Kollar, Y. Miyaoka and S. Mori : *Rational connected varieties*, J.Alg. Geom. **1** (1992), 429-448.
- [8] M. Miyanishi and D.-Q. Zhang : *Gorenstein log del Pezzo surfaces of rank one*, J. Alg., **118** (1988), 63-84.
- [9] M. Miyanishi and D.-Q. Zhang : *Gorenstein log del Pezzo surfaces, II*, J. Alg., **156** (1993), 183-193.
- [10] S.Mori : *On degree and genera of curves on smooth quartic surfaces in \mathbb{P}^3* , Nagoya Math. J. **96** (1984), 127-132.
- [11] M. Nori : *Zariski's conjecture and related problems*, Ann. Sci. Ecole Norm. Sup. **16** (1983), 305-344.
- [12] Miles Reid : *Projective morphisms according to Kawamata*, Preprint 1983.
- [13] B. Saint-Donat : *Projective models of K3-surfaces*, Amer. J. Math. **96** (1974), 602-639.
- [14] T. Sano : *Classification of \mathbb{Q} -Fano d -folds of Fano index greater than $d-2$* , Univ. of Tokyo Preprint 1992.
- [15] T. Sano : *On classification of \mathbb{Q} -Fano 3-folds whose Fano indices are one*, Univ. of Tokyo Preprint 1992.
- [16] Kil-Ho Shin : *3-dimensional Fano varieties with canonical singularities*, Tokyo J. Math. **12** (1989), 375-385.
- [17] D.-Q. Zhang : *Logarithmic del Pezzo surfaces with rational double and triple singular points*, Tohoku Math. J. **41** (1989), 399-452.
- [18] D.-Q. Zhang : *Algebraic surfaces with nef and big anti-canonical divisor*, Math. Proc. Cambridge Philo. Soc. **117** (1995), 161-163.
- [19] D.-Q. Zhang : *Log Fano threefolds of Fano index one and quotients of K3-surfaces*, in : Proc. Pacific Rim Geometry Conf. Singapore, Dec. 1994, to appear.

Department of Mathematics
National University of Singapore
Kent Ridge
SINGAPORE 0511
e-mail : matz dq @ nus. sg