

Title	Toroidal compactifications and Borel-Serre compactifications
Author(s)	Kato, Kazuya; Nakayama, Chikara; Usui, Sampei
Citation	
Version Type	A0
URL	https://hdl.handle.net/11094/84809
rights	
Note	

Osaka University Knowledge Archive : OUKA

<https://ir.library.osaka-u.ac.jp/>

Osaka University

Toroidal compactifications and Borel–Serre compactifications

Kazuya Kato, Chikara Nakayama, Sampei Usui

July 26, 2021

Abstract

We discuss connections of toroidal compactifications and Borel–Serre compactifications in view of the fundamental diagram of extended period domains. We give a complement to a work of Goresky–Tai.

0 Introduction

In this paper, as an application of the fundamental diagram of extended period domains introduced in our series [5]–[9] of papers, we give a complement to the work of Goresky and Tai [2] on the relation of the toroidal compactification and the reductive Borel–Serre compactification.

Let G be a reductive algebraic group over \mathbf{Q} and let $h_0 : S_{\mathbf{C}/\mathbf{R}} \rightarrow G_{\mathbf{R}}$ be a homomorphism satisfying certain conditions which are necessary to consider the period domain D associated to (G, h_0) . Then the main part of the associated fundamental diagram in [9] is the following.

$$\begin{array}{ccccccc}
\Gamma \backslash D_{\Sigma, [\text{val}]} & \leftarrow & D_{\Sigma, [\text{val}]}^{\#} & \xrightarrow{\psi} & D_{\text{SL}(2), \text{val}} & \xrightarrow{\eta} & D_{\text{BS}, \text{val}} \\
\downarrow & & \downarrow & & \downarrow & & \downarrow \\
\Gamma \backslash D_{\Sigma, [\cdot]} & \leftarrow & D_{\Sigma, [\cdot]}^{\#} & \xrightarrow{\psi} & D_{\text{SL}(2)} & & D_{\text{BS}} \\
\downarrow & & \downarrow & & & & \downarrow \\
\Gamma \backslash D_{\Sigma} & \leftarrow & D_{\Sigma}^{\#} & & & & D_{\text{BS}}^{\text{red}}
\end{array}$$

Here the notation is as in [9] Section 4. In particular, Σ is a weak fan and Γ is a neat semi-arithmetic subgroup (1.5) of $G'(\mathbf{Q})$ which is strongly compatible with Σ . (G' denotes the commutator group of G .)

MSC2020: Primary 14A21; Secondary 14D07, 32G20

Keywords: Hodge theory, moduli, toroidal compactification, Borel–Serre compactification

TOROIDAL COMPACTIFICATIONS AND BOREL–SERRE COMPACTIFICATIONS

In this diagram, all maps are continuous. All vertical maps and the maps to the left direction are surjective and the targets of these maps have the quotient topologies. The vertical arrows except the map $D_{\text{BS}} \rightarrow D_{\text{BS}}^{\text{red}}$ are proper.

Assume further that (G, h_0) comes from some Shimura data and that Γ is an arithmetic subgroup (1.5). Then we have the following two kinds of compactifications of the Shimura variety $\Gamma \backslash D$. On one hand, there is a Σ such that $\Gamma \backslash D_\Sigma$ is compact and this is a toroidal compactification of $\Gamma \backslash D$. On the other hand, we have the reductive Borel–Serre compactification $\Gamma \backslash D_{\text{BS}}^{\text{red}}$ of $\Gamma \backslash D$, which is a quotient of the Borel–Serre compactification $\Gamma \backslash D_{\text{BS}}$ of $\Gamma \backslash D$.

Now, Goresky and Tai constructed a space which connects, after replacing Σ by a finer subdivision, these two compactifications of $\Gamma \backslash D$ in the following fashion. That is, they show that, after replacing Σ by a sufficiently finer subdivision Σ' , there are a compact topological space T which contains $\Gamma \backslash D$ as a dense open subspace, and continuous surjective maps $f : T \rightarrow \Gamma \backslash D_{\Sigma'}$ and $g : T \rightarrow \Gamma \backslash D_{\text{BS}}^{\text{red}}$ such that f is a homotopy equivalence and such that f and g induce the identity map of $\Gamma \backslash D$:

$$\Gamma \backslash D_{\Sigma'} \xleftarrow{\text{hom. eq.}} T \longrightarrow \Gamma \backslash D_{\text{BS}}^{\text{red}}.$$

From this, the existence of the canonical maps

$$H^m(\Gamma \backslash D_{\text{BS}}^{\text{red}}, A) \rightarrow H^m(\Gamma \backslash D_\Sigma, A)$$

with A being an abelian group for any m , A being a group for $m = 0, 1$, and A being a set for $m = 0$ follows.

In this paper, we have the following complement to the work of Goresky and Tai (cf. 3.3). As we have shown in [9] 3.8, under the above assumption, there is a unique continuous map $D_{\text{SL}(2)} \rightarrow D_{\text{BS}}$ which makes the above fundamental diagram still commutative, and the composition $D_{\Sigma, [\cdot]}^\# \xrightarrow{\psi} D_{\text{SL}(2)} \rightarrow D_{\text{BS}}$ induces the surjection of the quotient spaces $\Gamma \backslash D_{\Sigma, [\cdot]} \rightarrow \Gamma \backslash D_{\text{BS}}^{\text{red}}$. On the other hand, as we will prove in this paper, the map $\Gamma \backslash D_{\Sigma, [\cdot]} \rightarrow \Gamma \backslash D_\Sigma$ in the fundamental diagram is proper, surjective, and a weak homotopy equivalence (Theorem 3.2 (3)):

$$\Gamma \backslash D_\Sigma \xleftarrow{\text{weak hom. eq.}} \Gamma \backslash D_{\Sigma, [\cdot]} \longrightarrow \Gamma \backslash D_{\text{BS}}^{\text{red}}.$$

That is, compared to [2], we do not need a subdivision of Σ and our intermediate space is a concrete one, that is, $\Gamma \backslash D_{\Sigma, [\cdot]}$. Thus we can give an alternative construction of the above canonical maps between cohomologies (Corollary 3.4). (Probably $\Gamma \backslash D_{\Sigma, [\cdot]} \rightarrow \Gamma \backslash D_\Sigma$ is in fact a homotopy equivalence. For this, it is enough to see that $\Gamma \backslash D_{\Sigma, [\cdot]}$ is triangulizable, which is plausible, though we don't try to show it.)

Further, even not necessarily in the situation of Shimura data, we have

$$\Gamma \backslash D_\Sigma \xleftarrow{\text{same cohomologies}} \Gamma \backslash D_{\Sigma, [\text{val}]} \longrightarrow \Gamma \backslash D_{\text{BS}}^{\text{red}},$$

where the first map induces the isomorphisms of cohomologies including H^1 with non-abelian coefficients, and we still have the above canonical maps between cohomologies.

The organization of this paper is as follows. Section 1 is a preparation. In Section 2, we prove the basic result that the fibers of the natural projection from the ratio space

are weakly contractible. Based on this result, the complement for Goresky–Tai alluded in the above is an easy consequence of the existence of our fundamental diagram, which is explained in Section 3. Section 4 is a valuative version of the results in Section 2. Section 5 is a generalization of the results in Section 3 to the case where (G, h_0) does not necessarily come from Shimura data.

K. Kato was partially supported by NFS grants DMS 1303421, DMS 1601861, and DMS 2001182. C. Nakayama was partially supported by JSPS Grants-in-Aid for Scientific Research (B) 23340008, (C) 16K05093, and (C) 21K03199. S. Usui was partially supported by JSPS Grants-in-Aid for Scientific Research (B) 23340008 and (C) 17K05200.

1 Preliminaries

In this section, we gather terminology and some well-known facts used in the following sections.

1.1. In this paper, we say that a continuous map $f : X \rightarrow Y$ of topological spaces *induces the isomorphisms of cohomologies* if for any locally constant sheaf F of sets (resp. groups, resp. abelian groups) on Y , the induced map $H^m(Y, F) \rightarrow H^m(X, f^{-1}F)$ is an isomorphism for $m = 0$ (resp. $m = 0, 1$, resp. $m \in \mathbf{Z}$). Note that we include the nonabelian coefficients here, which is essential in the following.

1.2. A topological space is said to be *cohomologically contractible* if the map to a point induces the isomorphisms of cohomologies in the above sense, i.e., its cohomology groups with abelian and nonabelian coefficients are all trivial.

A topological space is said to be *weakly contractible* if the map to a point is a weak homotopy equivalence, i.e., its π_n ($n \geq 0$) are all trivial.

A topological space is said to be *locally contractible* if each point has a fundamental system of contractible open neighborhoods.

A continuous map $f : X \rightarrow Y$ of topological spaces is said to be *proper* if it is universally closed and separated.

A topological space is said to be *compact* if the map to a point is proper. Hence a compact space is Hausdorff.

1.3. By the Hurewicz theorem, a continuous map f of locally contractible spaces induces the isomorphisms of cohomologies if and only if it is a weak homotopy equivalence (cf. [12] Chapter II, Section 3, Proposition 4).

In particular, a locally contractible space is cohomologically contractible if and only if it is weakly contractible.

1.4. A proper map $f : X \rightarrow Y$ induces the isomorphisms of cohomologies if each fiber is cohomologically contractible. In fact, let F be as in 1.1. By the proper base change theorem ([10] Theorem A2.1), $R^q f_* G$, where $G = f^{-1}F$ coincides with F if $q = 0$ and vanishes if otherwise. Hence f induces the isomorphisms of cohomologies by Leray spectral sequences and the exact sequence of pointed sets $H^1(Y, f_* G) \rightarrow H^1(X, G) \rightarrow H^0(Y, R^1 f_* G)$, where the first map is injective.

1.5. Let G be a linear algebraic group over \mathbf{Q} and let Γ be a subgroup of $G(\mathbf{Q})$. Then Γ is called an *arithmetic subgroup* of $G(\mathbf{Q})$ if there are $n \geq 1$ and an injective homomorphism $\rho : G \rightarrow \mathrm{GL}(n)$ such that Γ is a subgroup of $\{g \in G(\mathbf{Q}) \mid \rho(g) \in \mathrm{GL}(n, \mathbf{Z})\}$ of finite index. As in [9], we call Γ a *semi-arithmetic subgroup* of $G(\mathbf{Q})$ if there are $n \geq 1$ and an injective homomorphism $\rho : G \rightarrow \mathrm{GL}(n)$ such that $\rho(\Gamma) \subset \mathrm{GL}(n, \mathbf{Z})$.

2 Weak contractibility

In this section, we prove that the projection from the space of ratios is a weak homotopy equivalence.

2.1. Let S be a locally ringed space over $E = \mathbf{R}$ or \mathbf{C} with an fs log structure M_S satisfying the following conditions (i) and (ii).

(i) For every $s \in S$, the natural homomorphism $E \rightarrow \mathcal{O}_{S,s}/m_s$ is an isomorphism, where m_s is the maximal ideal of $\mathcal{O}_{S,s}$.

(ii) For every open set U of S and for every $f \in \mathcal{O}_S(U)$, the map $U \rightarrow E ; s \mapsto f(s)$ is continuous. Here $f(s)$ is the image of f in $\mathcal{O}_{S,s}/m_s = E$.

Then we have a topological space $S_{[\cdot]}$ over S , called the *space of ratios*. See [8] 4.2 and [9] 4.3.

In this section we will prove the following theorem, which is the basic result in this paper.

Theorem 2.2. *Let S be as above. Then the following hold:*

- (1) *All fibers of $S_{[\cdot]} \rightarrow S$ are weakly contractible and locally contractible.*
- (2) *The map $S_{[\cdot]} \rightarrow S$ induces the isomorphisms of cohomologies (1.1).*

Since $S_{[\cdot]} \rightarrow S$ is proper, by 1.3 and 1.4, (2) is reduced to (1).

(1) follows from the following theorem whose proof occupies the rest of this section. Note that the fiber of $S_{[\cdot]} \rightarrow S$ on $s \in S$ is identified with the space of ratios $R(\mathcal{S})$ of the sharp fs monoid $\mathcal{S} = (M_S/\mathcal{O}_S^\times)_s$, which is defined in [8] 4.1.

Theorem 2.3. *For a sharp fs monoid \mathcal{S} , the space of ratios $R(\mathcal{S})$ is weakly contractible and locally contractible.*

2.4. Let $\Phi = \{\mathcal{S}^{(i)} \mid 0 \leq i \leq n\}$, where $\mathcal{S}^{(i)}$ are faces of \mathcal{S} such that $\mathcal{S} = \mathcal{S}^{(0)} \supseteq \mathcal{S}^{(1)} \supseteq \dots \supseteq \mathcal{S}^{(n)} = \{1\}$. Then by [8] Corollary 4.2.17, the Φ -part $R(\mathcal{S})(\Phi)$ of $R(\mathcal{S})$, which is an open set of $R(\mathcal{S})$, has the following description as a topological space. For $0 \leq i \leq n-1$, take $a_i \in \mathcal{S}^{(i)}$ such that $a_i \notin \mathcal{S}^{(i+1)}$. Then $R(\mathcal{S})(\Phi)$ is homeomorphic to the subspace of $\prod_{i=0}^{n-1} \mathrm{Hom}(\mathcal{S}^{(i)}, \mathbf{R}_{\geq 0}^{\mathrm{add}})$ consisting of all elements $N = (N_i)_{0 \leq i \leq n-1}$ satisfying the following conditions (i), (ii), and (iii).

- (i) $N_i(a_i) = 1$ for $0 \leq i \leq n-1$.
 - (ii) For $0 \leq i \leq n-1$, the kernel of N_i coincides with $\mathcal{S}^{(j)}$ for some j such that $i < j \leq n$.
 - (iii) If $0 \leq i < j \leq n-1$, the restriction of N_i to $\mathcal{S}^{(j)}$ coincides with $N_i(a_j)N_j$.
- We will identify $R(\mathcal{S})(\Phi)$ with the set of these N .

2.5. Let $R(\mathcal{S})(\Phi)_{>0}$ be the subset of $R(\mathcal{S})(\Phi)$ consisting of all $(N_i)_{0 \leq i \leq n-1}$ such that $\text{Ker}(N_i) = 0$ for all i (this is equivalent to $\text{Ker}(N_0) = 0$).

Let H be the set of all homomorphisms $h : \mathcal{S} \rightarrow \mathbf{R}_{\geq 0}^{\text{add}}$ such that $\text{Ker}(h) = 0$. We have a surjection

$$\pi : H \rightarrow R(\mathcal{S})(\Phi)_{>0}$$

given by $h \mapsto N$, where N_i is the restriction of $h(a_i)^{-1}h$ to $\mathcal{S}^{(i)}$. For $h \in H$, we have $h = h(a_0)\pi(h)_0$. Hence π induces a homeomorphism $\{h \in H \mid h(a_0) = 1\} \rightarrow R(\mathcal{S})(\Phi)_{>0}$, whose inverse map is given by $N \mapsto N_0$.

2.6. For $0 \leq i \leq n-1$, fix a homomorphism $p_i : \mathcal{S} \rightarrow \mathcal{S}_{\mathbf{Q}_{\geq 0}}^{(i)}$ whose restriction to $\mathcal{S}^{(i)}$ is the inclusion map $\mathcal{S}^{(i)} \rightarrow \mathcal{S}_{\mathbf{Q}_{\geq 0}}^{(i)}$. The existence of such p_i is seen by the next lemma, which is probably well-known.

For an $N \in R(\mathcal{S})(\Phi)$ and a real number $t > 0$, define

$$\theta_t(N) := \sum_{i=0}^{n-1} t^i N_i \circ p_i \in H.$$

Lemma 2.7. *Let \mathcal{S} be a sharp fs monoid and let \mathcal{S}' be a face of \mathcal{S} . Then there is a homomorphism $\mathcal{S} \rightarrow \mathcal{S}'_{\mathbf{Q}_{\geq 0}}$ whose restriction to \mathcal{S}' is the inclusion map $\mathcal{S}' \rightarrow \mathcal{S}'_{\mathbf{Q}_{\geq 0}}$.*

Proof. By the induction on $\dim \mathcal{S}'$, we may assume that $\dim \mathcal{S}' = \dim \mathcal{S} - 1$. Take a direct decomposition $\mathcal{S}_{\mathbf{Q}} = \mathcal{S}'_{\mathbf{Q}} \oplus \mathbf{Q}$ with the first and the second projection being p and l respectively such that $l(\mathcal{S}) \subset \mathbf{Q}_{\geq 0}$. Let $a \in \mathcal{S}'$ and consider the homomorphism $p_a : \mathcal{S} \rightarrow \mathcal{S}'_{\mathbf{Q}}$; $x \mapsto p(x) + l(x)a$ (the semi-group law is written additively here). Then $p_a(x) = x$ for $x \in \mathcal{S}'$. Hence it is sufficient to prove that there is an a such that $p_a(\mathcal{S}) \subset \mathcal{S}'_{\mathbf{Q}_{\geq 0}}$.

Let x_1, \dots, x_n be a set of generators of \mathcal{S} . If $x_j \in \mathcal{S}'$, $p_a(x_j) = x_j \in \mathcal{S}'$. If $x_j \notin \mathcal{S}'$, $l(x_j) > 0$ so that if a is an inner point of \mathcal{S}' , $l(x_j)a$ is an inner point of $\mathcal{S}'_{\mathbf{Q}_{\geq 0}}$. Replacing a by ca for $c \gg 0$, we have $p_a(x_j) = p(x_j) + l(x_j)a \in \mathcal{S}'_{\mathbf{Q}_{\geq 0}}$ for all j . \square

Lemma 2.8. *$R(\mathcal{S})(\Phi)$ is contractible.*

Proof. Fix an $L \in H$. Consider the continuous map

$$f : R(\mathcal{S})(\Phi) \times (0, 1] \rightarrow R(\mathcal{S})(\Phi) \ ; \ f(N, t) = \pi(t^n L + (1-t)\theta_t(N)).$$

Then $f(N, 1) = \pi(L)$. It is sufficient to prove that f extends to a continuous map $f : R(\mathcal{S})(\Phi) \times [0, 1] \rightarrow R(\mathcal{S})(\Phi)$ such that $f(N, 0) = N$. Assume $M \in R(\mathcal{S})(\Phi)$ converges to $N \in R(\mathcal{S})(\Phi)$ and $t \in (0, 1)$ converges to 0. We prove that $f(M, t)$ converges to N . Let $0 \leq i \leq n-1$. Then the restriction of $t^n L + (1-t)\theta_t(M)$ to $\mathcal{S}^{(i)}$ coincides with $t^i(b(t)M_i + E(t))$, where $b(t) = (1-t) \sum_{k=0}^i t^{k-i} M_k(a_i) \geq (1-t)M_i(a_i) = 1-t$ and $E(t)$ is the restriction of $t^{n-i}L + (1-t) \sum_{k=i+1}^{n-1} t^{k-i} M_k \circ p_k$ to $\mathcal{S}^{(i)}$. Hence $\pi(t^n L + (1-t)\theta_t(M))_i = (M_i + b(t)^{-1}E(t))(1 + b(t)^{-1}E(t)(a_i))^{-1}$. This converges to N_i since $b(t) \geq 1-t$ and since M_i and $E(t)$ converge to N_i and 0, respectively. \square

2.9. We prove that $R(\mathcal{S})$ is weakly and locally contractible. First, by [9] Proposition 4.2.19, $R(\mathcal{S})$ is a topological manifold with boundaries. In particular, it is locally contractible. Next, $R(\mathcal{S})$ has an open covering by $\{R(\mathcal{S})(\Phi)\}_\Phi$, and for any Φ_1 and Φ_2 , we have $R(\mathcal{S})(\Phi_1) \cap R(\mathcal{S})(\Phi_2) = R(\mathcal{S})(\Phi_1 \cap \Phi_2)$. Together with this, Lemma 2.8 implies that $R(\mathcal{S})$ is cohomologically contractible. Hence, by 1.3, $R(\mathcal{S})$ is weakly contractible.

Remark 2.10. It is plausible that $R(\mathcal{S})$ is in fact contractible and even is a homeomorphic to a closed ball. At least, $R(\mathcal{S})$ is a compact Hausdorff topological manifold with boundaries whose interior is homeomorphic to an open ball. But the authors do not know if this implies that $R(\mathcal{S})$ is homeomorphic to a closed ball.

3 On the work of Goresky and Tai

3.1. Let G be a reductive algebraic group over \mathbf{Q} . Let $h_0 : S_{\mathbf{C}/\mathbf{R}} \rightarrow G_{\mathbf{R}}$ be a homomorphism as in [9] 1.2.13. Assume that h_0 is \mathbf{R} -polarizable ([9] 1.5.2). Let $D = D(G, h_0)$ be the period domain associated to (G, h_0) . Let Σ be a weak fan in $\text{Lie}(G')$ ([9] 4.1.7), where G' denotes the commutator group of G . Let Γ be a neat semi-arithmetic subgroup (1.5) of $G'(\mathbf{Q})$. Assume that Σ and Γ are strongly compatible.

Below, the *Shimura data case* ([1] 1.5) means the case where the Hodge type of $\text{Lie}(G_{\mathbf{R}})$ via h_0 is in $\{(1, -1), (0, 0), (-1, 1)\}$ (as in Shimura data).

Theorem 3.2. *Let the notation and the assumptions be as above. Then the following hold:*

- (1) *All fibers of $\Gamma \backslash D_{\Sigma, [\cdot]} \rightarrow \Gamma \backslash D_{\Sigma}$ are weakly contractible and locally contractible.*
- (2) *The map $\Gamma \backslash D_{\Sigma, [\cdot]} \rightarrow \Gamma \backslash D_{\Sigma}$ induces the isomorphisms of cohomologies (1.1).*
- (3) *In the Shimura data case (3.1), $\Gamma \backslash D_{\Sigma, [\cdot]} \rightarrow \Gamma \backslash D_{\Sigma}$ is also a weak homotopy equivalence.*

Proof. (1) is by Theorem 2.2 (1). (2) follows from (1) by 1.3 and 1.4. Then, by 1.3, to see (3), it is enough to show that, in the Shimura data case, $\Gamma \backslash D_{\Sigma, [\cdot]}$ is locally triangulizable so that locally contractible. Take a local chart of the fs log complex analytic space $\Gamma \backslash D_{\Sigma}$. Then, by [8] Proposition 4.2.14, $\Gamma \backslash D_{\Sigma, [\cdot]}$ is locally a fiber product of two analytic maps over a real manifold with singular corners. Hence it is locally a semianalytic subset of an Euclidean space so that locally triangulizable by a theorem of Hironaka. \square

3.3. As is explained in Introduction, this Theorem 3.2 gives a new interpretation to the main result of the work of Goresky and Tai on the relation between the toroidal compactification and the reductive Borel–Serre compactification, which we will explain.

Let the reductive Borel–Serre space $D_{\text{BS}}^{\text{red}}$ be the quotient of D_{BS} by the following equivalence relation. For $p_1 = (P_1, Z_1), p_2 = (P_2, Z_2) \in D_{\text{BS}}$, $p_1 \sim p_2$ if and only if $P_1 = P_2$ and $P_{1,u}Z_1 = P_{2,u}Z_2$, where $P_{i,u}$ denotes the unipotent radical of the parabolic subgroup P_i ($i = 1, 2$).

In the case where D is a Griffiths domain [3], this $D_{\text{BS}}^{\text{red}}$ is denoted by D_{BS}^b in [10] Section 9. If Γ is semi-arithmetic (resp. arithmetic), then $\Gamma \backslash D_{\text{BS}}^{\text{red}}$ is Hausdorff (resp. compact). The proof is similar to that in [10] Section 9.2.

In the Shimura data case, for an arithmetic Γ , $\Gamma \backslash D_{\text{BS}}^{\text{red}}$ is the well-known reductive Borel–Serre compactification of the symmetric domain $\Gamma \backslash D$ studied in [13].

Now assume that we are in the Shimura data case. Then we have the natural morphism $D_{\text{SL}(2)} \rightarrow D_{\text{BS}}$ in the fundamental diagram ([9] Theorem 3.8.2) and hence the CKS map $D_{\Sigma, [\cdot]}^{\#} \rightarrow D_{\text{SL}(2)}$ induces a continuous map $D_{\Sigma, [\cdot]}^{\#} \rightarrow D_{\text{BS}}$, which induces a continuous map $\Gamma \backslash D_{\Sigma, [\cdot]} \rightarrow \Gamma \backslash D_{\text{BS}}^{\text{red}}$.

On the other hand, the map $\Gamma \backslash D_{\Sigma, [\cdot]} \rightarrow \Gamma \backslash D_{\Sigma}$ is a weak homotopy equivalence (Theorem 3.2 (3)). Thus the formation of these maps are regarded as a variant of the construction by Goresky and Tai.

Corollary 3.4. *Let A be a set (resp. a group, resp. an abelian group). Then, in the Shimura data case (3.1), there is a canonical map*

$$H^m(\Gamma \backslash D_{\text{BS}}^{\text{red}}, A) \rightarrow H^m(\Gamma \backslash D_{\Sigma}, A)$$

for $m = 0$ (resp. $m = 0, 1$, resp. $m \in \mathbf{Z}$).

4 Valuative version

In this section, we consider the valuative version of the results in Section 2.

Theorem 4.1. *Let S be either an object of $\mathcal{B}'_{\mathbf{R}}(\log)$ ([6] 3.1, [8] 1.3, [9] 3.4.1) or an fs log topological space whose structural sheaf of rings is the sheaf of all continuous \mathbf{R} -valued functions. Let S_{val} (resp. $S_{[\text{val}]}$) be the topological space defined as in [8] 3.1.2 and [9] 4.4 (resp. [8] 4.3 and [9] 4.7). Then the following hold:*

- (1) *All fibers of $S_{\text{val}} \rightarrow S$ and all fibers of $S_{[\text{val}]} \rightarrow S$ are cohomologically contractible.*
- (2) *The map $S_{\text{val}} \rightarrow S$ and the map $S_{[\text{val}]} \rightarrow S$ induce the isomorphisms of cohomologies (1.1).*

Proof. Since $S_{\text{val}} \rightarrow S$ and $S_{[\text{val}]} \rightarrow S$ are proper, (2) is reduced to (1) by 1.4. Because $S_{[\text{val}]} \rightarrow S$ is the composition $S_{[\text{val}]} \rightarrow S_{[\cdot]} \rightarrow S$ and $S_{[\text{val}]} = (S_{[\cdot]})_{\text{val}}$ for the new log structure of $S_{[\cdot]}$, the $S_{[\text{val}]}$ case is reduced to the S_{val} case by Theorem 2.2 and the proper base change theorem.

Thus it suffices to show the S_{val} case of (1). It is sufficient to prove that for $f, g \in \mathcal{S}$, the log blowing-up B of S by (f, g) is contractible, where \mathcal{S} is a sharp fs monoid and S is a point with the ring \mathbf{R} and the log structure defined by the local homomorphism $\mathcal{S} \rightarrow \mathbf{R}_{\geq 0}$ (cf. [4] p.311). If either $f/g \in \mathcal{S}$ or $g/f \in \mathcal{S}$, $B = S$ and hence B is contractible. Otherwise, B is homeomorphic to the interval $[0, \infty]$ and hence B is contractible. \square

Remark 4.2. The S_{val} part of (1) gives an alternative proof of Theorem 2.2 without constructing homotopies, which we sketch here. It is enough to show that $S_{\text{val}} \rightarrow S_{[\cdot]}$ induces the isomorphisms of cohomologies, where S is as in the last paragraph in the proof of the theorem. By 1.4, it is enough to show that any fiber X of the last map is cohomologically contractible. This X is the closure of a stratum Y of S_{val} . For each log modification S' of S , the closure of the image of Y in S' is homeomorphic to a closed ball, and X is the inverse limit of these closures. Hence X is cohomologically contractible.

The above proof of Theorem 4.1 shows the following.

Theorem 4.3. *For an S as in Theorem 4.1, for every log modification of S , all fibers are cohomologically contractible.*

Remark 4.4. The fibers in Theorem 4.3 are in fact contractible because they are triangulizable by the theory of toric varieties. See [11]. One can ask if it is homeomorphic to a closed ball.

5 Relation to extended period domains

We can generalize 3.3 to the non-classical (without Shimura data) situation, where we may not have $D_{\mathrm{SL}(2)} \rightarrow D_{\mathrm{BS}}$. Here we only assume that G is reductive.

5.1. By Theorem 4.1 (1), all fibers of

$$\Gamma \backslash D_{\Sigma, [\mathrm{val}]} \rightarrow \Gamma \backslash D_{\Sigma, [\cdot]} \rightarrow \Gamma \backslash D_{\Sigma}$$

are cohomologically contractible. Hence, by 1.4, this map induces the isomorphisms of cohomologies (1.1).

5.2. On the other hand, we have continuous maps

$$\Gamma \backslash D_{\Sigma, [\mathrm{val}]} \rightarrow \Gamma \backslash D_{\mathrm{BS}}^{\mathrm{red}}$$

induced by the continuous map

$$D_{\Sigma, [\mathrm{val}]}^{\sharp} \rightarrow D_{\mathrm{SL}(2), \mathrm{val}} \rightarrow D_{\mathrm{BS}, \mathrm{val}} \rightarrow D_{\mathrm{BS}} \rightarrow D_{\mathrm{BS}}^{\mathrm{red}}$$

in the fundamental diagram.

Corollary 5.3. *Let A be a set (resp. a group, resp. an abelian group). Then, there is a canonical map*

$$H^m(\Gamma \backslash D_{\mathrm{BS}}^{\mathrm{red}}, A) \rightarrow H^m(\Gamma \backslash D_{\Sigma}, A)$$

for $m = 0$ (resp. $m = 0, 1$, resp. $m \in \mathbf{Z}$).

References

- [1] P. Deligne, *Travaux de Shimura*, Séminaire Bourbaki (1970/71), Exp. 389, Lecture Notes in Math., **244**, Springer (1971), 123–165.
- [2] M. Goresky and Y. Tai, *Toroidal and reductive Borel–Serre compactifications of locally symmetric spaces*, Amer. J. Math. **121** (5) (1999), 1095–1151.
- [3] P. Griffiths, *Periods of integrals on algebraic manifolds. I. Construction and properties of modular varieties*, Amer. J. Math. **90** (2) (1968), 568–626.

- [4] T. Kajiwara and C. Nakayama, *Higher direct images of local systems in log Betti cohomology*, J. Math. Sci. Univ. Tokyo **15** (2) (2008), 291–323.
- [5] K. Kato, C. Nakayama and S. Usui, *Classifying spaces of degenerating mixed Hodge structures*, I: Borel–Serre spaces, Adv. Stud. Pure Math. **54** (2009), 187–222.
- [6] K. Kato, C. Nakayama and S. Usui, *Classifying spaces of degenerating mixed Hodge structures*, II: Spaces of $SL(2)$ -orbits, Kyoto J. Math. **51** (1): Nagata Memorial Issue (2011), 149–261.
- [7] K. Kato, C. Nakayama and S. Usui, *Classifying spaces of degenerating mixed Hodge structures*, III: Spaces of nilpotent orbits, J. Algebraic Geometry **22** (2013), 671–772.
- [8] K. Kato, C. Nakayama and S. Usui, *Classifying spaces of degenerating mixed Hodge structures*, IV: The fundamental diagram, Kyoto J. Math. **58** (2018), 289–426.
- [9] K. Kato, C. Nakayama and S. Usui, *Classifying spaces of degenerating mixed Hodge structures*, V: Extended period domains and algebraic groups, preprint, submitted. <https://arxiv.org/abs/2107.03561>.
- [10] K. Kato and S. Usui, *Classifying spaces of degenerating polarized Hodge structures*, Ann. Math. Studies **169**, Princeton University Press (2009).
- [11] C. Nakayama, *Base change theorems for log analytic spaces*, preprint, submitted.
- [12] D. G. Quillen, *Homotopical algebra*, Lecture Notes in Mathematics, 43, Springer-Verlag, Berlin, New York, 1967.
- [13] S. Zucker, *L_2 cohomology of warped products and arithmetic groups*, Invent. Math. **70** (1982), 169–218.

Kazuya KATO
 Department of mathematics
 University of Chicago
 Chicago, Illinois, 60637, USA
kkato@math.uchicago.edu

Chikara NAKAYAMA
 Department of Economics
 Hitotsubashi University
 2-1 Naka, Kunitachi, Tokyo 186-8601, Japan
c.nakayama@r.hit-u.ac.jp

Sampei USUI
 Graduate School of Science
 Osaka University
 Toyonaka, Osaka, 560-0043, Japan
usui@math.sci.osaka-u.ac.jp