



Title	An elementary construction of the representations of $SL(2, GF(q))$
Author(s)	Silberger, Allan J.
Citation	Osaka Journal of Mathematics. 1969, 6(2), p. 329-338
Version Type	VoR
URL	<a href="https://doi.org/10.18910/8521">https://doi.org/10.18910/8521</a>
rights	
Note	

*The University of Osaka Institutional Knowledge Archive : OUKA*

<https://ir.library.osaka-u.ac.jp/>

The University of Osaka

## AN ELEMENTARY CONSTRUCTION OF THE REPRESENTATIONS OF $SL(2, GF(q))$

ALLAN J. SILBERGER

(Received February 10, 1969)

### 1. Introduction

Let  $GF(q)$  be a field containing  $q$  elements,  $q$  odd. Let  $\mathcal{G}$  denote  $GL(2, GF(q))$ , the group of non-singular two-by-two matrices with entries in  $GF(q)$ , and let  $G$  denote  $SL(2, GF(q))$ , the subgroup of  $\mathcal{G}$  consisting of matrices with determinant one. In this paper, assuming a knowledge of certain of the characters of  $\mathcal{G}$ , we construct all the irreducible unitary representations of  $G$ . Our construction involves essentially no technique beyond the theory of induced representations and the orthogonality relations on a finite group. For a similarly elementary computation of the characters of  $\mathcal{G}$  we refer the reader to [3]. In future papers we shall generalize the methods employed in this paper to construct the representations of the  $n \times n$  matrix groups  $GL(n, GF(q))$  and  $SL(n, GF(q))$ .

Kloosterman [2] was the first to describe all the irreducible matrix representations of  $SL(2, GF(q))$ . Weil in [5] generalizes and gives an alternative construction for Kloosterman's representations. In [4] Tanaka uses Weil's theory to construct representations and presents a complete and unified description of the representations of  $G$ . We also mention the paper [1] of Gelfand-Graev, which classifies but does not detail the actual construction of all the representations of  $G$ .

### 2. The representations

Let  $B$  be the upper unipotent,  $D$  the diagonal, and  $T$  the upper triangular subgroups of  $G$ . Then  $T = DB$ .  $G$  has order  $q(q^2 - 1)$  and contains an abelian subgroup  $R$  (unique up to conjugacy) of order  $q + 1$ . Except for plus-or-minus the identity of  $G$  elements of  $R$  have characteristic roots in  $GF(q^2) - GF(q)$ .  $R$  is isomorphic to the subgroup of  $GF(q^2)^\times$  comprised of elements of norm one.

The  $q + 4$  equivalence classes of irreducible representations of  $G$  break up roughly into two main classifications. The  $\frac{1}{2}(q + 5)$  representations of the

“principal series” all contain  $B$ -invariant vectors. Those  $\frac{1}{2}(q+3)$  inequivalent representations which do not contain  $B$ -invariant vectors we call discrete series.

More precisely, the principal series include:

- (1) The trivial representation of degree 1,  $U \equiv 1$ ;
- (2) A  $q$ -dimensional representation  $U_1^1$  which occurs with  $U \equiv 1$  in the induced representation  $\text{ind}_{T \uparrow G} 1$ ;
- (3)  $\frac{1}{2}(q-3)$  irreducible induced representations  $U^\alpha = \text{ind}_{T \uparrow G} \alpha$ , where  $\alpha$  is a one-dimensional representation of  $T$  which is not real-valued.  $U^\alpha$  has degree  $q+1$  and  $U^{\alpha'}$  is equivalent to  $U^\alpha$  if and only if  $\alpha' = \alpha$  or  $\alpha^{-1}$ .
- (4) Let  $\alpha = \text{sgn}$ , where  $\text{sgn} \equiv 1$  and  $\text{sgn}^2 \equiv 1$ . Then  $\text{ind}_{T \uparrow G} \text{sgn} = U^{\text{sgn}} = U_1^{\text{sgn}} + U_2^{\text{sgn}}$ , the direct sum of two inequivalent irreducible representations, each of degree  $\frac{1}{2}(q+1)$ .

The discrete series are as follows:

- (5) If  $\pi$  is a non-trivial character of  $R$ , then there is a representation  $U^\pi$  of  $G$  of degree  $q-1$  associated with  $\pi$ .  $U^\pi$  is characterized by the fact that it does *not* occur in  $\text{ind}_{R \uparrow G} \pi$ .  $U^\pi$  is irreducible if and only if  $\pi$  is not real-valued.  $U^\pi$  is equivalent to  $U^{\pi'}$  if and only if  $\pi' = \pi$  or  $\pi^{-1}$ , so there are  $\frac{1}{2}(q-1)$  inequivalent irreducible representations of degree  $q-1$ .
- (6) If  $\pi \equiv 1$ ,  $\pi^2 \equiv 1$ , then  $U^\pi = U_1^\pi + U_2^\pi$ , the direct sum of inequivalent representations of degree  $\frac{1}{2}(q-1)$ .

### 3. The construction of principal series

The construction of the representations of the principal series as induced representations is well-known. For completeness we discuss this problem in detail.

Let  $\alpha$  be a one-dimensional representation of  $T$ . Since  $B$  is the commutator subgroup of  $T$ ,  $\alpha(btb') = \alpha(t)$  for any  $b$  and  $b' \in B$  and  $t \in T$ .  $T/B$  is canonically  $D$ , so  $\alpha$  is the extension to  $T$  of a character of the abelian group  $D$ . The mapping which identifies  $d \in D$  with its upper diagonal entry regarded as an element of the multiplicative group  $GF(q)^\times$  is an isomorphism. In this section, when convenient, we regard  $\alpha$  as a function on  $GF(q)^\times$  via this identification. Let  $U^\alpha$  denote the representation of  $G$  induced from  $\alpha$ .

By the definition of  $U^\alpha$ ,  $G$  acts by right translation in the space  $V^\alpha$  which consists of complex-valued functions  $\psi$  on  $G$  satisfying

$$(3.1) \quad \psi(tg) = \alpha(t)\psi(g)$$

for all  $t \in T$  and  $g \in G$ . Any such function is determined by its restriction to a set of representatives of  $T \backslash G$ . Since two matrices in  $G$  with the same lower entries differ only by a left factor in  $B$ ,  $\psi \in V^\alpha$  implies  $\psi(g) = \psi(g_{21}, g_{22})$ ,  $g_{21}$  and

$g_{22}$  the lower entries of  $g \in G$ . Equation (3.1) entails

$$(3.2) \quad \psi(d^{-1}g_{21}, d^{-1}g_{22}) = \alpha(d)\psi(g_{21}, g_{22})$$

for  $d \in GF(q)^\times$ ,  $g_{21}$  and  $g_{22}$  as before, so  $\psi$  is actually determined by its values, which may be chosen arbitrarily, on a set of representatives for the projective line over  $GF(q)$ .

**Theorem 3.1.** *Let  $\alpha$  be a one-dimensional representation of  $T$ . Let  $U^\alpha$  be the representation of  $G$  induced from  $\alpha$ .  $U^\alpha$  is right translation in the space  $V^\alpha$  defined by relations (3.1) and (3.2).*

- (1) *The degree of  $U^\alpha$  is  $q+1$ .*
- (2)  *$U^\alpha$  is irreducible if and only if  $\alpha^2 \neq 1$ .*
- (3)  *$U^{\alpha'}$  is equivalent to  $U^\alpha$  if and only if  $\alpha' = \alpha$  or  $\alpha^{-1}$ .*
- (4)  *$U^1$  decomposes into the direct sum of an irreducible representation of degree  $q$  and the unique one-dimensional representation of  $G$ .*
- (5)  *$U^{\text{sgn}}$ , where  $\text{sgn} \neq 1$  but  $\text{sgn}^2 = 1$ , decomposes into the direct sum of two inequivalent representations of degree  $\frac{1}{2}(q+1)$ .*

*Proof.*

- (1) A set of representatives for the projective line over  $GF(q)$  (e.g.  $\{(0, 1), (-1, z) | z \in GF(q)\}$ ) has cardinality  $q+1$ . In view of the above remarks this proves that  $V^\alpha$  has dimension  $q+1$ .
- (2) The proofs of the remaining parts of this theorem depend upon an analysis of the commuting algebra of  $U^\alpha$ .

Let  $C^\alpha$  be the convolution algebra of all complex-valued functions  $f$  on  $G$  satisfying  $f(tgt') = \alpha(tt')f(g)$  for any  $t, t' \in T$  and  $g \in G$ . Then  $U^\alpha(g_0)(f * \psi) = f * U^\alpha(g_0)\psi$  for any  $g_0 \in G$  and  $\psi \in V^\alpha$ , since  $f \in C^\alpha$  acting from the left by convolution keeps  $V^\alpha$  stable and commutes with right translation. Frobenius' reciprocity theorem says precisely that  $C^\alpha$  is large enough to be the full commuting algebra of  $U^\alpha$ .

$f \in C^\alpha$  is determined by its values on a set of representatives for the double cosets  $T \backslash G / T$ , e.g.  $\left\{ e = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}, w = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} \right\}$ . Clearly,  $\dim C^\alpha \leq 2$ .  $\dim C^\alpha = 2$  if and only if  $f(w) \neq 0$  for some  $f \in C^\alpha$ , if and only if  $\alpha(t)f(w) = f(tw) = f(wt^{-1}) = \alpha^{-1}(t)f(w)$  for all  $t \in D$ . Thus  $\dim C^\alpha = 2$  if and only if  $\alpha^2(t) \equiv 1$ , so (2) is true.

(3) The space of intertwining operators between  $V^\alpha$  and  $V^{\alpha'}$ ,  $\alpha \neq \alpha'$ , is canonically the vector space  $T^{\alpha, \alpha'}$  of complex-valued functions on  $G$  satisfying  $f(tgt') = \alpha(t)f(g)\alpha'(t')$  for all  $t, t' \in T$  and  $g \in G$ . It is spanned by any function  $f$  which satisfies  $\alpha(t)f(w) = f(w)\alpha'(t^{-1})$  for all  $t \in D$ .  $f(w) \neq 0$  implies  $\alpha' = \alpha^{-1}$ .

(4)  $V^1$  contains the constant functions on  $G$  as a stable subspace. The orthogonal complement of this one dimensional module must be an irreducible  $q$ -dimensional representation space for  $G$ .

(5) By the analysis in (2) we know that  $U^{\text{sgn}}$  decomposes into the direct sum of two inequivalent representations,  $U_1^{\text{sgn}} + U_2^{\text{sgn}} = U^{\text{sgn}}$ . By Frobenius' reciprocity theorem  $\text{res}_{G \downarrow T} U_\nu^{\text{sgn}}$ , for  $\nu=1$  or  $2$ , contains  $\text{sgn}$  and no other one-dimensional representation of  $T$ . Since  $G/\{\pm e\}$  is a simple group,  $G$  has no non-trivial one-dimensional representations. Therefore, Lemma (4.3) implies that the degree of  $U_\nu^{\text{sgn}}$  is  $\frac{1}{2}(q+1)$ ,  $\nu=1$  or  $2$ .

REMARK. To complete our description of the representations of the principal series we need to be more specific about the  $G$ -stable subspaces  $V_1^{\text{sgn}}$  and  $V_2^{\text{sgn}}$  of  $V^{\text{sgn}}$ . Set  $\phi(-1, z) = \Phi(z)$  for  $z \in GF(q)$ , where  $\Phi$  is an additive character of  $GF(q)$ ; let  $\phi(0, 1) = 0$ . Then  $\phi$  extends uniquely to a function in  $V^{\text{sgn}}$  and  $U^{\text{sgn}}(b(u))\phi = \Phi(u)\phi$ , where  $u$  is the super diagonal entry of  $b(u) \in B$ . Moreover,  $U^{\text{sgn}}(d)\phi(-1, z) = \text{sgn}(d)\phi(-1, d^{-2}z) = \text{sgn}(d)\Phi(d^{-2}z)$  for all  $z \in GF(q)$ ,  $d \in D$  (identified with  $GF(q)^\times$ );  $U^{\text{sgn}}(d)\phi(0, 1) = 0$ . Let  $\Phi \neq 1$ . Then the  $\frac{1}{2}(q-1)$  functions  $\phi'$  which correspond to characters  $\Phi'$  such that  $\Phi'(d^{-2}z) = \Phi(z)$  for some  $d \in GF(q)^\times$  belong to  $V_\nu^{\text{sgn}}$ ; the other non-trivial additive characters of  $GF(q)$  must correspond to elements of  $V_{\nu'}^{\text{sgn}}$ ,  $1 \leq \nu \neq \nu' \leq 2$ .  $V_\nu^{\text{sgn}}$  also contains a vector  $\psi$  satisfying  $\psi(tgt') = \text{sgn}(tt')\psi(g)$  for all  $t, t' \in T$ ,  $g \in G$ . In fact  $\psi$  may be chosen to be an idempotent in  $C^{\text{sgn}}$ .

**Proposition 3.2.** *Set*

$$\begin{aligned} \psi(g) &= \frac{1}{2} \frac{|G|}{|T|} \text{sgn}(t), \quad \text{if } g = t \in T; \\ &= \frac{1}{2} \frac{|G|}{|T|} (q \text{sgn}(-1))^{-1/2} \text{sgn}(t), \end{aligned}$$

if  $g = twb$ , with  $t \in T$ ,  $b \in B$ , and  $w = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}$ .

Then  $\psi$  is an idempotent in the algebra  $C^{\text{sgn}}$ . There are two choices  $\psi, \psi'$  depending on the sign of  $[\text{sgn}(-1)]^{1/2}$ . Clearly,  $\psi + \psi'$  is the identity in  $C^{\text{sgn}}$ . The function  $\phi$  defined in the preceding remark and corresponding to the non-trivial character  $\Phi$  of  $GF(q)$  belongs to the same  $G$ -irreducible subspace of  $V^{\text{sgn}}$  as  $\psi$  if and only if  $\sum_{x \neq 0} \text{sgn}(x)\Phi(-x) = [q \text{sgn}(-1)]^{1/2}$  (with the same choice for the sign of the right hand side as in the definition of  $\psi$ ).

Proof. To show that  $\psi$  is an idempotent in  $C^{\text{sgn}}$  it suffices to show that  $\psi * \psi(e) = \psi(e)$  and  $\psi * \psi(w) = \psi(w)$ . We have

$$\begin{aligned} \psi * \psi(g) &= \frac{1}{|G|} \sum_{x \in G} \psi(x) \psi(x^{-1}g) = \frac{|T|}{|G|} \sum_{x \in G/T} \psi(x) \psi(x^{-1}g) \\ &= \frac{|T|}{|G|} \{ \psi(e) \psi(g) + \sum_{u \in GF(q)} \psi(w) \psi(w^{-1}b^{-1}(u)g) \}. \end{aligned}$$

Therefore,

$$\begin{aligned}\psi * \psi(e) &= \frac{|T|}{|G|} \cdot \frac{1}{4} \frac{|G|^2}{|T|^2} \{1 + [q \operatorname{sgn}(-1)]^{-1} \operatorname{sgn}(-1)q\} \\ &= \psi(e). \\ \psi * \psi(w) &= \frac{|T|}{|G|} \cdot \frac{1}{4} \frac{|G|^2}{|T|^2} \{[q \operatorname{sgn}(-1)]^{-1/2} + [q \operatorname{sgn}(-1)]^{-1/2} \\ &\quad + [q \operatorname{sgn}(-1)]^{-1} \sum_{\substack{u \in GF(q) \\ u \neq 0}} \operatorname{sgn}(-u)\}.\end{aligned}$$

The last term on the right, being a character sum, is zero. It arises from the relation

$$\begin{aligned} * \quad w^{-1}b^{-1}(u)w &= \begin{vmatrix} 0 & -1 \\ 1 & 0 \end{vmatrix} \begin{vmatrix} 1 & -u \\ 0 & 1 \end{vmatrix} \begin{vmatrix} 0 & 1 \\ -1 & 0 \end{vmatrix} \\ &= \begin{vmatrix} -u^{-1} & -1 \\ 0 & -u \end{vmatrix} \begin{vmatrix} 0 & 1 \\ -1 & 0 \end{vmatrix} \begin{vmatrix} 1 & u^{-1} \\ 0 & 1 \end{vmatrix}, \quad \text{if } u \neq 0.\end{aligned}$$

Thus,  $\psi * \psi(w) = \psi(w)$ .

Finally, since  $\psi$  is a minimal idempotent in  $C^{\operatorname{sgn}}$ ,  $\psi * \phi = \phi$ , if  $\psi \in V^{\operatorname{sgn}}$  and  $\phi \in V^{\operatorname{sgn}}$ . If  $\phi \notin V^{\operatorname{sgn}}$ , then  $\phi \in V^{\operatorname{sgn}}$ , so  $\psi * \phi = 0$ .

$$\begin{aligned}\psi * \phi(w) &= \frac{|T|}{|G|} \sum_{x \in G/T} \psi(x) \phi(x^{-1}w) \\ &= \frac{|T|}{|G|} \{ \psi(e) \phi(w) + \psi(w) \sum_{u \in GF(q)} \phi(w^{-1}b^{-1}(u)w) \}.\end{aligned}$$

Using relation  $*$  as well as the definitions of  $\psi$  and  $\phi$ , we obtain

$$\psi * \phi(w) = \phi(w) \left\{ \frac{1}{2} + \frac{1}{2} [q \operatorname{sgn}(-1)]^{-1/2} \sum_{u \neq 0} \operatorname{sgn}(-u) \Phi(u) \right\},$$

which implies the last part of the proposition.

#### 4. The construction of discrete series for $GL(2, GF(q))$

Let  $\Pi$  be a character of  $GF(q^2)^\times$  whose restriction to the elements of norm one is non-trivial. Then  $\Pi$  corresponds to a representation  $\mathcal{U}^\Pi$  of the discrete series of  $\mathcal{G} = GL(2, GF(q))$  (i.e.  $\operatorname{res}_{\mathcal{G} \downarrow B} \mathcal{U}^\Pi \neq 1$ ). It turns out that  $\operatorname{res}_{\mathcal{G} \downarrow \mathcal{I}} \mathcal{U}^\Pi$  is an irreducible representation of  $\mathcal{I}$ , the triangle subgroup of  $\mathcal{G}$ . To determine a space of functions which transforms under  $\mathcal{G}$  as  $\mathcal{U}^\Pi$  we find an irreducible representation  $m$  of  $\mathcal{I}$  such that  $m = \operatorname{res}_{\mathcal{G} \downarrow \mathcal{I}} \mathcal{U}^\Pi$ . Then, using the trace of  $\mathcal{U}^\Pi$  (which we assume known) we extend the matrix coefficients of  $m$  to  $\mathcal{G}$ . To determine the discrete series of  $G$  we study  $\operatorname{res}_{\mathcal{G} \downarrow G} \mathcal{U}^\Pi$ .

Let  $\mathcal{D}$  be the diagonal subgroup of  $\mathcal{G}$  and let  $\alpha$  be a character of  $\mathcal{D}$ . Ind  $\alpha$   $\mathcal{D} \uparrow \mathcal{I}$   $= M^\alpha$  is right translation in the space of complex-valued functions on  $\mathcal{I}$

which satisfy  $\psi(dt) = \alpha(d)\psi(t)$  for all  $d \in \mathcal{D}$  and  $t \in \mathcal{I}$ . Since  $B$  represents  $\mathcal{D} \setminus \mathcal{I}$ , we may consider  $M^\alpha$  as acting in a vector space  $B^\alpha$  of complex-valued functions on  $B$ . We write  $\psi \in B^\alpha$  as a function of the super diagonal entries of elements of  $B$ . Then

$$(4.1) \quad M^\alpha(db(u))\psi(x) = \alpha(d)\psi(d_{11}^{-1}d_{22}x+u)$$

for any  $d \in \mathcal{D}$  and  $b(u)$  the element of  $B$  with superdiagonal entry  $u \in GF(q)$ ,  $d_{11}$  and  $d_{22}$  the non-zero entries of  $d$ .

To see how  $M^\alpha$  decomposes take as an orthonormal basis of  $B^\alpha$  the  $q$  characters of  $B$ . The operators  $M^\alpha(b)$  for  $b \in B$  obviously diagonalize with respect to this basis. Let  $\Phi_0$  be the trivial character of  $B$ . Clearly  $\Phi_0$  transforms under  $M^\alpha$  as the one-dimensional representation  $\alpha$  of  $\mathcal{I}$ . Now let  $\Phi$  be a fixed non-trivial character of  $B$ . For  $i \in GF(q)^\times$  set  $\Phi_i(x) = \Phi(ix)$  for all  $x \in GF(q)$ . Then  $\Phi_i$  is a non-trivial character of  $B$  and every non-trivial character of  $B$  is of the form  $\Phi_i$  for some  $i \in GF(q)^\times$ . (4.1) entails that, except for scalar factors,  $\mathcal{D}$  acts transitively on the non-trivial characters of  $B$ . Since  $M^\alpha$  is completely reducible, we see that the  $(q-1)$ -dimensional subspace of  $B^\alpha$  spanned by the non-trivial characters of  $B$  must be irreducible. Call the resulting representation  $m_\alpha$ .

**Lemma 4.1.** *An irreducible representation of  $\mathcal{I}$  is either of degree one or  $q-1$ . An irreducible  $(q-1)$ -dimensional representation of  $\mathcal{I}$  is determined by its restriction to the center of  $\mathcal{I}$ .*

Proof. If an irreducible representation of  $\mathcal{I}$  is not one-dimensional, it is equivalent to a representation  $m_\alpha$  for some character  $\alpha$  of  $\mathcal{D}$ . Thus it is  $(q-1)$ -dimensional. By Frobenius' reciprocity theorem characters  $\alpha'$  which occur in  $\text{res}_{\mathcal{I} \downarrow \mathcal{D}} m_\alpha$  occur with multiplicity one. Since  $m_\alpha$  is irreducible, every  $\alpha'$  contained in  $\text{res}_{\mathcal{I} \downarrow \mathcal{D}} m_\alpha$  must have the same values on the center of  $\mathcal{I}$  (i.e. the scalars). There are  $q-1$  distinct characters of  $\mathcal{D}$  which agree on the scalars, so they must all occur in  $\text{res}_{\mathcal{I} \downarrow \mathcal{D}} m_\alpha$ . By Frobenius' theorem,  $m_\alpha$  is equivalent to  $m_{\alpha'}$ , for all such  $\alpha'$ .

**Lemma 4.2.** *Let  $\Phi$  be a non-trivial character of  $B$ . For  $i \in GF(q)^\times$  set  $\Phi_i(x) = \Phi(ix)$  for all  $x \in GF(q)$  (considered as super-diagonal entries of elements of  $B$ ). The matrix coefficients of the representation  $m_\alpha$  with respect to the basis for  $B^\alpha$  consisting of the  $q-1$  non-trivial characters  $\{\Phi_i\}_{i \in GF(q)^\times}$  of  $B$  are the  $(q-1)^2$  functions*

$$(4.2) \quad \begin{aligned} m_{ij}^\alpha(t) &= \langle m_\alpha(t)\Phi_j, \Phi_i \rangle, \quad i \text{ and } j \in GF(q)^\times, \\ &= \alpha(d)\Phi_j(u), \quad \text{if } \Phi_j(d_{11}^{-1}d_{22}x) = \Phi_i(x) \text{ for all } x \in GF(q); \\ &= 0, \quad \text{otherwise.} \end{aligned}$$

In (4.2)  $t=db(u)$ , where  $u \in GF(q)$  is the super-diagonal entry of the matrix  $b(u) \in B$  and  $d_{11}$  and  $d_{22}$  are the diagonal entries of  $d \in \mathcal{D}$ .

Proof. Immediate from equation (4.1).

**Lemma 4.3.** *Let  $m_\alpha$  be an irreducible representation of  $\mathcal{I}$  of degree  $q-1$ . Then  $\text{res}_{\mathcal{I} \downarrow B} m_\alpha$  decomposes into inequivalent representations of degree  $\frac{1}{2}(q-1)$ . Any  $\mathcal{I} \downarrow T$  irreducible representation of  $T$  is either one-dimensional or  $\frac{1}{2}(q-1)$ -dimensional.*

Proof.  $\text{Res}_{\mathcal{I} \downarrow B} m_\alpha$  decomposes simply; if  $\text{res}_{\mathcal{I} \downarrow T} m_\alpha$  decomposes, the component representations must be inequivalent. By (4.1)  $M^\alpha(d)\Phi(x) = \alpha(d)\Phi(d_{11}^{-1}x)$  for  $d \in D$ , so two characters  $\Phi$  and  $\Phi'$  of  $B$  occur in the restriction to  $B$  of the same irreducible subrepresentation of  $\text{res}_{\mathcal{I} \downarrow T} m_\alpha$  if and only if  $\Phi'(x) = \Phi(a^2x)$  for some  $a \in GF(q)^\times$  and all  $x \in GF(q)$ . Since half the characters of  $B$  satisfy this relation and half do not,  $\text{res}_{\mathcal{I} \downarrow T} m_\alpha$  contains two irreducible representations, each of degree  $\frac{1}{2}(q-1)$ . The last statement in Lemma (4.3) follows from the fact that any irreducible representation of  $T$  occurs in the restriction to  $T$  of some irreducible representation of  $\mathcal{I}$ .

**Lemma 4.4.** *Let  $G$  be a finite group and  $H$  a subgroup of  $G$ . Let  $U$  be a unitary representation of  $G$  whose degree is  $d$  and character is  $X$ . Assume  $\text{res}_{G \downarrow H} U$  is irreducible. Then, for any matrix coefficient  $u_{ij}$  of  $U$ ,  $1 \leq i, j \leq d$ , and any  $g \in G$*

$$u_{ij}(g) = \frac{d}{|H|} \sum_{h \in H} u_{ij}(h) X(h^{-1}g).$$

Proof.

$$\begin{aligned} \frac{d}{|H|} \sum_{h \in H} u_{ij}(h) X(h^{-1}g) &= \frac{d}{|H|} \sum_{h \in H} u_{ij}(h) \sum_{k=1}^d u_{k\bar{k}}(h^{-1}g) \\ &= \frac{d}{|H|} \sum_{h \in H} u_{ij}(h) \sum_{k=1}^d \sum_{l=1}^d u_{kl}(h^{-1}) u_{l\bar{k}}(g) \\ &= \frac{d}{|H|} \sum_{l, \bar{k}} u_{l\bar{k}}(g) \sum_{h \in H} u_{ij}(h) \bar{u}_{l\bar{k}}(h) \\ &= u_{ij}(g), \end{aligned}$$

by Schur's orthogonality relations on  $G$ .

Lemma (4.1) implies that for any representation  $\mathcal{U}^\pi$  of the discrete series of  $\mathcal{G}$ ,  $\text{res}_{\mathcal{G} \downarrow \mathcal{I}} \mathcal{U}^\pi$  is equivalent to an irreducible representation  $m_\alpha$ , where  $m_\alpha$  is, up to equivalence, the unique irreducible  $(q-1)$ -dimensional representation of  $\mathcal{I}$  which agrees with  $\mathcal{U}^\pi$  on the scalars. Since  $\mathcal{G} = \mathcal{I} \cup \mathcal{I}wB$ ,  $w = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}$ , it



suffices to compute the matrix coefficients for  $\mathcal{U}^\Pi$  at  $w$  in order to extend them from  $\mathcal{I}$  to all of  $\mathcal{G}$ . For this purpose we need the character  $X^\Pi$  of  $\mathcal{U}^\Pi$  (To find directions for the easy computation of  $X^\Pi$  consult [3], p. 227.). Figure 1 presents  $X^\Pi$ .

Conjugacy Classes on $G$	Values of $X^\Pi$
$\lambda \begin{vmatrix} 1 & 0 \\ 0 & 1 \end{vmatrix}$	$\Pi(\lambda)(q-1)$
$\lambda \begin{vmatrix} 1 & 0 \\ 1 & 1 \end{vmatrix}$	$-\Pi(\lambda)$
$\lambda \begin{vmatrix} t & 0 \\ 0 & 1 \end{vmatrix}^*$	0
$\begin{vmatrix} \varepsilon^q & 0 \\ 0 & \varepsilon \end{vmatrix} \sim \begin{vmatrix} \alpha & \beta \\ \beta\zeta & \alpha \end{vmatrix}^*$	$-(\Pi(\varepsilon) + \Pi(\varepsilon^q))$
$t, \lambda \in GF(q)^*, t \neq 1; \varepsilon = \alpha + \beta\sqrt{\zeta}, \alpha, \beta, \zeta \in GF(q)$ with $\zeta$ not a square and $\beta \neq 0$ . * Matrices with the same characteristic roots are conjugate.	

Figure 1.

**Lemma 4.5.** Let  $X^\Pi$  be the character of a representation  $\mathcal{U}^\Pi$  of the discrete series of  $\mathcal{G}$ . Let  $\alpha$  be a character of  $\mathcal{D}$  such that  $\alpha(\lambda) = \Pi(\lambda)$  for any scalar matrix  $\lambda \in \mathcal{D}$ . Then  $m_\alpha$  is equivalent to  $\text{res } \mathcal{U}^\Pi$ . Fix a non-trivial character  $\Phi$  of  $B$ . Let  $\{m_{ij}^\alpha\}_{i,j \in GF(q)^\times}$  be the matrix coefficients of  $m_\alpha$  with respect to the basis  $\{\Phi_i\}_{i \in GF(q)^\times}$  of  $B^\alpha$  (see Lemma (4.2) and relation (4.2)). The matrix coefficients  $m_{ij}^\alpha$  are the restrictions to  $\mathcal{I}$  of matrix coefficients  $u_{ij}^{\Pi,\alpha}$  of  $\mathcal{U}^\Pi$ . For  $w = \begin{vmatrix} 0 & 1 \\ -1 & 0 \end{vmatrix}$  and  $\delta(a, b)$  the diagonal matrix with diagonal entries  $\delta_{11} = a$  and  $\delta_{22} = b$ ,

$$(4.3) \quad u_{ij}^{\Pi,\alpha}(w) = -\alpha^{-1}(\delta(i, j))q^{-1} \sum_{\varepsilon: \varepsilon\varepsilon^q = ij} \Pi(\varepsilon)\Phi(\varepsilon + \varepsilon^q).$$

Proof. By Lemma (4.4) and relation (4.2)

$$\begin{aligned} u_{ij}^{\Pi,\alpha}(w) &= \frac{(q-1)}{|\mathcal{I}|} \sum_{t \in \mathcal{I}} m_{ij}^\alpha(t) X^\Pi(t^{-1}w) \\ &= q^{-1} \sum_{u \in GF(q)} \alpha(\delta) \Phi_j(u) X^\Pi(b^{-1}(u)\delta^{-1}w), \end{aligned}$$

where  $\delta = \delta(j, i)$  and  $b(u) \in B$  has super diagonal entry  $u$ . Use of the explicit formula for  $X^\Pi$  easily yields (4.3).

**Theorem 4.6.** Let  $\Pi$  be a character of  $GF(q^2)^\times$  whose restriction to the elements of norm one is not trivial. Let  $X^\Pi$  be the character of the irreducible representation of  $\mathcal{G}$  associated with  $\Pi$ . Let  $\alpha$  be any character of  $\mathcal{D}$  which agrees

with  $\Pi$  on the scalar matrices. Then  $m_\alpha$  is  $\text{res } \mathcal{U}^\Pi$  and  $B^\alpha$ , the representation space of  $m_\alpha$ , is a representation space for  $\mathcal{U}^\Pi$ . Fix a non-trivial character  $\Phi$  of  $B$  and write it as a function of the super-diagonal entries of elements of  $B$ . Take as a basis for  $B^\alpha$  the  $q-1$  non-trivial characters  $\{\Phi_i\}_{i \in GF(q)^\times}$ , where  $\Phi_i(x) = \Phi(ix)$  for all  $x \in GF(q)$ . Matrix coefficients for  $\mathcal{U}^\Pi$  acting in  $B^\alpha$  are as follows. For  $i, j \in GF(q)^\times$  set  $u_{ij}^{\Pi, \alpha} = \langle \mathcal{U}^\Pi(g) \Phi_j, \Phi_i \rangle$ . If  $g = db(u)$ , where  $d \in \mathcal{D}$  has diagonal entries  $d_{11}$  and  $d_{22}$  and  $b(u) \in B$  has super-diagonal entry  $u \in GF(q)$ , then

$$(4.4) \quad u_{ij}^{\Pi, \alpha}(g) = \alpha(d) \Phi_j(u), \quad \text{provided } d_{11}^{-1} d_{22} = j^{-1} i;$$

$$= 0, \quad \text{otherwise.}$$

If  $g = b(v)wdb(u)$ , where  $d \in \mathcal{D}$  has diagonal entries  $d_{11}$  and  $d_{22}$ ,  $w = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}$ , and  $b(u)$  and  $b(v) \in B$  have superdiagonal entries  $u$  and  $v$  respectively then

$$(4.5) \quad u_{ij}^{\Pi, \alpha}(g) = \Phi(iv + ju) [-\Pi(d_{11}) \alpha^{-1}(\delta(i, j)) q^{-1} \sum_{\varepsilon: \varepsilon \varepsilon^q = 1} \Pi(\varepsilon) \Phi(\varepsilon + \varepsilon^q)]$$

where  $\delta(i, j)$  is the diagonal matrix with upper entry  $i$  and lower entry  $j$  and  $l = ij d_{11}^{-1} d_{22}$ .

Proof. Relation (4.4) is the same as (4.2), so no proof is needed. To prove (4.5) note first that  $u_{ij}^{\Pi, \alpha}(b(v)gb(u)) = \Phi_i(v) u_{ij}^{\Pi, \alpha}(g) \Phi_j(u)$ . Moreover,  $u_{ij}^{\Pi, \alpha}(wd) = \alpha(d) u_{i, jd_{11}^{-1}d_{22}}^{\Pi, \alpha}(w)$ . Use of (4.3) to express  $u_{i, jd_{11}^{-1}d_{22}}^{\Pi, \alpha}(w)$  as an exponential sum leads to a proof of (4.5).

## 5. Discrete series of $G$

Let  $\Pi$  be a character of  $GF(q^2)^\times$  whose restriction to  $N^1$ , the elements of norm one in  $GF(q^2)^\times$ , is not trivial. Let  $\pi$  be  $\Pi$  restricted to  $N^1$ . Let  $\mathcal{U}^\Pi$  be the representation of the discrete series of  $\mathcal{G}$  associated with  $\Pi$ . Set  $U^\pi = \text{res } \mathcal{U}^\Pi$ . The trace  $X^\pi$  of  $U^\pi$  is the restriction to  $G$  of  $X^\Pi$ , so, up to equivalence,  $U^\pi$  depends only on the values of  $\Pi$  restricted to  $N^1$ . Furthermore,  $U^\pi$  and  $U^{\pi'}$  are equivalent if and only if  $\pi' = \pi$  or  $\pi^{-1}$ , since, if  $\pi'$  is the restriction to  $N^1$  of a character  $\Pi'$  of  $GF(q^2)^\times$ ,  $X^\pi = X^{\pi'}$  if and only if  $\pi' = \pi$  or  $\pi^{-1}$ .

We may take as representatives for the conjugacy classes in  $\mathcal{G}$  those representatives for conjugacy classes in  $\mathcal{G}$  which lie in  $G$  (see Figure 1.). However,  $\begin{pmatrix} 1 & 0 \\ 1 & 1 \end{pmatrix}$  and  $\begin{pmatrix} 1 & 0 \\ \zeta & 1 \end{pmatrix}$ ,  $\zeta$  a non-square, are not conjugate in  $G$ ; similarly  $-\begin{pmatrix} 1 & 0 \\ 1 & 1 \end{pmatrix}$  and  $-\begin{pmatrix} 1 & 0 \\ \zeta & 1 \end{pmatrix}$ .

**Theorem 5.1.** *Let  $\pi$  be a non-trivial character of  $N^1$  and let  $U^\pi$  be the corresponding representation of  $G$  defined above.  $U^\pi$  is irreducible if and only if  $\pi^2 \neq 1$ . If  $\pi^2 = 1$ ,  $U^\pi = U_1^\pi + U_2^\pi$ , the direct sum of inequivalent  $\frac{1}{2}(q-1)$ -dimensional representations.*

Proof. It suffices to show that  $|G|^{-1} \sum_{g \in G} |X^\pi(g)|^2 = 1$ , if  $\pi^2 \neq 1$ , and 2, otherwise. The computation is easy and we omit it. In the case that  $U^\pi$  is reducible, the components are  $\frac{1}{2}(q-1)$ -dimensional and inequivalent, since, according to Lemma (4.3), this statement holds already for  $\text{res}_{G \downarrow T} U^\pi$ . We may use Lemma (4.3) to obtain representation spaces for  $U_1^\pi$  and  $U_2^\pi$ .

There are  $q+4$  conjugacy classes in  $G$  and we have accounted for this many equivalence classes of irreducible representations, so our description of the irreducible representations of  $SL(2, GF(q))$  is complete.

BOWDOIN COLLEGE

---

### References

- [1] I.M. Gel'fand and M.I. Graev: *Categories of group representations and the problem of classifying irreducible representations*, Soviet Math. Dokl. **3** (1962), 1378–1381.
- [2] H.D. Kloosterman: *The behavior of general theta functions under the modular group and the characters of binary modular congruence groups I and II*, Ann. of Math. **47** (1946), I: 317–375 and II: 376–447.
- [3] R. Steinberg: *The representations of  $GL(3, q)$ ,  $GL(4, q)$ ,  $PGL(3, q)$ , and  $PGL(4, q)$* , Canad. J. Math. **3** (1951), 225–235.
- [4] S. Tanaka: *Construction and classification of irreducible representations of special linear group of the second order over a finite field*, Osaka J. Math. **4** (1967), 65–84.
- [5] A. Weil: *Sur certaines groupes d'opérateurs unitaires*, Acta Math. **111** (1964), 143–211.