

Title	Cobordism groups of immersions
Author(s)	Uchida, Fuichi
Citation	Osaka Journal of Mathematics. 1971, 8(2), p. 207-218
Version Type	VoR
URL	<a href="https://doi.org/10.18910/8652">https://doi.org/10.18910/8652</a>
rights	
Note	

*Osaka University Knowledge Archive : OUKA*

<https://ir.library.osaka-u.ac.jp/>

Osaka University

## COBORDISM GROUPS OF IMMERSIONS

FUICHI UCHIDA

(Received November 9, 1970)

### 1. Introduction

In the previous paper [3] we have introduced the cobordism groups  $I(n, k)$ ,  $E(n, k)$  and  $G(n, k)$  of immersions, embeddings and generic immersions of  $n$ -manifolds into  $(n+k)$ -manifolds respectively. Mainly we have considered the cobordism group  $G(n, k)$ , and we have obtained the following exact sequence:

$$0 \rightarrow E(n, k) \rightarrow G(n, k) \rightarrow B(n-k, k) \rightarrow 0$$

where the group  $B(n, k)$  is the cobordism group of bundles over manifolds with involution defined in [3]

In general  $E(n, k)$  is isomorphic to the bordism group  $\mathfrak{X}_{n+k}(MO(k))$ , so we have studied the cobordism group  $B(n, k)$  in [3], [4], in order to study the cobordism group  $I(n, k)$ , since  $I(n, k)$  is canonically isomorphic to  $G(n, k)$  in the meta-stable range (i.e.  $2k > n+1$ ). Especially, as one of the consequence of the previous paper, the forgetting homomorphism

$$\alpha_* : E(n, k) \rightarrow I(n, k)$$

is injective in the meta-stable range.

One of the results of the present paper is that the homomorphism

$$\alpha_* : E(n, k) \rightarrow I(n, k)$$

is injective without restriction of the meta-stable range.

First we will introduce the notion "completely regular ( $p$ )-immersion" and study some properties of completely regular ( $p$ )-immersions in sections 2, 3. We will define the cobordism group  $C(n, k; p)$  of completely regular ( $p$ )-immersions in section 4, and we will show, in section 5, that the forgetting homomorphism

$$\alpha_p : C(n, k; p) \rightarrow C(n, k; p+1)$$

is injective. Therefore the homomorphism

$$\alpha_* : E(n, k) \rightarrow I(n, k)$$

is injective.

Next, in section 6, we will study the oriented cobordism groups of completely regular ( $p$ )-immersions. In the final section 7, we will return to the unoriented case and study the cokernel of the homomorphism  $\alpha_p$ .

**2. Definitions and notations**

Let  $M, N$  be  $C^\infty$ -differentiable manifolds.

**2.1.** First we will give some definitions.

DEFINITION 2.1. A  $C^\infty$ -differentiable mapping  $f: M \rightarrow N$  is *proper* if  $f^{-1}(\partial N) = \partial M$  and there are collar neighborhoods

$$\begin{aligned} c &: \partial M \times [0, 1) \rightarrow M \\ c' &: \partial N \times [0, 1) \rightarrow N \end{aligned}$$

for which the following diagram is commutative:

$$\begin{array}{ccc} \partial M \times [0, 1) & \xrightarrow{c} & M \\ \downarrow (f|_{\partial M}) \times 1 & & \downarrow f \\ \partial N \times [0, 1) & \xrightarrow{c'} & N \end{array}$$

DEFINITION 2.2. Subspaces  $V_1, V_2, \dots, V_p$  of a vector space  $V$  are *in general position* if

$$\dim(V_{i_1} \cap \dots \cap V_{i_k}) = \sum_{r=1}^k \dim(V_{i_r}) - (k-1) \dim V$$

for  $1 \leq i_1 < i_2 < \dots < i_k \leq p$ .

DEFINITION 2.3. A proper immersion  $f: M \rightarrow N$  is *completely regular* if the subspaces  $df(M_{x_1}), \dots, df(M_{x_p})$  of  $N_y$  are in general position for  $x_1, \dots, x_p$  in  $M$  such that  $y = f(x_1) = \dots = f(x_p)$ , where  $M_x$  is a tangent space of  $M$  at  $x$ , and  $df: M_x \rightarrow N_{f(x)}$  is a differential of  $f$ .

DEFINITION 2.4. A completely regular immersion  $f: M \rightarrow N$  is *completely regular ( $p$ )-immersion* if  $f^{-1}(f(x))$  has at most  $p$ -elements for any point  $x$  in  $M$ .

**Lemma 2.5.** *Let  $V_1, \dots, V_p$  be subspaces of a vector space  $V$ , then the following conditions are equivalent:*

- (a)  $V_1, \dots, V_p$  are in general position,
- (b)  $V_{i_0} + (V_{i_1} \cap \dots \cap V_{i_k}) = V$  for mutually distinct indices  $i_0, i_1, \dots, i_k$ ,
- (c) orthogonal complements of  $V_1, \dots, V_p$  are linearly independent for some

inner product of  $V$ .

**2.2.** Next we will fix some notations.

$E(M, N)$ : the set of all proper embeddings from  $M$  to  $N$ ,

$I(M, N)$ : the set of all proper immersions from  $M$  to  $N$ ,

$CR(M, N)$ : the set of all proper completely regular immersions from  $M$  to  $N$ ,

$CR_{(p)}(M, N)$ : the set of all proper completely regular  $(p)$ -immersions from  $M$  to  $N$ ,

$C^\infty(M, N)$ : the set of all proper  $C^\infty$ -differentiable mappings from  $M$  to  $N$ .  
Then

$$E(M, N) = CR_{(1)}(M, N) \subset CR_{(2)}(M, N) \subset \dots \subset CR_{(p)}(M, N) \\ \subset CR_{(p+1)}(M, N) \subset \dots,$$

$$CR(M, N) = \bigcup_{p \geq 1} CR_{(p)}(M, N), \quad CR(M, N) \subset I(M, N) \subset C^\infty(M, N),$$

$$CR_{(p)}(M, N) = CR(M, N) \text{ for } p(\dim N - \dim M) > \dim M,$$

$$CR(M, N) = I(M, N) \text{ for } \dim M = \dim N.$$

### 3. Completely regular immersions

Let  $X$  be a set. Denote by  $X^{(p)}$  the  $p$ -fold cartesian product of  $X$ ,  $\Delta^p X$  the diagonal set of  $X^{(p)}$  and

$$\Delta_{(p)} X = \{(x_1, \dots, x_p) \in X^{(p)} \mid x_i = x_j \text{ for some } i < j\}.$$

Let  $f: X \rightarrow Y$  be a mapping. Denote by

$$f^{(p)} : X^{(p)} \rightarrow Y^{(p)}$$

the mapping defined by  $f^{(p)}(x_1, \dots, x_p) = (f(x_1), \dots, f(x_p))$ .

Then we have the following result from the definition of the transverse regularity condition.

**Lemma 3.1.** *Let  $M, N$  be  $C^\infty$ -differentiable manifolds without boundary and  $f: M \rightarrow N$  be a  $C^\infty$ -differentiable mapping. Then the  $C^\infty$ -differentiable mapping  $f^{(p)}: M^{(p)} - \Delta_{(p)} M \rightarrow N^{(p)}$  is transverse regular over the diagonal  $\Delta^p N$ , if and only if*

$$dfM_{x_1} + (dfM_{x_2} \cap \dots \cap dfM_{x_p}) = N_y,$$

for  $(x_1, \dots, x_p) \in M^{(p)} - \Delta_{(p)} M$  such that  $y = f(x_1) = \dots = f(x_p)$ .

As a corollary of this lemma we have the following result.

**Theorem 3.2.** *Let  $M, N$  be  $C^\infty$ -differentiable manifolds without boundary and  $f: M \rightarrow N$  be an immersion. Then*

(a)  $f$  is completely regular if and only if  $f^{(p)} : M^{(p)} - \Delta_{(p)}M \rightarrow N^{(p)}$  is transverse regular over the diagonal  $\Delta^p N$  for all  $p \geq 2$ ,

(b)  $f$  is completely regular ( $p$ )-immersion if and only if  $f^{(k)} : M^{(k)} - \Delta_{(k)}M \rightarrow N^{(k)}$  is transverse regular over the diagonal  $\Delta^k N$  for  $2 \leq k \leq p$  and  $f^{(p+1)}(M^{(p+1)} - \Delta_{(p+1)}M)$  does not meet the diagonal  $\Delta^{p+1}N$ .

**Corollary 3.3.**  $CR(M, N)$  and  $CR_{(p)}(M, N)$  are open subsets of  $I(M, N)$  with respect to the fine  $C^1$ -topology.

**Corollary 3.4.** Let  $f : M \rightarrow N$  be a completely regular ( $p$ )-immersion and  $X = \{x \in M \mid f^{-1}(f(x)) \text{ has just } p\text{-elements}\}$ . Then  $X$  is a closed submanifold of  $M$  with dimension  $\dim N - p(\dim N - \dim M)$ .

**Theorem 3.5.** Let  $M$  be a compact  $C^\infty$ -differentiable manifold and  $N$  a  $C^\infty$ -differentiable manifold. Then

(a) the set  $CR(M, N)$  is a dense open subset of  $I(M, N)$  with respect to the fine  $C^1$ -topology,

(b) let  $A$  be a closed subset of  $M$  and  $f : M \rightarrow N$  be an immersion, if the restriction of  $f$  over  $A$  is completely regular and  $f(A)$  does not intersect  $f(M - A)$ , then, as an arbitrarily closed  $C^1$ -approximation of  $f$ , there is a completely regular immersion  $g : M \rightarrow N$  such that  $g = f$  on  $A$ .

Proof. This is an immediate corollary of Theorem 3.2 and the generalized transversality theorem (Theorem 1.10[1]).

**Corollary 3.6.** Any immersion  $f : M \rightarrow N$  is differentiably homotopic to a completely regular immersion.

Proof. This follows from Theorem 3.5 and the fact that  $I(M, N)$  is locally contractible with respect to the fine  $C^1$ -topology (cf. [2]).

#### 4. Cobordism of immersions

**4.1.** A completely regular ( $p$ )-immersion of dimension  $(n, k)$  is a triple  $(f, M, N)$  consisting of two closed  $C^\infty$ -differentiable manifolds  $M, N$  of dimensions  $n, n+k$  respectively and a completely regular ( $p$ )-immersion  $f : M \rightarrow N$ . We identify  $(f, M, N)$  with  $(f', M', N')$  if and only if there are diffeomorphisms  $\varphi : M \rightarrow M'$  and  $\psi : N \rightarrow N'$  for which  $\psi f = f' \varphi$ .

A completely regular ( $p$ )-immersion  $(f, M, N)$  of dimension  $(n, k)$  will be said to be *cobordant to zero* if there exists a triple  $(F, V, W)$  where:

(1)  $V$  and  $W$  are compact  $C^\infty$ -differentiable manifolds of dimensions  $n+1, n+k+1$  respectively, and

(2)  $F : V \rightarrow W$  is a proper completely regular ( $p$ )-immersion such that  $(F \mid \partial V, \partial V, \partial W) = (f, M, N)$ .

Then we denote  $\partial(F, V, W)=(f, M, N)$ . Two completely regular ( $p$ )-immersions  $(f_0, M_0, N_0)$  and  $(f_1, M_1, N_1)$  of dimension  $(n, k)$  will be said to be *cobordant* if and only if the disjoint union  $(f_0, N_0, M_0)+(f_1, M_1, N_1)$  is cobordant to zero.

This cobordism relation is an equivalence relation and denote by  $\mathbf{C}(n, k; p)$  the set of equivalence classes under this relation of completely regular ( $p$ )-immersions of dimension  $(n, k)$ . As usual, an abelian group structure is imposed on  $\mathbf{C}(n, k; p)$  by disjoint union, which is called the cobordism group of completely regular ( $p$ )-immersions of dimension  $(n, k)$ , and every element of  $\mathbf{C}(n, k; p)$  is its own inverse.

**4.2.** In the above definition, if the term “completely regular ( $p$ )-immersion” is replaced by “embedding”, “immersion” and “completely regular immersion”, one may define the cobordism groups of embeddings  $\mathbf{E}(n, k)$ , immersions  $\mathbf{I}(n, k)$  and completely regular immersions  $\mathbf{C}(n, k)$ , of dimension  $(n, k)$  respectively.

By definition  $\mathbf{E}(n, k)=\mathbf{C}(n, k; 1)$  and there are natural forgetting homomorphisms

$$\begin{aligned} \alpha_p &: \mathbf{C}(n, k; p) \rightarrow \mathbf{C}(n, k; p+1) \\ \bar{\alpha}_p &: \mathbf{C}(n, k; p) \rightarrow \mathbf{I}(n, k) \\ \alpha_* &: \mathbf{E}(n, k) \rightarrow \mathbf{I}(n, k) \end{aligned}$$

such that  $\alpha_*=\bar{\alpha}_p \circ \alpha_{p-1} \circ \dots \circ \alpha_2 \circ \alpha_1$ .

REMARK. (a)  $\bar{\alpha}_p$  is an isomorphism for  $kp > n+1$ ,  
 (b)  $\alpha_1$  is injective, since  $\mathbf{C}(n, k; 2)=\mathbf{G}(n, k)$  the cobordism group of generic immersions defined in the previous paper ([3], Section 4).

In the next section, we will prove that the homomorphism  $\alpha_p$  is injective for all  $p \geq 1$ .

### 5. Splitting homomorphisms

**5.1.** Let  $f: M \rightarrow N$  be a proper completely regular ( $p$ )-immersion where  $M, N$  are compact  $C^\infty$ -differentiable manifolds of dimensions  $n, n+k$  respectively. Let

$$X = \{x \in M \mid f^{-1}(f(x)) \text{ has just } p\text{-elements}\}$$

and  $Y=f(X)$ . Then  $X$  and  $Y$  are closed submanifolds of  $M, N$  respectively, and  $\dim X=\dim Y=n-(p-1)k$ . Moreover  $f|X: X \rightarrow Y$  is a  $p$ -fold covering. Then

**Lemma 5.1.** *There are Riemannian metrics on  $M, N$  such that*

- (a) *the differential  $df: M_x \rightarrow N_{f(x)}$  is isometric for  $x \in X$ ,*
- (b) *the orthogonal complements of  $dfM_{x_1}, \dots, dfM_{x_p}$  in  $N_y$  are mutually orthogonal if  $y=f(x_1)=\dots=f(x_p)$ .*

Proof. Firstly, define a Riemannian metric satisfying (b) on a neighborhood in  $N$  of each point of  $Y$ . Define a Riemannian metric on  $N$  satisfying the condition (b) by making use of a  $C^\infty$ -differentiable partition of unity on  $N$ . Next, define a Riemannian metric on  $M$  as the induced metric by  $df$ .

**5.2.** Let  $\nu(X)$  and  $\nu(Y)$  be the normal bundles of the embeddings  $X \subset M$ ,  $Y \subset N$  respectively, with respect to the above Riemannian metrics. Denote by  $E(\nu(X))$ ,  $E(\nu(Y))$  the total spaces of these normal bundles, and  $E_\varepsilon(\nu(X))$ ,  $E_\varepsilon(\nu(Y))$  the set of all normal vectors with length  $\leq \varepsilon$ .

Then the differential  $df$  maps  $E(\nu(X))$  into  $E(\nu(Y))$  and the following diagram is commutative:

$$\begin{array}{ccc} E(\nu(X)) & \xrightarrow{df} & E(\nu(Y)) \\ \downarrow & & \downarrow \\ X & \xrightarrow{f|X} & Y \end{array}$$

where the vertical mappings are bundle projections. Then we have the following result by an elementary method.

**Lemma 5.2.** *There is a differentiably homotopic approximation  $g$  of  $f$  such that*

- (a)  $g=f$  on  $X$ ,
- (b)  $dg=df$  on  $E(\nu(X))$ ,
- (c) *the following diagram is commutative for some  $\varepsilon > 0$ :*

$$\begin{array}{ccc} E_{2\varepsilon}(\nu(X)) & \xrightarrow{dg} & E_{2\varepsilon}(\nu(Y)) \\ \downarrow \text{exp} & & \downarrow \text{exp} \\ M & \xrightarrow{g} & N \end{array}$$

where  $\text{exp}$  is the exponential mapping.

**5.3.** Under these notations, let

$$\begin{aligned} M_0 &= M - \text{exp}(\text{int } E_\varepsilon(\nu(X))), \\ N_0 &= N - \text{exp}(\text{int } E_\varepsilon(\nu(Y))), \\ \partial_1 M_0 &= \text{exp}(\partial E_\varepsilon(\nu(X))), \\ \partial_1 N_0 &= \text{exp}(\partial E_\varepsilon(\nu(Y))), \end{aligned}$$

Then  $\partial_1 M_0$  and  $\partial_1 N_0$  have fixed point free  $C^\infty$ -differentiable involutions  $a$ ,  $b$

induced from the bundle involutions of  $E(\nu(X))$ ,  $E(\nu(Y))$  respectively, and the following diagram is commutative:

$$\begin{array}{ccc}
 \partial_1 M_0 & \xrightarrow{g/\partial_1 M_0} & \partial_1 N_0 \\
 \downarrow a & & \downarrow b \\
 \partial_1 M_0 & \xrightarrow{g/\partial_1 M_0} & \partial_1 N_0
 \end{array}$$

Let  $M_1$  be the quotient space of  $M_0$  by the relation  $x=a(x)$  for  $x \in \partial_1 M_0$ . Let  $N_1$  be the quotient space of  $N_0$  by the relation  $y=b(y)$  for  $y \in \partial_1 N_0$ . Then  $M_1, N_1$  have naturally  $C^\infty$ -differentiable structures and a mapping  $g_1: M_1 \rightarrow N_1$  is induced from the mapping  $g|M_0: M_0 \rightarrow N_0$  which is proper completely regular  $(p-1)$ -immersion.

**Theorem 5.3.** *There is a homomorphism.*

$$\gamma_p : C(n, k; p) \rightarrow C(n, k; p-1)$$

such that  $\gamma_{p+1} \circ \alpha_p = \text{identity}$ .

*Pfoof.* By the above notations, the correspondence from  $(f, M, N)$  to  $(g_1, M_1, N_1)$  is cobordism invariant, and this defines a desired homomorphism.

**Corollary 5.4.** *The homomorphism  $\alpha_p: C(n, k; p) \rightarrow C(n, k; p+1)$  is injective for all  $p \geq 1$  and the image of  $\alpha_p$  is a direct summand.*

**Corollary 5.5.** *The homomorphism  $\alpha_*: E(n, k) \rightarrow I(n, k)$  is injective.*

*Proof.* If  $k > 0$ , then this follows from the above corollary. If  $k = 0$ , then this follows directly from the definition of  $E(n, k)$  and  $I(n, k)$ .

### 6. Oriented cobordism of immersions

**6.1.** By similar argument to the unoriented case one may define the oriented cobordism groups of completely regular  $(p)$ -immersions  $C^0(n, k; p)$ , embeddings  $E^0(n, k)$ , immersions  $I^0(n, k)$  and completely regular immersions  $C^0(n, k)$ , of dimension  $(n, k)$  respectively, where we consider orientation preserving mappings if  $k = 0$ .

By definition  $E^0(n, k) = C^0(n, k; 1)$  and there are forgetting homomorphisms

$$\begin{aligned}
 \alpha_p^0 &: C^0(n, k; p) \rightarrow C^0(n, k; p+1) \\
 \bar{\alpha}_p^0 &: C^0(n, k; p) \rightarrow I^0(n, k) \\
 \alpha_*^0 &: E^0(n, k) \rightarrow I^0(n, k)
 \end{aligned}$$

such that  $\alpha_*^0 = \bar{\alpha}_p^0 \circ \alpha_{p-1}^0 \circ \dots \circ \alpha_2^0 \circ \alpha_1^0$ .



REMARK. (a)  $\alpha_p^0$  is an isomorphism for  $kp > n + 1$ ,  
 (b)  $C^0(n, k; 2) = G^0(n, k)$  the oriented cobordism group of generic immersions defined in the previous paper ([3], Section 10).

We could not find such a homomorphism as  $\gamma_p$ , so we do not know in general whether the homomorphisms  $\alpha_p^0$  and  $\alpha_*^0$  are injective or not. In the following we give some partial results.

**6.2.** First, we consider the case of low codimensions. Let  $s$  be a point of  $k$ -sphere  $S^k$ . Let  $M, N$  be oriented closed  $C^\infty$ -differentiable manifolds of dimensions  $n, n+k$  respectively, and define a mapping

$$f : M \rightarrow N + M \times S^k$$

by  $f(x) = (x, s)$ , then  $f$  is an embedding. The function

$$\iota : \Omega_n \oplus \Omega_{n+k} \rightarrow E^0(n, k)$$

defined by  $\iota([M], [N]) = [f, M, N + M \times S^k]$  is a well-defined homomorphism.

**Lemma 6.1.** *The homomorphism*

$$\alpha_*^0 \circ \iota : \Omega_n \oplus \Omega_{n+k} \rightarrow I^0(n, k)$$

is injective and the image of  $\iota$  is a direct summand of  $E^0(n, k)$ .

Proof. Let  $\pi : I^0(n, k) \rightarrow \Omega_n \oplus \Omega_{n+k}$  be a homomorphism defined by  $\pi([f, M, N]) = ([M], [N])$ , then  $\pi \circ \alpha_*^0 \circ \iota = \text{identity}$ . Therefore we have the desired result.

**Proposition 6.2.** *The homomorphism*

$$\alpha_*^0 : E^0(n, k) \rightarrow I^0(n, k)$$

is injective for  $k=0$  and  $k=1$ .

Proof. In general  $E^0(n, k)$  is isomorphic to  $\Omega_{n+k}(MSO(k))$ . If  $k=1$ , then  $MSO(1)$  is homotopy equivalent to the circle and hence  $\Omega_*(MSO(1))$  is isomorphic to the tensor product  $H_*(MSO(1); Z) \otimes \Omega_*$ . Therefore  $E^0(n, 1)$  is isomorphic to  $\Omega_n \oplus \Omega_{n+1}$  and  $\alpha_*^0$  is injective by Lemma 6.1. If  $k=0$ , then the results follows directly from the definition of  $E^0(n, 0)$  and  $I^0(n, 0)$ .

**6.3.** If  $k > n + 1$ , then the homomorphism

$$\alpha_*^0 : E^0(n, k) \rightarrow I^0(n, k)$$

is isomorphic by Remark in 6.1. We consider the case  $k=n+1$  and  $k=n$ .

**Proposition 6.3.**

- (a)  $\alpha_*^0: E^0(n, n+1) \rightarrow I^0(n, n+1)$  is an isomorphism,
- (b)  $\alpha_*^0: E^0(n, n) \rightarrow I^0(n, n)$  is injective.

Proof. Since the homomorphism

$$\bar{\alpha}_2^0: C^0(n, k; 2) \rightarrow I^0(n, k)$$

is an isomorphism in both case except  $n=k=1$ , and  $E^0(1, 1)=0$ , it is sufficient to consider the homomorphism

$$\alpha_1^0: E^0(n, k) \rightarrow C^0(n, k; 2) = G^0(n, k).$$

There are exact sequences [3]:

$$\begin{aligned} G^0(2s-1, 2s-1) &\xrightarrow{\beta} B^-(0, 2s-1) \xrightarrow{\partial} E^0(2s-2, 2s-1) \xrightarrow{\alpha_1^0} G^0(2s-2, 2s-1) \rightarrow 0, \\ G^0(2s, 2s) &\xrightarrow{\beta} B^+(0, 2s) \xrightarrow{\partial} E^0(2s-1, 2s) \xrightarrow{\alpha_1^0} G^0(2s-1, 2s) \rightarrow 0. \end{aligned}$$

Now define a mapping

$$f: S_1^n + S_2^n \rightarrow S^n \times S^n$$

by  $f(x)=(x, s)$  for  $x \in S_1^n$  and  $f(y)=(s, y)$  for  $y \in S_2^n$ , where  $S_i^n$  is a copy of  $S^n$  and  $s \in S^n$  is a base point. Then  $f$  is a completely regular (2)-immersion with unique double point and therefore the homomorphisms  $\beta$  are onto in the above sequences. Consequently the homomorphism  $\alpha_1^0$  is an isomorphism.

Next, since  $B^-(1, 2s-1)=0$  in the exact sequence

$$B^-(1, 2s-1) \xrightarrow{\partial} E^0(2s-1, 2s-1) \xrightarrow{\alpha_1^0} G^0(2s-1, 2s-1) \xrightarrow{\beta} B^-(0, 2s-1) \rightarrow 0,$$

the homomorphism  $\alpha_1^0: E^0(2s-1, 2s-1) \rightarrow G^0(2s-1, 2s-1)$  is injective and not onto since  $B^-(0, 2s-1)=Z_2$ .

Lastly,  $B^+(1, 2s)=Z_2$  in the following exact sequence

$$G^0(2s+1, 2s) \xrightarrow{\beta} B^+(1, 2s) \xrightarrow{\partial} E^0(2s, 2s) \xrightarrow{\alpha_1^0} G^0(2s, 2s).$$

We will prove that  $\beta$  is onto, and it is sufficient to show the existence of a completely regular (2)-immersion  $(f, M, N)$  of dimension  $(2s+1, 2s)$  such that

$$X_f = \{x \in M \mid f^{-1}(f(x)) \text{ has 2-elements}\}$$

is diffeomorphic to the circle  $S^1$ . Let  $CP^s$  be the complex projective space and

$$f: S^1 \times CP^s \rightarrow S^1 \times CP^{2s}$$

a mapping defined by

$$\begin{aligned} & f(e^{2i\theta}, \langle z_0, z_1, \dots, z_s \rangle) \\ &= (e^{4i\theta}, \langle z_0 e^{i\theta}, z_1 \cos \theta, \dots, z_s \cos \theta, z_1 \sin \theta, \dots, z_s \sin \theta \rangle) \end{aligned}$$

where  $\langle z_0, z_1, \dots, z_s \rangle$  is a homogeneous coordinate of  $CP^s$ . Then  $f$  is a completely regular (2)-immersion and

$$X_f = S^1 \times \{ \langle 1, 0, \dots, 0 \rangle \}$$

REMARK. By direct calculation, if  $n+k \leq 7$  but  $(n, k) \neq (4, 2)$ , then the homomorphism

$$\alpha_*^0: E^0(n, k) \rightarrow I^0(n, k)$$

is injective, and the homomorphism

$$\alpha_1^0: E^0(4, 2) \rightarrow C^0(4, 2; 2) = G^0(4, 2)$$

is injective.

## 7. Bundles over covering spaces

7.1. Now we return to the unoriented case. The homomorphism

$$\alpha_p: C(n, k; p) \rightarrow C(n, k; p+1)$$

is injective and the image of  $\alpha_p$  is a direct summand by Corollary 5.4, so we study now the cokernel of  $\alpha_p$ . For this purpose, we introduce new cobordism groups as follows.

7.2. Let  $k, p$  be fixed non-negative integers. A pair of bundles over a covering space is a quadruple  $(\xi, \eta, h, \bar{h})$ , where

$$\xi: E(\xi) \rightarrow B(\xi), \quad \eta: E(\eta) \rightarrow B(\eta)$$

are  $C^\infty$ -differentiable vector bundles over compact  $C^\infty$ -differentiable manifolds with fibre dimensions  $pk, (p+1)k$  respectively,

$$\bar{h}: B(\xi) \rightarrow B(\eta)$$

is a  $(p+1)$ -fold covering which is a proper  $C^\infty$ -differentiable mapping, and

$$h: E(\xi) \rightarrow E(\eta)$$

is a  $C^\infty$ -differentiable mapping covering  $\bar{h}$ . The following must be satisfied:

- (1)  $h$  maps each fibre  $\xi_x$  over  $x \in B(\xi)$  linearly one to one into a fibre  $\eta_{\bar{h}(x)}$ ,
- (2) for each  $y$  in  $B(\eta)$  and  $x_0, x_1, \dots, x_p$  in  $B(\xi)$  such that  $y = \bar{h}(x_0) = \bar{h}(x_1) = \dots = \bar{h}(x_p)$ , subspaces  $h(\xi_{x_0}), \dots, h(\xi_{x_p})$  of a vector space  $\eta_y$  are in general position.

**7.3.** A quadruple  $(\xi, \eta, h, \bar{h})$  is identified with a quadruple  $(\xi', \eta', h', \bar{h}')$  if and only if there is a quadruple  $(a, \bar{a}, b, \bar{b})$  of  $C^\infty$ -diffeomorphisms

$$\begin{aligned} a &: E(\xi) \rightarrow E(\xi'), & \bar{a} &: B(\xi) \rightarrow B(\xi'), \\ b &: E(\eta) \rightarrow E(\eta'), & \bar{b} &: B(\eta) \rightarrow B(\eta'), \end{aligned}$$

such that  $b \circ h = h' \circ a$  and  $a, b$  are bundle mappings covering  $\bar{a}, \bar{b}$  respectively.

For a quadruple  $(\xi, \eta, h, \bar{h})$ , denote by  $\partial(\xi, \eta, h, \bar{h})$  a quadruplet consisting of the restrictions

$$\begin{aligned} \xi|_{\xi^{-1}(\partial B(\xi))} &: \xi^{-1}(\partial B(\xi)) \rightarrow \partial B(\xi), \\ \eta|_{\eta^{-1}(\partial B(\eta))} &: \eta^{-1}(\partial B(\eta)) \rightarrow \partial B(\eta), \\ h|_{\xi^{-1}(\partial B(\xi))} &: \xi^{-1}(\partial B(\xi)) \rightarrow \eta^{-1}(\partial B(\eta)), \\ \bar{h}|_{\partial B(\xi)} &: \partial B(\xi) \rightarrow \partial B(\eta). \end{aligned}$$

**7.4.** The cobordism group  $\mathbf{B}(n, k; p)$  of pairs of bundles over a covering spaces of  $n$ -manifold may be now defined. If  $B(\xi_0)$  and  $B(\xi_1)$  are closed  $n$ -manifolds, then a quadruple  $(\xi_0, \eta_0, h_0, \bar{h}_0)$  is cobordant to a quadruple  $(\xi_1, \eta_1, h_1, \bar{h}_1)$  if and only if there is a quadruple  $(\xi, \eta, h, \bar{h})$  as such that

$$\partial(\xi, \eta, h, \bar{h}) = (\xi_0, \eta_0, h_0, \bar{h}_0) + (\xi_1, \eta_1, h_1, \bar{h}_1)$$

where the symbol  $+$  denotes disjoint union. Then this cobordism relation is an equivalence relation. Denote by  $\mathbf{B}(n, k; p)$  the set of all cobordism classes. As usual an abelian group structure is imposed on  $\mathbf{B}(n, k; p)$  by disjoint union, then every element is its own inverse.

REMARK.  $\mathbf{B}(n, k; 1)$  is naturally isomorphic with the cobordism group  $\mathbf{B}(n, k)$  of bundles over manifolds with involution defined in the previous paper ([3], Section 3).

**7.5.** Now we define homomorphisms

$$\begin{aligned} \beta_p &: \mathbf{C}(n, k; p+1) \rightarrow \mathbf{B}(n-pk, k; p), \\ \pi_p &: \mathbf{B}(n, k; p) \rightarrow \mathbf{C}(n+pk, k; p+1). \end{aligned}$$

(7.5.1) Let  $a \in \mathbf{C}(n, k; p+1)$  be represented by a completely regular  $(p+1)$ -immersion  $f: M \rightarrow N$ . Let

$$X = \{x \in M \mid f^{-1}(f(x)) \text{ has just } (p+1)\text{-elements}\}$$

and  $Y = f(X)$ . Then there are Riemannian metrics on  $M, N$  satisfying the conditions of Lemma 5.1. Let  $\nu(X), \nu(Y)$  be the normal bundles of the embeddings  $X \subset M, Y \subset N$  respectively, with respect to these Riemannian metrics, and then the differential  $df$  maps  $E(\nu(X))$  into  $E(\nu(Y))$ . Define  $\beta_p(a)$  the cobordism class

of the quadruple  $(\nu(X), \nu(Y), df|E(\nu(X)), f|X)$ .

(7.5.2.) Let  $b \in \mathbf{B}(n, k; p)$  be represented by a quadruple  $(\xi, \eta, h, \bar{h})$ . Define  $\pi_p(b)$  the cobordism class of a completely regular  $(p+1)$ -immersion defined by the mapping

$$P(h \oplus 1) : P(\xi \oplus \theta^1) \rightarrow P(\eta \oplus \theta^1)$$

where  $\theta^1$  is the trivial line bundle,  $P(\xi \oplus \theta^1)$  and  $P(\eta \oplus \theta^1)$  are the total spaces of the associated projective space bundles, and  $P(h \oplus 1)$  is a mapping canonically induced from the mapping  $h$ .

**Theorem 7.1.** *There is an exact sequence:*

$$0 \longrightarrow \mathbf{C}(n, k; p) \xrightarrow{\alpha_p} \mathbf{C}(n, k; p+1) \xrightarrow{\beta_p} \mathbf{B}(n-pk, k; p) \longrightarrow 0.$$

Proof. The homomorphism  $\alpha_p$  is injective by Corollary 5.4, and the homomorphism  $\beta_p$  is surjective since  $\beta_p \circ \pi_p = \text{identity}$  by definition (cf. [3] Theorem A'). The exactness at  $\mathbf{C}(n, k; p+1)$  is proved by the handle attaching construction (cf. [3], Section 5), so we omit the details.

OSAKA UNIVERSITY

---

### References

- [1] A. Haefliger: *Plongements différentiables de variétés dans variétés*, Comment. Math. Helv. **36** (1961), 47-82.
- [2] J. Munkres: *Elementary Differential Topology*, Annals of Mathematics Studies, Princeton, 1963.
- [3] F. Uchida: *Exact sequences involving cobordism groups of immersions*, Osaka J. Math. **6** (1969), 397-408.
- [4] F. Uchida: *The structure of the cobordism groups  $\mathbf{B}(n, k)$  of bundles over manifolds with involution*, Osaka J. Math. **7** (1970), 193-202.