



|              |   |
|--------------|---|
| Title        | On the hypersurfaces of Hermitian symmetric spaces of compact type. II    |
| Author(s)    | Kimura, Yoshio  |
| Citation     | Osaka Journal of Mathematics. 1980, 17(2), p. 455-469                     |
| Version Type | VoR   |
| URL          | <a href="https://doi.org/10.18910/8739">https://doi.org/10.18910/8739</a> |
| rights       |   |
| Note         |   |

*The University of Osaka Institutional Knowledge Archive : OUKA*

<https://ir.library.osaka-u.ac.jp/>

The University of Osaka

Kimura, Y.  
Osaka J. Math.  
17 (1980), 455-469

## ON THE HYPERSURFACES OF HERMITIAN SYMMETRIC SPACES OF COMPACT TYPE II

YOSHIO KIMURA

(Received November 20, 1978)

### 1. Introduction.

Let  $M$  be an irreducible Hermitian symmetric space of compact type and let  $L$  be a holomorphic line bundle over  $M$ . Denote by  $\Omega^p(L)$  the sheaf of germs of  $L$ -valued holomorphic  $p$ -forms on  $M$ . In the previous paper [1] we have studied the cohomology groups  $H^q(M, \Omega^p(L))$  of  $M$  if  $M$  is of type *BDI*, *EIII* or *EVII*. This note is the continuation of [1], and we retain the notations introduced in [1]. In this note we study the cohomology groups  $H^q(M, \Omega^p(L))$  of  $M$  of type *AIII*, *CI* or *EIII* and show the following theorem.

**Theorem.** *Let  $M$  be an irreducible Hermitian symmetric space of compact type but not a complex projective space nor a complex quadric of even dimension. Let  $V$  be a hypersurface of  $M$  whose degree  $\geq 2$ . Then*

$$H^0(V, \Theta) = (0)$$

where  $\Theta$  is the sheaf of germs of holomorphic vector fields on  $V$ .

The author would like to express his gratitude to Professor S. Murakami, Professor M. Takeuchi and Doctor M. Numata for their useful suggestions and encouragements.

### 2. Proof of the Theorem.

Theorem 8 and Lemma 3 in the previous paper [1] is incorrect. The followings are true.

**Theorem 8.** *Let  $M$  be an irreducible Hermitian symmetric space of type *EIII*, *EVII* or a complex quadric of odd dimension (resp. a complex quadric of even dimension  $\geq 4$ ), and let  $V$  be a hypersurface of  $M$  whose degree is  $d$ . Then*

$$H^0(V, \Theta) = (0) \quad \text{if } d \geq 2 \text{ (resp. } d \geq 3\text{)}$$

**Lemma 3.** *Let  $M$  be an  $n$ -dimensional irreducible Hermitian symmetric space of compact type *EIII*, *EVII* or a complex quadric of odd dimension (resp. a*

complex quadric of even dimension  $\geq 4$ ). Then

$$H^q(M, \Omega^p(E_{-k\omega_j})) = (0), \quad H^{q+1}(M, \Omega^p(E_{-(k-d)\omega_j})) = (0)$$

for  $p+q=n+1$ ,  $k=pd-\lambda$  if  $2 \leq p \leq n-1$  and  $d \geq 2$  (resp.  $d \geq 3$ ).

In the proof of Theorem 8 in [1], we have to replace  $n$  by  $n-1$  since  $\dim V=n-1$ . Thus we need the above Lemma 3, which is verified by the computations in [1].

From the above theorem we may assume that  $M$  is of type *AIII*, *CI* or *DIII* but not a complex projective space nor a complex quadric. If we prove the following proposition for such a space  $M$  we get the Theorem the same way as in the proof of Theorem 8 in [1].

**Proposition 1.** If  $d \geq 2$

$$H^q(M, \Omega^p(E_{-k\omega_j})) = (0), \quad H^{q+1}(M, \Omega^p(E_{-(k-d)\omega_j})) = (0),$$

for  $p+q \geq n+1$ ,  $k=pd-\lambda$ .

By Theorems 1 and 2 in [1], we get Proposition 1 if we prove the following inequalities:

$$\begin{aligned} \#\{\beta \in \Delta(\mathfrak{n}^+); (\sigma\delta + (dn(\sigma) - \lambda)\omega_j, \beta) < 0\} &< n+1-n(\sigma), \\ \#\{\beta \in \Delta(\mathfrak{n}^+); (\sigma\delta + (dn(\sigma) - d - \lambda)\omega_j, \beta) < 0\} &< n+2-n(\sigma), \end{aligned}$$

for  $\sigma \in W^1$  and  $d \geq 2$ .

Since  $(\omega_j, \beta) > 0$  for  $\beta \in \Delta(\mathfrak{n}^+)$ , we only have to prove the inequalities in the case of  $d=2$ . Recall that  $\#\Delta(\mathfrak{n}^+) = n$ . We can restate the inequalities, in the case of  $d=2$ , as follows:

**Proposition 2.** For  $\sigma \in W^1$

$$\begin{aligned} \#\{\beta \in \Delta(\mathfrak{n}^+); (\sigma\delta, \beta) \geq ((\lambda - 2n(\sigma))\omega_j, \beta)\} &> n(\sigma) - 1, \\ \#\{\beta \in \Delta(\mathfrak{n}^+); (\sigma\delta, \beta) \geq ((\lambda + 2 - 2n(\sigma))\omega_j, \beta)\} &> n(\sigma) - 2. \end{aligned}$$

In the following we shall prove Proposition 2 in each case.

**2.1. The case that  $M$  is of type *AIII* but not a complex projective space, that is  $M = SU(l+1)/S(U(j) \times U(l+1-j))$ ,  $l \geq 3$  and  $2 \leq j \leq l-1$ .** We immediately see that  $n=j(l+1-j)$  and  $\lambda=l+1$ . The Dynkin diagram of  $\Pi$  is as follows:

$$\begin{array}{ccccccccccccc} \circ & \text{---} & \circ & \text{---} & \cdots & \text{---} & \circ & \text{---} & \cdots & \text{---} & \circ & \text{---} & \circ \\ \alpha_1 & & \alpha_2 & & & & \alpha_j & & & & \alpha_{l-1} & & \alpha_l \end{array}$$

Let  $\{\varepsilon_i; 1 \leq i \leq l+1\}$  be a usual basis of  $R^{l+1}$ . Then we have:

$$\begin{aligned}
\mathfrak{h}_0 &= \left\{ \sum_{i=1}^{l+1} a_i \varepsilon_i \in R^{l+1}; \sum_{i=1}^{l+1} a_i = 0 \right\}, \\
\Delta &= \{ \varepsilon_i - \varepsilon_k; 1 \leq i, k \leq l+1, i \neq k \}, \\
\Pi &= \{ \alpha_1 = \varepsilon_1 - \varepsilon_2, \alpha_2 = \varepsilon_2 - \varepsilon_3, \dots, \alpha_l = \varepsilon_l - \varepsilon_{l+1} \}, \\
\Delta(\mathfrak{n}^+) &= \{ \varepsilon_i - \varepsilon_k; 1 \leq i \leq j < k \leq l+1 \}, \\
2\delta &= l\varepsilon_1 + (l-2)\varepsilon_2 + (l-4)\varepsilon_3 + \dots + (l+2)\varepsilon_l - l\varepsilon_{l+1}, \\
\omega_j &= \varepsilon_1 + \dots + \varepsilon_j - \frac{j}{l+1} \sum_{i=1}^{l+1} \varepsilon_i.
\end{aligned}$$

An element  $\sigma \in W$  acts on  $R^{l+1}$  by  $\sigma \varepsilon_i = \varepsilon_{\sigma(i)}$  for  $1 \leq i \leq l+1$ , where  $\sigma$  in the index is a permutation of  $\{1, 2, \dots, l+1\}$ . We represent  $\sigma$  by

$$\begin{pmatrix} 1 & 2 & \dots & l+1 \\ \sigma(1) & \sigma(2) & \dots & \sigma(l+1) \end{pmatrix}.$$

Then

$$W^1 = \left\{ \sigma \in W; \sigma^{-1} = \begin{pmatrix} 1 & \dots & l+1 \\ \sigma^{-1}(1) & \dots & \sigma^{-1}(l+1) \end{pmatrix}, \sigma^{-1}(1) < \dots < \sigma^{-1}(j), \sigma^{-1}(j+1) < \dots < \sigma^{-1}(l+1) \right\}.$$

The index  $n(\sigma)$  of  $\sigma \in W^1$  is given by

$$n(\sigma) = \sum_{i=1}^j (\sigma^{-1}(i) - i)$$

(Takeuchi [2]). We see easily that

$$\begin{aligned}
(\omega_j, \beta) &= 1 \quad \text{for any } \beta \in \Delta(\mathfrak{n}^+), \\
(\sigma\delta, \varepsilon_i - \varepsilon_k) &= \sigma^{-1}(k) - \sigma^{-1}(i) \quad \text{for } 1 \leq i, k \leq l+1.
\end{aligned}$$

Therefore we have to prove that the following two inequalities are true for any  $\sigma \in W^1$

$$(1.1) \quad \#\{(i, k); 1 \leq i \leq j < k \leq l+1, \sigma^{-1}(k) - \sigma^{-1}(i) \geq l+1 - 2n(\sigma)\} > n(\sigma) - 1,$$

$$(1.2) \quad \#\{(i, k); 1 \leq i \leq j < k < l+1, \sigma^{-1}(k) - \sigma^{-1}(i) \geq l+3 - 2n(\sigma)\} > n(\sigma) - 2.$$

First we prove the inequality (1.1).

**Lemma 1.1.** *Let  $\sigma \in W^1$ . If  $n(\sigma) \geq l+1$ , the inequality (1.1) is true.*

Proof. Since  $n(\sigma) \geq l+1$ ,  $l+1 - 2n(\sigma) \leq -(l+1)$ . There exist no pair  $(i, k)$ ,  $i \neq k$ , which satisfies

$$\sigma^{-1}(k) - \sigma^{-1}(i) < -(l+1).$$

Therefore

$$\#\{(i, k); 1 \leq i \leq j < k \leq l+1, \sigma^{-1}(k) - \sigma^{-1}(i) \geq l+1 - 2n(\sigma)\} = n.$$

From the definition of the index  $n(\sigma) \leq n$ , it follows that  $n > n(\sigma) - 1$ . Q.E.D.

**Lemma 1.2.** *Let  $\sigma \in W^1$ . Assume that  $\sigma(1) \neq 1$  and  $\sigma(l+1) \neq l+1$ . Then  $n(\sigma) \geq l$ .*

Proof. By the assumption  $\sigma^{-1}(j) = l+1$  and  $\sigma^{-1}(i) - i \geq 1$ ,  $1 \leq i \leq j$ . Therefore

$$\begin{aligned} n(\sigma) &= \sum_{i=1}^j (\sigma^{-1}(i) - i) \\ &= \sigma^{-1}(j) - j + \sum_{i=1}^{j-1} (\sigma^{-1}(i) - i) \\ &\geq (l+1-j) + (j-1) \\ &= l. \end{aligned} \quad \text{Q.E.D.}$$

**Lemma 1.3.** *Let  $\sigma \in W^1$ . Assume that  $\sigma(1) \neq 1$  and  $\sigma(l+1) \neq l+1$ . Then the inequality (1.1) is true.*

Proof. By Lemmas 1.1 and 1.2 we may assume that  $n(\sigma) = l$ . Then such an element  $\sigma$  is unique and given by

$$\sigma^{-1} = \begin{pmatrix} 1 & \cdots & j-1 & j & j+1 & j+2 & \cdots & l+1 \\ 2 & \cdots & j & l+1 & 1 & j+1 & \cdots & l \end{pmatrix}.$$

The pair  $(i, k)$ ,  $1 \leq i \leq j < k \leq l+1$ , which satisfies

$$\sigma^{-1}(k) - \sigma^{-1}(i) < l+1 - 2n(\sigma) = 1-l$$

is  $(j, j+1)$ . Hence

$$\#\{(i, k); 1 \leq i \leq j < k \leq l+1, \sigma^{-1}(k) - \sigma^{-1}(i) \geq 1-l\} = n-1 > n(\sigma)-1. \quad \text{Q.E.D.}$$

**Lemma 1.4.** *If  $j=2$ , the inequality (1.1) is true for any  $\sigma \in W^1$ .*

Proof. From the definition of  $n(\sigma)$

$$(1.3) \quad n(\sigma) = \sigma^{-1}(1) + \sigma^{-1}(2) - 3.$$

If  $n(\sigma) = 0$ , the inequality (1.1) is clearly true. Let  $n(\sigma) = 1$ . Then  $\sigma^{-1}(1) = 1$ ,  $\sigma^{-1}(2) = 3$  and

$$\sigma^{-1}(l+1) - \sigma^{-1}(1) = l > l+1 - 2n(\sigma).$$

It follows that the inequality (1.1) is true. Let  $n(\sigma) = 2$ . It is easy to see that the inequality (1.1) is true.

By Lemma 1.1 we have already seen that if  $n(\sigma) \geq l+1$  the inequality is

true. Hence we only have to show that (1.1) is true under the following condition:

$$(1.4) \quad 5 < \sigma^{-1}(1) + \sigma^{-1}(2) < l + 4.$$

By (1.3)

$$l + 1 - 2n(\sigma) = l + 7 - 2(\sigma^{-1}(1) + \sigma^{-1}(2)).$$

Since  $\sigma^{-1}(k) \geq k - 2$  for  $2 < k \leq l + 1$ ,

$$\begin{aligned} & \#\{k; 2 < k \leq l + 1, \sigma^{-1}(k) - \sigma^{-1}(1) \geq l + 7 - 2(\sigma^{-1}(1) + \sigma^{-1}(2))\} \\ & \geq \min \{\sigma^{-1}(1) + 2\sigma^{-1}(2) - 7, l - 1\}. \end{aligned}$$

Similarly

$$\begin{aligned} & \#\{k; 2 < k \leq l + 1, \sigma^{-1}(k) - \sigma^{-1}(2) \geq l + 7 - 2(\sigma^{-1}(1) + \sigma^{-1}(2))\} \\ & \geq \min \{2\sigma^{-1}(1) + \sigma^{-1}(2) - 7, l - 1\}. \end{aligned}$$

Therefore

$$\begin{aligned} & \#\{(i, k); 1 \leq i \leq 2 < k \leq l + 1, \sigma^{-1}(k) - \sigma^{-1}(i) \geq l + 1 - 2n(\sigma)\} \\ & \geq \min \{3(\sigma^{-1}(1) + \sigma^{-1}(2)) - 14, l + 2\sigma^{-1}(1) + \sigma^{-1}(2) - 8, 2l - 2\}. \end{aligned}$$

It is easy to see that  $3(\sigma^{-1}(1) + \sigma^{-1}(2)) - 14$ ,  $l + 2\sigma^{-1}(1) + \sigma^{-1}(2) - 8$  and  $2l - 2$  are both larger than  $n(\sigma) - 1 = \sigma^{-1}(1) + \sigma^{-1}(2) - 4$  under the condition (1.4).

Q.E.D.

We get the following lemma in the similar way as above.

**Lemma 1.5.** *If  $j = l - 1$ , the inequality (1.1) is true for any  $\sigma \in W^1$ .*

We shall prove that the inequality (1.1) is true for any  $\sigma \in W^1$  by using induction on  $l$ . If  $l = 3$  so that  $j = 2$ , it follows, by Lemma 1.4, our assertion is true.

Let  $l = l_0 \geq 4$ . We can assume that  $3 \leq j = j_0 \leq l_0 - 2$  and either  $\sigma(1) = 1$  or  $\sigma(l_0 + 1) = l_0 + 1$  by Lemmas 1.3, 1.4 and 1.5.

Case 1:  $\sigma(1) = 1$ . Define the element  $\tau$  of  $W^1$ , which is considered as an element of  $W^1$  for  $l = l_0 - 1$  and  $j = j_0 - 1$ , by

$$\tau^{-1} = \begin{pmatrix} 1 & 2 & \cdots & l_0 \\ \sigma^{-1}(2) - 1 & \sigma^{-1}(3) - 1 & \cdots & \sigma^{-1}(l_0 + 1) - 1 \end{pmatrix}.$$

We immediately see that  $2 \leq j \leq l - 2$  and  $n(\tau) = n(\sigma)$ . By the assumption of the induction,

$$\#\{(i, k); 1 \leq i \leq j_0 - 1 < k \leq l_0, \tau^{-1}(k) - \tau^{-1}(i) \geq l_0 - 2n(\tau)\} > n(\tau) - 1.$$

Hence

$$(1.5) \quad \#\{(i, k); 2 \leq i \leq j_0 < k \leq l_0 + 1, \sigma^{-1}(k) - \sigma^{-1}(i) \geq l_0 - 2n(\sigma)\} > n(\sigma) - 1.$$

For any  $k, j_0 \leq k \leq l_0 + 1$ , if there exists  $i, 2 \leq i \leq j_0$ , which satisfies the following:

$$\sigma^{-1}(k) - \sigma^{-1}(i) = l_0 - 2n(\sigma),$$

such an integer  $i$  is unique and

$$\sigma^{-1}(k) - \sigma^{-1}(1) \geq l_0 + 1 - 2n(\sigma).$$

Hence (1.5) leads to (1.1).

Case 2:  $\sigma(l_0 + 1) = l_0 + 1$ . Define the element  $\tau \in W^1$ , which is considered as an element of  $W^1$  for  $l = l_0 - 1$  and  $j = j_0$ , by

$$\tau^{-1} = \begin{pmatrix} 1 & \cdots & l_0 \\ \sigma^{-1}(1) & \cdots & \sigma^{-1}(l_0) \end{pmatrix}.$$

Then  $3 \leq j \leq l - 1$  and  $n(\tau) = n(\sigma)$ . By the assumption of the induction,

$$(1.6) \quad \#\{(i, k); 1 \leq i \leq j_0 < k \leq l_0, \sigma^{-1}(k) - \sigma^{-1}(i) \geq l_0 - 2n(\sigma)\} > n(\sigma) - 1.$$

For any  $i, 1 \leq i \leq j_0$ , if there exists  $k, j_0 < k \leq l_0$ , which satisfies the following:

$$\sigma^{-1}(k) - \sigma^{-1}(i) = l_0 - 2n(\sigma),$$

such an integer  $k$  is unique and

$$\sigma^{-1}(l_0 + 1) - \sigma^{-1}(i) \geq l_0 + 1 - 2n(\sigma).$$

Hence (1.5) leads to (1.1).

Thus we have proved that the inequality (1.1) is true for any  $\sigma \in W^1$ .

In the following we shall prove that the inequality (1.2) is true for any  $\sigma \in W^1$ .

**Lemma 1.6.** *Let  $\sigma \in W^1$ . If  $n(\sigma) \geq l + 1$ , the inequality (1.2) is true.*

**Proof.** Since  $n(\sigma) \geq l + 1$

$$l + 3 - 2n(\sigma) \leq 1 - l.$$

If there exists a pair  $(i, k), i \neq k$ , which satisfies

$$\sigma^{-1}(k) - \sigma^{-1}(i) < 1 - l,$$

such a pair is unique. Therefore

$$\#\{(i, k); 1 \leq i \leq j \leq k \leq l + 1, \sigma^{-1}(k) - \sigma^{-1}(i) \geq l + 1 - 2n(\sigma)\} \geq n - 1 > n(\sigma) - 2.$$

Q.E.D.

**Lemma 1.7.** *Let  $\sigma \in W^1$ . Assume that  $\sigma(1) \neq 1$  and  $\sigma(l+1) \neq l+1$ . Then the inequality (1.2) is true.*

Proof. By Lemma 1.2 and 1.6 we may assume that  $n(\sigma) = l$ . Such an element  $\sigma$  is unique and represented by

$$\sigma^{-1} = \begin{pmatrix} 1 & \cdots & j-1 & j & j+1 & j+2 & \cdots & l+1 \\ 2 & \cdots & j & l+1 & 1 & j+1 & \cdots & l \end{pmatrix}.$$

The number of the pairs  $(i, k)$ ,  $l \leq i \leq j < k \leq l+1$ , which satisfies

$$\sigma^{-1}(k) - \sigma^{-1}(i) < l+3 - 2n(\sigma) = 3-l$$

is at most 2. Therefore

$$\#\{(i, k); 1 \leq i \leq j < k \leq l+1, \sigma^{-1}(k) - \sigma^{-1}(i) \geq l+3 - 2n(\sigma)\} \geq n-2.$$

Since  $n(\sigma) = l$ ,  $n(\sigma) < n$ . It follows that (1.2) is true. Q.E.D.

**Lemma 1.8.** *If  $j=2$ , the inequality (1.2) is true for any  $\sigma \in W^1$ .*

Proof. It is easy to see that (1.2) is true if  $n(\sigma) \leq 3$ . By Lemma 1.6 and (1.3), we only have to show that (1.2) is true under the condition:

$$(1.7) \quad 6 < \sigma^{-1}(1) + \sigma^{-1}(2) < l+4.$$

We get the following inequality in the same way as in the proof of Lemma 1.4.

$$\begin{aligned} & \#\{(i, k); 1 \leq i \leq 2 < k \leq l+1, \sigma^{-1}(k) - \sigma^{-1}(i) \leq l+3 - 2n(\sigma)\} \\ & \geq \min \{3(\sigma^{-1}(1) + \sigma^{-1}(2)) - 18, l+2\sigma^{-1}(1) + \sigma^{-1}(2) - 10, 2l-2\}. \end{aligned}$$

It is easy to see that  $3(\sigma^{-1}(1) + \sigma^{-1}(2)) - 18$ ,  $l+2\sigma^{-1}(1) + \sigma^{-1}(2) - 10$  and  $2l-2$  are both larger than  $n(\sigma) - 2 = \sigma^{-1}(1) + \sigma^{-1}(2) - 5$  under the condition (1.7). Q.E.D.


We get the following lemma in the similar way as above.

**Lemma 1.9.** *If  $j=l-1$ , the inequality (1.2) is true for any  $\sigma \in W^1$ .*

From Lemmas 1.7, 1.8 and 1.9, we can prove that the inequality (1.2) is true for any  $\sigma \in W^1$  in the same way as in the proof of the inequality (1.1).

**2.2. The case that  $M$  is of type  $CI$ , that is  $M = Sp(l)/U(l)$ .** If  $l=1$ ,  $M = P_1(C)$ . If  $l=2$ ,  $M$  is a complex quadroic of dimension 3. Hence we assume that  $l \geq 3$ .

In this case  $n = \frac{1}{2}l(l+1)$  and  $\lambda = l+1$ . The Dynkin diagram of  $\Pi$  is as follows:



where  $\alpha_i \odot$  shows  $\alpha_i = \alpha_l$ . Let  $\{\varepsilon_i; 1 \leq i \leq l\}$  be the basis of  $\mathfrak{h}_0$  which satisfies  $(\varepsilon_i, \varepsilon_j) = \delta_{ij}$ . Then we have:

$$\begin{aligned}\Delta &= \{\pm 2\varepsilon_i; 1 \leq i \leq l, \pm \varepsilon_i \pm \varepsilon_j; 1 \leq i < j \leq l\}, \\ \Pi &= \{\alpha_1 = \varepsilon_1 - \varepsilon_2, \dots, \alpha_{l-1} = \varepsilon_{l-1} - \varepsilon_l, \alpha_l = 2\varepsilon_l\}, \\ \Delta(\mathfrak{n}^+) &= \{2\varepsilon_i; 1 \leq i \leq l, \varepsilon_i + \varepsilon_j; 1 \leq i < j \leq l\}, \\ \delta &= l\varepsilon_1 + (l-1)\varepsilon_2 + \dots + \varepsilon_l, \\ \omega_l &= \varepsilon_1 + \dots + \varepsilon_l.\end{aligned}$$

An element  $\sigma \in W$  acts on  $\mathfrak{h}_0$  by  $\sigma \varepsilon_i = \pm \varepsilon_{\bar{\sigma}(i)}$  for  $1 \leq i \leq l$ , where  $\bar{\sigma}$  is a permutation of  $\{1, 2, \dots, l\}$ . We denote the element  $\sigma \in W$  by the symbol

$$\begin{pmatrix} 1 & 2 & \dots & l \\ \pm \bar{\sigma}(1) & \pm \bar{\sigma}(2) & \dots & \pm \bar{\sigma}(l) \end{pmatrix}$$

Then

$$\begin{aligned}W^1 = \left\{ \sigma \in W; \bar{\sigma}^{-1} = \begin{pmatrix} 1 & \dots & r & r+1 & \dots & l \\ \bar{\sigma}^{-1}(1) & \dots & \bar{\sigma}^{-1}(r) & -\bar{\sigma}^{-1}(r+1) & \dots & -\bar{\sigma}^{-1}(l) \end{pmatrix} \right. \\ \left. \text{for } 0 \leq r \leq l, \bar{\sigma}^{-1}(1) < \dots < \bar{\sigma}^{-1}(r), \bar{\sigma}^{-1}(r+1) > \dots > \bar{\sigma}^{-1}(l) \right\}.\end{aligned}$$

The index  $n(\sigma)$  of  $\sigma \in W^1$  is given by

$$n(\sigma) = \sum_{i=1}^r (\bar{\sigma}^{-1}(i) - i) +_{l+1-r} C_2$$

(Takeuchi [2]). We see easily that

$$\begin{aligned}(\omega_l, \beta) &= 2 \quad \text{for any } \beta \in \Delta(\mathfrak{n}^+), \\ (\sigma \delta, \varepsilon_i) &= \begin{cases} (l+1-\bar{\sigma}^{-1}(i)) & \text{if } 1 \leq i \leq r \\ -(l+1-\bar{\sigma}^{-1}(i)) & \text{if } r < i \leq l. \end{cases}\end{aligned}$$

Therefore we have to prove that the following inequalities are true for any  $\sigma \in W^1$

$$(2.1) \quad \#\{\beta \in \Delta(\mathfrak{n}^+); (\sigma \delta, \beta) \geq 2(l+1) - 4n(\sigma)\} > n(\sigma) - 1,$$

$$(2.2) \quad \#\{\beta \in \Delta(\mathfrak{n}^+); (\sigma \delta, \beta) \geq 2(l+3) - 4n(\sigma)\} > n(\sigma) - 2.$$

Since  $(\sigma \delta, \beta) \geq -2l$ ,  $\beta \in \Delta(\mathfrak{n}^+)$ , we immediately see that if  $n(\sigma) \geq l+1$  (resp.  $l+2$ ), the inequality (2.1) (resp. (2.2)) is true for any  $\sigma \in W^1$ .

**Lemma 2.1.** *Let  $\sigma \in W^1$ . If  $n(\sigma) \geq l$ , the inequality (2.1) is true.*

Proof. From the above notice we can assume that  $n(\sigma)=l$ . In this case

$$2(l+1)-4n(\sigma) = 2-2l.$$

It is easy to see that

$$\#\{\beta \in \Delta(\mathfrak{n}^+); (\sigma\delta, \beta) < 2-2l\} \leq 2.$$

Hence

$$\#\{\beta \in \Delta(\mathfrak{n}^+); (\sigma\delta, \beta) \geq 2-2l\} \geq l+1 - C_2 - 2 > l-1 = n(\sigma)-1.$$

Q.E.D.

**Lemma 2.2.** *Let  $\sigma \in W^1$ . If  $n(\sigma) \geq l$ , the inequality (2.2) is true.*

Proof. If  $n(\sigma) \geq l+1$ , the inequality is true in the same way as above. Therefore we may assume that  $n(\sigma)=l$ .

Case 1:  $l=3$ . If  $r=0$ ,  $n(\sigma)=6 \neq 3$ . Hence  $r>0$ , and  $\sigma$  is one of the following elements:

$$\begin{pmatrix} 1 & 2 & 3 \\ 1 & -3 & -2 \end{pmatrix}, \quad \begin{pmatrix} 1 & 2 & 3 \\ 2 & 3 & -1 \end{pmatrix}.$$

In each case (2.2) is true.

Case 2:  $l=4$ . If  $r \leq 1$ ,  $n(\sigma) \geq 6 > 4$ . Hence  $r \geq 2$ . It follows that  $(\sigma\delta, 2\epsilon_1)$ ,  $(\sigma\delta, \epsilon_1+\epsilon_2)$  and  $(\sigma\delta, 2\epsilon_2)$  are larger than  $2(l+3)-4n(\sigma)=-2$ . On the other hand  $n(\sigma)-2=2$ . Therefore (2.2) is true.

Case 3:  $l \geq 5$ . If  $\beta \in \Delta(\mathfrak{n}^+)$  satisfies

$$(\sigma\delta, \beta) < 2(l+3)-4n(\sigma) = 6-2l,$$

$\beta$  is one of the following 12 elements:

$$\begin{aligned} & 2\epsilon_l, \epsilon_l+\epsilon_{l-1}, \epsilon_l+\epsilon_{l-2}, \epsilon_l+\epsilon_{l-3}, \epsilon_l+\epsilon_{l-4}, \epsilon_l+\epsilon_{l-5}, \\ & 2\epsilon_{l-1}, \epsilon_{l-1}+\epsilon_{l-2}, \epsilon_{l-1}+\epsilon_{l-3}, \epsilon_{l-1}+\epsilon_{l-4}, 2\epsilon_{l-2}, \epsilon_{l-2}+\epsilon_{l-3}. \end{aligned}$$

On the other hand

$$\begin{aligned} & l+1 - C_2 - 12 - (l-2) \\ & = \frac{1}{2}\{(l(l+1)-20-2l\} \\ & = \frac{1}{2}(l^2+l-20) \\ & = \frac{1}{2}(l+4)(l-5) \geq 0. \end{aligned}$$

The equality holds only in the case  $l=5$ . But if  $l=5$ ,  $\epsilon_l+\epsilon_{l-5} \notin \Delta(\mathfrak{n}^+)$ . Therefore the inequality is true. Q.E.D.

**Lemma 2.3.** *Let  $\sigma \in W$ . If  $\sigma(1) \neq 1$ ,  $n(\sigma) \geq l$ .*

Proof. By the assumption,

$$\sum_{i=1}^r (\sigma(i) - i) \geq r.$$

Hence

$$\begin{aligned} n(\sigma) - l &\geq r + l_{+1} - C_2 - l \\ &= \frac{1}{2}(l - r - 1)(l - r) \geq 0. \end{aligned} \quad \text{Q.E.D.}$$

We shall prove that the inequality (2.1) is true for any  $\sigma \in W^1$  by using induction on  $l$ . Let  $l=3$ . If  $n(\sigma) \geq 3$ , the inequality is true by Lemma 2.1. If  $n(\sigma)=0$ , the inequality is also true for  $n(\sigma)-1 < 0$ . If  $n(\sigma)=1$  (resp. 2),

$$\sigma = \begin{pmatrix} 1 & 2 & 3 \\ 1 & 2 & -3 \end{pmatrix} \left( \text{resp. } \begin{pmatrix} 1 & 2 & 3 \\ 1 & 3 & -2 \end{pmatrix} \right),$$

and (2.1) is true.

Let  $l=l_0 > 3$ . By Lemmas 2.1 and 2.3, we may assume that  $\sigma(1)=1$ . Define the element  $\tau \in W^1$ , which is considered as an element of  $W^1$  for  $l=l_0-1$ , by

$$\tau^{-1} = \begin{pmatrix} 1 & \cdots & r-1 & r & \cdots & l_0-1 \\ \bar{\sigma}^{-1}(2)-1 & \cdots & \bar{\sigma}^{-1}(r)-1 & -(\bar{\sigma}^{-1}(r+1)-1) & \cdots & -\bar{\sigma}^{-1}(l_0)-1 \end{pmatrix}.$$

We easily see that  $n(\tau)=n(\sigma)$ . By the assumption of the induction,

$$\#\{\varepsilon_i + \varepsilon_j; 1 \leq i, j \leq l_0-1, (\tau\delta', \varepsilon_i + \varepsilon_j) \geq 2l_0 - 4n(\tau)\} > n(\tau) - 1,$$

where  $\delta' = (l_0-1)\varepsilon_1 + (l_0+2)\varepsilon_2 + \cdots + \varepsilon_{l_0-1}$ . It follows, by the fact that  $(\tau\delta', \varepsilon_{i-1}) = (\sigma\delta, \varepsilon_i)$  for  $2 \leq i \leq l_0$ , that

$$(2.3) \quad \#\{\varepsilon_i + \varepsilon_j; 2 \leq i, j \leq l_0, (\sigma\delta, \varepsilon_i + \varepsilon_j) \geq 2l_0 - 4n(\sigma)\} > n(\sigma) - 1.$$

**Lemma 2.4.** *Let*

$$\begin{aligned} s &= \#\{\varepsilon_i; 2 \leq i \leq l_0, \exists \varepsilon_j, 2 \leq j \leq l_0, j \neq i, \text{ such that} \\ &(\sigma\delta, \varepsilon_i + \varepsilon_j) = 2l_0 - 4n(\sigma) \text{ or } 2l_0 + 1 - 4n(\sigma)\}. \end{aligned}$$

*Then*

$$\begin{aligned} \#\{\varepsilon_i + \varepsilon_j; 2 \leq i < j \leq l_0, (\sigma\delta, \varepsilon_i + \varepsilon_j) = 2l_0 - 4n(\sigma) \text{ or} \\ 2l_0 + 1 - 4n(\sigma)\} \leq s - 1. \end{aligned}$$

Proof. Let  $\varepsilon_i, 2 \leq i \leq l_0$ , satisfy the condition that there exists  $\varepsilon_j, 2 \leq j \leq l_0, j \neq i$ , such that  $(\sigma\delta, \varepsilon_i + \varepsilon_j) = 2l_0 - n(\sigma)$  or  $2l_0 + 1 - n(\sigma)$ . For the element  $\varepsilon_i$

$$(2.4) \quad \begin{aligned} \#\{\varepsilon_i + \varepsilon_j; 2 \leq j \leq l_0, j \neq i, (\sigma\delta, \varepsilon_i + \varepsilon_j) = 2l_0 - 4n(\sigma) \\ \text{or } 2l_0 + 1 - 4n(\sigma)\} \leq 2. \end{aligned}$$

In this way we find at most  $2s$  ordered pairs  $(i, j)$ ,  $2 \leq i, j \leq l_0, j \neq i$ , which satisfies  $(\sigma\delta, \varepsilon_i + \varepsilon_j) = 2l_0 - 4n(\sigma)$  or  $2l_0 + 1 - 4n(\sigma)$ . On the other hand the distinct pairs  $(i, j)$  and  $(j, i)$  induce the same element  $\varepsilon_i + \varepsilon_j$ . Therefore

$$(2.5) \quad \#\{\varepsilon_i + \varepsilon_j; 2 \leq i < j \leq l_0, (\sigma\delta, \varepsilon_i + \varepsilon_j) = 2l - 4n(\sigma) \text{ or } 2l + 1 - 4n(\sigma)\} \leq s,$$

and the equality holds if and only if the equality in (2.4) holds for any  $\varepsilon_i, 2 \leq i \leq l_0$ , such that  $(\sigma\delta, \varepsilon_i + \varepsilon_j) = 2l - n(\sigma)$  or  $2l + 1 - n(\sigma)$ .

Define the integer  $i_0$  (resp.  $i_m$ ) by

$$\begin{aligned} & \min(\text{resp. max}) \{i; 2 \leq i \leq l_0, \exists j, 2 \leq j \leq l_0, j \neq i \text{ such that} \\ & (\sigma\delta, \varepsilon_i + \varepsilon_j) = l - 2n(\sigma) \text{ or } l + 1 - 2n(\sigma)\}. \end{aligned}$$

If the equalities in (2.4) for  $\varepsilon_{i_0}$  and  $\varepsilon_{i_m}$  hold, there exist the integers  $i$  and  $j$  such that

$$\begin{aligned} & (\sigma\delta, \varepsilon_{i_0} + \varepsilon_j) = l - 2n(\sigma) \text{ or } l + 1 - 2n(\sigma), \\ & (\sigma\delta, \varepsilon_i + \varepsilon_{i_m}) = l - 2n(\sigma) \text{ or } l + 1 - 2n(\sigma), \\ & i_0 < i \text{ and } j < i_m. \end{aligned}$$

Hence

$$(\sigma\delta, \varepsilon_i + \varepsilon_{i_m}) \leq (\sigma\delta, \varepsilon_{i_0} + \varepsilon_j) - 2.$$

This is impossible, and therefore, the equality in (2.5) does not hold. Q.E.D.

Let  $\varepsilon_i, 2 \leq i \leq l_0$ , satisfy that there exists  $\varepsilon_j, 2 \leq j \leq l_0, j \neq i$ , such that

$$(\sigma\delta, \varepsilon_i + \varepsilon_j) = 2l - 4n(\sigma) \text{ or } 2l + 1 - 4n(\sigma).$$

For this element  $\varepsilon_i$ ,

$$(\sigma\delta, \varepsilon_i + \varepsilon_1) \geq 2l + 2 - 4n(\sigma),$$

in all but the following case:

$$(\sigma\delta, \varepsilon_i + \varepsilon_2) = 2l - 4n(\sigma).$$

Therefore, by Lemma 2.4,

$$\begin{aligned} & \#\{\varepsilon_i + \varepsilon_j; 1 \leq i < j \leq l_0, (\sigma\delta, \varepsilon_i + \varepsilon_j) \geq 2l_0 + 2 - 4n(\sigma)\} \\ & \geq \#\{\varepsilon_i + \varepsilon_j; 2 \leq i < j \leq l_0, (\sigma\delta, \varepsilon_i + \varepsilon_j) \geq 2l_0 - 4n(\sigma)\}. \end{aligned}$$

There exist at most one element  $\varepsilon_i, 2 \leq i \leq l_0$ , such that

$$(\sigma\delta, 2\varepsilon_i) = 2l - 4n(\sigma) \text{ or } 2l + 1 - 4n(\sigma).$$

If such  $\varepsilon_i$  exists,


$$(\sigma\delta, 2\varepsilon_i) \geq 2l + 2 - 4n(\sigma).$$

Therefore the inequality (2.1) is true.

Thus we have proved that the inequality (2.1) is true for any  $\sigma \in W^1$ .

From Lemmas 2.2 and 2.3, we can prove that the inequality (2.2) is true for any  $\sigma \in W^1$  in the same way as above.

**2.3. The case that  $M$  is of type DIII, that is  $M = SO(2l)/U(l)$ .** If  $l=3$ ,  $M = P_3(C)$ . If  $l \geq 4$ ,  $M$  is a complex quadric of dimension 6. Hence we assume that  $l \geq 5$ . In this case  $n = \frac{1}{2}l(l-1)$  and  $\lambda = 2l-2$ . The Dinkin diagram of  $\Pi$  is as follows:



where  $\alpha_l \odot$  shows  $\alpha_j = \alpha_l$ . Let  $\{\varepsilon_i; 1 \leq i \leq l\}$  be the basis of  $\mathfrak{h}_0$  which satisfies  $(\varepsilon_i, \varepsilon_j) = \delta_{ij}$ . Then we have:

$$\begin{aligned}\Delta &= \{\pm \varepsilon_i \pm \varepsilon_j; 1 \leq i < j \leq l\}, \\ \Pi &= \{\alpha_1 = \varepsilon_1 - \varepsilon_2, \dots, \alpha_{l-1} = \varepsilon_{l-1} - \varepsilon_l, \alpha_l = \varepsilon_{l-1} + \varepsilon_l\}, \\ \Delta(\mathfrak{n}^+) &= \{\varepsilon_i + \varepsilon_j; 1 \leq i < j \leq l\}, \\ \delta &= (l-1)\varepsilon_1 + (l-2)\varepsilon_2 + \dots + \varepsilon_{l-1}, \\ \omega &= \frac{1}{2}(\varepsilon_1 + \dots + \varepsilon_l).\end{aligned}$$

An element  $\sigma \in W$  acts on  $\mathfrak{h}_0$  by  $\sigma \varepsilon_i = \pm \varepsilon_{\bar{\sigma}(i)}$  for  $1 \leq i \leq l$ , where  $\bar{\sigma}$  is a permutation of  $\{1, 2, \dots, l\}$ . We denote the element  $\sigma \in W$  by the symbol

$$\begin{pmatrix} 1 & 2 & \dots & l \\ \pm \bar{\sigma}(1) & \pm \bar{\sigma}(2) & \dots & \pm \bar{\sigma}(l) \end{pmatrix}.$$

Then

$$\begin{aligned}W^1 &= \left\{ \sigma \in W; \sigma^{-1} = \begin{pmatrix} 1 & \dots & r & r+1 & \dots & l \\ \bar{\sigma}^{-1}(1) & \dots & \bar{\sigma}^{-1}(r) & -\bar{\sigma}^{-1}(r+1) & \dots & -\bar{\sigma}^{-1}(l) \end{pmatrix}, \right. \\ &\quad \left. l-r \text{ is even, } \bar{\sigma}^{-1}(1) < \dots < \bar{\sigma}^{-1}(r), \bar{\sigma}^{-1}(r+1) > \dots > \bar{\sigma}^{-1}(l) \right\}.\end{aligned}$$

The index  $n(\sigma)$  of  $\sigma \in W^1$  is given by

$$n(\sigma) = \sum_{i=1}^r (\bar{\sigma}^{-1}(i) - i) +_{l-r} C_2$$

(Takeuchi [2]). We see easily that

$$\begin{aligned}(\omega_l, \beta) &= 1 \quad \text{for any } \beta \in \Delta(\mathfrak{n}^+), \\ (\sigma \delta, \varepsilon_i) &= \begin{cases} l - \bar{\sigma}^{-1}(i) & \text{if } 1 \leq i \leq r \\ -(l - \bar{\sigma}^{-1}(i)) & \text{if } r < i \leq l. \end{cases}\end{aligned}$$

Therefore we have to prove that the following inequalities are true for any  $\sigma \in W^1$ .

$$(3.1) \quad \#\{\beta \in \Delta(\mathfrak{n}^+); (\sigma\delta, \beta) \geq 2l-2-2n(\sigma)\} > n(\sigma)-1,$$

$$(3.2) \quad \#\{\beta \in \Delta(\mathfrak{n}^+); (\sigma\delta, \beta) \geq 2l-2n(\sigma)\} > n(\sigma)-2.$$

**Lemma 3.1.** *Let  $\sigma \in W^1$ . If  $n(\sigma) \geq 2l-3$ , the inequality (3.1) is true.*

Proof. By the assumption  $2l-2-2n(\sigma) \leq 4-2l$ . Let  $\beta$  be an element of  $\Delta(\mathfrak{n}^+)$  which satisfies that

$$(\sigma\delta, \beta) < 4-2l,$$

then  $\beta = \varepsilon_{l-1} + \varepsilon_l$ . Therefore

$$\#\{\beta \in \Delta(\mathfrak{n}^+); (\sigma\delta, \beta) \geq 2l-2-2n(\sigma)\} \geq n-1.$$

If the equality holds,  $n(\sigma) = 2l-3$  and  $n-n(\sigma) = \frac{1}{2}(l-2)(l-3) > 0$ . Q.E.D.

**Lemma 3.2.** *Let  $\sigma \in W^1$ . If  $n(\sigma) \geq 2l-3$ , the inequality (3.2) is true.*

Proof. If  $n(\sigma) \geq 2l-3$ , the inequality is true in the same way as above. Therefore we assume that  $n(\sigma) = 2l-3$ . The number of the elements  $\beta \in \Delta(\mathfrak{n}^+)$  such that

$$(\sigma\delta, \beta) < 2l-2n(\sigma) = 6-2l$$

is at most 4. Since  $l \geq 5$ ,

$$(n-4)-(n(\sigma)-2) = \frac{1}{2}l(l-1)-4-2l+5 = \frac{1}{2}l(l-5)+1 > 0.$$

Q.E.D.

**Lemma 3.3.** *If  $\bar{\sigma}^{-1}(1) \geq 3$ , then  $n(\sigma) \geq 2l-3$ .*

Proof. By the assumption

$$\sum_{i=1}^r (\bar{\sigma}^{-1}(i) - i) \geq 2r.$$

It follows that

$$\begin{aligned} & n(\sigma) - (2l-3) \\ & \geq 2r + l-rC_2 - (2l-3) \\ & = \frac{1}{2}(l-r-2)(l-r-3) \geq 0. \end{aligned}$$

Q.E.D.

We prove that the inequality (3.1) is true for all  $\sigma \in W^1$  by using induction on  $l$ . If  $l=5$ , we easily see that the inequality is true.

Let  $l=l_0 > 5$ . By Lemmas 3.1 and 3.3, we can assume that  $\bar{\sigma}^{-1}(1)=1$  or 2.

Case 1:  $\bar{\sigma}^{-1}(1)=1$ . Define the element  $\tau \in W^1$ , which is considered as an element of  $W^1$  for  $l=l_0-1$ , by

$$\tau^{-1} = \begin{pmatrix} 1 & \cdots & r-1 & r & \cdots & l_0-1 \\ \bar{\sigma}^{-1}(2)-1 & \cdots & \bar{\sigma}^{-1}(r)-1 & -(\bar{\sigma}^{-1}(r+1)-1) & \cdots & \bar{\sigma}^{-1}((l_0)-1) \end{pmatrix}.$$

Then  $n(\tau)=n(\sigma)$ . By the assumption of the induction,

$$\#\{\varepsilon_i + \varepsilon_j; 2 \leq i < j \leq l_0, (\sigma\delta, \varepsilon_i + \varepsilon_j) \geq 2l_0 - 4 - 2n(\sigma)\} > n(\sigma) - 1.$$

Let

$$\begin{aligned} s &= \#\{\varepsilon_i; 2 \leq i \leq l_0, \exists \varepsilon_j, 2 \leq j \neq i \leq l_0, \text{ such that} \\ &(\sigma\delta, \varepsilon_i + \varepsilon_j) = 2l_0 - 4 - 2n(\sigma) \text{ or } 2l_0 - 3 - 2n(\sigma)\}. \end{aligned}$$

Then, in the same way as in Lemma 2.4, we see that

$$\#\{\varepsilon_i + \varepsilon_j; 2 \leq i < j \leq l_0, (\sigma\delta, \varepsilon_i + \varepsilon_j) = 2l_0 - 4 - 2n(\sigma) \text{ or } 2l_0 - 3 - 2n(\sigma)\} \leq s - 1.$$

Let  $\varepsilon_i$  satisfy that there exists  $\varepsilon_j, 2 \leq j \leq l_0, j \neq i$ , such that

$$(\sigma\delta, \varepsilon_i + \varepsilon_j) = 2l_0 - 4 - 2n(\sigma) \text{ or } 2l_0 - 3 - 2n(\sigma).$$

Then

$$(\sigma\delta, \varepsilon_i + \varepsilon_1) \geq 2l_0 - 2 - 2n(\sigma)$$

in all but the following case:

$$(\sigma\delta, \varepsilon_i + \varepsilon_2) = 2l_0 - 4 - 2n(\tau) \text{ and } \bar{\sigma}^{-1}(2) = 2.$$

Therefore the inequality is true.

Case 2:  $\bar{\sigma}^{-1}(1)=2$ . By the definition of  $W^1$

$$\sigma^{-1} = \begin{pmatrix} 1 & 2 & \cdots & r & r+1 & \cdots & l_0 \\ 2 & \bar{\sigma}^{-1}(2) & \cdots & \bar{\sigma}^{-1}(r) & -\bar{\sigma}^{-1}(r+1) & \cdots & -1 \end{pmatrix}.$$

Define the element  $\sigma' \in W^1$  by

$$(\sigma')^{-1} = \begin{pmatrix} 1 & 2 & \cdots & r & r+1 & \cdots & l_0-1 & l_0 \\ 1 & \bar{\sigma}^{-1}(2) & \cdots & \bar{\sigma}^{-1}(r) & -\bar{\sigma}^{-1}(r+1) & \cdots & -\bar{\sigma}^{-1}(l_0-1) & -2 \end{pmatrix}.$$

Then  $n(\sigma')=n(\sigma)-1$ . Define another element  $\tau \in W^1$ , which is considered for  $l=l_0-1$ , by

$$\tau^{-1} = \begin{pmatrix} 1 & \cdots & r & r+1 & \cdots & l_0-1 \\ \bar{\sigma}^{-1}(2)-1 & \cdots & \bar{\sigma}^{-1}(r)-1 & -(\bar{\sigma}^{-1}(r+1)-1) & \cdots & -1 \end{pmatrix}.$$

Then  $n(\tau)=n(\sigma')$ .

Assume that the inequality (3.2) is true for  $\tau$ . If we notice that  $(\overline{\sigma'})^{-1}(2) > 2$ , we get the following inequality in the same way as in case 1.

$$\#\{\beta \in \Delta(\mathfrak{n}^+); (\sigma'\delta, \beta) \geq 2l_0 - 2 - 2n(\sigma')\} > n(\sigma') .$$

Clearly

$$(\sigma\delta, \beta) \geq (\sigma'\delta, \beta) - 2 \quad \text{for any } \beta \in \Delta(\mathfrak{n}^+) .$$

Hence if  $\beta \in \Delta(\mathfrak{n}^+)$  satisfies that

$$(\sigma'\delta, \beta) \geq 2l_0 - 2 - 2n(\sigma') ,$$

then

$$(\sigma\delta, \beta) \geq 2l_0 - 2 - 2n(\sigma) .$$

Therefore

$$\#\{\beta \in \Delta(\mathfrak{n}^+); (\sigma\delta, \beta) \geq 2l_0 - 2 - 2n(\sigma)\} > n(\sigma) - 1 .$$

Thus we have proved that the inequality (3.1) is true for any  $\sigma \in W^1$ . We can prove that the inequality (3.2) is true in the same way as above.

### References

- [1] Y. Kimura: *On the hypersurfaces of Hermitian symmetric spaces of compact type*, Osaka J. Math. **16** (1979), 97–119.
- [2] M. Takeuchi: *Cell decompositions and Morse equalities on certain symmetric spaces*, J. Fac. Sci. Univ. Tokyo **12** (1965), 81–192.

Kobe University of Commerce  
Tarumi, Kobe 655, Japan

