

Title	GENUS ZERO PALF STRUCTURES ON THE AKBULUT-YASUI PLUGS
Author(s)	Ukida, Takuya
Citation	Osaka Journal of Mathematics. 2022, 59(2), p. 363-368
Version Type	VoR
URL	https://doi.org/10.18910/87482
rights	
Note	

Osaka University Knowledge Archive : OUKA

<https://ir.library.osaka-u.ac.jp/>

Osaka University

GENUS ZERO PALF STRUCTURES ON THE AKBULUT-YASUI PLUGS

TAKUYA UKIDA

(Received June 11, 2019, revised January 13, 2021)

Abstract

We construct a genus zero PALF structure on each of plugs introduced by Akbulut and Yasui and describe the monodromy as a positive factorization in the mapping class group of a fiber.

1. Introduction

The problem of classifying all differential structures defined on a given 4-manifold is an important problem in understanding the overall picture of a 4-manifold.

Akbulut and Yasui [4] introduced corks and plugs. Corks and plugs are compact Stein surfaces. Matveyev, Curtis-Freedman-Hsiang-Stong, and Akbulut-Matveyev's theorem show that the study of exotic manifold pairs constructed using cork is important for the classification problem of the differential structure of 4-manifolds.

Theorem 1.1 (Matveyev [12], Curtis-Freedman-Hsiang-Stong [7], Akbulut-Matveyev [2]). *For every homeomorphic but non-diffeomorphic pair of simply connected closed 4-manifolds, one is obtained from the other by removing a contractible 4-manifold and gluing it via an involution on the boundary. Such a contractible 4-manifold has since been called a Cork. Furthermore, corks and their complements can always be made compact Stein 4-manifolds.*

It is shown by Akbulut and Yasui [6] using cork that an infinite number of exotic Stein surface pairs embedded in X exist for any four-dimensional two-handle body X with $b_2(X) \geq 1$.

The plug generalizes the Gluck twist. The plug is also used to make exotic manifolds, as well as cork.

On the other hand, Loi and Piergallini [11] proved that every compact Stein surface admits a positive allowable Lefschetz fibration over D^2 (a PALF for short). Therefore we can investigate compact Stein surfaces in terms of positive factorizations in mapping class groups (see also Akbulut and Ozbagci [3], Akbulut and Arıkan [1]).

Since corks and plugs are Stein surfaces, the study of the relationship between Stein surfaces and mapping class groups using PALFs plays an important role in classifying differential structures on 4-manifolds.

If a PALF is created from a Stein surface by the existing method ([11], [3], [1]), its genus will be large, and it will be complicated and difficult to handle as a mapping class

group element.

Gompf [8] indicates that the Stein surface is compatible with Kirby calculus. In this paper, we use Kirby calculus to construct PALFs on Akbulut-Yasui plugs realizing the smallest possible fiber genera.

One planar (i.e. genus zero) PALF on the Akbulut cork was made in the previous paper of the author [14], but in this paper, we made an infinite number of planar PALFs on the Akbulut-Yasui plugs. Being planar is also playing an important role in [10].

In this paper, we construct a genus zero PALF structure on each of plugs introduced by Akbulut and Yasui [4] and describe the monodromy as a positive factorization in the mapping class group of a fiber.

Theorem 1.2. *For any $m \geq 1, n \geq 2$, the Akbulut-Yasui plug $(W_{m,n}, f_{m,n})$ admits a genus zero PALF structure. The monodromy of the PALF is described by the factorization $t_{\alpha_{2n+m}} \cdots t_{\alpha_1}$, where t_α is a right-handed Dehn twist along a simple closed curve α on a fiber and $\alpha_{2n+m}, \dots, \alpha_1$ are simple closed curves shown in Figure 2.*




Fig. 1. Kirby diagram for $W_{m,n}$




Fig. 2. Vanishing cycles of a genus zero PALF on $W_{m,n}$.

Note that the genus of a PALF on the manifold $W_{m,n}$ in a known way (cf. [3] and [1]) is much more than zero. We obtained similar results for the Akbulut cork W_1 [14]. In the present paper, we construct a genus zero PALF on an infinite number of the Akbulut-Yasui plugs.

2. Preliminaries

2.1. Mapping class groups. In this subsection, we review a precise definition of the mapping class groups of surfaces with boundary and that of Dehn twists along simple closed curves on surfaces.

DEFINITION 2.1. Let F be a compact oriented connected surface with boundary. Let $\text{Diff}^+(F, \partial F)$ be the group of all orientation-preserving self-diffeomorphisms of F fixing the boundary ∂F point-wise. Let $\text{Diff}_0^+(F, \partial F)$ be the subgroup of $\text{Diff}^+(F, \partial F)$ consisting of self-diffeomorphisms isotopic to the identity. The quotient group $\text{Diff}^+(F, \partial F) / \text{Diff}_0^+(F, \partial F)$ is called the mapping class group of F and it is denoted by $\text{Map}(F, \partial F)$.

DEFINITION 2.2. A *positive (or right-handed) Dehn twist* along a simple closed curve α , $t_\alpha : F \rightarrow F$ is a diffeomorphism obtained by cutting F along α , twisting 360° to the right and regluing.

2.2. PALF.

DEFINITION 2.3. Let M^4 and B^2 be compact oriented smooth manifolds of dimensions 4 and 2. Let $f : M \rightarrow B$ be a smooth map. f is called a *positive Lefschetz fibration* over B if it satisfies the following conditions (1) and (2):

- (1) There are finitely many critical values b_1, \dots, b_m of f in the interior of B and there is a unique critical point p_i on each fiber $f^{-1}(b_i)$, and
- (2) The map f is locally written as $f(z_1, z_2) = z_1^2 + z_2^2$ with respect to some local complex coordinates around p_i and b_i compatible with the orientations of M and B .

DEFINITION 2.4. A positive Lefschetz fibration is called *allowable* if its all vanishing cycles are homologically non-trivial on the fiber. A positive allowable Lefschetz fibration over D^2 with bounded fibers is called a *PALF* for short.

The following Lemma is useful to prove Theorem 1.2.

Lemma 2.5 (cf. Akbulut-Ozbagci [3, Remark 1]). *Suppose that a 4-manifold X admits a PALF. If a 4-manifold Y is obtained from X by attaching a Lefschetz 2-handle, then Y also admits a PALF.*

The Lefschetz 2-handle is defined as follows.

DEFINITION 2.6. Suppose that X admits a PALF. A *Lefschetz 2-handle* is a 2-handle attached along a homologically non-trivial simple closed curve in the boundary of X with framing -1 relative to the product framing induced by the fiber structure.

2.3. Stein surfaces. In this subsection, we recall a definition of the Stein surfaces. The question of which smooth 4-manifolds admit Stein structures can be completely reduced to a problem in handlebody theory.

DEFINITION 2.7. A complex manifold is called a *Stein manifold* if it admits a proper bi-holomorphic embedding to \mathbb{C}^n .

DEFINITION 2.8. Let W be a compact manifold with boundary. The manifold W is called a *Stein domain* if it satisfies following condition: There is a Stein manifold X and a plurisubharmonic function $\varphi : X \rightarrow [0, \infty)$ such that $W = \varphi^{-1}([0, a])$ for a regular value a of φ .

DEFINITION 2.9. A Stein manifold or a Stein domain is called a *Stein surface* if its complex dimension is 2.

2.4. Plugs. In this subsection, we give the definition of the plug.

DEFINITION 2.10 (AKBULUT-YASUI [4, Definition 2.2.]). Let P be a compact Stein 4-manifold with boundary and $\tau : \partial P \rightarrow \partial P$ an involution on the boundary, which cannot extend to any self-homeomorphism of P . We call (P, τ) a *Plug* of X , if $P \subset X$ and X keeps its homeomorphism type and changes its diffeomorphism type when removing P and gluing it via τ . We call (P, τ) a *Plug* if there exists a smooth 4-manifold X such that (P, τ) is a plug of X .

DEFINITION 2.11 (AKBULUT-YASUI [4, Definition 2.3.]). Let $W_{m,n}$ be a smooth 4-manifold given by Figure 1. Let $f_{m,n} : \partial W_{m,n} \rightarrow \partial W_{m,n}$ be the obvious involution obtained from first surgering $S^1 \times D^3$ to $D^2 \times S^2$ in the interiors of $W_{m,n}$, then surgering the other imbedded $D^2 \times S^2$ back to $S^1 \times D^2$ (i.e. replacing the dot in Figure 1).

Theorem 2.12 (Akbulut-Yasui [4, Theorem 2.5(2)]). *For $m \geq 1$ and $n \geq 2$, the pair $(W_{m,n}, f_{m,n})$ is a plug.*

3. Proof of Theorem 1.2.

In this section, we give the proof of Theorem 1.2.

Proof of Theorem 1.2. Let $F_{m,n}$ be the compact oriented surface of genus zero with $2n + m$ boundary components and $\alpha_1, \dots, \alpha_{2n+m}$ the curves on $F_{m,n}$ shown in Figure 4 (a). Note that Figure 2 and Figure 4 (a) show the same PALF. We denote the right-handed Dehn twists along $\alpha_1, \dots, \alpha_{2n+m}$ by $t_{\alpha_1}, \dots, t_{\alpha_{2n+m}}$, respectively. Let $f : X_{m,n} \rightarrow D^2$ be a Lefschetz fibration over D^2 with monodromy representation $(t_{\alpha_{2n+m}}, \dots, t_{\alpha_1})$. Since each curve α_i is homologically non-trivial on $F_{m,n}$, we see that f is a PALF with fiber $F_{m,n}$.

We now show that $X_{m,n}$ is diffeomorphic to $W_{m,n}$.

The Kirby diagram for $X_{m,n}$ corresponding to the monodromy representation $(t_{\alpha_{2n+m}}, \dots,$




Fig. 3




Fig.4

t_{α_1}) is given by Figure 4 (b). We slide the -1 -framed 2-handles over -1 -framed 2-handles and erase canceling 1-handle/2-handle pairs to get Figure 4 (c). We get Figure 4 (d) by sliding the $-m$ -framed 2-handle over -1 -framed 2-handles and sliding the $-n$ -framed 2-handle over -1 -framed 2-handles and erasing canceling 1-handle/2-handle pairs.

The Kirby diagram for $W_{m,n}$ is given by Figure 3 (a). We slide the 0-framed 2-handle under the 1-handle to get Figure 3 (b).

Since Figure 3 (b) and Figure 4 (d) are the same, we conclude that $X_{m,n}$ is diffeomorphic to $W_{m,n}$, which implies the theorem. \square

ACKNOWLEDGEMENTS. The author would like to thank Hisaaki Endo for his helpful comments and his encouragement. The author wishes to thank Kouichi Yasui and Yuichi Yamada for their useful comments.

References

- [1] S. Akbulut and M.F. Arıkan: *A note on Lefschetz fibrations on compact Stein 4-manifolds*, Commun. Contemp. Math. **14** (2012), 1250035, 14pp.
- [2] S. Akbulut and R. Matveyev: *A convex decomposition theorem for 4-manifolds*, Internat. Math. Res. Notices. (1998), 371–381.
- [3] S. Akbulut and B. Ozbagci: *Lefschetz fibration on compact Stein surfaces*, Geom. Topol. **5** (2001), 319–334.
- [4] S. Akbulut and K. Yasui: *Corks, Plugs and exotic structures*, J. Gökova Geometry Topology, GGT **2** (2008), 40–82.
- [5] S. Akbulut and K. Yasui: *Small exotic Stein manifolds*, Comment. Math. Helv. **85** (2010), 705–721.
- [6] S. Akbulut and K. Yasui: *Cork twisting exotic Stein 4-manifolds*, J. Differential Geom. **93** (2013), 1–36.
- [7] C.L. Curtis, M.H. Freedman, W.C. Hsiang, and R. Stong: *A decomposition theorem for h-cobordant smooth simply-connected compact 4-manifolds*, Invent. Math. **123** (1996), 343–348.
- [8] R. Gompf: *Handlebody construction of Stein surfaces*, Ann. of Math. **148** (1998), 619–693.
- [9] R.E. Gompf and A.I. Stipsicz: *4-manifolds and Kirby calculus*, Graduate Studies in Math. **20**, American Mathematical Society, Providence, RI, 1999.
- [10] Ç. Karakurt, T. Oba, T. Ukida: *Planar Lefschetz fibrations and Stein structures with distinct Ozsváth-Szabó invariants on corks*, Topology Appl. **221** (2017), 630–637.
- [11] A. Loi and R. Piergallini: *Compact Stein surfaces with boundary as branched covers of B^4* , Invent. Math. **143** (2001), 325–348.
- [12] R. Matveyev: *A decomposition of smooth simply-connected h-cobordant 4-manifolds*, J. Differential Geom. **44** (1996), 571–582.
- [13] B. Ozbagci and A.I. Stipsicz: *Surgery on contact 3-manifolds and Stein surfaces*, Bolyai Society Mathematical Studies, **13**, Springer-Verlag, Berlin; János Bolyai Mathematical Society, Budapest, 2004.
- [14] T. Ukida: *A genus zero Lefschetz fibration on the Akbulut cork*, Topology Appl. **214** (2016), 127–136.

Department of Creative Technology Engineering
National Institute of Technology, Anan College
265 Aoki, Minobayashi-chou, Anan-shi
Tokushima 774–0017
Japan
e-mail: ukida@anan-nct.ac.jp