

Title	A characterization of QF-algebras
Author(s)	Harada, Manabu
Citation	Osaka Journal of Mathematics. 1983, 20(1), p. 1-4
Version Type	VoR
URL	https://doi.org/10.18910/8764
rights	
Note	

Osaka University Knowledge Archive : OUKA

<https://ir.library.osaka-u.ac.jp/>

Osaka University

A CHARACTERIZATION OF QF-ALGEBRAS

MANABU HARADA

(Received February 19, 1981)

We have defined a new class of rings in [3] which we call self mini-injective rings and we have noted that there exist artinian rings in the new class which are not quasi-Frobenius rings (briefly QF-rings).

We shall show in this note that if a ring R is an algebra over a field with finite dimension, then a self mini-injective algebra is a QF-algebra.

Throughout this note we assume a ring R contains the identity and every module is a unitary right R -module. We shall refer for the definitions of mini-injectives and the extending property, etc. to [3].

Let K be a field and R a K -algebra with finite dimension over K .

Theorem 1 (cf. [3], Theorems 13 and 14). *Let R be as above. Then the following conditions are equivalent.*

- 1) R is self mini-injective as a right R -module.
- 2) R is self mini-injective as a left R -module.
- 3) Every projective right R -module has the extending property of direct decomposition of the socle.
- 4) Every projective left R -module has the extending property of direct decomposition of the socle.
- 5) R is a QF-algebra.

Proof. R is self-injective as a left or right R -module if and only if R is a QF-algebra by [2]. In this case R is self-injective as both a right and left R -module by [1]. It is clear from [3], Theorem 3 and Proposition 8 that 1), 2) are equivalent to 3), 4), respectively. Hence, we may assume R is a basic algebra by [4] and [6].

1) \rightarrow 5). Let $R = \sum_{i=1}^n e_i R$ be the standard decomposition, namely $\{e_i\}$ is a set of mutually orthogonal primitive idempotents and $e_i R \approx e_{i'} R$ if $i \neq i'$. Since R is right self mini-injective, R is right QF-2 by [3], Proposition 8 and $S(e_i R) \approx S(e_{i'} R)$ for $i \neq i'$ by [3], Theorem 5, where $S(\)$ means the socle. Now $e_i R$ is uniform as a right R -module and so the injective envelope $E(e_i R)$ of $e_i R$ is indecomposable. We put $M^* = \text{Hom}_K(M, K)$ for a K -module M . Then $E(e_i R)^*$

is indecomposable and projective as a left R -module. Hence, $E(e_i/R)^* \approx Re_{i'}$ and $E(e_i R) \approx (Re_{i'})^*$. From the fact $E(e_i R) \approx E(e_j R)$ for $i \neq j$, a mapping $\pi: i \rightarrow i'$ is a permutation on $\{1, 2, \dots, n\}$. Accordingly, $\sum_{i=1}^n [E(e_i R): K] = \sum_{i=1}^n [Re_{\pi(i)}: K] = [R: K]$. Therefore, $E(R) = \sum_{i=1}^n \oplus E(e_i R) = R$. The remaining part is clear.

In the above proof we have used only the facts that R is right QF-2 and $S(e_i R) \approx S(e_j R)$ if $i \neq j$. Hence, we have

Theorem 2. *Let R be a K -algebra as above. If R is right QF-2 and $S(eR) \approx S(e'R)$ if $eR \approx e'R$ then R is QF, where e and e' are primitive idempotents, where J is the Jacobson radical of R .*

Corollary. *Let R be the K -algebra as above. We assume R/J is a simple algebra. Then R is a QF-algebra if and only if R is a right QF-2 algebra.*

We note that the above facts are not true for right and left artinian rings (see [3], Example 2).

Next we shall consider a characterization of a right artinian and self mini-injective ring.

Theorem 3. *Let R be a right artinian ring. Then the following conditions are equivalent.*

- 1) R is self mini-injective as a right R -module.
- 2) R satisfies
 - i) if $e_1 R \approx e_2 R$, any minimal right ideal in $e_1 R$ is not isomorphic to one in $e_2 R$.
 - ii) there exists a minimal right ideal I contained in $e_1 J^k$ ($e_1 J^{k+1} = 0$) such

that $\text{End}_R(I) = \{a \in \overline{e_1 R e_1} \mid aI \subseteq I\}$, i.e. $\text{End}_R(I)$ is extended to $\text{End}_R(e_1 R)$ and $S(e_1 R) = e_1 R e_1 I$ for each e_i , where the e_i is primitive idempotent and $S(\)$ is socle and $\bar{R} = R/J$.

Proof. 1) \rightarrow 2). It is clear from [3], Theorem 5. 2) \rightarrow 1). The second part of ii) implies that each minimal right ideal I' in $e_1 R$ is isomorphic to I . We assume $I \approx e_2 \bar{R}$ and $I = xR$, $I' = x'R$. Then we may assume $xe_2 = x$ and $x'e_2 = x'$. We obtain from ii) that $x' = x'e_2 = \sum y_i x r_i$; $y_i \in e_1 R e_1$, $r_i \in R e_2$. Now $x r_i e_2 = x e_2 r_i e_2 = x e_2 r_i e_2$. Since a mapping $xz \rightarrow x e_2 r_i e_2 z$ is an R -endomorphism of I , there exists an element a_i in $\overline{e_1 R e_1}$ with $a_i x = x e_2 r_i e_2$ from ii). Hence, $x' = (\sum y_i a_i) x = \bar{b} x$, where $\bar{b} = \sum y_i a_i$. We quote the proof of [3], Proposition 9. Since $\bar{b} \neq 0$, $x = \bar{b}^{-1} x'$. Put $g(x'z) = xz$; $z \in R$. Let f be any element in $\text{Hom}_R(I, I')$. Then $gf \in \text{End}_R(I)$. Hence, there exists a in $\overline{e_1 R e_1}$ such that $gf(x) = ax$ by ii). Therefore, $f(x) = g^{-1}(ax) = \bar{b} a x$ and f is extended to an element in $\text{End}_R(e_1 R)$. We know similarly that $\text{End}_R(I') = \bar{b}^{-1} \text{End}_R(I) \bar{b} = \{\bar{c} \in \overline{e_1 R e_1} \mid \bar{c} I' \subseteq I'\}$.

Hence, I' satisfies ii). Let $h \in \text{Hom}_R(I', R)$ and $R = \sum_{i=1}^n \oplus e_i R$. Let $\pi_i: R \rightarrow e_i R$ be the projection. If $\pi_i h \neq 0$, $e_i R \approx e_1 R$ by i). We assume $\pi_i h = h_i \neq 0$ for $i=1, 2, \dots, t$ and $h_j = 0$ for $j > t$. Since $e_1 R \approx e_i R$ for $i \leq t$, there exists $c_i \in e_i R e_i$ and $d_i \in e_i R e_1$ such that $c_i d_i = e_1$ and $d_i c_i = e_i$. Using d_i and c_i , we know as above that any element in $\text{Hom}_R(I', h_i(I'))$ is extended to an element in $\text{Hom}_R(e_1 R, e_i R)$ for $i \leq t$. Take $p_i \in R$ such that $p_i x' = h_i(x')$. Then $h(x') = \sum h_i(x') = (\sum p_i)x'$. Hence, R is right self mini-injective by [3], Theorem 2.

REMARK. The above three conditions in Theorem 3, 2) are independent.

Corollary 1. *Let R be a right artinian and right self mini-injective. Then R is a right QF-2 if and only if $\text{End}_R(I) = \overline{e_1 R e_1}$ in ii) of Theorem 3.*

Corollary 2. *Let R be a right artinian ring and e a primitive idempotent. We assume that i) R is right QF-2, ii) any monomorphism of $\overline{e R e}$ into itself as a division ring is isomorphic for each e (e.g. algebraic extension of the prime field) and iii) $S(eR) \approx S(e'R)$ if $eR \approx e'R$. Then R is right self mini-injective.*

Proof. We may assume R is basic. Since $S(eR) \supset eJ^k \neq 0$ ($eJ^{k+1} = 0$), $S(eR) = eJ^k$ by i). Put $S(eR) = uR$. $eJ^k \subset eJ^{k+1} = 0$ and so uR is a left $\overline{e R e}$ -module. We assume $uR \approx \overline{e'R}$. Since R is basic, $\overline{e'R} = \overline{e'R e'}$. Hence, $\overline{u e' R e'} = uR$. Let \bar{x} be in $\overline{e R e}$. Then $\bar{x}u = u\bar{y}$; $\bar{y} \in \overline{e'R e'}$. It is clear that the mapping $\bar{x} \rightarrow \bar{y}$ gives us a monomorphism of the division ring $\overline{e R e}$ into $\overline{e'R e'}$ as a division ring. Repeating this procedure, we can find a chain $e, e', \dots, e^{(t)}$ of primitive idempotents. We know from iii) that $e^{(s)} = e$ for some s (cf. [3], the proof of Proposition 8). Hence, $\overline{e R e} = \overline{u e' R e'}$ by ii). Therefore, R is right self mini-injective by Theorem 3.

We do not know any example of a right QF-2 and right self mini-injective ring which is not QF.

References

- [1] S. Eilenberg and T. Nakayama: *On the dimension of modules and algebras* II, Nagoya Math. J. **9** (1956), 1-16.
- [2] C. Faith: *Rings with ascending condition on annihilators*, ibid. **27** (1966), 179-191.
- [3] M. Harada: *On self mini-injective rings*, Osaka J. Math. **19** (1982), 587-597.
- [4] M. Oshima: *Notes on basic rings*, Math. J. Okayama Univ. **2** (1952-53), 103-110.
- [5] K. Morita: *Duality for modules and its applications to the theory of rings with*

minimum condition, Sci. Rep. Tokyo Kyoiku Daigaku Sect. A **6** (1958), 83–142.

Department of Mathematics
Osaka City University
Sumiyoshi-ku, Osaka 558
Japan