



Title	Clinical implications of serum autotoxin in regular follow up after pediatric living donor liver transplantation for biliary atresia
Author(s)	Ueno, Takehisa; Takase, Koki; Toyama, Chiyoshi et al.
Citation	Journal of Pediatric Surgery. 2022, 57, p. 1215-1220
Version Type	AM
URL	https://hdl.handle.net/11094/87646
rights	© 2022. This manuscript version is made available under the Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International License.
Note	

The University of Osaka Institutional Knowledge Archive : OUKA

<https://ir.library.osaka-u.ac.jp/>

The University of Osaka

Table 2. Receiver operating characteristic (ROC) of fibrosis markers.

	AUC	Value	Sensitivity, %	Specificity, %
Autotaxin (mg/L)	0.86	0.897	100	74
APRI	0.63	0.90	30	100
AST to ALT ratio	0.56	1.32	70	54
Platelet count($10^4/\text{mm}^3$)	0.58	13.4	40	83
FIB-4 index	0.61	0.67	50	91
Type IV collagen 7s (mg/L)	0.82	6.6	67	94
Hyaluronic acid(ng/ml)	0.72	19	100	53

Table 2. Receiver operating characteristic of fibrosis markers. Summary of AUC, optimal cutoff point, sensitivity, and specificity for each fibrotic marker: Histological findings were classified by METAVIR score. APRI: the AST to Platelet Ratio Index; AST: aspartate aminotransferase; ALT: alanine aminotransferase; FIB-4: Fibrosis-4 index