



Title	A generalization of Magnus' theorem
Author(s)	Nakai, Yoshikazu; Baba, Kiyoshi
Citation	Osaka Journal of Mathematics. 1977, 14(2), p. 403-409
Version Type	VoR
URL	https://doi.org/10.18910/8846
rights	
Note	

The University of Osaka Institutional Knowledge Archive : OUKA

<https://ir.library.osaka-u.ac.jp/>

The University of Osaka

A GENERALIZATION OF MAGNUS' THEOREM

YOSHIKAZU NAKAI* AND KIYOSHI BABA

(Received May 4, 1976)

Let $f(x, y)$ and $g(x, y)$ be polynomials in two variables with integral coefficients. O.H. Keller raised the problem in [1]: If the functional determinant $\partial(f, g)/\partial(x, y)$ is equal to 1, then is it possible to represent x and y as polynomials of f and g with integral coefficients? This problem drew many mathematicians' attention and several attempts have been made by enlarging the coefficient domain to the complex number field \mathbb{C} . But no success has been reported yet. On the other hand A. Magnus studied the volume preserving transformation of complex planes and obtained a result which is relevant to Keller's problem ([2]). From his results it is immediately deduced that Keller's problem is answered affirmatively provided one of $f(x, y)$ and $g(x, y)$ has prime degree. For the proof Magnus used recursive formulas. But these formulas are complicated and not easy to handle. In this paper we shall give a simple proof of his theorem based on the notion of quasi-homogeneity for generalized polynomials. Moreover we shall go one step further than he did. Our results ensure that Keller's problem is valid provided one of $f(x, y)$ and $g(x, y)$ has degree 4 or larger degree is of the form $2p$ with an odd prime p . Since a complete solution of Keller's problem is not found yet our paper will be of some interest and worth-while publication.

1. Quasi-homogeneous generalized polynomials

Let x and y be two indeterminates. We shall set $\tilde{A} = \sum_{i,j \in \mathbb{Z}} C x^i y^j$ where \mathbb{C} is the complex number field and \mathbb{Z} is the ring of rational integers. \tilde{A} is a graded ring and the polynomial ring $\mathbb{C}[x, y]$ is a graded subring. Hereafter we shall call an element $f(x, y)$ of \tilde{A} a generalized polynomial or simply a g -polynomial. We shall denote by $S(f)$ the set of lattice points (i, j) in the real two space \mathbb{R}^2 such that the monomial $x^i y^j$ appears in $f(x, y)$ with a non-zero coefficient. $S(f)$ will be called the *support* of $f(x, y)$. A g -polynomial $f(x, y)$ is called a homogeneous g -polynomial or a g -form if $S(f)$ lies in the straight line of the form $X + Y = m$ where $m \in \mathbb{Z}$ and is called the degree of the g -form $f(x, y)$.

*) Supported by Takeda Science Foundation.

We shall use the symbol $S[f]$ to denote the set of monomials $x^i y^j$ such that the lattice point (i, j) is in $S(f)$.

Proposition 1. *Let $f(x, y)$ and $g(x, y)$ be non-constant g -forms of degrees m and n respectively such that the functional determinant $\partial(f, g)/\partial(x, y)$ is equal to zero. We shall define an integer d by the rule: (a) d is equal to the GCD of $|m|$ and $|n|$ if one of m and n is positive, (b) d is equal to the negative of GCD $(|m|, |n|)$ if both of m and n are negative. We shall set $m/d=m'$ and $n/d=n'$. Then we have the following:*

(i) *If one of m and n is zero, so is the other and $f(x, y)$ and $g(x, y)$ are g -polynomials in one variable (y/x) .*

(ii) *If $mn < 0$, then both of $f(x, y)$ and $g(x, y)$ are monomials and there exist a monomial $h(x, y)$ of degree d such that $f=c_1 h^{m'}$, and $g=c_2 h^{n'}$ where $c_i (i=1, 2)$ are constants.*

(iii) *If $mn > 0$, there exists a g -form $h(x, y)$ of degree d such that $f=c_1 h^{m'}$ and $g=c_2 h^{n'}$.*

Proof. Assume first $m=0$ and $n \neq 0$. It follows from $\partial(f, g)/\partial(x, y)=0$ that we have $\partial f/\partial x = \partial f/\partial y = 0$. This is against the assumption. Since a g -form of degree zero is necessarily of the form $\sum_{i \in \mathbb{Z}} a_i (y/x)^i$ we get the assertion (i). To prove (ii) we assume $m > 0$ and $n < 0$ and let $f_1 = f^{-n}$ and $g_1 = g^m$. Then $\partial(f_1, g_1)/\partial(x, y) = 0$. Since the degrees of f_1 and g_1 differ only in sign we see immediately that $f_1 \frac{\partial g_1}{\partial x} + g_1 \frac{\partial f_1}{\partial x} = 0$, or equivalently, $\partial(f_1 g_1)/\partial x = 0$. Similarly we have $\partial(f_1 g_1)/\partial y = 0$. Hence $f_1 g_1$ must be a constant. But such a case can occur only when f_1 , hence f , is a monomial because g_1 is a g -polynomial. The rest follows easily from this. The proof of (iii) will be carried out by a similar device and the detailed proof will be omitted.

DEFINITION. A g -polynomial $f(x, y)$ is called a quasi homogeneous g -polynomial (or simply a quasi g -form) if the support $S(f)$ of $f(x, y)$ is contained in the straight line. When the equation of that straight line has the form $Y + \alpha X = \lambda$. We shall say that the quasi g -form $f(x, y)$ is (α) -homogeneous of degree λ .

It should be noticed that if α is an irrational number, monomials only can be (α) -homogeneous g -forms.

Proposition 2. *Let $f(x, y)$ and $g(x, y)$ be (α) -homogeneous g -forms of positive degrees λ and μ respectively such that $\partial(f, g)/\partial(x, y) = 0$. Assume that α is a rational number q/p with coprime integers $p (> 0)$ and q . Let $d = \text{GCD}(p\lambda, p\mu)$. Then there exists an (α) -homogeneous g -form $h(x, y)$ of degree d/p such that $f=c_1 h^{m'}$ and $g=c_2 h^{n'}$ where $m'=p\lambda/d$, $n'=p\mu/d$ and $c_i (i=1, 2)$ are constants.*

Proof. Let u, v be new indeterminates and let $x=u^p$ and $y=v^q$. Then $F(u, v)=f(u^p, v^q)$ and $G(u, v)=g(u^p, v^q)$ are g -forms of degrees $p\lambda$ and $p\mu$ respectively. The rest follows easily from Proposition 1.

Let γ be an arbitrary real number. Then we can define a grading on \bar{A} in the following way. Let λ be a real number and let \bar{A}_λ be the vector space over \mathbf{C} generated by the set of g -monomials $x^i y^j$ such that $j+\gamma i=\lambda$. Then we have $\bar{A}=\bigoplus_{\lambda} \bar{A}_\lambda$ where the sum is extended over all real numbers contained in the additive subgroup of \mathbf{R} generated by 1 and γ . In case $\gamma=1$ we have the standard grading and its degree function is the ordinary function. The term "homogeneous" is reserved for this standard grading.

Proposition 3. Let $f(x, y)$ and $g(x, y)$ be g -polynomials in x and y such that $\partial(f, g)/\partial(x, y) \in \mathbf{C}$. Let α be any real number and let $f=\bigoplus f_\lambda$ and $g=\bigoplus g_\mu$ be the direct sum decomposition by the (α) -grading. Then we have

$$\sum_{\substack{\lambda+\mu=s \\ 1+\alpha\neq s}} \frac{\partial(f_\lambda, g_\mu)}{\partial(x, y)} = 0.$$

The proof is immediate and will be omitted.

2. Magnus' Theorem

For future reference we shall give Magnus' Theorem in a slightly different formulation from Magnus' original one.

Theorem 1. Let $f(x, y)$ and $g(x, y)$ be polynomials in two variables x and y with complex coefficients and let m and n be the degrees of $f(x, y)$ and $g(x, y)$ respectively. Assume that the functional determinant $\partial(f, g)/\partial(x, y)$ is a nonzero constant. If $\text{Min}(m, n) > 1$, then we have $\text{GCD}(m, n) > 1$.

Proof. Assume that $\text{GCD}(m, n)=1$. Let f_m and g_n be the degree forms of f and g respectively. From proposition 1, there is a linear form, say h , such that $f_m=\varepsilon_1 h^m$ and $g_n=\varepsilon_2 h^n$. Without loss of generalities we can assume that $h=x$ and $\varepsilon_i=1$. We shall pick up a point $P=(p_1, p_2)$ in $S(f)$ in the following way. Let L be the line defined by the equation $X=m$ and let L rotate around the point $M=(m, 0)$ counterclockwise until L meets a point in $S(f)$ other than M . Let l be the line thus obtained. The point in $S(f) \cap l$ with the smallest X -coordinate is the desired point P . Pick up a point $Q=(q_1, q_2)$ in $S(g)$ in a similar way.

Now assume we have either $(m >) p_2 > 0$ or $(n >) q_2 > 0$. Then we easily verify that one of the following situation takes place.

- (1) The lines MP and NQ are not parallel where $N=(n, 0)$.
- (2) The three points P, Q and the origin are not collinear.

If the case (a) occurs let

$$Y + aX = am, \quad Y + bx = bn$$

be the equations of the lines MP and NQ respectively. Then we have $a \neq b$. If $a > b$ let γ be a real number such that $a > \gamma > b$. If we choose γ near enough to a , then $x^{p_1}y^{p_2}$ will have the highest (γ) -degree in $S[f]$ and x^n will have the highest (γ) -degree in $S[g]$. Hence by Proposition 3, $\partial(x^{p_1}y^{p_2}, x^n)/\partial(x, y) = np_2x^{n+p_1-1}y^{p_2-1} = 0$. But this is impossible. Similarly we have a contradiction if $a < b$.

Now assume the lines MP and NQ are parallel, i.e., $a = b$ then we have the case (2), i.e., $p_2q_1 \neq q_1p_2$. Let $\gamma = a - \varepsilon$ with $\varepsilon < 0$. If we choose ε small enough, then $x^{p_1}y^{p_2}$ will have the highest (γ) -degree in $S[f]$ and $x^{q_1}y^{q_2}$ will have the highest (γ) -degree in $S[g]$. But this contradicts Proposition 3 because we have $q_1p_2 \neq q_2p_1$.

Thus we have seen that $p_2 = q_2 = 0$, i.e., $f(x, y)$ and $g(x, y)$ are polynomials in x alone. But this is impossible because $\partial(f, g)/\partial(x, y)$ is a non-zero constant, and the proof of Theorem 1 is complete.

For the sake of reference we shall call the method adopted in this proof "*the method of rotation of lines around the points M and N* ".

3. A generalization of Magnus' Theorem

Theorem 2. *Under the same notations and assumptions as Theorem 1, we have the following: If $\text{Min}(m, n) > 2$, then we have $\text{GCD}(m, n) > 2$.*

Proof. Assume that $\text{Min}(m, n) > 2$ and $\text{GCD}(m, n) = 2$ and we shall draw a contradiction. Let f_m and g_n be degree forms of f and g respectively. From Proposition 2 it follows that there exists a quadratic form $h(x, y)$ such that $f_m = ah^{m'}$ and $g_n = bh^{n'}$, where $m = 2m'$ and $n = 2n'$. There are two possibilities.

(I) h is a product of two independent linear forms. In this case we can assume without loss of generalities that $f_m = (xy)^{m'}$ and $g_n = (xy)^{n'}$. Apply the method of rotation of lines around the points $M_1 = (m', m')$ and $N_1 = (n', n')$. Then we can easily see that any point (i, j) in $S(f)$ satisfies the condition $j \leq m'$, and any point (s, t) in $S(g)$ satisfies the condition $t \leq n'$.

Now consider the (0)-grading in A. The degree forms of f and g are respectively of the forms

$$\begin{aligned} f_m^{(0)} &= y^{m'}(a_0 + a_1x + \cdots + a_{m'-1}x^{m'-1} + x^{m'}) \\ g_n^{(0)} &= y^{n'}(b_0 + b_1x + \cdots + b_{n'-1}x^{n'-1} + x^{n'}) \end{aligned}$$

From Propositions 2 and 3 there is a linear form $c + x$ such that

$$f_m^{(0)} = y^{m'}(c + x)^{m'} \quad \text{and} \quad g_n^{(0)} = y^{n'}(c + x)^{n'}$$

If we set $x_1 = c + x$ and consider f and g as polynomials in new variables x_1 and

$y_1=y$, then the support $S_1(f)$ have no point (i, j) with $j \geq m'$ except the point (m', m') . Similarly $S_1(g)$ have no point (s, t) with $t \geq n'$. Apply again the method of rotation of lines around the points M_1 and N_1 . Then we can see finally that no point (i, j) with $i < j$ is in $S(f)$ and no point (s, t) with $s < t$ is in $S(g)$. This means that $f(x, y)$ and $g(x, y)$ lack the terms $y^s (s \geq 1)$. This is impossible because of the assumption $\partial(f, g)/\partial(x, y)$ is an element of C^* .

(II) h is a power of a linear form: In this case we can assume as before that the degree forms are of the forms $f_m = x^m$ and $g_n = x^n$ respectively. Then we can see, following the method of rotations of lines around the point $M=(m, 0)$ and $N(n, 0)$, that $S(f)$ is contained in the region defined by the inequality $Y + \frac{1}{2}X \leq \frac{m}{2}$ and (g) is in the region $Y + \frac{1}{2}X \leq \frac{n}{2}$. Consider $(1/2)$ -grading and apply Propositions 2 and 3. Then we see that degree forms of f and g by this grading are

$$(ay + x^2)^{m'} \text{ and } (ay + x^2)^{n'}$$

respectively. If $a=0$ we can proceed further and we see that no point (i, j) with $j > 0$ is in $S(f)$ and no point (s, t) with $t > 0$ is in $S(g)$. This is a contradiction. Hence we must have $a \neq 0$. Then apply de Jonquiere transformation

$$Y_1 = ay + x^2, x_1 = x.$$

Since we have

$$f(x, y) = (ay + x^2)^{m'} + \sum_{j+i/2 < m'} a_{ij} x^i y^j$$

and

$$g(x, y) = (ay + x^2)^{n'} + \sum_{j+i/2 < n'} b_{ij} x^i y^j$$

We easily see that

$$f_1(x_1, y_1) = y_1^{m'} + \sum_{j+i/2 < m'} a'_{ij} x_1^i y_1^j$$

and

$$g_1(x_1, y_1) = y_1^{n'} + \sum_{j+i/2 < n'} b'_{ij} x_1^i y_1^j$$

where

$$f(x_1, y_1) = f(x_1, a^{-1}(y_1 - x_1^2)) \text{ and } g_1(x_1, y_1) = g(x_1, a^{-1}(y_1 - x_1^2)).$$

By the method of (clockwise) rotation of lines around the points $(0, m')$ and $(0, n')$ applied to the pair of polynomials $f_1(x_1, y_1)$ and $g_1(x_1, y_1)$, we see that

$S(f_1)$ is in the half plane $X+Y \leq m'$ and $S(g_1)$ is in the half plane $X+Y \leq n'$. This means that $f_1(x_1, y_1)$ is of degree m' and $g_1(x_1, y_1)$ is of degree n' . Moreover $\frac{\partial(f_1, g_1)}{\partial(x_1, y_1)} = a^{-1} \frac{\partial(f, g)}{\partial(x, y)}$ is in \mathbf{C}^* . Since $\text{Min}(m, n) > 2$, we have $\text{Min}(m', n') > 1$. Moreover $\text{GCD}(m', n') = 1$. This is the situation negated in Theorem 1.

4. Application to Keller's problem

Theorem 3. *Let $f(x, y)$ and $g(x, y)$ be polynomials of degrees m and n respectively with complex coefficients and assume that the functional determinant $\partial(f, g)/\partial(x, y)$ is a non-zero constant. Then we have $\mathbf{C}[x, y] = \mathbf{C}[f, y], g(x, y)]$ in the following three cases:*

- (1) m or n is a prime number;
- (2) m or n is 4;
- (3) $m = 2p \geq n$ where p is an odd prime.

Proof. In any case it follows from Theorems 1 and 2 that smaller degree, say n , divides larger degree m . Then from Proposition 2 and 3 the degree forms f_m and g_n are related like this, $f_m = \varepsilon g_n^{m/n}$. Then

$$f_1 = f - (\varepsilon^{n/m} g)^{m/n}$$

has lower degree than f and $\partial(f_1, g)/\partial(x, y) = \partial(f, g)/\partial(x, y)$ is a non-zero constant. Thus we can use induction on the sum $m+n$ of degrees to get the conclusion. q.e.d.

Keller's Original problem is also settled in these three cases cited in Theorem 3 because of the following

Proposition 4. *Let $f(x, y)$ and $g(x, y)$ be the polynomials in x and y with integer coefficients such that the functional determinant is equal to 1 and $\mathbf{C}[f, g] = \mathbf{C}[x, y]$. Then we have necessarily $\mathbf{Z}[f, g] = \mathbf{Z}[x, y]$.*

Proof. It suffices to prove that x and y are in $\mathbf{Z}[f, g]$. By assumption we have

$$x = \sum c_{ij} f^i g^j, c_{ij} \in \mathbf{C}.$$

If we set

$$f(x, y) = f_{10}x + f_{01}y + \cdots$$

$$g(x, y) = g_{10}x + g_{01}y + \cdots$$

then the assumption implies that $f_{10}g_{01} - f_{01}g_{10} = 1$. Apply the unimodular transformation of variables

$$x' = f_{10}x + f_{01}y$$

$$y' = g_{10}x + g_{01}y.$$

Then $\mathbf{Z}[x, y] = \mathbf{Z}[x', y']$ and $f = x' + (\text{higher degree terms})$ and $g = y' + (\text{higher degree terms})$. Hence to prove the assertion we can assume without loss of generalities that linear parts of f and g are x and y respectively. We shall define a linear order in the set (i, j) of lattice points in \mathbf{R}^2 by the way: $(i, j) > (i', j')$ if (i) $i + j > i' + j'$ or (ii) $i + j = i' + j'$ and $i > i'$. We shall show that every c_{ij} is in \mathbf{Z} by induction on the linear order just defined. Assume every $c_{i'j'}$ with $(i', j') < (i, j)$ is integer. Then the coefficients of the polynomial

$$c_{ij}f^i g^j + c_{i+1j-1}f^{i+1}g^{j-1} + \dots + c_{0i+j+1}g^{i+j+1} + \dots$$

are integers. In this polynomial $x^i y^j$ appears once with the coefficient c_{ij} . Hence c_{ij} must be an integer. Similarly y is in $\mathbf{Z}[f, g]$ and the assertion is proved completely.

OSAKA UNIVERSITY

References

- [1] O.H. Keller: *Ganze Cremona-Transformationen*, Monatsh. Math. und Phys. **47** (1939), 299–306.
- [2] A. Magnus: *On polynomial solutions of a differential equation*, Math. Scand. **3** (1955), 255–260.

