

Title	NEF CONES OF PROJECTIVE BUNDLES OVER SURFACES AND SESHADRI CONSTANTS
Author(s)	Misra, Snehajit; Ray, Nabanita
Citation	Osaka Journal of Mathematics. 2022, 59(3), p. 639–651
Version Type	VoR
URL	https://doi.org/10.18910/88490
rights	
Note	

The University of Osaka Institutional Knowledge Archive : OUKA

https://ir.library.osaka-u.ac.jp/

The University of Osaka

NEF CONES OF PROJECTIVE BUNDLES OVER SURFACES AND SESHADRI CONSTANTS

SNEHAJIT MISRA and NABANITA RAY

(Received February 22, 2021, revised May 26, 2021)

Abstract

In this article, we give a description of the closed cone of curves of the projective bundle $\mathbb{P}(E)$ over a smooth projective variety X. Using duality, we then calculate the nef cone of divisors in $\mathbb{P}(E)$ over some special surfaces X and for some special bundles on X. As an application, we also calculate the Seshadri constants of semistable ample vector bundles with vanishing discriminant on some special ruled surfaces at special points.

1. Introduction

A fundamental invariant of a projective variety X is its cone of nef divisors, denoted by Nef(X). It encodes all the information about embeddings of X in projective spaces. Also, knowledge of these cones can be used to study positivity questions, interpolations problems, Seshadri constants etc. Nef cones of a wide class of varieties have been calculated over the last few years (see [12], [5], [14], [10]).

In his paper [13], Yoichi Miyaoka studied the nef cone of a projective bundle $\mathbb{P}(E)$ over a smooth complex projective curve *C*, where *E* is any semistable vector bundle of rank *r* on *C*. More generally, it is showed in [5] that Nef($\mathbb{P}(E)$) is determined by the numerical data $\mu_{\min}(E)$ in the Harder-Narasimhan filtration of *E*. In these cases, the Picard number of $\mathbb{P}(E)$ is 2, and hence the nef cones are generated by two extremal rays in a two dimensional space. In general, when the Picard number is at least 3, the nef cones might not be a finite polyhedron, and hence are not so easy to calculate. For example if *E* is a rank 2 bundle obtained by the Serre construction from the ideal sheaf of 10 very general points on \mathbb{P}^2 , then the positivity of *E* is related to the Nagata conjecture. Thus one has to settle for special bundles *E* on special varieties *X*, even when dimension of *X* is 2. Motivated by this, we consider a projective bundle $\mathbb{P}(E) \longrightarrow X$. In section 3, we prove the following:

Theorem 1.1. Let E be a vector bundle of rank $r \ge 2$ on a smooth complex projective variety X. For an irreducible curve C in X together with its normalization $\eta_c : \tilde{C} \longrightarrow C$, consider the following fibre product diagram:



²⁰²⁰ Mathematics Subject Classification. Primary 14C20, 14E30, 14E25, 14J60; Secondary 14J26.

where *i* and *j* are inclusions. We define $\psi_c := j \circ \tilde{\eta_c}$. Then the closed cone of curves in $\mathbb{P}(E)$ is given by

$$\overline{\operatorname{NE}}(\mathbb{P}(E)) = \sum_{C \in \Gamma} (\psi_c)_* \Big(\overline{\operatorname{NE}}(\mathbb{P}(\eta_c^*(E|_C))) \Big),$$

where Γ is the set of all irreducible curves in *X*.

Consequently, we calculate the nef cones $Nef(\mathbb{P}(E))$ of some special projective bundles $\mathbb{P}(E)$ over some special smooth surfaces X (see Corollary 3.4, Corollary 3.5) by applying duality to Theorem 1.1. An important finding in section 3 is the following:

Corollary 1.2. Let X be a smooth complex projective surface with $\overline{\text{NE}}(X) = \{a_1[C_1] + a_2[C_2] + \cdots + a_n[C_n] \mid a_i \in \mathbb{R}_{\geq 0}\}$ for some irreducible curves C_1, C_2, \cdots, C_n in X. If E is a semistable vector bundle of rank $r \geq 2$ on X with vanishing discriminant, i.e. $\Delta(E) := 2rc_2(E) - (r-1)c_1^2(E) = 0$, then

 $\operatorname{Nef}(\mathbb{P}(E)) = \left\{ y_0 \xi + \pi^* \gamma \mid \gamma \in N^1(X)_{\mathbb{R}}, y_0 \ge 0, \ y_0 \mu(E|_{C_j}) + (\gamma \cdot C_j) \ge 0 \text{ for all } 1 \le j \le n \right\},$ where ξ denotes the numerical equivalence classes of the tautological bundle $\mathcal{O}_{\mathbb{P}(E)}(1)$.

We also give several examples to illustrate our results. In most of these examples, the projective bundles $\mathbb{P}(E)$ have Picard number at least 3, and the nef cones Nef($\mathbb{P}(E)$) are all finite polyhedra.

A vector bundle *V* on *X* is called ample if $\mathcal{O}_{\mathbb{P}(V)}(1)$ is ample on $\mathbb{P}(V)$. The Seshadri constant $\varepsilon(V, x)$ of an ample vector bundle *V* on a projective variety *X* at a closed point $x \in X$ is introduced in [7]. Let us consider the following pullback diagram under the blow up map $\rho_x : \tilde{X}_x = \text{Bl}_x(X) \longrightarrow X$

Let $\tilde{\xi}_x$ be the numerical equivalence class of the tautological bundle $\mathcal{O}_{\mathbb{P}(\rho_x^*V)}(1)$, and $\tilde{E}_x := \tilde{\rho}_x^{-1}(F_x)$, where F_x is the class of the fibre of the map π' over the point x. Then the Seshadri constant of V at $x \in X$ is defined as

$$\varepsilon(V, x) := \sup \{ \lambda \in \mathbb{R}_{>0} \mid \tilde{\xi}_x - \lambda \tilde{E}_x \text{ is nef} \}.$$

Note that the Seshadri constant $\varepsilon(V, x)$ is not equal to the Seshadri constant of the tautological line bundle $\mathcal{O}_{\mathbb{P}(V)}(1)$ on $\mathbb{P}(V)$. These interesting constants have been extensively studied by many authors especially for line bundles on algebraic surfaces (see [12], Chapter 5 and [1]). In [7], it is shown that if *V* is an ample vector bundle on a smooth projective curve *C*, then the Seshadri constant $\varepsilon(V, x) = \mu_{\min}(V)$ at $x \in C$. Recently, Seshadri constants for equivariant bundles have been calculated on toric varieties in [9] and on certain homogeneous varieties in [3].

As an application of our results in section 3, we calculate the Seshadri constant of semistable ample vector bundles with vanishing discriminant on some special ruled surfaces at special points (see Theorem 4.2).

2. Preliminaries

Throughtout this article, all the varieties are assumed to be over the field of complex numbers \mathbb{C} . In this section, we recall the definition and duality property of nef cone as well as the definition of semistability of vector bundles on smooth projective varieties. Let *X* be a smooth complex projective variety of dimension *n* with a fixed ample line bundle *H* on it.

2.1. Harder-Narasimhan Filtration: A non-zero torsion-free coherent sheaf G on X is said to be *H*-semistable if

$$\mu_H(\mathcal{F}) = \frac{c_1(\mathcal{F}) \cdot H^{n-1}}{\operatorname{rank}(\mathcal{F})} \le \mu_H(\mathcal{G}) = \frac{c_1(\mathcal{G}) \cdot H^{n-1}}{\operatorname{rank}(\mathcal{G})}$$

for all subsheaves \mathcal{F} of \mathcal{G} . For every vector bundle E on X, there is a unique filtration

$$0 = E_0 \subsetneq E_1 \subsetneq E_2 \subsetneq \cdots \subsetneq E_{k-1} \subsetneq E_k = E$$

of subbundles of *E*, called the Harder-Narasimhan filtration of *E*, such that E_i/E_{i-1} is *H*-semistable torsion free sheaf for each $i \in \{1, 2, \dots, k\}$ and $\mu_H(E_i/E_{i-1}) > \mu_H(E_{i+1}/E_i)$ for each $i \in \{1, 2, \dots, k-1\}$. We define $Q_k := E_k/E_{k-1}$ and $\mu_{\min}(E) := \mu_H(Q_k) = \mu_H(E_k/E_{k-1})$.

2.2. Nef cone. Let $\text{Div}^0(X) := \{D \in \text{Div}(X) \mid D \cdot C = 0 \text{ for all curves } C \text{ in } X\}$ be the subgroup of Div(X) consisting of numerically trivial divisors. The quotient $\text{Div}(X)/\text{Div}^0(X)$ is called the Néron Severi group of X, and is denoted by $N^1(X)_{\mathbb{Z}}$. The Néron Severi group $N^1(X)_{\mathbb{Z}}$ is a free abelian group of finite rank. Its rank, denoted by $\rho(X)$ is called the Picard number of X. In particular, $N^1(X)_{\mathbb{R}} := N^1(X)_{\mathbb{Z}} \otimes \mathbb{R} = (\text{Div}(X)/\text{Div}^0(X)) \otimes \mathbb{R}$ is called the real Néron Severi group.

A Cartier divisor D on X (with \mathbb{Z}, \mathbb{Q} or \mathbb{R} coefficients) is called nef if $D \cdot C \ge 0$ for all irreducible curves $C \subseteq X$. The intersection product being independent of numerical equivalence class, one can talk about nef classes in $N^1(X)_{\mathbb{R}}$. The convex cone of all nef classes in $N^1(X)_{\mathbb{R}}$ is called the nef cone of X, and is denoted by Nef(X).

A formal sum $\gamma = \sum_{i} a_i C_i$, where $a_i \in \mathbb{R}$ and $C_i \subseteq X$ are irreducible curves in X, is called a real 1-cycle in X. The \mathbb{R} -vector space of all real 1-cycle in X is denoted by $Z_1(X)_{\mathbb{R}}$. Two elements $\gamma_1, \gamma_2 \in Z_1(X)_{\mathbb{R}}$ are called numerically equivalent, denoted by $\gamma_1 \equiv \gamma_2$ if $D \cdot \gamma_1 = D \cdot \gamma_2$ for every divisor D in X. We denote $N_1(X)_{\mathbb{R}} = Z_1(X)_{\mathbb{R}} / \equiv$. The cone of curves in X, denoted by NE(X) is the cone spanned by the effective 1-cycle classes in $N_1(X)_{\mathbb{R}}$. If Cis an effective curve, then we denote the numerical equivalence class of curves by [C].

The intersection pairing $N^1(X)_{\mathbb{R}} \times N_1(X)_{\mathbb{R}} \longrightarrow \mathbb{R}$ induces a perfect pairing. The closed cone of curves $\overline{NE}(X)$, which is defined to be the closure of NE(X), is known to be the dual to the nef cone Nef(X), i.e.

$$NE(X) = \{ \gamma \in N_1(X)_{\mathbb{R}} \mid \alpha \cdot \gamma \ge 0 \text{ for all } \alpha \in Nef(X) \}.$$

3. Main Results

We quickly recall our set up. Let *E* be a vector bundle of rank ≥ 2 on a smooth complex projective variety *X*, and $\pi : \mathbb{P}(E) \longrightarrow X$ be the projection. In this section, we give a description of the closed cone $\overline{NE}(\mathbb{P}(E))$ of curves in $\mathbb{P}(E)$ in terms of the closed cone of curves

 $\overline{NE}(\mathbb{P}(E|_C))$ for every irreducible curve *C* in *X*. Proof of Theorem 1.1. For an irreducible curve *C* in *X* together with its normalization $\eta_c : \tilde{C} \longrightarrow C$, we recall the following fibre product diagram:

where *i* and *j* are inclusions and $\psi_c := j \circ \tilde{\eta_c}$. Let $\xi \in N^1(\mathbb{P}(E))_{\mathbb{R}}$ be the numerical equivalence class of the tautological line bundle $\mathcal{O}_{\mathbb{P}(E)}(1)$ on $\mathbb{P}(E)$. Then

$$N^1(\mathbb{P}(E))_{\mathbb{R}} = \mathbb{R}\xi \oplus \pi^* N^1(X)_{\mathbb{R}}$$

We fix the notations $\xi_{\tilde{c}}$ and $f_{\tilde{c}}$ for the numerical equivalence classes of the tautological line bundle $\mathcal{O}_{\mathbb{P}(\eta_c^*(E|_C))}(1)$ of $\mathbb{P}(\eta_c^*(E|_C))$ and a fibre of the map $\tilde{\pi_c}$ respectively. Then $N^1(\mathbb{P}(\eta_c^*(E|_C))_{\mathbb{R}})$ is generated by $\xi_{\tilde{c}}$ and $f_{\tilde{c}}$. The map ψ_c induces the map

 $\psi_c^*: N^1(\mathbb{P}(E))_{\mathbb{R}} \longrightarrow N^1(\mathbb{P}(\eta_c^*(E|_C))_{\mathbb{R}})$

between the Néron Severi groups such that

$$\psi_c^*(\xi) = \xi_{\tilde{c}} \text{ and } \psi_c^*(\pi^*L) = (L \cdot C)f_{\tilde{c}}, \text{ for any } L \in N^1(X)_{\mathbb{R}}$$

As a consequence, we get the pushforward map

$$(\psi_c)_* : N_1(\mathbb{P}(\eta_c^*(E|_C))_{\mathbb{R}} \longrightarrow N_1(\mathbb{P}(E))_{\mathbb{R}})$$

Hence

$$\overline{\sum_{C\in\Gamma} (\psi_c)_* \left(\overline{\operatorname{NE}}(\mathbb{P}(\eta_c^*(E|_C))) \right)} \subseteq \overline{\operatorname{NE}}(\mathbb{P}(E)),$$

where the sum is taken over the set Γ of all irreducible curves in X. Now to prove the reverse inequality, we consider the numerical equivalence class $[\bar{C}] \in \text{NE}(\mathbb{P}(E))$ of an irreducible curve \bar{C} in $\mathbb{P}(E)$ which is not contained in any fibre of π . Denote $\pi(\bar{C}) = C$. Then, $\bar{C} \subseteq$ $\mathbb{P}(E|_C)$. Then there exists a unique irreducible curve $C' \subseteq \mathbb{P}(\eta_c^*(E|_C))$ such that $\tilde{\eta}_c(C') = \bar{C}$ and $(\psi_c)_*([C']) = [\bar{C}]$. Also, the numerical equivalence classes of curves in a fibre of π_c maps to the numerical classes of curves in a fibre of π by $(\psi_c)_*$. Hence, we have

$$\overline{\operatorname{NE}}(\mathbb{P}(E)) \subseteq \overline{\sum_{C \in \Gamma} (\psi_c)_* (\overline{\operatorname{NE}}(\mathbb{P}(\eta_c^*(E|_C)))))}.$$

This completes the proof.

Corollary 3.1. Let X be a smooth complex projective surface with $\overline{\text{NE}}(X) = \{a_1[C_1] + a_2[C_2] + \cdots + a_n[C_n] \mid a_i \in \mathbb{R}_{\geq 0}\}$ for some irreducible curves C_1, C_2, \cdots, C_n in X. If E is a semistable vector bundle of rank $r \geq 2$ on X with vanishing discriminant, i.e. $\Delta(E) := 2rc_2(E) - (r-1)c_1^2(E) = 0$, then

 $\operatorname{Nef}(\mathbb{P}(E)) = \{ y_0\xi + \pi^*\gamma \mid \gamma \in N^1(X)_{\mathbb{R}}, y_0 \ge 0, y_0\mu(E|_{C_j}) + (\gamma \cdot C_j) \ge 0 \text{ for all } 1 \le j \le n \},$ where ξ denotes the numerical equivalence class of the tautological bundle $\mathcal{O}_{\mathbb{P}(E)}(1)$.

Proof. Let *C* be an irreducible curve in *X* such that $[C] = \sum_{i=1}^{n} x_i[C_i] \in N_1(X)_{\mathbb{R}}$ for some $x_i \in \mathbb{R}_{\geq 0}$. As *E* is semistable with vanishing discriminant, applying Theorem 1.2 in [2] to the map $i \circ \eta_c : \tilde{C} \longrightarrow C \hookrightarrow X$, we get that $\eta_c^*(E|_C)$ is also semistable bundle on \tilde{C} for any irreducible curve *C* in *X*. Using Theorem 3.1 in [13], we have

$$\operatorname{Nef}(\mathbb{P}(\eta_c^*(E|_C))) = \left\{ a(\xi_{\tilde{c}} - \mu(\eta_c^*(E|_C))f_{\tilde{c}}) + bf_{\tilde{c}} \mid a, b \in \mathbb{R}_{\geq 0} \right\}.$$

We define $l_c := \deg(\eta_c^*(E|_C)) - \mu(\eta_c^*(E|_C)).$

Applying duality, we then get

$$\overline{\operatorname{NE}}(\mathbb{P}(\eta_c^*(E|_C))) = \left\{ a(\xi_{\tilde{c}}^{r-2}f_{\tilde{c}}) + b(\xi_{\tilde{c}}^{r-1} - l_c\xi_{\tilde{c}}^{r-2}f_{\tilde{c}}) \mid a, b \in \mathbb{R}_{\geq 0} \right\}.$$

Therefore, by Theorem 1.1 we have

$$\overline{\operatorname{NE}}(\mathbb{P}(E)) = \sum_{C \in \Gamma} (\psi_c)_* \left(\overline{\operatorname{NE}}(\mathbb{P}(\eta_c^*(E|_C))) \right) = \sum_{C \in \Gamma} \left[\mathbb{R}_{\geq 0}(\xi^{r-2}F) + \mathbb{R}_{\geq 0}(\xi^{r-1}\pi^*[C] - l_c\xi^{r-2}F) \right],$$

where *E* is the numerical equivalence class of the fibre of the projection map $\pi : \mathbb{P}(E)$.

where *F* is the numerical equivalence class of the fibre of the projection map $\pi : \mathbb{P}(E) \longrightarrow X$.

Also,
$$\mu(\eta_c^*(E|_C)) = \frac{\deg(E|_C)}{r} = \frac{\sum_{i=1}^{n} x_i \deg(E|_{C_i})}{r} = \sum_{i=1}^{n} x_i \mu(\eta_{c_i}^*(E|_{C_i})),$$

 $l_{c_i} = \deg(\eta_{c_i}^*(E|_C)) - \mu(\eta_{c_i}^*(E|_{C_i})) = \frac{r-1}{r} \deg(E|_{C_i}) \text{ for every } 1 \le i \le n, \text{ and } l_c = \sum_{i=1}^{n} x_i l_{c_i}.$

This shows that $\overline{NE}(\mathbb{P}(E))$ is generated by

$$\xi^{r-2}F$$
, $\xi^{r-1}\pi^*[C_i] - \frac{r-1}{r}\deg(E|_{C_i})\xi^{r-2}F \mid 1 \le i \le n$,

and the nef cone

$$Nef(\mathbb{P}(E)) = \{ y_0 \xi + \pi^* \gamma \mid \gamma \in N^1(X)_{\mathbb{R}}, y_0 \ge 0, y_0 \mu(E|_{C_j}) + (\gamma \cdot C_j) \ge 0 \text{ for all } 1 \le j \le n \}.$$

REMARK 1. Let X be a smooth irregular complex projective surface X, i.e., $q = H^1(X, \mathcal{O}_X) \neq 0$, and the closed cone of curves $\overline{NE}(X)$ is a finite polyhedron generated by classes of irreducible curves. Examples of such surfaces X include ruled surface over a smooth curve of genus greater than 0 with closed cone of curves generated by irreducible curve classes, very general abelian surfaces with Picard number 1, del Pezzo surfaces etc. Then there exists non-split extension of the form

$$0 \longrightarrow \mathcal{O}_C \longrightarrow E \longrightarrow \mathcal{O}_C \longrightarrow 0.$$

In this case, *E* is a semistable bundle of rank 2 with vanishing discriminant. Moreover, for any positive integer *r*, the vector bundles of the forms $E^{\oplus r} \oplus \mathcal{O}_C$ and $E^{\oplus r}$ are examples of semistable bundles of ranks 2r + 1 and 2r respectively with vanishing discriminant. In this way, one can produce examples of semistable bundles with vanishing discriminant of any rank on such *X*.

Also, for a semistable bundle V on X with discriminant 0, and a nonsplit extension of the form $0 \longrightarrow V \longrightarrow E \longrightarrow V \longrightarrow 0$, E is also semistable with vanishing discriminant. In all these cases, one can calculate the nef cones using Corollary 3.1.

EXAMPLE 3.2. Let $\rho : X = \mathbb{P}(W) \longrightarrow C$ be a ruled surface on a smooth complex projective curve *C* defined by a normalized bundle (in the sense of Hartshorne [see [8], Ch 5]) *W* with $\mu_{\min}(W) = \deg(W)$. Then $\overline{NE}(X) = \{a\zeta + bf \mid a, b \in \mathbb{R}_{\geq 0}\}$ is a finite polyhedron generated by classes of two irreducible curves, where $\zeta = [\mathcal{O}_X(\sigma)]$ (here σ is the normalized section such that $\mathcal{O}_{\mathbb{P}(W)}(1) \cong \mathcal{O}_X(\sigma)$, and *f* is the numerical equivalence class of a fibre of ρ). Let *V* be a semistable (resp. stable) vector bundle of rank *r* and degree *d* on *C*. Then the pullback bundle $E := \rho^*(V)$ is also semistable (resp. stable) on *X* with discriminant

$$\Delta(E) = \Delta(\rho^* V) = 2rc_2(\rho^* V) - (r-1)c_1^2(\rho^* V) = 0 - (r-1)df \cdot df = 0.$$

Therefore, one can calculate the nef cone Nef($\mathbb{P}(E)$) in this case using Corollary 3.1. Note that in this case $\mathbb{P}(E) \cong \mathbb{P}(W) \times_C \mathbb{P}(V)$, and nef cone of the fiber product Nef($\mathbb{P}(W) \times_C \mathbb{P}(V)$) is already known due to [11]. Our result in this case matches with the result in [11].

For example, any indecomposable normalized rank 2 bundle *W* over an elliptic curve *C* is one of the following type (see Proposition 2.15, Chapter V, [8]):

- Either (i) $0 \longrightarrow \mathcal{O}_C \longrightarrow W \longrightarrow \mathcal{O}_C \longrightarrow 0$,
- or (ii) $0 \longrightarrow \mathcal{O}_C \longrightarrow W \longrightarrow \mathcal{O}_C(p) \longrightarrow 0$ for some closed point $p \in C$.

In both the cases, $deg(W) \ge 0$, and hence W is semistable.

We consider the ruled surface $\rho : X = \mathbb{P}(W) \longrightarrow C$ over the elliptic curve *C* defined by the nonsplit extension $0 \longrightarrow \mathcal{O}_C \longrightarrow W \longrightarrow \mathcal{O}_C \longrightarrow 0$. Let us define $[C_1] := [\mathcal{O}_X(\sigma)] = [\mathcal{O}_{\mathbb{P}(W)}(1)]$, and $[C_2] := f$ = numerical class of a fibre of the map ρ . Then $\overline{\text{NE}}(X) = \{a[C_1] + b[C_2] \mid a, b \in \mathbb{R}_{\geq 0}\}$.

Let V be the bundle on C which sits in the nonsplit extension $0 \longrightarrow \mathcal{O}_C \longrightarrow V \longrightarrow \mathcal{O}_C(p) \longrightarrow 0$ for some closed point p in C. Then, $E := \rho^*(V)$ is a stable bundle on X with $\Delta(E) = 0$. Note that, in this example, the Picard number of $\mathbb{P}(E)$ is 3.

Now,

$$deg(E|_{C_1}) = c_1(\rho^*V) \cdot \zeta = deg(V)\zeta \cdot f = 1; \quad deg(E|_{C_2}) = c_1(\rho^*V) \cdot f = deg(V)f \cdot f = 0.$$

$$b_{11} := C_1 \cdot C_1 = \zeta^2 = deg(W) = 0; \quad b_{12} = b_{21} := C_1 \cdot C_2 = \zeta \cdot f = 1;$$

$$b_{22} := C_2 \cdot C_2 = f^2 = 0.$$

$$y_0\mu(E|_{C_1}) + y_1b_{11} + y_2b_{21} = \frac{y_0}{2} + y_2; \quad y_0\mu(E|_{C_2}) + y_1b_{12} + y_2b_{22} = y_1.$$

Therefore, for the projective bundle $\pi : \mathbb{P}(E) \longrightarrow X$, the nef cone is

Nef($\mathbb{P}(E)$) = { $y_0\xi + y_1(\pi^*\zeta) + y_2(\pi^*f) \mid y_0 \ge 0, y_1 \ge 0, \frac{y_0}{2} + y_2 \ge 0$ }.

EXAMPLE 3.3. Let $\rho : X = \mathbb{P}(W) \longrightarrow C$ be a ruled surface over a smooth elliptic curve *C* defined by the rank two bundle $W = \mathcal{O}_C \oplus \mathcal{O}_C$. Then $\overline{\text{NE}}(X) = \{a\zeta + bf \mid a, b \in \mathbb{R}_{\geq 0}\}$, where $\zeta = [\mathcal{O}_{\mathbb{P}(W)}(1)] \in N^1(X)$ and *f* is the numerical equivalence class of a fibre of ρ .

Let $\rho_x : \tilde{X}_x = Bl_x(X) \longrightarrow X$ be the blow up of X at a closed point x in a section σ such that $\mathcal{O}_X(\sigma) \cong \mathcal{O}_{\mathbb{P}(W)}(1)$, and E_x be the exceptional divisor. Then we claim that $\overline{NE}(\tilde{X}_x) = \{a[C_1]+b[C_2]+c[C_3] \mid a, b, c \in \mathbb{R}_{\geq 0}\}$, where $[C_1] = \rho_x^*f - E_x$, $[C_2] = \rho_x^*\zeta - E_x$ and $[C_3] = E_x$.

To prove our claim enough to show that if *V* is an irreducible curve in *X* which is not a fibre i.e. $[V] = a\zeta + bf$; a > 0 and $b \ge 0$, then $\text{mult}_x V = r \le a$.

Let *F* be a fibre passing through *x*. Then $mult_xF = 1$. Hence

$$a = V \cdot F = \sum_{P \in V \cap F} (V.F)_P \ge (V.F)_x \ge r$$
 (see [8], Chapter V, Proposition 1.4)

Now we also have

 $\begin{aligned} b_{11} &:= C_1 \cdot C_1 = (\rho_x^* f - E_x) \cdot (\rho_x^* f - E_x) = E_x^2 = -1; \\ b_{22} &:= C_2 \cdot C_2 = (\rho_x^* \zeta - E_x) \cdot (\rho_x^* \zeta - E_x) = \deg(W) - 1 = -1; \\ b_{33} &:= C_3 \cdot C_3 = E_x^2 = -1. \\ b_{12} &= b_{21} &:= C_1 \cdot C_2 = (\rho_x^* f - E_x) \cdot (\rho_x^* \zeta - E_x) = \zeta \cdot f + E_x^2 = 0; \\ b_{13} &= b_{31} &:= C_1 \cdot C_3 = (\rho_x^* f - E_x) \cdot E_x = 1; \\ b_{23} &= b_{32} &:= C_2 \cdot C_3 = (\rho_x^* \zeta - E_x) \cdot E_x = 1; \\ \text{Let } E &= \rho_x^* (\rho^*(V)), \text{ where } V \text{ is the semistable bundle on } C \text{ given by the nonsplit extension} \\ 0 &\longrightarrow \mathcal{O}_C \longrightarrow V \longrightarrow \mathcal{O}_C(p) \longrightarrow 0 \text{ for some closed point } p \in C. \text{ Then } E \text{ is also semistable bundle with } \triangle(E) = 0. \text{ Note that, in this example, the Picard number of } \mathbb{P}(E) \text{ is } 4. \end{aligned}$

 $\begin{aligned} &\deg(E|_{C_1}) = c_1(\rho_x^*(\rho^*(V))) \cdot (\rho_x^*f - E_x) = 0 ; \deg(E|_{C_2}) = c_1(\rho_x^*(\rho^*(V))) \cdot (\rho_x^*\zeta - E_x) = 1; \\ &\deg(E|_{C_3}) = c_1(\rho_x^*(\rho^*(V))) \cdot E_x = 0. \\ &y_0\mu(E|_{C_1}) + y_1b_{11} + y_2b_{21} + y_3b_{31} = y_3 - y_1 ; \\ &y_0\mu(E|_{C_2}) + y_1b_{12} + y_2b_{22} + y_3b_{32} = \frac{y_0}{2} + y_3 - y_2 ; \\ &y_0\mu(E|_{C_3}) + y_1b_{13} + y_2b_{23} + y_3b_{33} = y_1 + y_2 - y_3. \end{aligned}$

Therefore, the nef cone of the projective bundle $\pi : \mathbb{P}(E) \longrightarrow \tilde{X}_x$ is

$$\operatorname{Nef}(\mathbb{P}(E)) = \left\{ y_0 \xi + \sum_{i=1}^3 y_i \pi^*(C_i) \mid y_0 \ge 0, y_3 - y_1 \ge 0, \frac{y_0}{2} + y_3 - y_2 \ge 0, y_1 + y_2 - y_3 \ge 0 \right\}.$$

Corollary 3.4. Let X be a smooth complex projective surface such that $\overline{NE}(X) = \{a_1[C_1] + a_2[C_2] + \dots + a_n[C_n] \mid a_1, \dots, a_n \in \mathbb{R}_{\geq 0}\}$ for some irreducible curves C_1, C_2, \dots, C_n of X. Let $E = L_1 \oplus L_2 \oplus \dots \oplus L_r$ be a completely decomposable vector bundle of rank r on X. Let ξ be the numerical class of $\mathcal{O}_{\mathbb{P}(E)}(1)$. Then the nef cone

$$\operatorname{Nef}(\mathbb{P}(E)) = \left\{ y_0 \xi + \pi^* \gamma \mid \gamma \in N^1(X)_{\mathbb{R}}, y_0 \ge 0, y_0 \min\{C_i \cdot L_j \mid 1 \le j \le r\} + (\gamma \cdot C_i) \ge 0 \\ for all \ 1 \le i \le n \right\}.$$

Proof. Let *C* be an irreducible curve in *X* numerically equivalent to $\sum_{i=1}^{n} x_i C_i$ for $x_i \in \mathbb{R}_{\geq 0}$ for all *i* and let $\eta_c : \tilde{C} \longrightarrow C$ be its normalization. Then, by Lemma 2.1 in [5]

 $\operatorname{Nef}(\mathbb{P}(\eta_c^*(E|_C))) = \left\{ a(\xi_{\tilde{c}} - \mu_{\min}(\eta_c^*(E|_C))f_{\tilde{c}}) + bf_{\tilde{c}} \mid a, b \in \mathbb{R}_{\geq 0} \right\},$ $\operatorname{Nef}(\mathbb{P}(\eta_{c_i}^*(E|_{C_i}))) = \left\{ a(\xi_{\tilde{c}_i} - \mu_{\min}(\eta_{c_i}^*(E|_{C_i}))f_{\tilde{c}_i}) + bf_{\tilde{c}_i} \mid a, b \in \mathbb{R}_{\geq 0} \right\} \text{ for each } i \in \{1, 2, \cdots, n\}.$ For any curve $C \subset X$, we define

 $l_c = \deg(\eta_c^*(E|_C)) - \min\{C \cdot L_j \mid 1 \le j \le r\} = \deg(\eta_c^*(E|_C)) - \mu_{\min}(\eta_c^*(E|_C))$ Applying duality, we also have

 $\overline{\mathrm{NE}}(\mathbb{P}(\eta_{c}^{*}(E|_{C}))) = \left\{ a(\xi_{\tilde{c}}^{r-2}f_{\tilde{c}}) + b(\xi_{\tilde{c}}^{r-1} - l_{c}\xi_{\tilde{c}}^{r-2}f_{\tilde{c}}) \mid a, b \in \mathbb{R}_{\geq 0} \right\},\$ $\overline{\mathrm{NE}}(\mathbb{P}(\eta_{c_{i}}^{*}(E|_{C_{i}}))) = \left\{ a(\xi_{\tilde{c}_{i}}^{r-2}f_{\tilde{c}_{i}}) + b(\xi_{\tilde{c}_{i}}^{r-1} - l_{c_{i}}\xi_{\tilde{c}_{i}}^{r-2}f_{\tilde{c}_{i}}) \mid a, b \in \mathbb{R}_{\geq 0} \right\} \text{ for each } i \in \{1, 2, \cdots, n\}.$

Observe that

 $\sum_{i=1}^{n} x_i \min\{C_i \cdot L_j \mid 1 \le j \le r\} \le \min\{\sum_{i=1}^{n} x_i(C_i \cdot L_j) \mid 1 \le j \le r\} = \mu_{\min}(\eta_c^*(E|_C)),$ and deg $(\eta_c^*(E|_C)) = \sum_{i=1}^{n} x_i \deg(\eta_{c_i}^*(E|_{C_i})).$ Therefore

$$l_c \le \sum_{i=1}^n x_i l_{c_i}.$$

Recall from Theorem 1.1 that

$$\overline{\mathrm{NE}}(\mathbb{P}(E)) = \sum_{C \in \Gamma} \left[\mathbb{R}_{\geq 0}(\xi^{r-2}F) + \mathbb{R}_{\geq 0}\{\xi^{r-1}\pi^*[C] - (\deg(\eta^*_c(E|_C)) - \mu_{\min}(\eta^*_c(E|_C)))\xi^{r-2}F\} \right]$$
$$= \sum_{C \in \Gamma} \left[\mathbb{R}_{\geq 0}(\xi^{r-2}F) + \mathbb{R}_{\geq 0}\{\xi^{r-1}\pi^*[C] - l_c\xi^{r-2}F\} \right]$$
$$= \sum_{C \equiv \sum_i x_i C_i} \left[\mathbb{R}_{\geq 0}(\xi^{r-2}F) + \mathbb{R}_{\geq 0}\{\sum_{i=1}^n x_i(\xi^{r-1}\pi^*[C_i] - l_{c_i}\xi^{r-2}F) + (\sum_{i=1}^n x_il_{c_i} - l_c)\xi^{r-2}F\} \right]$$

This shows that $\overline{NE}(\mathbb{P}(E))$ is generated by

$$\xi^{r-2}F, (\xi^{r-1}\pi^*[C_i] - l_{c_i}\xi^{r-2}F) \mid 1 \le i \le r \}.$$

Therefore

 $\operatorname{Nef}(\mathbb{P}(E)) = \left\{ y_0 \xi + \pi^* \gamma \mid \gamma \in N^1(X)_{\mathbb{R}}, y_0 \ge 0, y_0 \mu_{\min}(\eta_{c_i}^*(E|_{C_i})) + (\gamma \cdot C_i) \ge 0 \ \forall 1 \le i \le n \right\}.$

But $\mu_{\min}(\eta_{c_i}^*(E|_{C_i})) = \min\{C_i \cdot L_j \mid 1 \le j \le r\}$ for each $i \in \{1, 2, \dots, n\}$. Hence the result follows.

A particular case of Corollary 3.4 is the following.

Corollary 3.5. Let X be a smooth complex projective surface with Picard number 1 and L_X be an ample generator of the real Néron Severi group $N^1(X)_{\mathbb{R}}$. Let $E = M_1 \oplus M_2 \oplus \cdots \oplus M_r$ be a completely decomposable vector bundle of rank $r \ge 2$ on X such that $M_i \equiv a_i L_X \in N^1(X)_{\mathbb{R}}$ for each $i \in \{1, 2, \dots, r\}$ and $a_1 \le a_2 \le a_3 \le \cdots \le a_r$. Then

 $\operatorname{Nef}(\mathbb{P}(E)) = \{ y_0 \xi + y_1 \pi^* L_X \mid y_0 \ge 0, y_0 a_1 + y_1 \ge 0 \}.$

In particular, if $X = \mathbb{P}^2$, and $E = \mathcal{O}_{\mathbb{P}^2}(a_1) \oplus \mathcal{O}_{\mathbb{P}^2}(a_2) \oplus \cdots \oplus \mathcal{O}_{\mathbb{P}^2}(a_r)$ a completely decomposable bundle on \mathbb{P}^2 with $a_1 \leq a_2 \leq a_3 \leq \cdots \leq a_r$, then

$$Nef(\mathbb{P}(E)) = \{y_0\xi + y_1\pi^*H \mid y_0 \ge 0, y_0a_1 + y_1 \ge 0\},\$$

where *H* is the numerical equivalence class of $\mathcal{O}_{\mathbb{P}^2}(1)$ in $N^1(\mathbb{P}^2)_{\mathbb{R}}$.

Proof. Let *m* be the least positive integer such that $H^0(mL_X) \neq 0$ and $C_0 \in |mL_X|$ be an irreducible curve. Then we have

$$\overline{\operatorname{NE}}(X) \subseteq \mathbb{R}_{\geq 0} L_X = \mathbb{R}_{\geq 0} \frac{1}{m} [C_0] \subseteq \operatorname{Nef}(X).$$

Hence $\overline{\text{NE}}(X) = \mathbb{R}_{\geq 0}[C_0]$. Now $\min\{C_0 \cdot L_j \mid 1 \le j \le r\} = \min\{ma_j L_X^2 \mid 1 \le j \le r\} = ma_1 L_X^2$.

Thus by the previous Corollary 3.4 we have

$$Nef(\mathbb{P}(E)) = \{y_0\xi + y_1\pi^*L_X \mid y_0 \ge 0, y_0 \min\{C_0 \cdot L_j \mid 1 \le j \le r\} + (y_1L_X \cdot C_0) \ge 0\} \\ = \{y_0\xi + y_1\pi^*L_X \mid y_0 \ge 0, y_0a_1 + y_1 \ge 0\}.$$

646

-		
Т		
-		

REMARK 2. We known that \mathbb{P}^2 is a toric variety and any line bundle is equivariant. Then the second part of Corollary 3.5, $\mathbb{P}(E)$ is toric variety where $E = \mathcal{O}_{\mathbb{P}^2}(a_1) \oplus \mathcal{O}_{\mathbb{P}^2}(a_2) \oplus \cdots \oplus \mathcal{O}_{\mathbb{P}^2}(a_r)$. For more detailed description of the fan structure of $\mathbb{P}(E)$ check [15], page 53. If Δ is the fan of $\mathbb{P}(E)$, then it is clear from the construction that the support of Δ has full dimension. The mori cone of curves of $\mathbb{P}(E)$ is (see [4], Theorem 6.3.20)

$$\overline{\mathrm{NE}}(\mathbb{P}(E)) = \sum_{\tau \text{ a wall of } \Delta} \mathbb{R}_{\geq 0}[V(\tau)].$$

Hence the dual of the mori cone gives the Nef cone of $\mathbb{P}(E)$.

EXAMPLE 3.6. Let $\phi_x : X = \tilde{\mathbb{P}}_x^2 = \operatorname{Bl}_x \mathbb{P}^2 \longrightarrow \mathbb{P}^2$ be the blow-up of \mathbb{P}^2 at a closed point $x \in \mathbb{P}^2$ with exceptional divisor $\phi_x^{-1}(x) = E_x$. Then, $\overline{\operatorname{NE}}(X) = \{a[C_1] + b[C_2] \mid a, b \in \mathbb{R}_{\geq 0}\}$, where $[C_1] = [\phi_x^*(\mathcal{O}_{\mathbb{P}^2}(1)) - E_x]$, and $[C_2] = [E_x]$. Then, $b_{11} = C_1 \cdot C_1 = 0$, $b_{12} = b_{21} = C_1 \cdot C_2 = 1$, $b_{22} = C_2 \cdot C_2 = -1$.

Let us consider the rank 2 bundle $E = \phi_x^*(\mathcal{O}_{\mathbb{P}^2}(1)) \oplus \mathcal{O}_X(E_x)$ on X. Then, $\mathbb{P}(E)$ has Picard number 3. Fix the notations $L_1 = \phi_x^*(\mathcal{O}_{\mathbb{P}^2}(1))$, $L_2 = \mathcal{O}_X(E_x)$, and $a_{ij} = C_i \cdot L_j$ for $1 \le i \le 2$, $1 \le j \le 2$. Then, $a_{11} = 1$, $a_{12} = 1$, $a_{21} = 0$, $a_{22} = -1$, $\deg(E|_{C_1}) = 2$, $\deg(E|_{C_2}) = -1$. $\mu_{\min}(\eta_{c_1}^*(E|_{c_1})) = \min\{a_{1j} \mid 1 \le j \le 2\} = 1$, $\mu_{\min}(\eta_{c_2}^*(E|_{c_2})) = \min\{a_{2j} \mid 1 \le j \le 2\} = -1$.

Therefore, $y_0\xi + \sum_{i=1}^2 y_i(\pi^*[C_i]) \in N^1(\mathbb{P}(E))_{\mathbb{R}}$ is in Nef($\mathbb{P}(E)$) if and only if $y_0 \ge 0$, $y_0\mu_{\min}(\eta^*_{c_1}(E|_{c_1})) + \sum_{i=1}^2 y_ib_{i1} = y_0 + y_2 \ge 0$ and $y_0\mu_{\min}(\eta^*_{c_2}(E|_{c_2})) + \sum_{i=1}^2 y_ib_{i2} = y_1 - y_0 - y_2 \ge 0$, i.e.

Nef(
$$\mathbb{P}(E)$$
) = $\left\{ y_0 \xi + \sum_{i=1}^2 y_i(\pi^*[C_i]) \mid y_0 \ge 0, y_0 + y_2 \ge 0 \text{ and } y_1 - y_0 - y_2 \ge 0 \right\}.$

Also note that X is a toric variety. The fan structure of X is given in [4], Definition 3.3.17. Similarly, as it is mentioned in Remark 2, one can give the toric description of the Nef cone of X.

EXAMPLE 3.7. Let $\rho : X = \mathbb{P}(W) \longrightarrow C$ be a ruled surface over a smooth curve *C*, defined by an unstable normalized rank 2 bundle $W = \mathcal{O}_C \oplus M$ for some line bundle *M* on *C* with $\deg(M) = l < 0$. Then, $\overline{NE}(X) = \{a\zeta + bf \mid a, b \in \mathbb{R}_{\geq 0}\}$, where $[\mathcal{O}_{\mathbb{P}(W)}(1)] = \zeta \in N^1(X)_{\mathbb{R}}$ and *f* is the numerical class of a fibre of ρ .

Let $\rho_x : \tilde{X}_x = Bl_x(X) \longrightarrow X$ be the blow up of X at a closed point x in the section σ such that $\mathcal{O}_X(\sigma) \cong \mathcal{O}_{\mathbb{P}(W)}(1)$, and E_x be the exceptional divisor. Then, $\overline{NE}(\tilde{X}_x) = \{a[C_1] + b[C_2] + c[C_3] \mid a, b, c \in \mathbb{R}_{\geq 0}\}$, where $[C_1] = \rho_x^* f - E_x$, $[C_2] = \rho_x^* \zeta - E_x$, $[C_3] = E_x$ (see Example 3.3). The intersection products,

$$b_{11} := C_1 \cdot C_1 = (\rho_x^* f - E_x) \cdot (\rho_x^* f - E_x) = E_x^2 = -1; \\ b_{22} := C_2 \cdot C_2 = (\rho_x^* \zeta - E_x) \cdot (\rho_x^* \zeta - E_x) = \deg(W) - 1 = l - 1; \\ b_{12} = b_{21} := C_1 \cdot C_2 = (\rho_x^* f - E_x) \cdot (\rho_x^* \zeta - E_x) = \zeta \cdot f + E_x^2 = 0; \\ b_{13} = b_{31} := C_1 \cdot C_3 = (\rho_x^* f - E_x) \cdot E_x = 1; \\ b_{23} = b_{32} := C_2 \cdot C_3 = (\rho_x^* \zeta - E_x) \cdot E_x = 1; \\ b_{23} = b_{23} := C_2 \cdot C_3 = (\rho_x^* \zeta - E_x) \cdot E_x = 1; \\ b_{23} = b_{23} := C_2 \cdot C_3 = (\rho_x^* \zeta - E_x) \cdot E_x = 1; \\ b_{23} = b_{23} := C_2 \cdot C_3 = (\rho_x^* \zeta - E_x) \cdot E_x = 1; \\ b_{23} = b_{23} := C_2 \cdot C_3 = (\rho_x^* \zeta - E_x) \cdot E_x = 1; \\ b_{23} = b_{23} := C_2 \cdot C_3 = (\rho_x^* \zeta - E_x) \cdot E_x = 1; \\ b_{23} = b_{23} := C_2 \cdot C_3 = (\rho_x^* \zeta - E_x) \cdot E_x = 1; \\ b_{23} = b_{23} := C_2 \cdot C_3 = (\rho_x^* \zeta - E_x) \cdot E_x = 1; \\ b_{23} = b_{23} := C_2 \cdot C_3 = (\rho_x^* \zeta - E_x) \cdot E_x = 1; \\ b_{23} = b_{23} := C_2 \cdot C_3 = (\rho_x^* \zeta - E_x) \cdot E_x = 1; \\ b_{23} = b_{23} := C_2 \cdot C_3 = (\rho_x^* \zeta - E_x) \cdot E_x = 1; \\ b_{23} = b_{23} := C_2 \cdot C_3 = (\rho_x^* \zeta - E_x) \cdot E_x = 1; \\ b_{23} = b_{23} := C_2 \cdot C_3 = (\rho_x^* \zeta - E_x) \cdot E_x = 1; \\ b_{23} = b_{23} := C_2 \cdot C_3 = (\rho_x^* \zeta - E_x) \cdot E_x = 1; \\ b_{23} = b_{23} := C_2 \cdot C_3 = (\rho_x^* \zeta - E_x) \cdot E_x = 1; \\ b_{23} = b_{23} := C_2 \cdot C_3 = (\rho_x^* \zeta - E_x) \cdot E_x = 1; \\ b_{23} = b_{23} := C_2 \cdot C_3 = (\rho_x^* \zeta - E_x) \cdot E_x = 1; \\ b_{23} = b_{23} := C_2 \cdot C_3 = (\rho_x^* \zeta - E_x) \cdot E_x = 1; \\ b_{23} = b_{23} := C_2 \cdot C_3 = (\rho_x^* \zeta - E_x) \cdot E_x = 1; \\ b_{23} = b_{23} := C_2 \cdot C_3 = (\rho_x^* \zeta - E_x) \cdot E_x = 1; \\ b_{23} = b_{23} := C_2 \cdot C_3 = (\rho_x^* \zeta - E_x) \cdot E_x = 1; \\ b_{23} = b_{23} := C_2 \cdot C_3 = (\rho_x^* \zeta - E_x) \cdot E_x = 1; \\ b_{23} = b_{23} := C_2 \cdot C_3 = (\rho_x^* \zeta - E_x) \cdot E_x = 1; \\ b_{23} = b_{23} := C_2 \cdot C_3 = (\rho_x^* \zeta - E_x) \cdot E_x = 1; \\ b_{23} = b_{23} := C_2 \cdot C_3 = (\rho_x^* \zeta - E_x) \cdot E_x = 1; \\ b_{23} = b_{23} := C_2 \cdot C_3 = (\rho_x^* \zeta - E_x) \cdot E_x = 1; \\ b_{23} = b_{23} := C_2 \cdot C_3 = (\rho_x^* \zeta - E_x) \cdot E_x = 1; \\ b_{23} = b_{23} := C_2 \cdot C_3 = (\rho_x^* \zeta - E_x) \cdot E_x = 1; \\ b_{23} = b_{23} := C_2 \cdot C_3 = (\rho_x^* \zeta - E_x) \cdot E_x = 1; \\ b_{23}$$

 $b_{13} = b_{31} := C_1 \cdot C_3 = (\rho_x^* f - E_x) \cdot E_x = 1 ; b_{23} = b_{32} := C_2 \cdot C_3 = (\rho_x^* \zeta - E_x) \cdot E_x = 1.$ Let us consider the rank 2 bundle $E = \mathcal{O}_{\tilde{X}_x} \oplus \mathcal{O}_{\tilde{X}_x}(nE_x)$ on \tilde{X}_x , for some $n \in \mathbb{Z}$ with $n \ge 0$. Then, $\mathbb{P}(E)$ has Picard number 4 in this case. Fix the notations $L_1 := \mathcal{O}_{\tilde{X}_x}, L_2 := \mathcal{O}_{\tilde{X}_x}(nE_x)$, and $a_{ij} = C_i \cdot L_j$ for $1 \le i \le 3, 1 \le j \le 2$. Then, $a_{11} = 0, a_{12} = n, a_{21} = 0, a_{22} = n, a_{31} = 0, a_{32} = -n$. Hence $\mu_{\min}(\eta_{c_1}^*(E|_{C_1})) = \min\{a_{1j} \mid 1 \le j \le 2\} = 0 ; \mu_{\min}(\eta_{c_2}^*(E|_{C_2})) = \min\{a_{2j} \mid 1 \le j \le 2\} = 0$
$$\begin{split} 1 &\leq j \leq 2\} = 0 \;; \; \mu_{\min}(\eta_{c_3}^*(E|_{C_3})) = \min\{a_{3j} \mid 1 \leq j \leq 2\} = -n. \\ \text{Also, } \deg(E|_{C_1}) &= (\mathcal{O}_{\bar{X}_x} \oplus nE_x) \cdot (\rho_x^*f - E_x) = n. \\ \text{Similarly, } \deg(E|_{C_2}) = n \;; \; \deg(E|_{C_3}) = -n. \\ \text{Therefore, } y_0 \xi + \sum_{i=1}^3 y_i (\pi^*[C_i]) \in N^1(\mathbb{P}(E))_{\mathbb{R}} \text{ is in Nef}(\mathbb{P}(E)) \text{ if and only if } y_0 \geq 0 \;, \\ y_0 \mu_{\min}(\eta_{c_1}^*(E|_{C_1})) + \sum_{i=1}^3 y_i b_{i1} = -y_1 + y_3 \geq 0, \\ y_0 \mu_{\min}(\eta_{c_2}^*(E|_{C_2})) + \sum_{i=1}^3 y_i b_{i2} = (l-1)y_2 + y_3 \geq 0 \text{ and} \\ y_0 \mu_{\min}(\eta_{c_3}^*(E|_{C_3})) + \sum_{i=1}^3 y_i b_{i3} = -ny_0 + y_1 + y_2 - y_3 \geq 0. \end{split}$$

4. Seshadri Constants

We begin this section by the following easy observation.

Lemma 4.1. Let $Z \subseteq X$ be a closed subvariety in a smooth complex projective variety X. For a vector bundle V on X and for any point $x \in Z$

$$0 < \varepsilon(V, x) \le \varepsilon(V|_Z, x).$$

Proof. We have the following commutative diagram.

$$\begin{array}{ccc} \operatorname{Bl}_{x} Z & \stackrel{j}{\longrightarrow} & \operatorname{Bl}_{x} X \\ & \downarrow^{\psi_{x}} & \qquad \downarrow^{\rho_{x}} \\ Z & \stackrel{i}{\longrightarrow} & X \end{array}$$

where *j* and *i* are inclusions. Note that the exceptional divisor in $Bl_x Z$ is the restriction of the exceptional divisor in $Bl_x X$, as $Bl_x Z$ is the strict transform of Z in $Bl_x X$.

Also we have the following commutative diagram.

In the above diagram we have the inclusion $\mathbb{P}(\psi_x^*(V|_Z)) \hookrightarrow \mathbb{P}(\rho_x^*V)$ and

$$\mathcal{O}_{\mathbb{P}(\rho_x^*V)}(1)|_{\mathbb{P}(\psi_x^*(V|_Z))} = \mathcal{O}_{\mathbb{P}(\psi_x^*(V|_Z))}(1).$$

As restriction of nef divisor is nef, we conclude from the definition of Seshadri constant that

$$0 < \varepsilon(V, x) \le \varepsilon(V|_Z, x).$$

We recall from [8,Chapter 5] that a vector bundle W of rank 2 on a smooth projective curve C is said to be normalized if $H^0(W) \neq 0$, but $H^0(W \otimes L) = 0$ for all line bundle L on C with deg(L) < 0. We notice that a normalized bundle W is semistable if and only if deg(W) ≥ 0 .

Theorem 4.2. Let ρ : $Y = \mathbb{P}(W) \longrightarrow C$ be a ruled surface over a smooth curve defined by a normalized rank 2 bundle W such that $\mu_{\min}(W) = \deg(W)$. Let σ be a section of ρ such that $\mathcal{O}_Y(\sigma) \cong \mathcal{O}_{\mathbb{P}(W)}(1)$ and f denotes a fibre of the map ρ . If V is a semistable ample bundle with discriminant $\Delta(V) = 0$, then the Seshadri constant at a closed point $y \in \sigma$ is given by

$$\varepsilon(V, y) = \min\{\mu(V|_{\sigma}), \mu(V|_{f})\}.$$

Proof. Consider the following fibre product diagram.

Since $\mu_{\min}(W) = \deg(W)$, we have $\overline{NE}(Y) = \{a[\sigma] + b[f] \mid a, b \in \mathbb{R}_{\geq 0}\}$. We denote the exceptional divisor of the map ρ_y by E_y . We also note that

$$NE(X) = \{a[C_1] + b[C_2] + c[C_3] \mid a, b, c \in \mathbb{R}_{\ge 0}\},\$$

where $C_1 \equiv (\rho_y^* \sigma - E_y)$, $C_2 \equiv (\rho_y^* f - E_y)$ and $C_3 \equiv E_y$ (see Example 3.3). Note that all three curves C_1 , C_2 and C_3 are smooth. We define $E := \rho_u^*(V)$.

Now, if *V* is a semistable ample bundle with $\triangle(V) = 0$, then the pullback bundle $E = \rho_y^*(V)$ is also semistable bundle with $\triangle(E) = 0$ by Theorem 1.2 in [2]. Let ξ_y be the numerical equivalence class of the tautological bundle $\mathcal{O}_{\mathbb{P}(\rho_y^*V)}(1)$, and $\tilde{E}_y := \rho_y^{-1}(F_y)$, where F_y is the class of the fibre of the map π' over the point *y*. Therefore, applying Corollary 3.1 we get that for a positive real number $\lambda > 0$, $\xi_y - \lambda \tilde{E}_y$ is nef if and only if

 $\mu(E|_{C_1}) - \lambda(C_3 \cdot C_1) = (\mu(V|_f) - \lambda) \ge 0, \ \mu(E|_{C_2}) - \lambda(C_3 \cdot C_2) = (\mu(V|_{\sigma}) - \lambda) \ge 0, \text{ and} \\ \mu(E|_{C_3}) - \lambda(C_3 \cdot C_3) = \lambda \ge 0.$

Therefore, the Seshadri constant at a closed point $y \in \sigma$ is given by

$$\varepsilon(V,y) = \sup \{\lambda > 0 \mid \tilde{\xi}_y - \lambda \tilde{E}_y \text{ is nef } \} = \min \{\mu(V|_{\sigma}), \mu(V|_f) \}.$$

REMARK 3. Let $X : \mathbb{P}(\mathcal{O}_C \oplus \mathcal{L}) \longrightarrow C$ be a ruled surface defined by a normalized rank 2 bundle $W = \mathcal{O}_C \oplus \mathcal{L}$ with deg $(\mathcal{L}) < 0$. The above result generalizes known results about Seshadri constants of ample line bundles on such ruled surfaces (see Theorem 4.1 in [6]).

EXAMPLE 4.3. Let $V = L_1 \oplus L_2 \oplus \cdots \oplus L_r$ is a completely decomposable ample bundle on \mathbb{P}^2 . Consider the following commutative fibred diagram.

We define $E := \phi_x^*(V)$. Let *l* be a line in \mathbb{P}^2 passing through *x* and $\phi_x^*(l) = H_x$. We denote the exceptional divisor of the map ϕ_x by E_x . Then we have $\overline{NE}(X) = \{a[C_1] + b[C_2] \mid a, b \in \mathbb{R}_{\geq 0}\}$, where $C_1 \equiv H_x - E_x$, $C_2 \equiv E_x$. Note that both C_1 and C_2 are smooth curves. We denote the numerical equivalence class of the tautological bundle $\mathcal{O}_{\mathbb{P}(E)}(1)$ by ξ_x , and define $\tilde{E}_x := \pi^*(E_x)$.

Using Corollary 3.4 we get that for a positive real number $\lambda > 0$, $\tilde{\xi}_x - \lambda \tilde{E}_x = \tilde{\xi}_x - \lambda \pi^* [C_2]$ is nef if and only if

 $\lambda \ge 0, \text{ and } \min_{1 \le i \le r} \{L_i \cdot l\} - \lambda = \min_{1 \le i \le r} \{\deg(L_i)\} - \lambda = \mu_{\min}(V) - \lambda \ge 0.$ Hence, the Seshadri constant of $V = L_1 \oplus L_2 \oplus \cdots \oplus L_r$ at $x \in \mathbb{P}^2$ is $\varepsilon(V, x) = \sup \{\lambda > 0 \mid \tilde{\xi}_x - \lambda \tilde{E}_x \text{ is nef }\} = \mu_{\min}(V) = \min_{1 \le i \le r} \{\deg(L_i)\}.$

A completely decomposable ample vector bundle on \mathbb{P}^2 is a special case of a torus equivariant vector bundle over a toric variety. The Seshadri constant of a nef toric vector bundle on a toric variety is calculated in [9, Proposition 3.2] which also agrees with our result.

Acknowledgements. The authors would like to thank Prof. Indranil Biswas, Prof. D S Nagaraj and Dr. Krishna Hanumanthu for several useful discussions. The authors would also like to thank the referee for his/her valuable comments and useful suggestions towards the overall improvement of the content of this article. This work is supported financially by a fellowship from TIFR, Mumbai under DAE, Government of India.

References

- T. Bauer, S. Di Rocco, B. Harbourne, M. Kapustka, A. Knutsen, W. Syzdek and T. Szemberg: A primer on Seshadri constants; in Interactions of classical and numerical algebraic geometry, Contemp. Math. 496, Amer. Math. Soc., Providence, RI 2009, 33–70.
- [2] I. Biswas and U. Bruzzo: On semistable principal bundles over a complex projective manifold, Int. Math. Res. Not. IMRN (2008), Article ID rnn035, 28pp.
- [3] I. Biswas, K. Hanumanthu and D.S. Nagaraj: *Positivity of vector bundles on homogeneous varieties*, Internat. J. Math. **31** (2020), 2050097, 11pp.
- [4] D.A. Cox, J.B. Little and H.K. Schenck: Toric Varieties, Graduate Studies in Mathematics 124, American Mathematical Society, Providence, RI, 2011.
- [5] M. Fulger: The cones of effective cycles on projective bundles over curves, Math. Z. 269 (2011), 449–459.
- [6] L.F. Garciá: Seshadri constants on ruled surfaces: the rational and elliptic cases, Manuscripta Math. 119 (2006), 483–505.
- [7] C. Hacon: Remarks on Seshadri constants of vector bundles, Ann. Inst. Fourier (Grenoble) 50 (2000), 767–780.
- [8] R. Hartshorne: Algebraic Geometry, Graduate Text in Mathematics 52, Springer-Verlag, New York-Heidelberg, 1977.
- [9] M. Hering, M. Mustatá and S. Payne: *Positivity properties of toric vector bundles*, Ann. Inst. Fourier (Grenoble) 60 (2010), 607–640.
- [10] R. Karmakar, S. Misra and N. Ray: Nef and Pseudoeffective cones of product of projective bundles over a curve, Bull. Sci. Math. 151 (2019), 1–12.
- [11] R. Karmakar and S. Misra: Nef cone and Seshadri constants of product of projective bundles over curves, J. Ramanujan Math. Soc. 35 (2020), 317–325.
- [12] R. Lazarsfeld: Positivity in Algebraic Geometry I, Springer-Verlag, Berlin, 2004.
- [13] Y. Miyaoka: The Chern classes and Kodaira dimension of a minimal variety; in Algebraic Geometry, Sendai, 1985, Adv. Stud. Pure Math. 10, North-Holland, Amsterdam, 1987, 449–476.
- [14] R. Muñoz, G. Occhetta and L.E. Solá Conde: On rank 2 vector bundles on Fano manifolds, Kyoto J. Math. 54 (2014), 167–197.
- [15] T. Oda: Convex Bodies and Algebraic Geometry, Springer-Verlag Berlin, 1988.

Snehajit Misra Tata Institute of Fundamental Research (TIFR) Homi Bhabha Road, Mumbai 400005 India e-mail: smisra@math.tifr.res.in

Nabanita Ray Tata Institute of Fundamental Research (TIFR) Homi Bhabha Road, Mumbai 400005 India e-mail: nray@math.tifr.res.in