



Title	A STUDY OF A FUCHSIAN SYSTEM OF RANK 8 IN 3 VARIABLES AND THE ORDINARY DIFFERENTIAL EQUATIONS AS ITS RESTRICTIONS
Author(s)	Ebisu, Akihito; Haraoka, Yoshishige; Kaneko, Masanobu et al.
Citation	Osaka Journal of Mathematics. 2023, 60(1), p. 153-206
Version Type	VoR
URL	https://doi.org/10.18910/89997
rights	
Note	

The University of Osaka Institutional Knowledge Archive : OUKA

<https://ir.library.osaka-u.ac.jp/>

The University of Osaka

A STUDY OF A FUCHSIAN SYSTEM OF RANK 8 IN 3 VARIABLES AND THE ORDINARY DIFFERENTIAL EQUATIONS AS ITS RESTRICTIONS

AKIHIKO EBISU, YOSHISHIGE HARAOKA, MASANOBU KANEKO, HIROYUKI OCHIAI,
TAKESHI SASAKI and MASAAKI YOSHIDA

(Received May 11, 2020, revised October 26, 2021)

Abstract

A Fuchsian system of rank 8 in 3 variables with 4 parameters is found. The singular locus consists of six planes and a cubic surface. The restriction of the system onto the intersection of two singular planes is an ordinary differential equation of order four with three singular points. A middle convolution of this equation turns out to be the tensor product of two Gauss hypergeometric equations, and another middle convolution sends this equation to the Dotsenko-Fateev equation. Local solutions of these ordinary differential equations are found. Their coefficients are *sums* of products of the Gamma functions. These sums can be expressed as special values of the generalized hypergeometric series ${}_4F_3$ at 1.

Contents

Introduction	155
Part I. A Fuchsian system of rank 8 in 3 variables and its restrictions	156
1. A Fuchsian system of rank 8 in 3 variables $Z_3(A)$	156
1.1. Outline of the proof of Theorem 1.1	158
2. Restriction of $Z_3(A)$ onto the diagonal $t_1 = t_2 = t_3$	158
2.1. $Z_{\Delta 8}(A)$	158
2.2. $Z_{\Delta 6}(A)$	159
2.3. $Z_{\Delta 4}(A)$	159
3. Restriction of $Z_3(A)$ onto the plane $t_3 = 1$ and $Z_2(A)$	159
3.1. Equation $Z_2(A)$	159
3.2. Outline of the proof of Theorem 3.1	161
3.3. Restriction of $Z_2(A)$ onto the diagonal $t_1 = t_2$	161
4. Restriction of $Z_2(A)$ onto the line $t_2 = 1$ and $Z(A)$	161
4.1. Equation $Z(A)$	161
4.2. A small change $\tilde{Z}(A)$ of $Z(A)$	162
4.3. Invariants of ordinary differential operators	163
5. $\tilde{Z}(A)$ is related to the tensor product of two Gauss equations	164
5.1. Definition and fundamental properties of addition and middle convolution	164

5.2. A middle convolution connects $\tilde{Z}(A)$ with the tensor product of two Gauss equations	165
6. Relation between $\tilde{Z}(A)$ and the Dotsenko-Fateev equation	168
6.1. The Dotsenko-Fateev equation	168
6.2. A middle convolution and an addition send $\tilde{Z}(A)$ to the Dotsenko-Fateev equation	169
7. Table of related differential equations	170
8. Explicit expressions of matrix 1-forms	172
8.1. 8×8 -matrix form $\omega = M_1 dt_1 + M_2 dt_2 + M_3 dt_3$	172
8.2. 6×6 -matrix form $\omega_6 = N_1 dt_1 + N_2 dt_2$	174
9. Tensor product of two Gauss equations	175
9.1. Tensor product without apparent singularities	177
9.2. Tensor product without apparent singularities Case 1	177
9.3. Tensor product without apparent singularities Case 2	180
Part II. Local solutions of ordinary differential equations related to the Dotsenko-Fateev equation	182
10. Local solutions of $Z(A)$ at $x = 0$ with exponent 0 and $2A_0$, and those at infinity	182
10.1. Invariants of linear difference equations	183
10.2. Recurrence relation $Rc_0(A)$ for the coefficients of a holomorphic solution of $Z(A)$ at $x = 0$	184
10.3. 3-term relation for the special values (at 1) of balanced terminating hypergeometric series ${}_4F_3$	185
10.4. Local solutions at zero I: solving $Rc_0(A)$	186
10.5. Other local solutions expressed in terms of $f^{(0,0)}$	189
10.6. Local solutions at infinity I: using invariants of the difference equations	190
10.7. Correspondence of solutions via the Riemann-Liouville transformation	192
10.8. Partial correspondence of local solutions at $x = 0, 1$	193
10.9. Local solutions at infinity II: using middle convolution	193
11. Local solutions of $Z(A)$ at $x = 0$ with exponent $A_0 \pm 1/2$	194
11.1. Recurrence relation $Rc_1(A)$	194
11.2. Special values of non-terminating ${}_4F_3$ at $x = 1$	194
11.3. Difference equation $Rc^{(1)}(\alpha)$: an extension of $Rc^{(0)}(\hat{\alpha})$	195
11.4. Local solutions at zero II: solving $Rc_1(A)$	196
12. Local solutions of $Q(A)$	199
12.1. Holomorphic solution $f_Q^{(0,0)}(A; x)$ to $Q(A)$ at $x = 0$	200
12.2. Local solution to $Q(A)$ at $x = 0$ with exponent $A_0 - A_2$	201
12.3. Pfaff transforms of the solutions of $Q(A)$	202
12.4. Local solutions of $Q(A)$	202
13. Local solutions of the Dotsenko-Fateev equation	202
13.1. Local solutions of the Dotsenko-Fateev equation at $x = 0$	203
13.2. Local solutions of the Dotsenko-Fateev equation at $x = 1$	204
13.3. Solutions of the Dotsenko-Fateev equation at $x = \infty$	205
References	205

Introduction

In Part I, we find a Fuchsian system $Z_3(A)$ of rank 8 in 3 variables (t_1, t_2, t_3) with 4 parameters $A = (A_0, A_1, A_2, A_3)$. The singular locus consists of six planes and a cubic surface (the Fricke surface):

$$t_i = \pm 1 \quad (i = 1, 2, 3), \quad 1 - t_1^2 - t_2^2 - t_3^2 + 2t_1t_2t_3 = 0.$$

In general, we can have ordinary differential equations as we like, we have only to give polynomials to coefficients, but to have a system in more than 2 variables of finite and non-zero rank, the coefficients must satisfy the integrability condition (a system of non-linear differential equations). Very few examples are known (most of them belong to the so-called hypergeometric family). Our system $Z_3(A)$, which is not hypergeometric, is an important example.

The restriction of the system onto the plane, say $t_3 = 1$, is a system $Z_2(A)$ of rank 6 in 2 variables. Its singular locus consists of

$$t_i = \pm 1, \quad (i = 1, 2) \quad t_1 = t_2,$$

which is equal to that of the well-known Appell's hypergeometric system F_1 .

In the future, we would like to study these systems; power series solutions, integral representations of the solutions, etc.

To understand these systems we begin with studying its several restrictions. The restriction of the system onto the diagonal $t_1 = t_2 = t_3$ is an ordinary differential equation $Z_{\Delta 8}$ of order 8.

The restriction of the system $Z_2(A)$ onto the line, say, $t_2 = 1$ is an ordinary differential equation $Z(A)$ of order 4 with three singular points $t_1 = \pm 1$ and ∞ . This equation $Z(A)$ has not been studied so far, to the authors knowledge.

While studying local solutions of $Z(A)$, which are fully presented in Part II, we find a power-series solution to $Z(A)$ at $t = 1$, which is very similar to the product of two Gauss hypergeometric series. This leads to the discovery that a middle convolution sends the equation $Z(A)$ to the tensor product of two Gauss hypergeometric equations, with special parameters. We also find that another middle convolution sends $Z(A)$ to the Dotsenko-Fateev equation.

In Part II, we study local solutions for the ordinary differential equation $Z(A)$ and for several related ones around their singular points. We see the relation between $Z(A)$ and the tensor product of two specific Gauss hypergeometric equations. At a singular point of $Z(A)$, say $t = 1$, the coefficients of the holomorphic solution to $Z(A)$ satisfy a 3-term difference equation $Rc_0(A)$. On the other hand, ${}_4F_3(*; 1)$, special values at the unit argument of the terminating generalized hypergeometric series ${}_4F_3$, satisfy a linear difference equation of order 2, if the parameters are carefully chosen. Comparing the invariant of this difference equation with that of $Rc_0(A)$, we find solutions of $Rc_0(A)$ expressed in terms of ${}_4F_3(*; 1)$. From the observation that the special values ${}_4F_3(*; 1)$ appear as the coefficients of the product of two Gauss hypergeometric series, we notice that its product has relevance to the holomorphic solution to $Z(A)$ at $t = 1$, which leads to the discovery stated above.

For most local solutions of the ordinary differential equations related to $Z(A)$, we can make use of middle convolutions connecting the equation and the tensor product of two

Gauss equations to get explicit expressions for the solutions. But in these cases also, we present a way to get them by using the difference equations for ${}_4F_3(*; 1)$, because this method gives various expressions.

The coefficients of hypergeometric-type series are products of the Gamma functions. However for our equation $Z(A)$ and the related ones including the Dotsenko-Fateev equation, the coefficients of local solutions are *sums* of products of the Gamma functions. These sums can be expressed as special values ${}_4F_3(*; 1)$.

Solutions of the ordinary differential equations we studied in this paper admit Euler integral representations, which will be discussed elsewhere.

Part I. A Fuchsian system of rank 8 in 3 variables and its restrictions

In §1, we find a Fuchsian system $Z_3(A)$ of rank 8 in 3 variables (t_1, t_2, t_3) with 4 parameters $A = (A_0, A_1, A_2, A_3)$.

In §2, the restriction $Z_{\Delta 8}(A)$ of the system $Z_3(A)$ onto the diagonal $t_1 = t_2 = t_3$ is studied.

In §3, the restriction of the system $Z_3(A)$ onto the plane $t_3 = 1$ is studied and the system $Z_2(A)$ is found. The restriction of the system $Z_2(A)$ onto the diagonal $t_1 = t_2$ is also studied.

In §4, the restriction of the system $Z_2(A)$ onto the line $t_2 = 1$ is studied and the system $Z(A)$ is found. This is an ordinary differential equation of order 4 with three singular points $t_1 = \pm 1$ and ∞ .

§5 gives a relation between $Z(A)$ and the tensor product of two specific Gauss hypergeometric equations.

§6 gives a relation between $Z(A)$ and the Dotsenko-Fateev equation.

§7 illustrates the relation among the differential equations appeared above.

In §8, a Pfaffian form of $Z_3(A)$ is presented.

§9 studies the tensor products of two Gauss hypergeometric equations.

1. A Fuchsian system of rank 8 in 3 variables $Z_3(A)$

We treat ideals of the ring of differential operators $\mathbf{C}[a_0, \dots, t_1, \dots, \partial/\partial t_1, \dots]$. We often call a set of generators of an ideal simply as a system, which sometimes also means the corresponding system of differential equations, after introducing an unknown, say F, u, \dots .

In 2017, Don Zagier showed us a system generated by the differential operator

$$(1 - t_1^2)\partial_{11} + 2(t_3 - t_1 t_2)\partial_{12} + (1 - t_2^2)\partial_{22} + a_0 t_1 \partial_1 + a_0 t_2 \partial_2$$

and those obtained by a succession of the cyclic permutation $1 \rightarrow 2 \rightarrow 3 \rightarrow 1$ with a parameter a_0 , where $\partial_1 = \partial/\partial t_1$, $\partial_{12} = \partial^2/\partial t_1 \partial t_2$, etc. This system in 3 variables (t_1, t_2, t_3) is Fuchsian of rank 8, and is highly reducible. Hoping to have less reducible system of rank 8, we considered a bit general system with more parameters and got the following result.

Theorem 1.1. *The system generated by the operator*

$$E_3 = (1 - t_1^2)\partial_{11} + 2(t_3 - t_1 t_2)\partial_{12} + (1 - t_2^2)\partial_{22} + a_{31} t_1 \partial_1 + a_{32} t_2 \partial_2 + a_{33} t_3 \partial_3 + a_{30}$$

and those obtained by a succession of the cyclic permutation $1 \rightarrow 2 \rightarrow 3 \rightarrow 1$ with constants

a_{ij} ($i = 1, 2, 3$, $j = 0, 1, 2, 3$) is of rank 8 if and only if

$$a_{11} = a_{22} = a_{33} = 0, \quad a_{12} = a_{13} = a_{21} = a_{23} = a_{31} = a_{32} (=: a_0).$$

Set $a_1 = a_{10}$, $a_2 = a_{20}$, $a_3 = a_{30}$. Then the operators E_1 , E_2 and E_3 are given as

$$\begin{aligned} E_1 &= (1 - t_2^2)\partial_{22} + 2(t_1 - t_2 t_3)\partial_{23} + (1 - t_3^2)\partial_{33} + a_0 t_2 \partial_2 + a_0 t_3 \partial_3 + a_1, \\ E_2 &= (1 - t_3^2)\partial_{33} + 2(t_2 - t_3 t_1)\partial_{31} + (1 - t_1^2)\partial_{11} + a_0 t_3 \partial_3 + a_0 t_1 \partial_1 + a_2, \\ E_3 &= (1 - t_1^2)\partial_{11} + 2(t_3 - t_1 t_2)\partial_{12} + (1 - t_2^2)\partial_{22} + a_0 t_1 \partial_1 + a_0 t_2 \partial_2 + a_3, \end{aligned}$$

with parameters $a = (a_0, a_1, a_2, a_3)$. We often use parameters $A = (A_0, A_1, A_2, A_3)$ related to a by

$$a_0 = 2A_0 - 3, \quad a_i = A_i^2 - (A_0 - 1)^2 \quad i = 1, 2, 3,$$

and name the system as $Z_3(A)$.

By using $b_i = (a_1 + a_2 + a_3)/2 - a_i$ ($i = 1, 2, 3$) as parameters, F as unknown, and writing $F_1 = \partial_1 F$, $F_{12} = \partial_{12} F$, etc, this system can be also written as

$$\begin{aligned} (t_1^2 - 1)F_{11} &= (t_3 - t_1 t_2)F_{12} + (t_2 - t_3 t_1)F_{13} - (t_1 - t_2 t_3)F_{23} + a_0 t_1 F_1 + b_1 F, \\ (t_2^2 - 1)F_{22} &= (t_1 - t_2 t_3)F_{23} + (t_3 - t_1 t_2)F_{21} - (t_2 - t_3 t_1)F_{31} + a_0 t_2 F_2 + b_2 F, \\ (t_3^2 - 1)F_{33} &= (t_2 - t_3 t_1)F_{31} + (t_1 - t_2 t_3)F_{32} - (t_3 - t_1 t_2)F_{12} + a_0 t_3 F_3 + b_3 F. \end{aligned}$$

Proposition 1.2. *The system $Z_3(A)$ is Fuchsian, and the singular locus in the finite space consists of six planes and a cubic surface:*

$$t_i = \pm 1 \quad (i = 1, 2, 3), \quad 1 - t_1^2 - t_2^2 - t_3^2 + 2t_1 t_2 t_3 = 0.$$

The local exponents along the divisors are given as

$$t_i = \pm 1 : \quad 0, 1, 2, 3, 4, 5, 1/2 \pm A_i,$$

$$\text{the cubic surface} : \quad 0, 1, 2, 3, A_0, A_0 + 1, A_0 + 2, A_0 + 3,$$

$$\begin{aligned} t_i = \infty : \quad 1 - A_0 \pm A_j, \quad 1 - A_0 \pm A_k, \quad 2 - A_0 \pm A_j, \quad 2 - A_0 \pm A_k \\ (\{i, j, k\} = \{1, 2, 3\}). \end{aligned}$$

The local exponents along a divisor are defined as those of the ordinary differential equation obtained by restricting the system onto a curve intersecting the divisor transversely at an ordinary point of the divisor.

The singularities are known from the matrix 1-form ω in the next subsection. If we restrict the system onto a generic line $t_2 = \text{constant}$, $t_3 = \text{constant}$, we get an ordinary differential equation of order 8 in $t := t_1$ with polynomial coefficients:

$$(t + 1)^2(t - 1)^2(1 - t^2 - t_2^2 - t_3^2 + 2t_2 t_3 t)^4 P(t) \frac{d^8 F}{dt^8} + \dots = 0,$$

where $P(t)$ is of degree 16, the number of apparent singular points, the local exponents at each points are 0, 1, 2, 3, 4, 5, 6 and 8. Though we omit the explicit expression of the coefficients of the ordinary equation above, we find the local exponents at the singular points as in the proposition.

REMARK 1.3 (SYMMETRY). The system $Z_3(A)$ is invariant under

$$(t_1, t_2, t_3) \rightarrow (\varepsilon_1 t_1, \varepsilon_2 t_2, \varepsilon_3 t_3), \quad \varepsilon_i = \pm 1, \quad \varepsilon_1 \varepsilon_2 \varepsilon_3 = 1,$$

$$A_j \rightarrow -A_j \quad (j = 1, 2, 3),$$

$$(t_1, t_2, t_3, A_1, A_2, A_3) \rightarrow (t_{\sigma(1)}, t_{\sigma(2)}, t_{\sigma(3)}, A_{\sigma(1)}, A_{\sigma(2)}, A_{\sigma(3)}),$$

where σ is a permutation of $\{1, 2, 3\}$.

1.1. Outline of the proof of Theorem 1.1. Several integrable systems of partial differential equations with many variables are known; for example Appell-Lauricella's hypergeometric system F_A in n variables. The rank of F_A is known to be 2^n . The form of the equations tells immediately the rank does not exceed 2^n . But it would be quite difficult to prove that the rank is exactly 2^n by manipulating the differential equations; this is proved by finding 2^n linearly independent hypergeometric series at a singular point.

In our case, no local solutions are known; so, we are forced to check honestly the integrability condition. We transform the system $Z_3(A)$ into a Pfaffian form of size 8, and show the integrability.

Let F be the unknown, $F_{i..k}$ the partial derivative of F by t_i, t_j, \dots, t_k , and set

$$e = {}^{\text{tr}}(F, F_1, F_2, F_3, F_{12}, F_{13}, F_{23}, DF_{123}), \quad D := -1 + t_1^2 + t_2^2 + t_3^2 - 2t_1 t_2 t_3.$$

A computation shows that the derivatives $F_{i..k}$ can be written as linear combinations of $F, F_1, F_2, F_3, F_{12}, F_{13}, F_{23}$ and F_{123} , and thus we get a Pfaffian system of the form

$$de = \omega e,$$

where ω is an 8×8 -matrix 1-form given in §8.1. The integrability condition of the system is written as

$$d\omega = \omega \wedge \omega,$$

and, by computation, we get Theorem 1.1.

2. Restriction of $Z_3(A)$ onto the diagonal $t_1 = t_2 = t_3$

Let $F(t_1, t_2, t_3)$ be a solution of $Z_3(A)$. The function $F(t, t, t)$ satisfies a Fuchsian ordinary differential equation. In this section, its singular points and the exponents are described. Proofs are omitted.

2.1. $Z_{\Delta 8}(A)$. For generic parameters $A = (A_0, \dots, A_3)$ the function $F(t, t, t)$ satisfies an ordinary differential equation $Z_{\Delta 8}(A)$ of order 8 with regular singular points at $-1, -1/2, 1, \infty$ and apparent singular points at -2 and other 8 points. The local exponents are given as

$$t = -1 : \quad 0, 1, \frac{1}{2} \pm A_1, \frac{1}{2} \pm A_2, \frac{1}{2} \pm A_3,$$

$$t = -\frac{1}{2} : \quad 0, 1, 2, 3, A_0, A_0 + 1, A_0 + 2, A_0 + 3,$$

$$t = 1 : \quad 0, 2A_0, A_0 - \frac{1}{2}, A_0 + \frac{1}{2}, A_0 + \frac{3}{2}, A_0 + \frac{5}{2}, A_0 + \frac{7}{2}, A_0 + \frac{9}{2},$$

$$t = \infty : \quad \frac{1}{2}(3 - 3A_0 \pm A_1 \pm A_2 \pm A_3),$$

$$t = -2 : \quad 0, 1, 3, 4, 5, 6, 8, 9,$$

$$t = \text{other 8 points} : \quad 0, 1, 2, 3, 4, 5, 6, 8.$$

2.2. $Z_{\Delta 6}(A)$. If $A_3 = A_2$ then $F(t, t, t)$ satisfies an ordinary differential equation $Z_{\Delta 6}$ of order 6 with regular singular points at $-1, -1/2, 1, \infty$ and apparent singular points at -2 and other 4 points. The local exponents are given as

$$\begin{aligned} t = -1 : & \quad 0, 1, \frac{1}{2} \pm A_1, \frac{1}{2} \pm A_2, \\ t = -\frac{1}{2} : & \quad 0, 1, 2, A_0, A_0 + 1, A_0 + 2, \\ t = 1 : & \quad 0, 2A_0, A_0 - \frac{1}{2}, A_0 + \frac{1}{2}, A_0 + \frac{3}{2}, A_0 + \frac{5}{2}, \\ t = \infty : & \quad \frac{1}{2}(3 - 3A_0 \pm A_1 \pm 2A_2), \frac{1}{2}(3 - 3A_0 \pm A_1), \\ t = -2 : & \quad 0, 1, 3, 4, 5, 6, \\ t = \text{other 4 points} : & \quad 0, 1, 2, 3, 4, 6. \end{aligned}$$

2.3. $Z_{\Delta 4}(A)$. If $A_3 = A_2 = A_1$ then $F(t, t, t)$ satisfies an ordinary differential equation $Z_{\Delta 4}$ of order 4 with regular singular points at $-1, -1/2, 1, \infty$ and only one apparent singular point at -2 . The local exponents are given as

$$\begin{aligned} t = -1 : & \quad 0, 1, \frac{1}{2} \pm A_1, \\ t = -\frac{1}{2} : & \quad 0, 1, A_0, A_0 + 1, \\ t = 1 : & \quad 0, 2A_0, A_0 - \frac{1}{2}, A_0 + \frac{1}{2}, \\ t = \infty : & \quad \frac{1}{2}(3 - 3A_0 \pm 3A_1), \frac{1}{2}(3 - 3A_0 \pm A_1), \\ t = -2 : & \quad 0, 1, 3, 4. \end{aligned}$$

3. Restriction of $Z_3(A)$ onto the plane $t_3 = 1$ and $Z_2(A)$

3.1. Equation $Z_2(A)$. The restriction $Z_3(A)|_{t_3=1}$ of $Z_3(A)$ onto the plane $t_3 = 1$ is, by definition, generated by the operators P , where

$$P(t_1, t_2, \partial_1, \partial_2) + (t_3 - 1)Q, \quad \partial_i := \partial/\partial t_i$$

belongs to $Z_3(A)$ for some operator $Q = Q(t_1, t_2, t_3, \partial_1, \partial_2, \partial_3)$. We find two such operators P_1 and P_2 as follows. Since

$$E_3 = (1 - t_1^2)\partial_{11} + 2(1 - t_1 t_2)\partial_{12} + (1 - t_2^2)\partial_{22} + a_0(t_1 \partial_1 + t_2 \partial_2) + a_3 + 2(t_3 - 1)\partial_{12},$$

we cut off the last term, and define P_1 as

$$P_1 := (1 - t_1^2)\partial_{11} + 2(1 - t_1 t_2)\partial_{12} + (1 - t_2^2)\partial_{22} + a_0(t_1 \partial_1 + t_2 \partial_2) + a_3.$$

We next express E_1 and E_2 as

$$\begin{aligned} E_1 &= G_1 + (t_3 - 1)R_1 + 2(t_1 - t_2)\partial_{23} + a_0\partial_3, \\ E_2 &= G_2 + (t_3 - 1)R_2 + 2(t_2 - t_1)\partial_{13} + a_0\partial_3, \end{aligned}$$

where

$$G_1 = (1 - t_2^2)\partial_{22} + a_0 t_2 \partial_2 + a_1, \quad R_1 = -2t_2 \partial_{23} - (1 + t_3) \partial_{33} + a_0 \partial_3;$$

G_2 and R_2 are given by exchanging 1 and 2 in G_1 and R_1 , respectively. Differentiate these:

$$\begin{aligned} E_{1,1} &= G_{1,1} + (t_3 - 1)R_{1,1} + 2\partial_{23} + 2(t_1 - t_2)\partial_{123} + a_0 \partial_{13}, \\ E_{2,2} &= G_{2,2} + (t_3 - 1)R_{2,2} + 2\partial_{13} + 2(t_2 - t_1)\partial_{123} + a_0 \partial_{23}, \end{aligned}$$

where $E_{1,1} := \partial_1 E_1$, $G_{1,1} := \partial_1 G_1$, etc, for example,

$$G_{1,1} = (1 - t_2^2)\partial_{122} + a_0 t_2 \partial_{12} + a_1 \partial_1.$$

We have

$$\begin{aligned} E_1 - E_2 &\equiv G_1 - G_2 + 2(t_1 - t_2)(\partial_{23} + \partial_{13}), \\ E_{1,1} + E_{2,2} &\equiv G_{1,1} + G_{2,2} + (2 + a_0)(\partial_{23} + \partial_{13}) \end{aligned}$$

modulo $(t_3 - 1)$, and so

$$2(t_1 - t_2)(E_{1,1} + E_{2,2}) - (2 + a_0)(E_1 - E_2) \equiv 2(t_1 - t_2)(G_{1,1} + G_{2,2}) - (2 + a_0)(G_1 - G_2).$$

Now we define the second operator P_2 by the right hand-side of this identity:

$$\begin{aligned} P_2 := & 2(t_1 - t_2)\{(1 - t_2^2)\partial_{122} + a_0 t_2 \partial_{12} + a_1 \partial_1 + (1 - t_1^2)\partial_{112} + a_0 t_1 \partial_{12} + a_2 \partial_2\} \\ & - (2 + a_0)\{(1 - t_2^2)\partial_{22} + a_0 t_2 \partial_2 + a_1 - (1 - t_1^2)\partial_{11} - a_0 t_1 \partial_1 - a_2\}. \end{aligned}$$

Though we have no rigorous proof that P_1 and P_2 generate the ideal $Z_3(a)|_{t_3=1}$, we study the system $Z_2(A)$ in (t_1, t_2) generated by P_1 and P_2 .

Theorem 3.1. *The system $Z_2(A) := \langle P_1, P_2 \rangle$ is of rank 6. The singular locus in $\mathbb{P}^1 \times \mathbb{P}^1$ is given by*

$$t_i = \pm 1, \infty \quad (i = 1, 2), \quad t_1 = t_2.$$

Proposition 3.2. *The local exponents along the divisors above are given as*

$$\begin{aligned} t_1 = \pm 1 : & 0, 1, 2, 3, \frac{1}{2} \pm A_1, \\ t_2 = \pm 1 : & 0, 1, 2, 3, \frac{1}{2} \pm A_2, \\ t_1 = t_2 : & 0, 1, 2A_0, 2A_0 + 1, A_0 \pm A_3, \\ t_i = \infty : & 1 - A_0 \pm A_j, 2 - A_0 \pm A_j, 1 - A_0 \pm A_3 \quad (\{i, j\} = \{1, 2\}). \end{aligned}$$

If we restrict the system $Z_2(A)$ further onto a generic line $t_2 = \text{constant}$, we get an ordinary differential equation of order 6 in $t := t_1$ with polynomial coefficients:

$$(t + 1)^2(t - 1)^2(t - t_2)^4 P(t) \frac{d^6 F}{dt^6} + \dots = 0,$$

where $P(t)$ is of degree 6, the number of apparent singular points, whose local exponents are 0, 1, 2, 3, 4 and 6. Though we omit the explicit expression of the coefficients of the ordinary equation above, we find the local exponents at the singular points as in the proposition.

REMARK 3.3. Any set of six independent solutions defines a map from (t_1, t_2) -space into the five dimensional projective space, whose image is regarded as a surface. We remark that the operator P_1 implies that the second jet-space of the surface is always degenerate; the system $Z_2(A)$ is not general in this sense among those systems of rank 6.

3.2. Outline of the proof of Theorem 3.1. Using unknown F , we rewrite the system in Pfaffian form relative to a frame

$$e_6 = {}^{\text{tr}}(F, F_1, F_2, (t_1 - t_2)F_{11}, (t_1 - t_2)F_{12}, (t_1 - t_2)^2 F_{112}).$$

This time, by using $P_1 = 0$ and $P_2 = 0$, and their higher-order derivatives, we can see that the derivatives $F_{ij,k}$, $i, j, k = 1, 2$, can be written in terms of $F, F_1, F_2, F_{11}, F_{12}$ and F_{112} . Thus, we get a Pfaffian form ω_6 such that $de_6 = \omega_6 e_6$. It is a straightforward computation to see that the integrability condition $d\omega_6 = \omega_6 \wedge \omega_6$ holds. The 6×6 -matrix 1-form ω_6 is listed in §8.2.

3.3. Restriction of $Z_2(A)$ onto the diagonal $t_1 = t_2$. Change the coordinates from (t_1, t_2) to (t, s) by $t_1 = t, t_2 = t + s$. Then the operator P_1 becomes

$$\partial_{11} - t^2 \partial_{11} + a_0 t \partial_1 + a_3 + s \{-s \partial_{22} - 2t(\partial_{12} - \partial_{22}) + a_0 \partial_2 - 2t \partial_{22}\}.$$

Thus the restriction of $Z_2(A)$ onto the diagonal $s = 0$ is the ordinary differential equation

$$(1 - t^2)F_{11} + a_0 t F_1 + a_3 F = 0, \quad F_1 = dF/dt.$$

The local exponents at $t = -1, 1$ and ∞ are

$$0, A_0 - \frac{1}{2}; \quad 0, A_0 - \frac{1}{2} \quad \text{and} \quad 1 - A_0 \pm A_3,$$

respectively.

4. Restriction of $Z_2(A)$ onto the line $t_2 = 1$ and $Z(A)$

4.1. Equation $Z(A)$.

Express P_1 and P_2 as

$$\begin{aligned} P_1 &\equiv Q_1 + 2(1 - t_1)\partial_{12} + a_0 \partial_2, \\ P_2 &\equiv Q_2 + 2(t_1 - 1)\{a_0 \partial_{12} + (1 - t_1^2)\partial_{112} + a_0 t_1 \partial_{12} + a_2 \partial_2\} - (2 + a_0)a_0 \partial_2 \\ &= Q_2 + 2a_0(t_1^2 - 1)\partial_{12} - 2(t_1^2 - 1)(t_1 - 1)\partial_{112} + \{2a_2(t_1 - 1) - (2 + a_0)a_0\}\partial_2 \end{aligned}$$

mod $(1 - t_2)$, where

$$\begin{aligned} Q_1 &= (1 - t_1^2)\partial_{11} + a_0 t_1 \partial_1 + a_3, \\ Q_2 &= 2(t_1 - 1)a_1 \partial_1 - (2 + a_0)\{a_1 - (1 - t_1^2)\partial_{11} - a_0 t_1 \partial_1 - a_2\}. \end{aligned}$$

Differentiate P_1 , and we have

$$\begin{aligned} P_{1,1} := \partial_1 P_1 &= Q_{1,1} - 2\partial_{12} - 2(t_1 - 1)\partial_{112} + a_0 \partial_{12} \\ &= Q_{1,1} + (a_0 - 2)\partial_{12} - 2(t_1 - 1)\partial_{112}, \end{aligned}$$

where $Q_{1,1} := \partial_1 Q_1, \partial_{112} := \partial_1 \partial_{12}$. Set

$$P_3 := P_2 - (t_1^2 - 1)P_{1,1} = Q_2 - (t_1^2 - 1)Q_{1,1} + (t_1^2 - 1)(a_0 + 2)\partial_{12} + \{2a_2(t_1 - 1) - (2 + a_0)a_0\}\partial_2,$$

and differentiate:

$$\begin{aligned} P_{3,1} &= Q_{2,1} - 2t_1 Q_{1,1} - (t_1^2 - 1)Q_{1,1,1} \\ &\quad + 2t_1(a_0 + 2)\partial_{12} + (t_1^2 - 1)(a_0 + 2)\partial_{112} + 2a_2 \partial_2 + \{2a_2(t_1 - 1) - (2 + a_0)a_0\}\partial_{12}. \end{aligned}$$

By using P_1, P_3 and $P_{1,1}$, express $\partial_2, \partial_{12}$ and ∂_{112} in terms of $Q_1, Q_{1,1}$. Substitute these expressions into $P_{3,1}$, and we get an ordinary differential operator $Z(a)$ of order four.

Proposition 4.1. *The equation $Z(a)$ is irreducible.*

Proof. If $Z(a)$ factors as Z_1Z_2 , then the local exponents of Z_2 at $t = \pm 1$ and ∞ are subsets of those of $Z(a)$. Riemann relation says that the sum of the local exponents of Z_2 is an integer. The Riemann scheme of $Z(a)$ below shows that this can not happen if Z_2 is of order 1 or 3. Assume that the order of Z_2 is 2, and let k be the number of apparent singular points. Then Riemann relation says that the sum of the local exponents of Z_2 is equal to $1 + k$. On the other hand the Riemann scheme shows that the sum is greater than $3 + k$. \square

This assures that $Z(a)$ is the restriction of $Z_2(a)$ onto $t_2 = 1$.

Theorem 4.2. *The restriction $Z(a)$ of $Z_2(a)$ onto the line $t_2 = 1$ is given by*

$$Z(a) := p_0\partial^4 + p_1\partial^3 + p_2\partial^2 + p_3\partial + p_4,$$

where $\partial = d/dt$, $t = t_1$, and

$$\begin{aligned} p_0 &= 2(t+1)^2(t-1)^3, \\ p_1 &= -4(t+1)(t-1)^2\{(2+a_0)+(a_0-2)t\}, \\ p_2 &= 2(t-1)\{(a_0^2-2a_1+6a_0+2+a_2+a_3)+(3a_0^2+4a_0-4+2a_1)t \\ &\quad +(a_0^2-4a_0+2-a_2-a_3)t^2\}, \\ p_3 &= (-4a_0^2-8a_0+4a_0a_1+4a_1-(2a_0+4)(a_2+a_3)) \\ &\quad +(-2a_0^3-6a_0^2-4a_0a_1-4a_1+4(a_2+a_3))t+2a_0(a_0+a_2+a_3)t^2, \\ p_4 &= 2a_2a_3t+(a_1-a_2-a_3)(a_0+2)^2-2a_2a_3. \end{aligned}$$

This equation has one accessory parameter; the local exponents do not change if we add a constant to p_4 .

We denote the operator $Z(a)$ with parameters (A_0, A_1, A_2, A_3) by $Z(A)$, the explicit form of which will be given in §10. The Riemann scheme of $Z(A)$ is given as

$$\left\{ \begin{array}{ccc} t = 1 & t = -1 & t = \infty \\ 0 & \frac{1}{2} - A_1 & 1 - A_0 + A_2 \\ A_0 - \frac{1}{2} & 0 & 1 - A_0 - A_2 \\ A_0 + \frac{1}{2} & 1 & 1 - A_0 + A_3 \\ 2A_0 & \frac{1}{2} + A_1 & 1 - A_0 - A_3 \end{array} \right\}.$$

REMARK 4.3 (SYMMETRY). $Z(A)$ is invariant under

$$A_j \rightarrow -A_j \quad (j = 1, 2, 3) \quad \text{and} \quad A_2 \leftrightarrow A_3.$$

REMARK 4.4. The Dotsenko-Fateev equation (§6.1) appears as the restriction on a divisor of the Appell's equation F_4 in two variables (see [5]).

4.2. A small change $\tilde{Z}(A)$ of $Z(A)$. To increase symmetry, we introduce an operator $\tilde{Z}(A)$ as¹

$$\tilde{Z}(A) := \text{Ad}((t-1)^{-A_0+\frac{1}{2}})Z(A) = (t-1)^{-A_0+\frac{1}{2}} \circ Z(A) \circ (t-1)^{A_0-\frac{1}{2}}.$$

¹Equivalent to changing the unknown z of the equation $Z(A)$ to a new unknown w by $z = (t-1)^{A_0-\frac{1}{2}}w$

(Ad stands for addition which will be recalled in §5.1.)² We further change the variable t , used for $Z(A)$ and $\tilde{Z}(A)$ etc, into the new variable

$$x = \frac{1-t}{2}.$$

In x -coordinate, the differential operator $\tilde{Z}(A)$ changes into³

$$(1) \quad \tilde{Z}(A) = x^2(x-1)^2\partial^4 + m_1(x)\partial^3 + m_2(x)\partial^2 + m_3(x)\partial + m_4(x), \quad \partial := d/dx,$$

where

$$\begin{aligned} m_1 &= 4(x-1)x(2x-1), \\ m_2 &= \frac{1}{4} \left(4A_0^2x - 4A_0^2 - 4A_1^2x - 4A_2^2x^2 + 4A_2^2x - 4A_3^2x^2 + 4A_3^2x + 58x^2 - 58x + 9 \right), \\ m_3 &= \frac{1}{2} \left(2A_0^2 - 2A_1^2 - 4A_2^2x + 2A_2^2 - 4A_3^2x + 2A_3^2 + 10x - 5 \right), \\ m_4 &= \left(A_2 - \frac{1}{2} \right) \left(A_2 + \frac{1}{2} \right) \left(A_3 - \frac{1}{2} \right) \left(A_3 + \frac{1}{2} \right). \end{aligned}$$

The local exponents do not change if we add a constant to m_3 ; the constant term of m_3 is called the *accessory parameter*.

REMARK 4.5 (SYMMETRY). $\tilde{Z}(A)$ is invariant under

$$A_j \rightarrow -A_j \quad (j = 0, 1, 2, 3) \quad \text{and} \quad A_2 \longleftrightarrow A_3$$

and

$$(x, A_0, A_1) \longleftrightarrow (1-x, A_1, A_0).$$

The Riemann scheme of $\tilde{Z}(A)$ is given as

$$\left\{ \begin{array}{ccc} x = 0 & x = 1 & x = \infty \\ \frac{1}{2} - A_0 & \frac{1}{2} - A_1 & \frac{1}{2} + A_2 \\ 0 & 0 & \frac{1}{2} - A_2 \\ 1 & 1 & \frac{1}{2} + A_3 \\ \frac{1}{2} + A_0 & \frac{1}{2} + A_1 & \frac{1}{2} - A_3 \end{array} \right\}.$$

4.3. Invariants of ordinary differential operators. For a differential operator $L = p_0\partial^4 + p_1\partial^3 + p_2\partial^2 + p_3\partial + p_4$, the operator

$$L^* = \partial^4 \circ p_0 - \partial^3 \circ p_1 + \partial^2 \circ p_2 - \partial \circ p_3 + p_4, \quad \partial = d/dx$$

is called the *adjoint operator*.

Proposition 4.6. *The equation $\tilde{Z}(A)$ is self-adjoint.*

To explain the meaning of this proposition, we recall some differential invariants of ordinary differential operators. An ordinary differential operator

$$\partial^4 + Q_1\partial^3 + Q_2\partial^2 + Q_3\partial + Q_4$$

²Strictly speaking, $\tilde{Z}(A) = \frac{1}{2}(t-1)^{-1}\text{Ad}((t-1)^{-A_0+\frac{1}{2}})Z(A)$.

³The equations $Z(A)$ and $\tilde{Z}(A)$ rewritten in the new variable x will be denoted by the same notation.

is transformed into the operator of the form

$$(2) \quad \partial^4 + q_2\partial^2 + q_3\partial + q_4$$

which has no third-order term, by multiplying a non-zero function to the dependent variable. The coefficients q_i are given as

$$\begin{aligned} q_2 &= Q_2 - \frac{3}{2}Q'_1 - \frac{3}{8}Q_1^2, \\ q_3 &= Q_3 - \frac{1}{2}Q_1Q_2 + \frac{1}{8}Q_1^3 - Q''_1, \\ q_4 &= Q_4 - \frac{1}{4}Q_1Q_3 + \frac{1}{16}Q_1^2Q_2 - \frac{3}{256}Q_1^4 - \frac{1}{4}Q_2Q'_1 + \frac{3}{32}Q_1^2Q'_1 + \frac{3}{16}(Q'_1)^2 - \frac{1}{4}Q'''_1. \end{aligned}$$

It is known ([7]) that, for an appropriate choice of the dependent variable and the coordinate $y = y(x)$, the operator (2) can be transformed further into an operator

$$(3) \quad \partial^4 + r_3\partial + r_4, \quad \partial = d/dy.$$

Though r_3 and r_4 are not unique, the forms

$$\theta_3 := r_3 dy^{\otimes 3} = (q_3 - q'_2) dx^{\otimes 3}, \quad \theta_4 := \left(r_4 - \frac{1}{2}r'_3 \right) dy^{\otimes 4} = \left(q_4 - \frac{1}{2}q'_3 - \frac{9}{100}q_2^2 + \frac{1}{5}q''_2 \right) dt^{\otimes 4}$$

are unique and are called the fundamental invariants of the operator (2).

By an easy calculation, we see that the adjoint operator of (2) is

$$\partial^4 + q_2\partial^2 + (2q'_2 - q_3)\partial + q_4 + q''_2 - q'_3.$$

Hence, we have:

Lemma 4.7. *The operator (2) is self-adjoint if and only if $\theta_3 = 0$.*

REMARK 4.8. The property that $\theta_3 \equiv 0$ is rephrased geometrically as follows: Let z_1, \dots, z_4 be linearly independent solutions of the equation and let us consider $z = [z_1, \dots, z_4]$ as a curve in the projective space \mathbf{P}^3 . Then, we can see that, when $\theta_3 \equiv 0$, the curve formed by the tangent vectors to this curve z , which lies in the 5-dimensional projective space of all lines in \mathbf{P}^3 , is degenerate in the sense that it lives in a 4-dimensional hyperplane.

5. $\tilde{Z}(A)$ is related to the tensor product of two Gauss equations

In §10.4, we study local solutions of $\tilde{Z}(A)$ at $x = 0$ and find that they are closely related to the product of two specific Gauss hypergeometric series. In this section we show that an addition and a middle convolution connect $\tilde{Z}(A)$ with the tensor product of the two Gauss equations. We begin with introducing two important operations for differential operators.

Detailed study of the tensor product of two Gauss equations in general is made in the last section of Part 1.

5.1. Definition and fundamental properties of addition and middle convolution. For a differential operator P in x and a function f in x , the **addition** by f is defined as

$$\text{Ad}(f)P := f \circ P \circ f^{-1},$$

which is already appeared in §4.2; multiplying a non-zero function f to the dependent variable to get a new one.

For a differential operator P in x and a complex number μ , the **middle convolution** $mc_\mu P$ with parameter μ is defined symbolically (cf. Definition 2.3 in [6]) as

$$mc_\mu P := \partial^{-\mu} \circ P \circ \partial^\mu, \quad \partial = \frac{d}{dx}.$$

Actual procedure is as follows: Write the operator P in the form

$$\sum p_{ij}x^i\partial^j.$$

Set $r = \max\{i - j, p_{ij} \neq 0\}$, and multiply ∂^r to P from the left, then substitute

$$x^i\partial^i = \theta(\theta - 1)\cdots(\theta - i + 1), \quad \theta = x\partial$$

to express $\partial^r P$ as a linear combination of $\{\theta^i\partial^j\}$. Then replace θ by $\theta - \mu$, and finally divide the operator by ∂ from the left as many times as possible to obtain $mc_\mu P$. Fundamental properties:

$$mc_{\mu+\mu'} = mc_\mu \circ mc_{\mu'}, \quad mc_{-\mu} = mc_\mu^{-1}, \quad mc_\mu\theta = \theta - \mu, \quad mc_\mu\partial = \partial.$$

5.2. A middle convolution connects $\tilde{Z}(A)$ with the tensor product of two Gauss equations.

Recall the Gauss equation:

$$E\begin{pmatrix} a, b \\ c \end{pmatrix} = x(x-1)\partial^2 + ((a+b+1)x - c)\partial + ab.$$

Let us consider the tensor product

$$K = K(A) = K(A_{-++}, A_{-+-}, 1 - A_0, A_{---}, A_{--+}, 1 - A_0)$$

of the two Gauss equations

$$E\begin{pmatrix} A_{-++}, A_{-+-} \\ 1 - A_0 \end{pmatrix} \quad \text{and} \quad E\begin{pmatrix} A_{---}, A_{--+} \\ 1 - A_0 \end{pmatrix},$$

where

$$A_{\varepsilon_0, \varepsilon_1, \varepsilon_2, \varepsilon_3} := \frac{\varepsilon_0 A_0 + \varepsilon_1 A_1 + \varepsilon_2 A_2 + \varepsilon_3 A_3 + 1}{2} \quad \varepsilon_j = \pm.$$

It is, by definition, the differential equation satisfied by the *product of the solutions* of the two Gauss equations⁴, and is given as follows (see §9):

$$K(A) = x^3(x-1)^2\partial^4 + k_1(x)\partial^3 + k_2(x)\partial^2 + k_3(x)\partial + k_4(x), \quad \partial = d/dx,$$

where

$$\begin{aligned} k_1 &= (1-x)x^2(4A_0x - 4A_0 - 10x + 5), \\ k_2 &= x(6A_0^2x^2 - 11A_0^2x + 5A_0^2 - 24A_0x^2 + 33A_0x - 9A_0 \\ &\quad - A_1^2x - A_2^2x^2 + A_2^2x - A_3^2x^2 + A_3^2x + 25x^2 - 25x + 4), \end{aligned}$$

⁴Do not confuse this and the *product of two operators*

$$\begin{aligned}
k_3 &= \frac{1}{2}(-8A_0^3x^2 + 12A_0^3x - 4A_0^3 + 36A_0^2x^2 - 39A_0^2x + 6A_0^2 \\
&\quad + 4A_0A_1^2x + 4A_0A_2^2x^2 - 4A_0A_2^2x + 4A_0A_3^2x^2 - 4A_0A_3^2x - 56A_0x^2 \\
&\quad + 42A_0x - 2A_0 - 3A_1^2x - 6A_2^2x^2 + 3A_2^2x - 6A_3^2x^2 + 3A_3^2x + 30x^2 - 15x), \\
k_4 &= \frac{1}{2}(2A_0^4x - 2A_0^4 - 8A_0^3x + 5A_0^3 - 2A_0^2A_1^2 - 2A_0^2A_2^2x \\
&\quad + 2A_0^2A_2^2 - 2A_0^2A_3^2x + 2A_0^2A_3^2 + 12A_0^2x - 4A_0^2 + A_0A_1^2 + 4A_0A_2^2x \\
&\quad - A_0A_2^2 + 4A_0A_3^2x - A_0A_3^2 - 8A_0x + A_0 + 2A_2^2A_3^2x - 2A_2^2x - 2A_3^2x + 2x).
\end{aligned}$$

To increase symmetry, we make a change (addition) as:

$$L(A) = \text{Ad}(x^{-A_0})K(A) = x^{-A_0} \circ K(A) \circ x^{A_0},$$

which can be expressed as

$$L(A) = x^2(x-1)^2\partial^4 + \ell_1(x)\partial^3 + \ell_2(x)\partial^2 + \ell_3(x)\partial + \ell_4(x), \quad \partial = d/dx$$

where

$$\begin{aligned}
\ell_1 &= 5(x-1)x(2x-1), \\
\ell_2 &= A_0^2x - A_0^2 - A_1^2x - A_2^2x^2 + A_2^2x - A_3^2x^2 + A_3^2x + 25x^2 - 25x + 4, \\
\ell_3 &= -\frac{3}{2}(-A_0^2 + A_1^2 + 2A_2^2x - A_2^2 + 2A_3^2x - A_3^2 - 10x + 5), \\
\ell_4 &= (A_2 - 1)(A_2 + 1)(A_3 - 1)(A_3 + 1).
\end{aligned}$$

The constant term ℓ_{30} of $\ell_3(x) = \ell_{31}x + \ell_{30}$ is the accessory parameter. The Riemann scheme of $L(A)$ is given as

$$\begin{aligned}
&x^{-A_0} \left\{ \begin{array}{ccc} x=0 & x=1 & x=\infty \\ 0 & 0 & A_{-++} \\ A_0 & -A_1 & A_{-++} \end{array} \right\} \times \left\{ \begin{array}{ccc} x=0 & x=1 & x=\infty \\ 0 & 0 & A_{----} \\ A_0 & A_1 & A_{-++} \end{array} \right\} \\
&= x^{-A_0} \left\{ \begin{array}{ccc} x=0 & x=1 & x=\infty \\ 0 & 0 & -A_0 - A_2 + 1 \\ A_0 & -A_1 & -A_0 + A_3 + 1 \\ A_0 + 1 & A_1 & -A_0 - A_3 + 1 \\ 2A_0 & 1 & -A_0 + A_2 + 1 \end{array} \right\} = \left\{ \begin{array}{ccc} x=0 & x=1 & x=\infty \\ -A_0 & -A_1 & 1 + A_2 \\ 0 & 0 & 1 - A_3 \\ 1 & 1 & 1 + A_3 \\ A_0 & A_1 & 1 - A_2 \end{array} \right\}.
\end{aligned}$$

REMARK 5.1 (SYMMETRY). L is invariant under

$$A_j \rightarrow -A_j \quad (j = 0, 1, 2, 3) \quad \text{and} \quad A_2 \longleftrightarrow A_3$$

and

$$(x, A_0, A_1) \longleftrightarrow (1-x, A_1, A_0).$$

This operator L is connected with the operator $\tilde{Z}(A)$, introduced in §4.2, by the middle convolution as follows.

Theorem 5.2. $mc_{-\frac{1}{2}}\tilde{Z}(A) = L(A)$.

We first express the operator $\tilde{Z}(A)$ defined in §4.2 (1) as a polynomial in $\theta = x\partial_x$ and

$\partial = \partial_x$:

$$\begin{aligned}\tilde{Z}(A) &= (\theta + \frac{1}{2} - A_2)(\theta + \frac{1}{2} + A_2)(\theta + \frac{1}{2} - A_3)(\theta + \frac{1}{2} + A_3) \\ &\quad - (\theta + 1)(2\theta^2 + 4\theta + A_1^2 - A_0^2 - A_2^2 - A_3^2 + \frac{5}{2})\partial \\ &\quad + (\theta + \frac{3}{2} + A_0)(\theta + \frac{3}{2} - A_0)\partial^2.\end{aligned}$$

Then, replacing θ by $\theta + 1/2$, we have

$$\begin{aligned}mc_{-1/2}\tilde{Z}(A) &= (\theta + 1 - A_2)(\theta + 1 + A_2)(\theta + 1 - A_3)(\theta + 1 + A_3) \\ &\quad - (\theta + \frac{3}{2})(2\theta^2 + 6\theta + A_1^2 - A_0^2 - A_2^2 - A_3^2 + 5)\partial \\ &\quad + (\theta + 2 + A_0)(\theta + 2 - A_0)\partial^2,\end{aligned}$$

which turns out to be $L(A)$. This expression leads to

Corollary 5.3. *Assume A_2 is a positive integer: $A_2 = m + 1$ for $m \geq 0$, (or a negative integer $A_2 = -m - 1$, $m \geq 0$; namely, A_2 is a non-zero integer). Then, the equation $L(A)u = 0$ has a polynomial solution of degree m .*

Proof. For an integer k , we see that

$$L(A)x^k = p_k x^k + q_{k-1} x^{k-1} + r_{k-2} x^{k-2},$$

where

$$\begin{aligned}p_k &= ((k+1)^2 - A_2^2)((k+1)^2 - A_3^2), \\ q_{k-1} &= -k(2k+1)(k^2 + k - 2 + \alpha), \quad \alpha = (A_1^2 - A_0^2 - A_2^2 - A_3^2 + 5)/2, \\ r_{k-2} &= k(k-1)(k^2 - A_0^2),\end{aligned}$$

and $q_1 = r_2 = r_1 = 0$. This implies that the operator $L(A)$ sends the space of polynomials of degree smaller than or equal to m , into itself. Since $p_m = 0$ by assumption, the image is a proper subspace; hence, there exists a polynomial solution. \square

Let $u = \sum_{k=0}^m a_k x^k$ be such a solution. Then, we see

$$\begin{aligned}a_{m-1}p_{m-1} + a_m q_{m-1} &= 0, \\ a_{m-2}p_{m-2} + a_{m-1}q_{m-2} + a_m r_{m-2} &= 0, \\ &\dots, \\ a_k p_k + a_{k+1} q_k + a_{k+2} r_k &= 0, \\ &\dots, \\ a_0 p_0 + a_1 q_0 + a_2 r_0 &= 0.\end{aligned}$$

If $A_3 \notin \mathbf{Z}$, for a given value a_m , the other a_k are uniquely determined as follows. Let M be a lower triple-triangular matrix defined as

$$M = \begin{pmatrix} p_{m-1} & & & & & \\ q_{m-2} & p_{m-2} & & & & \\ r_{m-3} & q_{m-3} & p_{m-3} & & & \\ & r_{m-4} & q_{m-4} & p_{m-4} & & \\ & & \ddots & \ddots & \ddots & \\ & & & r_0 & q_0 & p_0 \end{pmatrix},$$

and $a = {}^{\text{tr}}(a_{m-1}, a_{m-2}, \dots, a_0)$ be a column vector. Then, the linear equations above are written as

$$Ma = {}^{\text{tr}}(-a_m q_{m-1}, -a_m r_{m-2}, 0, \dots, 0).$$

Hence, multiplying the inverse of M yields the solution a ; note that, since $A_3 \notin \mathbf{Z}$, the determinant of M is non-vanishing. First few terms of a are given as

$$a_m := 1, \quad a_{m-1} = \frac{-m(m^2 + m - 2 + \alpha)}{m^2 - A_3^2},$$

$$a_{m-2} = \frac{m(m-1) \left\{ (2m-1)(m^2 + m - 2\alpha)((m-1)^2 + m - 3 + \alpha) - (m^2 - A_0^2)(m^2 - A_3^2) \right\}}{2(2m+1)((m-1)^2 - A_3^2)(m^2 - A_3^2)}.$$

6. Relation between $\tilde{Z}(A)$ and the Dotsenko-Fateev equation

6.1. The Dotsenko-Fateev equation. The Dotsenko-Fateev operator ([2]) is an operator of order 3 defined as

$$S = S(a, b, c, g) = x^2(x-1)^2\partial^3 + s_1\partial^2 + s_2\partial + s_3, \quad \partial := d/dx$$

where

$$s_1 = -(-1+x)x(3ax + 3bx + 6cx + 2gx - 3a - 3c - g),$$

$$\begin{aligned} s_2 = & 2a^2x^2 + 4abx^2 + 12acx^2 + 3agx^2 + 2b^2x^2 + 12bcx^2 + 3bgx^2 + 12c^2x^2 + 8cgx^2 \\ & + g^2x^2 - 4a^2x - 4abx - 16acx - 4agx + ax^2 - 8bcx - 2bgx + bx^2 - 12c^2x - 8cgx \\ & + 6cx^2 - g^2x + gx^2 + 2a^2 + 4ac + ag - 2ax + 2c^2 + cg - 6cx - gx + a + c, \end{aligned}$$

$$s_3 = -c(2a + 2 + 2b + 2c + g)(2ax + 2bx + 4cx + 2gx - 2a - 2c - g + 2x - 1).$$

The constant term of s_3 is the accessory parameter. The Riemann scheme is

$$\left\{ \begin{array}{ccc} x = 0 & x = 1 & x = \infty \\ 0 & 0 & -2c \\ a + c + 1 & b + c + 1 & -a - b - 2c - g - 1 \\ 2a + 2c + g + 2 & 2b + 2c + g + 2 & -2a - 2b - 2c - g - 2 \end{array} \right\}.$$

REMARK 6.1 (SYMMETRY). The adjoint operator of $S(a, b, c, g)$ is given by S with the change:

$$(a, b, c, g) \rightarrow (-1 - a, -1 - b, -1 - c, -g).$$

6.2. A middle convolution and an addition send $\tilde{Z}(A)$ to the Dotsenko-Fateev equation. The equation $\tilde{Z}(A) = x^2(x-1)^2\partial^4 + \dots$ has the Riemann scheme

$$\left\{ \begin{array}{ccc} x=0 & x=1 & x=\infty \\ 0 & 0 & \frac{1}{2} - A_2 \\ 1 & 1 & \frac{1}{2} + A_2 \\ \frac{1}{2} - A_0 & \frac{1}{2} - A_1 & \frac{1}{2} - A_3 \\ \frac{1}{2} + A_0 & \frac{1}{2} + A_1 & \frac{1}{2} + A_3 \end{array} \right\}.$$

Proposition 6.2. *A middle convolution with parameter $-\frac{1}{2} - A_2$ sends $\tilde{Z}(A)$ to the equation defined by*⁵

$$Q(A) = mc_{-\frac{1}{2} - A_2}(\tilde{Z}(A)) = x^2(x-1)^2\partial^3 + q_1(x)\partial^2 + q_2(x)\partial + q_3(x)$$

of order 3, where

$$\begin{aligned} q_1 &= (2A_2 + 3)(x-1)x(2x-1), \\ q_2 &= A_0^2x - A_0^2 - A_1^2x + 5A_2^2x^2 - 5A_2^2x + A_2^2 + 12A_2x^2 - 12A_2x + 2A_2 \\ &\quad - A_3^2x^2 + A_3^2x + 7x^2 - 7x + 1, \\ q_3 &= \frac{1}{2}(2A_2 + 1)(A_0^2 - A_1^2 + 2A_2^2x - A_2^2 + 4A_2x - 2A_2 - 2A_3^2x + A_3^2 + 2x - 1). \end{aligned}$$

The Riemann scheme of $Q(A)$ is

$$\left\{ \begin{array}{ccc} x=0 & x=1 & x=\infty \\ 0 & 0 & 1 + 2A_2 \\ -A_0 - A_2 & -A_1 - A_2 & 1 + A_2 - A_3 \\ A_0 - A_2 & A_1 - A_2 & 1 + A_2 + A_3 \end{array} \right\}.$$

The coefficients are determined by the local exponents except q_{30} , where $q_3(x) = q_{31}x + q_{30}$, which is the accessory parameter.

REMARK 6.3 (SYMMETRY). $Q(A)$ is invariant under

$$A_j \rightarrow -A_j \quad (j = 0, 1, 3) \quad \text{and} \quad (x, A_0, A_1) \longleftrightarrow (1-x, A_1, A_0).$$

Moreover, the change $(x, A_0, A_1, A_2, A_3) \rightarrow (\frac{1}{x}, A_3, A_1, A_2, A_0)$ takes $Q(A)$ into $\text{Ad}(x^{1+2A_2})Q(A)$.

REMARK 6.4. The symmetry of $Q(A)$ under $x \rightarrow 1/x$ and its Riemann scheme determine the accessory parameter as above, and so characterize the equation $Q(A)$.

Proposition 6.5. *The operator $R(A)$ defined by*

$$R(A) := \text{Ad}(x^{A_0+A_2}(x-1)^{A_1+A_2})Q(A)$$

⁵It is known (cf. [6]) that if we choose the parameter μ of a middle convolution as

$$\mu + 1 = \text{one of the local exponents of } \tilde{Z}(A) \text{ at } \infty,$$

say, $\mu = \frac{1}{2} - A_2 - 1 = -\frac{1}{2} - A_2$, the resulting equation is of order 3; for generic parameter it is of order 4.

has an expression:

$$R(A) = R(A_0, A_1, A_2, A_3) = x^2(x-1)^2\partial^3 + r_1(x)\partial^2 + r_2(x)\partial + r_3(x),$$

where

$$\begin{aligned} r_1 &= (1-x)x(3A_0x - 3A_0 + 3A_1x + 2A_2x - A_2 - 6x + 3), \\ r_2 &= 3A_0^2x^2 - 5A_0^2x + 2A_0^2 + 6A_0A_1x^2 - 6A_0A_1x + 4A_0A_2x^2 \\ &\quad - 6A_0A_2x + 2A_0A_2 - 9A_0x^2 + 12A_0x - 3A_0 + 3A_1^2x^2 - A_1^2x + 4A_1A_2x^2 \\ &\quad - 2A_1A_2x - 9A_1x^2 + 6A_1x + A_2^2x^2 - A_2^2x - 6A_2x^2 + 6A_2x - A_2 - A_3^2x^2 \\ &\quad + A_3^2x + 7x^2 - 7x + 1, \\ r_3 &= -\frac{1}{2}(2A_0x - 2A_0 + 2A_1x - 2x + 1)(A_0 + A_1 + A_2 - A_3 - 1)(A_0 + A_1 + A_2 + A_3 - 1). \end{aligned}$$

The Riemann scheme of $R(A)$ is

$$\left\{ \begin{array}{lll} x=0 & x=1 & x=\infty \\ 0 & 0 & 1-A_0-A_1 \\ 2A_0 & 2A_1 & 1-A_0-A_1-A_2-A_3 \\ A_0+A_2 & A_1+A_2 & 1-A_0-A_1-A_2+A_3 \end{array} \right\},$$

and the constant term of r_3 is the accessory parameter.

Proposition 6.6. *Change the parameters $\{A_0, A_1, A_2, A_3\}$ to $\{a, b, c, g\}$ by*

$$A_0 = \frac{2a+2c+g+2}{2}, \quad A_1 = \frac{2b+2c+g+2}{2}, \quad A_2 = \frac{-g}{2}, \quad A_3 = \frac{2a+2b+g+2}{2}.$$

Then $R(A_0, A_1, A_2, A_3)$ exactly coincides with the Dotsenko-Fateev equation $S(a, b, c, g)$.

REMARK 6.7 (SYMMETRY). $R(A)$ is invariant only under

$$A_j \rightarrow -A_j \quad (j=3) \quad \text{and} \quad (x, A_0, A_1) \longleftrightarrow (1-x, A_1, A_0).$$

7. Table of related differential equations

Though the equations $Z(A)$, $K(A)$ and $R(A)$ have origin in the Zagier system $Z_3(A)$, the hypergeometric equations and the Dotsenko-Fateev equation, respectively, the equations $\tilde{Z}(A)$, $L(A)$ and $Q(A)$ are more accessible. They are related as in the table below:

$$\begin{array}{ccc} Z_3(A) & & E_1, E_2 \\ \text{rest } (t_3 = 1) & \downarrow & \downarrow \otimes \\ Z_2(A) & & K(A) = E_1 \otimes E_2 \\ \text{rest } (t_2 = 1) & \downarrow & \downarrow \text{Ad}_1 \\ Z(A) & & L(A) \end{array}$$

$$\begin{array}{ccccc}
\text{Ad}_2 & \downarrow & & \downarrow & mc_{\frac{1}{2}} \\
\tilde{Z}(A) & \sim & \tilde{Z}(A) & \text{in } x = \frac{1-t}{2} \\
& & \downarrow & & mc_{-\frac{1}{2}-A_2} \\
& & Q(A) & & \\
& & \downarrow & & \text{Ad}_3 \\
& & R(A) & = & S(a, b, c, g) : DF
\end{array}$$

Here E_1 and E_2 are Gauss hypergeometric equations:

$$E_1 = E\left(\begin{matrix} A_{-++}, A_{-++} \\ 1 - A_0 \end{matrix}; x\right), \quad E_2 = E\left(\begin{matrix} A_{---}, A_{---} \\ 1 - A_0 \end{matrix}; x\right).$$

The additions Ad_j are given as

$$\text{Ad}_1 = \text{Ad}(x^{-A_0}), \quad \text{Ad}_2 = \text{Ad}((t-1)^{\frac{1}{2}-A_0}), \quad \text{Ad}_3 = \text{Ad}(x^{A_0+A_2}(x-1)^{A_1+A_2})$$

are used just for cosmetic changes. Since a middle convolution is additive and invertible, from

$$mc_{-\frac{1}{2}-A_2}(\tilde{Z}) = Q, \quad \text{Ad}_3(Q) = R, \quad \text{Ad}_1(K) = L, \quad mc_{\frac{1}{2}}(L) = \tilde{Z},$$

we have

$$Q = mc_{-\frac{1}{2}-A_2}(\tilde{Z}) = \left(mc_{-\frac{1}{2}-A_2} \circ mc_{\frac{1}{2}}\right)(L) = mc_{-A_2}(L),$$

and

$$R = \text{Ad}_3(Q) = (\text{Ad}_3 \circ mc_{-A_2})(L) = (\text{Ad}_3 \circ mc_{-A_2} \circ \text{Ad}_1)(K),$$

and conversely,

$$K = \left(\text{Ad}_1^{-1} \circ mc_{A_2} \circ \text{Ad}_3^{-1}\right)(R).$$

Relation of the system of parameters: a_* , A_* , $A_{\pm\pm\pm\pm}$ and (a, b, c, g) :

$$a_0 = 2A_0 - 3, \quad a_i = A_i^2 - (A_0 - 1)^2, \quad i = 1, 2, 3,$$

$$A_{\varepsilon_0, \varepsilon_1, \varepsilon_2, \varepsilon_3} := (\varepsilon_0 A_0 + \varepsilon_1 A_1 + \varepsilon_2 A_2 + \varepsilon_3 A_3 + 1)/2, \quad \varepsilon_j = \pm,$$

$$a = \frac{A_{-++} - 1}{2}, \quad b = \frac{A_{-++} - 1}{2}, \quad c = \frac{A_{-++} - 1}{2}, \quad g = -2A_2,$$

$$A_0 = \frac{2a + 2c + g + 2}{2}, \quad A_1 = \frac{2b + 2c + g + 2}{2}, \quad A_2 = \frac{-g}{2}, \quad A_3 = \frac{2a + 2b + g + 2}{2}.$$

8. Explicit expressions of matrix 1-forms

8.1. 8×8 -matrix form $\omega = M_1 dt_1 + M_2 dt_2 + M_3 dt_3$. In §1.1 the system $Z_3(a)$ is transformed into the Pfaffian form $de = \omega e$ with the frame

$$e = {}^{\text{tr}}(F, F_1, F_2, F_3, F_{12}, F_{13}, F_{23}, DF_{123}), \quad D = -1 + t_1^2 + t_2^2 + t_3^2 - 2t_1 t_2 t_3.$$

We express in this subsection the 8×8 -matrix 1-form $\omega = M_1 dt_1 + M_2 dt_2 + M_3 dt_3$. We use parameters

$$b_1 = (-a_1 + a_2 + a_3)/2, \quad b_2 = (a_1 - a_2 + a_3)/2, \quad b_3 = (a_1 + a_2 - a_3)/2.$$

$$M_1 = \begin{pmatrix} 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ \frac{b_1}{(t_1^2-1)} & \frac{a_0 t_1}{(t_1^2-1)} & 0 & 0 & \frac{-(t_1 t_2 - t_3)}{(t_1^2-1)} & \frac{-(t_1 t_3 - t_2)}{(t_1^2-1)} & \frac{t_2 t_3 - t_1}{(t_1^2-1)} & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & \frac{p152}{(t_1^2-1)D} & \frac{p153}{(t_1^2-1)D} & \frac{p154}{(t_1^2-1)D} & \frac{p155}{(t_1^2-1)D} & \frac{1}{(t_1^2-1)} & \frac{p157}{(t_1^2-1)D} & \frac{(t_2 - t_3 t_1)}{(t_1^2-1)D} \\ 0 & \frac{p162}{(t_1^2-1)D} & \frac{p163}{(t_1^2-1)D} & \frac{p164}{(t_1^2-1)D} & \frac{1}{(t_1^2-1)} & \frac{p166}{(t_1^2-1)D} & \frac{p167}{(t_1^2-1)D} & \frac{-(t_1 t_2 - t_3)}{(t_1^2-1)D} \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & \frac{1}{D} \\ \frac{m181}{(t_1^2-1)D} & \frac{m182}{D} & \frac{m183}{(t_1^2-1)D} & \frac{m184}{(t_1^2-1)D} & \frac{m185}{(t_1^2-1)} & \frac{m186}{(t_1^2-1)} & \frac{m187}{(t_1^2-1)D} & \frac{m188}{(t_1^2-1)D} \end{pmatrix},$$

$$p152 = -b_2(t_1 t_2 - t_3) + b_3 t_1(t_1 t_3 - t_2),$$

$$p153 = b_1(t_2^2 - 1) - (b_1 + b_3)t_1(t_2 t_3 - t_1),$$

$$p154 = b_2(t_2 t_3 - t_1) - b_1 t_3(t_1 t_3 - t_2),$$

$$p155 = -a_0(t_1^2 - 1)(t_2 t_3 - t_1) + (t_1 t_2 - t_3)(t_1 t_3 - t_2),$$

$$p157 = -(1 + a_0)(t_2 t_3 - t_1)(t_1 t_3 - t_2),$$

$$p162 = \sigma_{23} \circ p152 = -b_3(t_1 t_3 - t_2) + b_2 t_1(t_1 t_2 - t_3),$$

$$p163 = \sigma_{23} \circ p154 = b_3(t_2 t_3 - t_1) - b_1 t_2(t_1 t_2 - t_3),$$

$$p164 = \sigma_{23} \circ p153 = b_1(t_3^2 - 1) - (b_1 + b_2)t_1(t_2 t_3 - t_1),$$

$$p166 = \sigma_{23} \circ p155 = p155 = -a_0(t_1^2 - 1)(t_2 t_3 - t_1) + (t_1 t_2 - t_3)(t_1 t_3 - t_2),$$

$$p167 = \sigma_{23} \circ p157 = -(1 + a_0)(t_2 t_3 - t_1)(t_1 t_2 - t_3),$$

where we use the permutation σ_{23} : $(\sigma_{23} \circ P)(t_1, t_2, t_3, b_1, b_2, b_3) = P(t_1, t_3, t_2, b_1, b_3, b_2)$.

$$m181 = -b_2 b_3(t_1^2 - 1)(t_2 t_3 - t_1) + b_1(b_2 + b_3)(t_1 t_2 - t_3)(t_1 t_3 - t_2),$$

$$m182 = b_3(1 - t_3^2) + b_2(1 - t_2^2),$$

$$m183 = a_0 b_1 t_2(t_1 t_2 - t_3)(t_1 t_3 - t_2) - a_0 b_3(t_1 t_3 - t_2)(t_2 t_3 - t_1) + b_1 t_3(t_1^2 - 1)(t_2^2 - 1),$$

$$m184 = a_0 b_1 t_3(t_1 t_3 - t_2)(t_1 t_2 - t_3) - a_0 b_2(t_1 t_2 - t_3)(t_2 t_3 - t_1) + b_1 t_2(t_1^2 - 1)(t_3^2 - 1),$$

$$m185 = a_0(t_2 - t_3 t_1) + b_3(t_2 - t_3 t_1) + b_2(t_2 t_1^2 - t_3 t_1),$$

$$m186 = a_0(t_3 - t_1 t_2) + b_3(t_1^2 t_3 - t_1 t_2) + b_2(t_3 - t_1 t_2),$$

$$m187 = a_0^2(t_2 t_3 - t_1)(t_1 t_3 - t_2)(t_1 t_2 - t_3) - (a_0 + b_2 + b_3)t_1(t_2 t_3 - t_1)D + b_1(t_1^2 - 1)D \\ + (t_1^2 - 1)(t_2^2 - 1)(t_3^2 - 1),$$

$$m188 = a_0(-t_1 t_2^2 + 2t_2 t_3 - t_3^2 t_1 + t_1^3 - t_1) + 2(t_1^2 - 1)(t_2 t_3 - t_1).$$

Note that $m181, m182, m187$ and $m188$ are σ_{23} -invariant, and $\sigma_{23} \circ m183 = m184, \sigma_{23} \circ m185 = m186$.

$$M_2 = \begin{pmatrix} 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ \frac{b_2}{(t_2^2 - 1)} & 0 & \frac{a_0 t_2}{(t_2^2 - 1)} & 0 & \frac{-(t_1 t_2 - t_3)}{(t_2^2 - 1)} & \frac{t_1 t_3 - t_2}{(t_2^2 - 1)} & \frac{-(t_2 t_3 - t_1)}{(t_2^2 - 1)} & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & \frac{p252}{(t_2^2 - 1)D} & \frac{p253}{(t_2^2 - 1)D} & \frac{p254}{(t_2^2 - 1)D} & \frac{p255}{(t_2^2 - 1)D} & \frac{p256}{(t_2^2 - 1)D} & \frac{1}{(t_2^2 - 1)} & \frac{-(t_2 t_3 - t_1)}{(t_2^2 - 1)D} \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & \frac{1}{D} \\ 0 & \frac{p272}{(t_2^2 - 1)D} & \frac{p273}{(t_2^2 - 1)D} & \frac{p274}{(t_2^2 - 1)D} & \frac{1}{(t_2^2 - 1)} & \frac{p276}{(t_2^2 - 1)D} & \frac{p277}{(t_2^2 - 1)D} & \frac{-(t_1 t_2 - t_3)}{(t_2^2 - 1)D} \\ \frac{m281}{(t_2^2 - 1)D} & \frac{m282}{(t_2^2 - 1)D} & \frac{m283}{D} & \frac{m284}{(t_2^2 - 1)D} & \frac{m285}{(t_2^2 - 1)} & \frac{m286}{(t_2^2 - 1)D} & \frac{m287}{(t_2^2 - 1)} & \frac{m288}{(t_2^2 - 1)D} \end{pmatrix},$$

$$p252 = \sigma_{12} \circ p153,$$

$$p253 = \sigma_{12} \circ p152,$$

$$p254 = \sigma_{12} \circ p154,$$

$$p255 = \sigma_{12} \circ p155,$$

$$p256 = \sigma_{12} \circ p157 = p157,$$

$$m281 = \sigma_{12} \circ m181,$$

$$m282 = \sigma_{12} \circ m183,$$

$$m283 = \sigma_{12} \circ m182,$$

$$m284 = \sigma_{12} \circ m184,$$

$$p272 = \sigma_{12} \circ p163,$$

$$p273 = \sigma_{12} \circ p162,$$

$$p274 = \sigma_{12} \circ p164,$$

$$p276 = \sigma_{12} \circ p167,$$

$$p277 = \sigma_{12} \circ p166 = p255,$$

$$m285 = \sigma_{12} \circ m185,$$

$$m286 = \sigma_{12} \circ m187,$$

$$m287 = \sigma_{12} \circ m186,$$

$$m288 = \sigma_{12} \circ m188,$$

where $(\sigma_{12} \circ P)(t_1, t_2, t_3, b_1, b_2, b_3) = P(t_2, t_1, t_3, b_2, b_1, b_3)$.

$$M_3 = \begin{pmatrix} 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ \frac{b_3}{(t_3^2 - 1)} & 0 & 0 & \frac{a_0 t_3}{(t_3^2 - 1)} & \frac{t_1 t_2 - t_3}{(t_3^2 - 1)} & \frac{-(t_1 t_3 - t_2)}{(t_3^2 - 1)} & \frac{-(t_2 t_3 - t_1)}{(t_3^2 - 1)} & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & \frac{1}{D} \\ 0 & \frac{p362}{(t_3^2 - 1)D} & \frac{p363}{(t_3^2 - 1)D} & \frac{p364}{(t_3^2 - 1)D} & \frac{p365}{(t_3^2 - 1)D} & \frac{p366}{(t_3^2 - 1)D} & \frac{1}{(t_3^2 - 1)} & \frac{-(t_2 t_3 - t_1)}{(t_3^2 - 1)D} \\ 0 & \frac{p372}{(t_3^2 - 1)D} & \frac{p373}{(t_3^2 - 1)D} & \frac{p374}{(t_3^2 - 1)D} & \frac{p375}{(t_3^2 - 1)D} & \frac{1}{(t_3^2 - 1)} & \frac{p377}{(t_3^2 - 1)D} & \frac{-(t_1 t_2 - t_3)}{(t_3^2 - 1)D} \\ \frac{m381}{(t_3^2 - 1)D} & \frac{m382}{(t_3^2 - 1)D} & \frac{m383}{(t_3^2 - 1)D} & \frac{m384}{D} & \frac{m385}{(t_3^2 - 1)D} & \frac{m386}{(t_3^2 - 1)} & \frac{m387}{(t_3^2 - 1)} & \frac{m388}{(t_3^2 - 1)D} \end{pmatrix},$$

$$\begin{aligned}
p362 &= \sigma_{13} \circ p164, & p372 &= \sigma_{13} \circ p154, \\
p363 &= \sigma_{13} \circ p163, & p373 &= \sigma_{13} \circ p153, \\
p364 &= \sigma_{13} \circ p162, & p374 &= \sigma_{13} \circ p152, \\
p365 &= \sigma_{13} \circ p167 = p167, & p375 &= \sigma_{13} \circ p157, \\
p366 &= \sigma_{13} \circ p166, & p377 &= \sigma_{13} \circ p155 = p366, \\
m381 &= \sigma_{13} \circ m181, & m385 &= \sigma_{13} \circ m187, \\
m382 &= \sigma_{13} \circ m184, & m386 &= \sigma_{13} \circ m186, \\
m383 &= \sigma_{13} \circ m183, & m387 &= \sigma_{13} \circ m185, \\
m384 &= \sigma_{13} \circ m182, & m388 &= \sigma_{13} \circ m188,
\end{aligned}$$

where $(\sigma_{13} \circ)P(t_1, t_2, t_3, b_1, b_2, b_3) = P(t_3, t_2, t_1, b_3, b_2, b_1)$.

REMARK 8.1. All the poles of the entries of M_i are simple.

8.2. 6×6-matrix form $\omega_6 = N_1 dt_1 + N_2 dt_2$. In §3.1 the system $Z_2(A)$ is transformed into the Pfaffian form $de_6 = \omega_6 e_6$ with the frame

$$e_6 = \text{tr}(F, F_1, F_2, (t_1 - t_2)F_{11}, (t_1 - t_2)F_{12}, (t_1 - t_2)^2 F_{112}).$$

We express in this subsection the 6×6 -matrix 1-form $\omega_6 = N_1 dt_1 + N_2 dt_2$.

$$N_1 = \begin{pmatrix} 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & \frac{1}{t_1 - t_2} & 0 & 0 \\ 0 & 0 & 0 & 0 & \frac{1}{t_1 - t_2} & 0 \\ \frac{-b_1(2+a_0)}{t_1^2 - 1} & \frac{n142}{t_1^2 - 1} & \frac{-(b_1+b_3)(t_1 - t_2)}{t_1^2 - 1} & \frac{n144}{(t_1^2 - 1)(t_1 - t_2)} & \frac{n145}{(t_1^2 - 1)(t_1 - t_2)} & \frac{1 - 2t_1 t_2 + t_1^2}{(t_1^2 - 1)(t_1 - t_2)} \\ 0 & 0 & 0 & 0 & \frac{1}{t_1 - t_2} & \frac{1}{t_1 - t_2} \\ \frac{n161}{t_1^2 - 1} & \frac{n162}{(t_1^2 - 1)(t_1 - t_2)} & \frac{n163}{(t_1^2 - 1)(t_1 - t_2)} & \frac{n164}{(t_1^2 - 1)(t_1 - t_2)} & \frac{n165}{(t_1^2 - 1)(t_1 - t_2)} & \frac{n166}{(t_1^2 - 1)(t_1 - t_2)} \end{pmatrix}$$

$$\begin{aligned}
n142 &= -a_0(t_1 + t_2) - a_0^2 t_1 + (b_1 - b_3)(t_1 - t_2), \\
n144 &= t_1^2 + 2t_1 t_2 - 3 + a_0(2t_1^2 - t_1 t_2 - 1), \\
n145 &= 2(t_2^2 - 1) - a_0(t_1^2 - 2t_1 t_2 + 1), \\
n161 &= (2 + a_0)b_1 - (b_1 + b_2)(b_1 + b_3), \\
n162 &= -a_0(b_1 + b_3)t_1(t_1 - t_2) + a_0(2 + a_0)t_1(t_1 - t_2) - a_0b_2(t_1^2 - 2t_1 t_2 + 1) + 2b_2(t_2^2 - 1) \\
&\quad + 2b_3(t_1 - t_2)^2, \\
n163 &= -a_0b_3t_2(t_1 - t_2) + 2b_3(t_1 - t_2)^2 + 2b_1(1 - 2t_1 t_2 + t_1^2) + a_0b_1(1 - t_1 t_2), \\
n164 &= 2 - 2t_1^2 + a_0(1 - t_1^2) + 2b_3t_1(t_1 - t_2) + b_2(1 - 2t_1 t_2 + t_1^2) - b_1(1 - t_1^2), \\
n165 &= 2 - 2t_2^2 + a_0(3t_1^2 - 4t_1 t_2 - t_2^2 + 2) + a_0^2(1 - t_1 t_2) + b_3(t_1^2 - t_2^2) + b_1(t_1^2 + t_2^2 - 2), \\
n166 &= -2t_1^2 + 4t_1 t_2 - 2 + a_0(t_1^2 + t_1 t_2 - 2),
\end{aligned}$$

$$N_2 = \begin{pmatrix} 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & \frac{1}{t_1 - t_2} & 0 \\ \frac{b_1 + b_2}{t_2^2 - 1} & \frac{a_0 t_1}{t_2^2 - 1} & \frac{a_0 t_2}{t_2^2 - 1} & \frac{-(t_1^2 - 1)}{(t_2^2 - 1)(t_1 - t_2)} & \frac{2(1 - t_1 t_2)}{(t_2^2 - 1)(t_1 - t_2)} & 0 \\ 0 & 0 & 0 & \frac{-1}{t_1 - t_2} & 0 & \frac{-1}{t_1 - t_2} \\ \frac{b_1(2 + a_0)}{t_2^2 - 1} & \frac{n252}{t_2^2 - 1} & \frac{(b_1 + b_3)(t_1 - t_2)}{t_2^2 - 1} & \frac{-(2 + a_0)(t_1^2 - 1)}{(t_2^2 - 1)(t_1 - t_2)} & \frac{n255}{(t_2^2 - 1)(t_1 - t_2)} & \frac{-(t_1^2 - 1)}{(t_2^2 - 1)(t_1 - t_2)} \\ \frac{n261}{t_2^2 - 1} & \frac{n262}{(t_2^2 - 1)(t_1 - t_2)} & \frac{n263}{(t_2^2 - 1)(t_1 - t_2)} & \frac{n264}{(t_2^2 - 1)(t_1 - t_2)} & \frac{n265}{(t_2^2 - 1)(t_1 - t_2)} & \frac{-2(t_2^2 - 1) + a_0(2 - t_1^2 - t_2^2)}{(t_2^2 - 1)(t_1 - t_2)} \end{pmatrix}$$

$$\begin{aligned}
n252 &= (2 + a_0)a_0 t_1 + (b_2 + b_3)(t_1 - t_2), \\
n255 &= -t_2^2 - 2t_1 t_2 + 3 + a_0(t_1^2 - t_2^2 + 1 - t_1 t_2), \\
n261 &= (2 + a_0)a_0 b_1 + (b_1 + b_3)(b_1 + b_2), \\
n262 &= (2 + a_0)a_0^2 t_1(t_1 - t_2) + a_0 b_1 t_1(t_1 - t_2) + a_0 b_2(1 - 2t_1 t_2 + t_1^2) \\
&\quad + a_0 b_3(2t_1^2 - 3t_1 t_2 + t_2^2) + 2b_2(1 - t_2^2), \\
n263 &= a_0 b_3 t_1(t_1 - t_2) - 2b_1(1 - t_2^2) + a_0 b_1(t_2^2 + t_1^2 - 1 - t_1 t_2), \\
n264 &= (2a_0 + a_0^2 + b_1 + b_3)(1 - t_1^2) - (b_2 + b_3)(1 - t_2^2), \\
n265 &= a_0(t_2^2 - 2t_1 t_2 + 1) + a_0^2 t_1(t_1 - t_2) - 2b_3 t_2(t_1 - t_2) + 2b_1(1 - t_1 t_2).
\end{aligned}$$

REMARK 8.2. All the poles of the entries of N_i are simple.

Interested readers may refer to our list of data in Maple format⁶ for the matrix 1-forms in this section as well as related ordinary differential equations in Sections 1-3.

9. Tensor product of two Gauss equations

Consider two differential equations

$$z_1'' = S_1 z_1 \quad \text{and} \quad z_2'' = S_2 z_2 \quad (z' := dz/dx),$$

with dependent variables z_1 and z_2 . If $S_1 \neq S_2$, the product $w = z_1 z_2$ satisfies the fourth-order differential equation $K_{S_1, S_2} w = 0$, where

$$K_{S_1, S_2} := \partial^4 + f_3 \partial^3 + f_2 \partial^2 + f_1 \partial + f_0, \quad (\partial := d/dx)$$

and

$$\begin{aligned}
f_3 &:= -\frac{S'_1 - S'_2}{S_1 - S_2}, \quad f_2 := -2(S_1 + S_2), \\
f_1 &:= -\frac{S_1 S'_1 - S_2 S'_2 + 5(S_1 S'_2 - S'_1 S_2)}{S_1 - S_2}, \\
f_0 &:= -S_1'' - S_2'' + (S_1 - S_2)^2 + \frac{(S'_1)^2 - (S'_2)^2}{S_1 - S_2}.
\end{aligned}$$

If $S_1 = S_2 = S$, $w = z_1^2$ satisfies the third-order equation $K_S w = 0$, where

⁶<http://www.math.kobe-u.ac.jp/OpenXM/Math/Z12data/z12.html>

$$(4) \quad K_S = \partial^3 - 4S\partial - 2S'.$$

For two Gauss equations

$$y_j'' + p_j y_j' + q_j y_j = 0, \quad p_j = \frac{c_j - (a_j + b_j + 1)x}{x(1-x)}, \quad q_j = -\frac{a_j b_j}{x(1-x)}, \quad (j = 1, 2)$$

we let

$$y_j = \lambda_j z_j, \quad \lambda_j = x^{-c_j/2} (x-1)^{(c_j-a_j-b_j-1)/2}.$$

Then, z_j satisfies the equation

$$z_j'' = S_j z_j, \quad S_j = -q_j + \frac{1}{4} p_j^2 + \frac{1}{2} p_j',$$

with the Riemann scheme

$$\left\{ \begin{array}{ccc} x = 0 & x = 1 & x = \infty \\ c_j/2 & (a_j + b_j - c_j + 1)/2 & (a_j - b_j - 1)/2 \\ 1 - c_j/2 & (c_j - a_j - b_j + 1)/2 & (b_j - a_j - 1)/2 \end{array} \right\}.$$

From the equation K_{S_1, S_2} satisfied by $z_1 z_2$, we get the differential equation

$$K = K(a_1, b_1, c_1, a_2, b_2, c_2) = \text{Ad}(\lambda_1 \lambda_2) K_{S_1, S_2}$$

satisfied by $y_1 y_2 = (\lambda_1 \lambda_2) z_1 z_2$. Though we omit the explicit form of K , if $(a_1, b_1, c_1) \neq (a_2, b_2, c_2)$, it is of order four and has *generically* two apparent singular points say $\{x_1, x_2\}$ other than $\{0, 1, \infty\}$, and the Riemann scheme is given as

$$\begin{aligned} & \left\{ \begin{array}{ccc} x = 0 & x = 1 & x = \infty \\ 0 & 0 & a_1 \\ 1 - c_1 & c_1 - a_1 - b_1 & b_1 \end{array} \right\} \times \left\{ \begin{array}{ccc} x = 0 & x = 1 & x = \infty \\ 0 & 0 & a_2 \\ 1 - c_2 & c_2 - a_2 - b_2 & b_2 \end{array} \right\} \\ & = \left\{ \begin{array}{ccccc} x = 0 & x = 1 & x = \infty & x = x_1 & x = x_2 \\ 0 & 0 & a_1 + a_2 & 0 & 0 \\ 1 - c_1 & c_1 - a_1 - b_1 & a_1 + b_2 & 1 & 1 \\ 1 - c_2 & c_2 - a_2 - b_2 & b_1 + a_2 & 2 & 2 \\ 2 - c_1 - c_2 & c_1 + c_2 - a_1 - b_1 - a_2 - b_2 & b_1 + b_2 & 4 & 4 \end{array} \right\}. \end{aligned}$$

Note that though the Gauss equations have no accessory parameters, K has one. The two apparent singular points $\{x_1, x_2\}$ are the roots of the following quadratic form:

$$\begin{aligned} App(x) := & (a_1 + a_2 - b_1 - b_2)(a_1 - a_2 - b_1 + b_2)x^2 \\ & + 2(2a_1 b_1 - 2a_2 b_2 + (1 - a_1 - b_1)c_1 - (1 - a_2 - b_2)c_2)x + (c_1 - c_2)(c_1 + c_2 - 2). \end{aligned}$$

If for example, $c_1 = c_2$, then App is divisible by x , and so we set $x_2 = 0$, and if the other parameters remain generic, the Riemann scheme becomes

$$\left\{ \begin{array}{cccc} x = 0 & x = 1 & x = \infty & x = x_1 \\ 0 & 0 & a_1 + a_2 & 0 \\ 1 - c_1 & c_1 - a_1 - b_1 & a_1 + b_2 & 1 \\ 2 - c_1 & c_1 - a_2 - b_2 & b_1 + a_2 & 2 \\ 2 - 2c_1 & 2c_1 - a_1 - b_1 - a_2 - b_2 & b_1 + b_2 & 4 \end{array} \right\}.$$

9.1. Tensor product without apparent singularities. There are several choices of parameters that the tensor product has no apparent singularities, that is the cases App reduces to constant times

$$x^2, \quad (x-1)^2, \quad 1, \quad \text{and} \quad x-1, \quad x, \quad x(x-1),$$

corresponding to

$$\{x_1, x_2\} \rightarrow 0, 1, \infty, \quad \text{and} \quad \{x_1, x_2\} \rightarrow \{0, \infty\}, \{1, \infty\}, \{0, 1\},$$

respectively. Thanks to the symmetry of the Gauss equations on the three singular points, we consider only two cases: x^2 and $x(x-1)$. The first case occurs only when

$$(1.1) \quad \{c_1 = c_2, a_1 = (2a_2b_2 - a_2c_2 + b_1c_2 - b_2c_2)/(2b_1 - c_2)\}, \quad \text{or}$$

$$(1.2) \quad \{c_1 = 2 - c_2, a_1 = (2a_2b_2 - a_2c_2 - b_1c_2 - b_2c_2 + 2b_1 + 2c_2 - 2)/(2b_1 + c_2 - 2)\};$$

and the second case,

$$(2.1) \quad \{c_1 = c_2, a_1 = -b_1 + 2c_2 - a_2 - b_2\}, \quad \text{or} \quad (\text{this is used below})$$

$$(2.2) \quad \{c_1 = c_2, a_1 = a_2 - b_1 + b_2\}, \quad \text{or}$$

$$(2.3) \quad \{c_1 = 2 - c_2, a_1 = -b_1 - 2c_2 + 2 + a_2 + b_2\}, \quad \text{or}$$

$$(2.4) \quad \{c_1 = 2 - c_2, a_1 = -a_2 - b_1 - b_2 + 2\}.$$

For the first case, two of the local exponents at $x = 0$ differ by 2, and for the second case, two of the local exponents at $x = 0$ and at $x = 1$ differ by 1. Thanks to the adjoint symmetry (cf. §4.3)

$$G(\alpha, \beta, \gamma) \longleftrightarrow G(1 - \alpha, 1 - \beta, 2 - \gamma)$$

of the Gauss equation⁷, we study only two cases (1.1) and (2.1); in these cases we have

$$c_1 = c_2 =: c.$$

For each case, in the following, we consider the renormalized equation (addition by x^{c-1})

$$L := \text{Ad}(x^{c-1})(K) = x^{c-1} \circ K \circ x^{1-c}.$$

After cancelling the common factor (denoted also by L), it is of the form

$$L = \begin{cases} x^3(x-1)^3\partial^4 + \dots & \text{in case (1.1),} \\ x^2(x-1)^2\partial^4 + \dots & \text{in case (2.1).} \end{cases}$$

9.2. Tensor product without apparent singularities Case 1. In this subsection we assume

$$a_1 = (2a_2b_2 - a_2c_2 + b_1c_2 - b_2c_2)/(2b_1 - c_2), \quad c_1 = c_2,$$

and study the middle convolution of L . The local exponents of L are given as follows:

$$x = 0 : [0, 2, c_2 - 1, -c_2 + 1],$$

⁷Adjoint equation of $G(\alpha, \beta, \gamma)$ is $G(1 - \alpha, 1 - \beta, 2 - \gamma)$.

$$\begin{aligned} x = 1 : & [0, c_1 - a_1 - b_1, c_2 - a_2 - b_2, c_1 - a_1 - b_1 + c_2 - a_2 - b_2], \\ x = \infty : & [a_1 + a_2 - c_2 + 1, a_1 + b_2 - c_2 + 1, b_1 + b_2 - c_2 + 1, b_1 + a_2 - c_2 + 1], \end{aligned}$$

where $a_1 = (2a_2b_2 - a_2c_2 + b_1c_2 - b_2c_2)/(2b_1 - c_2)$ and $c_1 = c_2$ should be assumed. We follow the recipe of making the middle convolution: we consider $L6 := \partial^2 \circ L$, and express it in terms of $(x\partial_x)^i \circ (\partial_x)^j$ with constant coefficients and replace $x\partial_x$ by $x\partial_x - m$, where m is a constant (parameter of middle convolution). The resulting operator $M6$ is of order 6, with parameter m , written as

$$M6 = cm_6\partial^6 + cm_5\partial^5 + cm_4\partial^4 + cm_3\partial^3 + cm_2\partial^2 + cm_1\partial^1 + cm_0,$$

where

$$\begin{aligned} cm_6 &= (2b_1 - c_2)^2 x^3 (x - 1)^3, \\ cm_5 &= x^2 (x - 1)^2 (2b_1 - c_2) (4xb_1^2 - 8xb_1c_2 + 4xa_2b_1 + 44xb_1 \\ &\quad - 12xb_1m + 4xb_2b_1 + 4c_2^2x + 4a_2b_2x - 4c_2b_2x - 4c_2a_2x - 22c_2x \\ &\quad + 6xmc_2 - 20b_1 + 6b_1m + 10c_2 - 3mc_2), \\ cm_4 &= x(x - 1)P_2(x), \quad cm_3 = P_3(x), \quad cm_2 = P_2(x), \quad cm_1 = (m - 1)P_1(x), \\ cm_0 &= (m - 1)(m - 2)(b_1 + b_2 - c_2 + 1 - m)(b_1 + a_2 - c_2 + 1 - m) \\ &\quad \times (-2b_1m + mc_2 + 2b_1 - c_2 - b_1c_2 + c_2^2 + 2a_2b_1 - 2c_2a_2 + 2a_2b_2 - c_2b_2) \\ &\quad \times (-2b_1m + mc_2 + 2b_1 - c_2 - b_1c_2 + c_2^2 - c_2a_2 + 2b_2b_1 - 2c_2b_2 + 2a_2b_2), \end{aligned}$$

where $P_k(x)$ denotes symbolically a polynomial of degree k in x . The local exponents of $\partial^2 \circ L$ are

$$\begin{aligned} x = 0 : & [0, 1, 2, 2, c_2 - 1, -c_2 + 1], \\ x = 1 : & [0, 1, 2, c_2 - a_2 - b_2, c_1 - a_1 - b_1, c_1 - a_1 - b_1 + c_2 - a_2 - b_2], \\ x = \infty : & [1, 2, 1 + b_1 + b_2 - c_2, 1 + b_1 + a_2 - c_2, 1 + a_1 + a_2 - c_2, 1 + a_1 + b_2 - c_2], \end{aligned}$$

and those of $M6$ are

$$\begin{aligned} x = 0 : & [0, 1, 2, m + 2, c_2 - 1 + m, 1 - c_2 + m], \\ x = 1 : & [0, 1, 2, m + c_2 - a_2 - b_2, m + c_1 - a_1 - b_1, m + c_1 - a_1 - b_1 + c_2 - a_2 - b_2], \\ x = \infty : & [-m + 1, -m + 2, -m + 1 + b_1 + b_2 - c_2, -m + 1 + b_1 + a_2 - c_2, \\ &\quad -m + 1 + a_1 + a_2 - c_2, -m + 1 + a_1 + b_2 - c_2]. \end{aligned}$$

The difference is

$$[0, 0, 0, m, m, m], \quad [0, 0, 0, m, m, m], \quad [-m, -m, -m, -m, -m, -m].$$

Though L would be generically irreducible, since the local exponents at $x = 0$ are $[0, 2, c_2 - 1, -c_2 + 1]$, the resulting $M6$ might be reducible (cf. [6]). In fact $M6$ breaks as

$$M6 = Y \circ M5, \quad M5 = u_5\partial^5 + u_4\partial^4 + u_3\partial^3 + u_2\partial^2 + u_1\partial + u_0,$$

where Y is a first order operator, and u_0, \dots, u_5 are polynomials in x and the parameters, and u_5 is given as

$$u_5 = (2b_1 - c_2)^2 x^2 (x-1)^3 (b_1 - c_2 + b_2)(b_1 - c_2 + a_2)(x + (m-1)\lambda),$$

where

$$\lambda = -\frac{2b_1 - c_2}{(b_1 - c_2 + b_2)(b_1 - c_2 + a_2)}.$$

This shows that $M5$ has one apparent singularity at $p := -(m-1)\lambda$. The local exponents of $M5$ are

$$\begin{aligned} x = 0 : & [0, 1, 2, c_2 - 1 + m, -c_2 + 1 + m], \\ x = 1 : & [0, 1, -b_2 + c_2 + m - a_2, c_1 - a_1 - b_1 + m, c_1 - a_1 - b_1 + c_2 - a_2 - b_2 + m], \\ x = \infty : & [1 - m, b_1 + b_2 - c_2 + 1 - m, b_1 + a_2 - c_2 + 1 - m, a_1 + b_2 - c_2 + 1 - m, \\ & a_1 + a_2 - c_2 + 1 - m], \\ x = p : & [0, 1, 2, 3, 5]. \end{aligned}$$

9.2.1. When $M6$ is divisible from the left by a first-order operator. Note first that the operator $L = x^3(x-1)^3\partial^4 + \dots$ can be written as

$$L = x^2 Q_1(\theta, \partial) + \{\lambda(\theta-1) + x\} Q_2(\theta, \partial), \quad \theta = x\partial.$$

Since

$$\partial^2 x^2 = (\theta+1)(\theta+2), \quad \partial^2 \{\lambda(\theta-1) + x\} = \{(\theta+1)(\lambda\theta+1) + 1\}\partial,$$

$L6 = \partial^2 L$ is written in terms of θ and ∂ :

$$L6 = (\theta+1)(\theta+2)Q_1(\theta, \partial) + \{(\theta+1)(\lambda\theta+1) + 1\}\partial Q_2(\theta, \partial).$$

So $M6$ can be obtained from $L6$ by replacing θ by $\theta - m$:

$$M6 = (\theta+1-m)(\theta+2-m)Q_1(\theta-m, \partial) + \{(\theta+1-m)(\lambda\theta+1) + 1\}\partial Q_2(\theta-m, \partial).$$

On the other hand we have the following formulae:

$$\begin{aligned} (\theta+1-m)(\theta+2-m) &= \left(\theta+1-m + \frac{x}{x+u}\right) \left(\theta+2-m - \frac{x}{x+u}\right) \quad \text{for any constant } u, \\ (\theta+1-m)(\lambda\theta+1) + 1 &= \left(\theta+1-m + \frac{x}{x+(m-1)\lambda}\right) \left(\lambda\theta+1 - \frac{\lambda}{x+(m-1)\lambda}\right). \end{aligned}$$

Applying these by putting $u = (m-1)\lambda$, we see that $M6$ can be divisible from the left by

$$Y = \theta + 1 - m + \frac{x}{x + (m-1)\lambda}.$$

9.2.2. When $M5$ has no apparent singular point. Further to forget the singularity p , we assume, for example, $b_1 = c_2 - b_2$. Then we also have $a_1 = c_2 - a_2$, namely, the two Gauss are equal up to a Euler transformation: $F(a, b, c; x) = (1-x)^{c-a-b} F(c-a, c-b, c; x)$. But $M5$ remains to be an equation of order 5 (see §9.3.1 for an analogous phenomenon), with the local exponents

$$\begin{aligned} x = 0 : & [0, 1, 2, c_2 - 1 + m, -c_2 + 1 + m], \\ x = 1 : & [0, 1, m, b_2 + a_2 - c_2 + m, c_2 - b_2 + m - a_2], \\ x = \infty : & [2 - m, 1 - m, 1 - m, 1 - b_2 + a_2 - m, 1 - a_2 + b_2 - m], \end{aligned}$$

and is reducible of type [14]⁸. We have $[1] = -x(x-1)^2(x(x-1)\partial - (m-6)x-2)$, and

$$x^2(x-1)^2[4] = x^2(x-1)^2\partial^4 + m_3\partial^3 + m_2\partial^2 + m_1\partial^1 + m_0,$$

where

$$\begin{aligned} m_3 &= -x(2x-1)(x-1)(2m-5), \\ m_2 &= 2c_2b_2x + 2a_2b_2x^2 + 2c_2a_2x + 3 + 25x^2 - x^2b_2^2 - 24mx^2 - x^2a_2^2 + 6m^2x^2 \\ &\quad - 2c_2x - 4a_2b_2x - c_2^2 + 2c_2 - 4m + m^2 - 24x - 6m^2x + 24mx, \\ m_1 &= (-3 + 2m)(-2m^2x + m^2 - 3m + 6mx - c_2b_2 + x^2b_2^2 - c_2a_2 + 2 - 2a_2b_2x + 2a_2b_2 \\ &\quad + c_2 + x^2b_2^2 - 5x), \\ m_0 &= -(-1 + m)^2(a_2 - 1 + m - b_2)(a_2 + 1 - m - b_2). \end{aligned}$$

The local exponents are

$$\begin{aligned} x = 0 : & [0, 1, c_2 - 1 + m, -c_2 + 1 + m], \\ x = 1 : & [0, 1, -a_2 + c_2 - b_2 + m, -c_2 + b_2 + a_2 + m], \\ x = \infty : & [1 - m, 1 - m, -a_2 + 1 - m + b_2, a_2 + 1 - m - b_2]. \end{aligned}$$

9.3. Tensor product without apparent singularities Case 2.

In this section we assume

$$a_1 = -b_1 + c_1 + c_2 - a_2 - b_2, \quad c_1 = c_2 =: c,$$

and study the middle convolution of L . This case happens to connect Gauss equations and the equation $\tilde{Z}(A)$, because the above assumption exactly fits the parameter change (see §5.2):

$$a_1 = A_{-++}, \quad b_1 = A_{-++}, \quad a_2 = A_{---}, \quad b_2 = A_{---}, \quad c = 1 - A_0.$$

The middle convolution $ML4(m)$ of L with parameter m is now computed without multiplying ∂ from the left, and we get

$$ML4(m) := x^2(x-1)^2\partial^4 + m\ell_3\partial^3 + m\ell_2\partial^2 + m\ell_1\partial + m\ell_0,$$

where

$$\begin{aligned} m\ell_3 &= -x(2x-1)(x-1)(2m-5), \\ m\ell_2 &= 3 + 2b_1xb_2 + 2b_1xa_2 + 25x^2 - 24x + 2c - c^2 - 4m + 2xb_1^2 + 2ca_2x^2 \\ &\quad + 2cb_2x^2 - 2a_2b_2x - 2cx + 2c^2x - 2x^2b_1^2 - x^2a_2^2 - x^2b_2^2 - 2c^2x^2 \\ &\quad - 4cb_1x + 6m^2x^2 - 24mx^2 - 6m^2x + 24mx + m^2 - 2x^2b_1b_2 - 2x^2b_1a_2 + 4x^2b_1c, \\ m\ell_1 &= (-3 + 2m)(m^2 - 2m^2x - 3m + 6mx + 2 + 2c^2x + 2b_1xa_2 + 2b_1xb_2 + c - c^2 + a_2b_2 \\ &\quad - 5x - b_1^2 + 2xb_1^2 - 2ca_2x + 2b_1c - 2cb_2x - b_1a_2 - b_1b_2 - 4cb_1x + xb_2^2 + xa_2^2), \\ m\ell_0 &= (b_1 - 1 + m + b_2 - c)(b_1 + 1 - m + b_2 - c)(b_1 + 1 - c - m + a_2)(b_1 - 1 - c + m + a_2). \end{aligned}$$

The local exponents of $ML4(m)$ are

⁸An operator P decomposes (is reducible) for example of type [14] means P can be written as $P_1 \circ P_2$, where P_1 is of order 1 and P_2 is of order 4. [4] part means P_2 . The decomposition is not necessarily an irreducible decomposition, which is not unique.

$$\begin{aligned}
x = 0 : & [0, 1, c - 1 + m, 1 - c + m], \\
x = 1 : & [0, 1, b_2 - c + m + a_2, -b_2 - a_2 + c + m], \\
x = \infty : & [-b_1 + 1 + c - b_2 - m, b_1 + 1 + b_2 - c - m, -b_1 + 1 + c - m - a_2, \\
& \quad 1 + b_1 - c - m + a_2].
\end{aligned}$$

Recall that $L(A) = \text{Ad}(x^{-A_0})(K(A))$. Since $c - 1 = -A_0$, our L just agrees with $L(A)$. The middle convolution $ML4(m)$ of $L = L(A)$ is given by

$$\begin{aligned}
m\ell_3 &= -2x(2x - 1)(x - 1)(-2 + (m - 1/2)), \\
m\ell_2 &= 29/2x^2 - 29/2x - 3(m - 1/2) + 9/4 - 6x(m - 1/2)^2 + 18x(m - 1/2) \\
&\quad + 6x^2(m - 1/2)^2 - 18x^2(m - 1/2) + (m - 1/2)^2 - A_0^2 + xA_0^2 + xA_3^2 - xA_1^2 \\
&\quad + xA_2^2 - x^2A_2^2 - x^2A_3^2, \\
m\ell_1 &= -1/2(-1 + (m - 1/2))(10x + 8(m - 1/2) - 5 + 8x(m - 1/2)^2 - 16x(m - 1/2) \\
&\quad - 4(m - 1/2)^2 + 2A_0^2 + 2A_3^2 + 2A_2^2 - 2A_1^2 - 4xA_3^2 - 4xA_2^2), \\
m\ell_0 &= 1/16(2A_2 - 1 + 2(m - 1/2))(-2A_2 - 1 + 2(m - 1/2))(-2A_3 - 1 + 2(m - 1/2)) \\
&\quad \times (2A_3 - 1 + 2(m - 1/2)).
\end{aligned}$$

Thus we rediscovered Theorem 5.2: $ML4(1/2) = \tilde{Z}(A)$.

9.3.1. K_{S_1, S_2} when $A_2 = 0$. The quadratic form App giving the two apparent singularities of K and K_{S_1, S_2} reduces to $-4A_2A_3x(x - 1)$; note that

$$S_1 - S_2 = -\frac{A_2A_3}{x(x - 1)}.$$

The coefficients of $K_{S_1, S_2} = \partial^4 + f_3dx^3 + \dots$ are given as

$$\begin{aligned}
f_3 &:= (2x - 1)/(x(-1 + x)), \\
f_2 &:= (-A_2^2x^2 - A_3^2x^2 + A_0^2x - A_1^2x + A_2^2x + A_3^2x - A_0^2 + x^2 - x + 1)/(x^2(-1 + x)^2), \\
f_1 &:= -(-2A_2^2x^3 - 2A_3^2x^3 + 5A_0^2x^2 - 5A_1^2x^2 + 3A_2^2x^2 + 3A_3^2x^2 - 9A_0^2x + A_1^2x \\
&\quad - A_2^2x - A_3^2x + 2x^3 + 4A_0^2 - 3x^2 + 9x - 4)/(2x^3(-1 + x)^3), \\
f_0 &:= (2A_2^2A_3^2x^4 - 4A_2^2A_3^2x^3 + 2A_2^2A_3^2x^2 - 2A_2^2x^4 - 2A_3^2x^4 + 6A_0^2x^3 \\
&\quad - 6A_1^2x^3 + 4A_2^2x^3 + 4A_3^2x^3 - 15A_0^2x^2 + 3A_1^2x^2 - 3A_2^2x^2 - 3A_3^2x^2 + 2x^4 + 13A_0^2x \\
&\quad - A_1^2x + A_2^2x + A_3^2x - 4x^3 - 4A_0^2 + 15x^2 - 13x + 4)/(2x^4(-1 + x)^4).
\end{aligned}$$

When $A_2 = 0$, K_{S_1, S_2} decomposes of type [13]:

$$\begin{aligned}
[1] &= \partial + (2x - 1)/(x(-1 + x)), \\
[3] &= \partial^3 + (-A_3^2x^2 + A_0^2x - A_1^2x + A_3^2x - A_0^2 + x^2 - x + 1)\partial/(x^2(-1 + x)^2) \\
&\quad - (-2A_3^2x^3 + 3A_0^2x^2 - 3A_1^2x^2 + 3A_3^2x^2 - 5A_0^2x + A_1^2x - A_3^2x + 2x^3 \\
&\quad + 2A_0^2 - 3x^2 + 5x - 2)/(2x^3(-1 + x)^3).
\end{aligned}$$

When $A_1 = A_2 = 0$ ($(a_1, b_1, c_1) = (a_2, b_2, c_2)$), the 3rd order operator [3] above is nothing but the equation (4): $K_S = \partial^3 - 4S\partial - 2S'$.

Part II. Local solutions of ordinary differential equations related to the Dotsenko-Fateev equation

In Part I, we found a Fuchsian system of rank 8 in 3 variables with 4 parameters, and an ordinary differential equation $Z(A)$ of order 4 with three singular points by restricting the system on a projective line. In Part II, we study the ordinary differential equation $Z(A)$ and several related ones, around their singular points.

In §10, we study a linear difference equation $Rc_0(A)$ of order 2, which is the recurrence relation satisfied by the coefficients of a power series solution to $Z(A)$ at $x = 0$.

§10.1 introduces the invariant of such a difference equation.

§10.3 introduces special values ${}_4F_3(*; 1)$ of terminating generalized hypergeometric series ${}_4F_3$ at 1 satisfying a linear difference equation $Rc^{(0)}$ of order 2.

In §10.4, by studying the invariant of this difference equation and that of $Rc_0(A)$, we find solutions of $Rc_0(A)$ expressed in terms of ${}_4F_3(*; 1)$. This expression is very near to the product of two Gauss hypergeometric series. This observation leads to the discovery: A middle convolution sends $Z(A)$ to the product of two Gauss hypergeometric equation.

In §10.5 and 6, invariants of the difference equations are used to get local solutions of $Z(A)$ at $x = 0, 1$ and ∞ .

In §10.7, Riemann-Liouville transformation is recalled.

In §10.8 and 9, we get local solutions of $Z(A)$ at $x = 0, 1$ and ∞ by using middle convolution.

In §11, we study a linear difference equation $Rc_1(A)$ of order 2, which is the recurrence relation satisfied by the coefficients of a solution to $Z(A)$ at $x = 0$ with exponents $A_0 \pm 1/2$.

§11.2 introduces special values ${}_4F_3(*; 1)$ of non-terminating generalized hypergeometric series ${}_4F_3$ at 1 satisfying a linear difference equations of order 2.

§11.3 studies the difference equation $Rc^{(1)}$ satisfied by this special values.

In §11.4, by studying the invariant of this difference equation and that of $Rc_1(A)$, we find solutions of $Rc_1(A)$ expressed in terms of non-terminating ${}_4F_3(*; 1)$.

§12 and 13 give a set of local solutions of $Q(A)$ and the Dotsenko-Fateev equation.

10. Local solutions of $Z(A)$ at $x = 0$ with exponent 0 and $2A_0$, and those at infinity

The explicit form of the differential equation in question is given as

$$Z(A) = p_0 \partial^4 + p_1 \partial^3 + p_2 \partial^2 + p_3 \partial + p_4, \quad \partial = d/dx,$$

where

$$p_0 = x^3(x - 1)^2,$$

$$p_1 = -2x^2(x - 1)(2A_0x - 2A_0 - 5x + 3),$$

$$p_2 = x(p_{22}x^2 + p_{21}x + p_{20}),$$

$$p_{22} = 6A_0^2 - A_2^2 - A_3^2 - 24A_0 + 25,$$

$$p_{21} = -11A_0^2 - A_1^2 + A_2^2 + A_3^2 + 36A_0 - 59/2,$$

$$p_{20} = (10A_0 - 9)(2A_0 - 3)/4,$$

$$\begin{aligned}
p_3 &= p_{32}x^2 + p_{31}x + p_{30}, \\
p_{32} &= -(2A_0 - 3)(2A_0^2 - A_2^2 - A_3^2 - 6A_0 + 5), \\
p_{31} &= (A_0 - 1)(6A_0^2 + 2A_1^2 - 2A_2^2 - 2A_3^2 - 16A_0 + 11), \\
p_{30} &= -((2A_0 - 3)(2A_0 - 1)^2)/4, \\
p_4 &= p_{41}x + p_{40}, \\
p_{41} &= (A_0 - 1 + A_3)(A_0 - 1 - A_3)(A_0 - 1 + A_2)(A_0 - 1 - A_2), \\
p_{40} &= -(2A_0 - 1)^2(A_0^2 + A_1^2 - A_2^2 - A_3^2 - 2A_0 + 1)/4,
\end{aligned}$$

its Riemann scheme is given as

$$\left\{ \begin{array}{ccc} x = 0 & x = 1 & x = \infty \\ 0 & \frac{1}{2} - A_1 & 1 - A_0 + A_2 \\ A_0 - \frac{1}{2} & 0 & 1 - A_0 - A_2 \\ A_0 + \frac{1}{2} & 1 & 1 - A_0 + A_3 \\ 2A_0 & \frac{1}{2} + A_1 & 1 - A_0 - A_3 \end{array} \right\}.$$

In this section we give the two local solutions around the point $x = 0$, whose exponents are 0 and $2A_0$. The other local solutions around $x = 0$, whose exponents are $A_0 \pm 1/2$, will be given in the next section. The coefficients of the power series satisfy a 3-term recurrence relation (a homogeneous linear difference equation of order 2), say $Rc_0(A)$.

On the other hand, under some condition, special values of the terminating generalized hypergeometric series ${}_4F_3$ at 1 satisfy a linear difference equation of order 2. By making use of this fact, we solve the equation $Rc_0(A)$ in terms of the special values of ${}_4F_3$ at 1. The result suggests a relation between $Z(A)$ and the tensor product of two Gauss hypergeometric differential equations: they are connected by a middle convolution. This method will be applied to other equations at other singular points.

The local exponents of the equation $\tilde{Z}(A)$ at $x = 0$ are

$$0, \quad 1, \quad 1/2 + A_0, \quad 1/2 - A_0;$$

the holomorphic solution of $Z(A)$ at $x = 0$ corresponds to the solution of $\tilde{Z}(A)$ with local exponent $1/2 - A_0$, so the solution of $Z(A)$ at $x = 0$ with exponent $2A_0$ corresponds to the solution of $\tilde{Z}(A)$ with exponent $1/2 + A_0$, which can be obtained from that with exponent $1/2 - A_0$ by changing the sign of A_0 .

But for the other exponents $A_0 \pm 1/2$ of $Z(A)$ at $x = 0$ ($\{0, 1\}$ for $\tilde{Z}(A)$), we need to make use of non-terminating series ${}_4F_3$. These are studied in the next section.

10.1. Invariants of linear difference equations. In this subsection, we introduce the invariants of linear difference equations following [3]. Let us consider a homogeneous linear difference equation of order 2

$$P : C_n = p_1(n)C_{n-1} + p_2(n)C_{n-2},$$

where p_1 and p_2 are rational functions in n . The quantity

$$H := \frac{p_1(n)p_1(n+1)}{p_2(n+1)}$$

is called the **invariant** of the difference equation P . Consider another such equation

$$Q : D_n = q_1(n)D_{n-1} + q_2(n)D_{n-2}.$$

The two equations are said to be **essentially the same** if there is a homogeneous linear difference equation of order 1:

$$\lambda(n-1) = \mu(n)\lambda(n), \quad \mu(n) : \text{a rational function in } n$$

and a solution $\lambda(n)$ so that

$$\{C_n\} = \lambda(n)\{D_n\},$$

where $\{C_n\}$ is the set of solutions of P , and $\{D_n\}$ that of solutions of Q .

Proposition 10.1. *The two equations P and Q are essentially the same if and only if the two invariants agree.*

In fact, substituting

$$C_n = \lambda(n)D_n, \quad C_{n-1} = \lambda(n-1)D_{n-1}, \quad C_{n-2} = \lambda(n-2)D_{n-2},$$

into P and equalizing with Q , we have

$$\frac{q_1(n)}{p_1(n)} = \mu(n), \quad \frac{q_2(n)}{p_2(n)} = \mu(n)\mu(n-1),$$

and, by eliminating μ ,

$$\frac{p_1(n)p_1(n-1)}{p_2(n)} = \frac{q_1(n)q_1(n-1)}{q_2(n)}.$$

On the other hand if we assume the last identity, we can trace back the argument up to $C_n = \lambda(n)D_n$.

Corollary 10.2. *For two essentially the same equations P and Q , the spaces of solutions $\{C_n\}$ and $\{D_n\}$ are related as*

$$\{C_n\} = \frac{1}{w^{n+1}} \cdot \frac{\prod_j \Gamma(n + v_j + 1)}{\prod_i \Gamma(n + u_i + 1)} \cdot \{D_n\},$$

where the rational function $\frac{q_1(n)}{p_1(n)}$ is factorized as

$$\frac{q_1(n)}{p_1(n)} = w \cdot \frac{\prod_i (n + u_i)}{\prod_j (n + v_j)}, \quad w : \text{independent of } n.$$

10.2. Recurrence relation $Rc_0(A)$ for the coefficients of a holomorphic solution of $Z(A)$ at $x = 0$. Let

$$f^{(0,0)}(A; x) := \sum_{n=0}^{\infty} C_n x^n, \quad C_0 = 1$$

be the (normalized) power series solution to $Z(A)$ at $x = 0$. Substituting this expression into $Z(A)$, we see that the coefficients C_n satisfy the following recurrence relation $Rc_0(A)$ ($C_{-1} = 0, C_0 = 1$):

$$Rc_0(A) : \begin{aligned} C_n &= \frac{\{2(n-A_0)-1\}^2 \{2n^2-4A_0n+A_0^2+A_1^2-A_2^2-A_3^2+1-2(n-A_0)\}}{n(n-2A_0)(2n-2A_0-1)(2n-2A_0+1)} C_{n-1} \\ &\quad - \frac{4(n-A_0-A_2-1)(n-A_0+A_2-1)(n-A_0-A_3-1)(n-A_0+A_3-1)}{n(n-2A_0)(2n-2A_0-1)(2n-2A_0+1)} C_{n-2}. \end{aligned}$$

Thus we see that the invariant of $Rc_0(A)$ is given as

$$H_0(n; A) = \frac{-\{4(n-A_0)^2-1\} \left\{ (2n^2-4A_0n+A_0^2+A_1^2-A_2^2-A_3^2+1)^2 - 4(n-A_0)^2 \right\}}{4n(n-2A_0)(n-A_0-A_2)(n-A_0+A_2)(n-A_0-A_3)(n-A_0+A_3)}.$$

REMARK 10.3 (SYMMETRY). $Rc_0(A)$ is invariant under

$$A_j \rightarrow -A_j \ (j=1, 2, 3) \quad \text{and} \quad A_2 \leftrightarrow A_3.$$

10.3. 3-term relation for the special values (at 1) of balanced terminating hypergeometric series ${}_4F_3$. We consider the special value (at 1) of the generalized hypergeometric series:

$${}_4F_3(\alpha; 1) = \sum_{k=0}^{\infty} \frac{(\alpha_0)_k(\alpha_1)_k(\alpha_2)_k(\alpha_3)_k}{(\beta_1)_k(\beta_2)_k(\beta_3)_k} \frac{1}{k!}, \quad \alpha = \begin{pmatrix} \alpha_0, \alpha_1, \alpha_2, \alpha_3 \\ \beta_1, \beta_2, \beta_3 \end{pmatrix}.$$

Here we assume α_0 is a non-positive integer (the series terminates), and also assume it is *balanced*:

$$\beta_1 + \beta_2 + \beta_3 - \alpha_0 - \alpha_1 - \alpha_2 - \alpha_3 = 1.$$

Set

$$e_1 := \begin{pmatrix} 1, 0, 0, 0 \\ 1, 0, 0 \end{pmatrix}, \quad e_2 := \begin{pmatrix} 0, 1, 0, 0 \\ 0, 1, 0 \end{pmatrix}, \quad e_{12} := \begin{pmatrix} 1, 1, 0, 0 \\ 1, 1, 0 \end{pmatrix}.$$

We have

Proposition 10.4 ([1] (§3.7), see also [8]).

$$(5) \quad {}_4F_3(\alpha; 1) = U_1^{(0)}(\alpha) {}_4F_3(\alpha + e_1; 1) + V_1^{(0)}(\alpha) {}_4F_3(\alpha + e_{12}; 1),$$

$$(6) \quad {}_4F_3(\alpha; 1) = U_2^{(0)}(\alpha) {}_4F_3(\alpha + e_2; 1) + V_2^{(0)}(\alpha) {}_4F_3(\alpha + e_{12}; 1),$$

where

$$(7) \quad U_1^{(0)}(\alpha) := \frac{-(\beta_1 - \alpha_1)(\beta_1 + \beta_2 - \alpha_2 - \alpha_3)}{\beta_1(\beta_3 - \alpha_0 - 1)}, \quad V_1^{(0)}(\alpha) := \frac{-\alpha_1(\beta_2 - \alpha_2)(\beta_2 - \alpha_3)}{\beta_1\beta_2(\beta_3 - \alpha_0 - 1)},$$

$$(8) \quad U_2^{(0)}(\alpha) := \frac{-(\beta_2 - \alpha_0)(\beta_1 + \beta_2 - \alpha_2 - \alpha_3)}{\beta_2(\beta_3 - \alpha_1 - 1)}, \quad V_2^{(0)}(\alpha) := \frac{-\alpha_0(\beta_1 - \alpha_2)(\beta_1 - \alpha_3)}{\beta_1\beta_2(\beta_3 - \alpha_1 - 1)}.$$

Perform a change $\alpha \mapsto \alpha + e_{12}$ in (5), and we have

$$(9) \quad {}_4F_3(\alpha + e_{12}; 1) = U_1^{(0)}(\alpha + e_{12}) {}_4F_3(\alpha + 2e_1 + e_2; 1) + V_1^{(0)}(\alpha + e_{12}) {}_4F_3(\alpha + 2e_{12}; 1),$$

and a change $\alpha \mapsto \alpha + e_1$ in (6) to get

$$(10) \quad {}_4F_3(\alpha + e_1; 1) = U_2^{(0)}(\alpha + e_1) {}_4F_3(\alpha + e_{12}; 1) + V_2^{(0)}(\alpha + e_1) {}_4F_3(\alpha + 2e_1 + e_2; 1).$$

Eliminating ${}_4F_3(\alpha + 2e_1 + e_2; 1)$ and ${}_4F_3(\alpha + e_1; 1)$ from (5), (9) and (10), we eventually get

Lemma 10.5.

$${}_4F_3(\alpha; 1) = q_1^{(0)}(\alpha) {}_4F_3(\alpha + e_{12}; 1) + q_2^{(0)}(\alpha) {}_4F_3(\alpha + 2e_{12}; 1),$$

where

$$\begin{aligned} q_1^{(0)}(\alpha) &= U_1^{(0)}(\alpha) U_2^{(0)}(\alpha + e_1) + V_1^{(0)}(\alpha) + \frac{U_1^{(0)}(\alpha) V_2^{(0)}(\alpha + e_1)}{U_1^{(0)}(\alpha + e_{12})}, \\ q_2^{(0)}(\alpha) &= \frac{-U_1^{(0)}(\alpha) V_1^{(0)}(\alpha + e_{12}) V_2^{(0)}(\alpha + e_1)}{U_1^{(0)}(\alpha + e_{12})}. \end{aligned}$$

If we change α into

$$(n; \hat{\alpha}) := \begin{pmatrix} -n, \alpha_1 - n, \alpha_2, \alpha_3 \\ \beta_1 - n, \beta_2 - n, \beta_3 \end{pmatrix} = -n e_{12} + \hat{\alpha}, \quad \text{where } \hat{\alpha} = \begin{pmatrix} 0, \alpha_1, \alpha_2, \alpha_3 \\ \beta_1, \beta_2, \beta_3 \end{pmatrix}$$

in the lemma above, we get

Proposition 10.6.

$$D(n; \hat{\alpha}) := {}_4F_3 \left(\begin{matrix} -n, \alpha_1 - n, \alpha_2, \alpha_3 \\ \beta_1 - n, \beta_2 - n, \beta_3 \end{matrix}; 1 \right) \quad \text{with } \beta_1 + \beta_2 + \beta_3 - \alpha_1 - \alpha_2 - \alpha_3 = 1$$

satisfies the homogeneous linear difference equation

$$(11) \quad Rc^{(0)}(\hat{\alpha}) : D_n = q_1^{(0)}(n; \hat{\alpha}) D_{n-1} + q_2^{(0)}(n; \hat{\alpha}) D_{n-2}.$$

The invariant of the difference equation $Rc^{(0)}(\hat{\alpha})$ is

$$H^{(0)}(n; \hat{\alpha}) := \frac{q_1^{(0)}(n; \hat{\alpha}) q_1^{(0)}(n+1; \hat{\alpha})}{q_2^{(0)}(n+1; \hat{\alpha})}.$$

REMARK 10.7 (SYMMETRY). $Rc^{(0)}(\hat{\alpha})$ is invariant under

$$(12) \quad \alpha_2 \leftrightarrow \alpha_3 \quad \text{and} \quad \beta_1 \leftrightarrow \beta_2.$$

10.4. Local solutions at zero I: solving $Rc_0(A)$.

Proposition 10.8. *The two invariants $H_0(n; A)$ and $H^{(0)}(n; \hat{\alpha})$ of the difference equations $Rc_0(A)$ and $Rc^{(0)}(\hat{\alpha})$ agree, as functions in n , if and only if*

$$(13) \quad \hat{\alpha} = \begin{pmatrix} 0, \alpha_1, \alpha_2, \alpha_3 \\ \beta_1, \beta_2, \beta_3 \end{pmatrix} = \begin{pmatrix} 0, A_0, A_{-++}, A_{-+-} \\ A_{++-}, A_{++}, 1 - A_0 \end{pmatrix},$$

$$(14) \quad \text{or} \quad = \begin{pmatrix} 0, A_0 - A_2, A_{-++}, A_{-+-} \\ A_{++-}, A_{++}, 1 - A_0 - A_2 \end{pmatrix},$$

up to the symmetries of the two difference equations. Thus $Rc_0(A)$ and $Rc^{(0)}(\hat{\alpha})$ are essentially the same in these cases.

This can be obtained by solving the system

$$H_0(n; A) = H^{(0)}(n; \hat{\alpha}) \quad n = 1, 2, \dots$$

with unknown $\hat{\alpha}$. Actual process is as follows: The numerator of $H_0(n; A) - H^{(0)}(n; \hat{\alpha})$ is a polynomial in n of degree 14. The coefficient of the top term decomposes as

$$\text{constant} \times (\alpha_1 - \beta_1 - \beta_2 + 1 - A_1)(\alpha_1 - \beta_1 - \beta_2 + 1 + A_1).$$

If we set $\alpha_1 = \beta_1 + \beta_2 - 1 \pm A_1$, then the second top term decomposes as

$$\text{constant} \times (2A_1 - 1)(2A_1 + 1)(\alpha_2 + \alpha_3 - \beta_1 - \beta_2 + 2A_0).$$

Since we assume A_1 is a free parameter, we have

$$(15) \quad \alpha_2 + \alpha_3 - \beta_1 - \beta_2 + 2A_0 = 0.$$

Eliminating β_2 and β_3 from $H^{(0)}(n; \hat{\alpha})$ by making use of this equality and the balance condition

$$(16) \quad \beta_1 + \beta_2 + \beta_3 - \alpha_1 - \alpha_2 - \alpha_3 = 1$$

in Proposition 10.6, we get an expression

$$H^{(0)}(n; \hat{\alpha}) = \frac{-\gamma^{(0)} + O((n - A_0)^2)}{\gamma^{(0)} + O((n - A_0)^2)},$$

where

$$(17) \quad \gamma^{(0)} := A_0^2(\alpha_1 - A_0)^2(\alpha_2 - \beta_1 + A_0)^2(\alpha_3 - \beta_1 + A_0)^2.$$

Now, assume $\gamma^{(0)} \neq 0$. Then, $H^{(0)}(A_0; \hat{\alpha}) = -1$, whereas $H_0(A_0; A)$ is parameter dependent:

$$H_0(A_0; A) = \frac{-\left(A_0^2 - A_1^2 + A_2^2 + A_3^2 - 1\right)^2}{4A_0^2 A_2^2 A_3^2}.$$

So it does not happen that $H_0(n; A) = H^{(0)}(n; \hat{\alpha})$ in the case $\gamma^{(0)} \neq 0$. Therefore, $\gamma^{(0)}$ must be equal to 0. Thanks to the symmetry of $Rc^{(0)}(\hat{\alpha})$ (cf. Remark 10.3), we have only to study two cases:

Case 1 : $\alpha_1 = A_0$ and Case 2 : $\beta_1 = \alpha_3 + A_0$.

Case 1: Eliminating β_2 and β_3 from $H^{(0)}(n; \hat{\alpha})$ by (15) and (16), we have

$$H^{(0)}(n; \hat{\alpha}) = \frac{-\{4(n - A_0)^2 - 1\} \epsilon_1^{(0)}(n) \epsilon_1^{(0)}(n + 1)}{4n(n - 2A_0)(n + \alpha_2 - \beta_1)(n - \alpha_2 + \beta_1 - 2A_0)(n + \alpha_3 - \beta_1)(n - \alpha_3 + \beta_1 - 2A_0)},$$

where $\epsilon_1^{(0)}(n) = 2(n^2 - 2A_0n + 2A_0\beta_1 + \alpha_2\alpha_3 + \alpha_2\beta_1 + \alpha_3\beta_1 - \beta_1^2 - \alpha_2 - \alpha_3 + 1 - n)$.

Equating this and $H_0(n; A)$, we conclude that $\hat{\alpha}$ satisfies

- $\epsilon_1^{(0)}(0) = A_0^2 + A_1^2 - A_2^2 - A_3^2 + 2A_0 + 1$,
- $\{\pm(-\alpha_2 + \beta_1 - A_0) - A_0, \pm(-\alpha_3 + \beta_1 - A_0) - A_0\} = \{-A_0 \pm A_2, -A_0 \pm A_3\}$.

These lead to (13), up to the symmetries of $Rc_0(A)$ and $Rc^{(0)}(\hat{\alpha})$.

Case 2: In the same way we have

$$H^{(0)}(n; \hat{\alpha}) = \frac{-\{4(n - A_0)^2 - 1\} \epsilon_2^{(0)}(n) \epsilon_2^{(0)}(n+1)}{4n(n - 2A_0)(n - \alpha_1)(n + \alpha_1 - 2A_0)(n - \alpha_2 + \alpha_3 - A_0)(n + \alpha_2 - \alpha_3 - A_0)},$$

$$\text{where } \epsilon_2^{(0)}(n) = 2(n^2 - 2A_0n + 2A_0^2 - A_0\alpha_1 + 2A_0\alpha_2 + 2A_0\alpha_3)$$

$$- \alpha_1\alpha_2 - \alpha_1\alpha_3 + 2\alpha_2\alpha_3 + \alpha_1 - \alpha_2 - \alpha_3 + 1 - A_0 - n),$$

and the conditions

- $\epsilon_2^{(0)}(0) = A_0^2 + A_1^2 - A_2^2 - A_3^2 + 2A_0 + 1$,
- $\{\pm(\alpha_1 - A_0) - A_0, \pm(\alpha_2 - \alpha_3) - A_0\} = \{-A_0 \pm A_2, -A_0 \pm A_3\}$,

which lead to (14), up to the symmetries of $Rc_0(A)$ and $Rc^{(0)}(\hat{\alpha})$.

These conclude the proof the proposition.

This proposition together with Corollary 10.2 implies

Proposition 10.9.

$$\begin{aligned} (\text{Solutions of } Rc_0(A)) &= \frac{(A_{----})_n (A_{-++})_n}{n! \left(\frac{3}{2} - A_0\right)_n} (\text{Solutions of } Rc^{(0)}(\hat{\alpha}) \text{ with } \hat{\alpha} \text{ as (13)}) \\ &= \frac{(1 - A_0 - A_2)_n (A_{--+})_n (A_{-++})_n}{n! (1 - A_0)_n \left(\frac{3}{2} - A_0\right)_n} (\text{Solutions of } Rc^{(0)}(\hat{\alpha}) \text{ with } \hat{\alpha} \text{ as (14)}). \end{aligned}$$

Picking the solutions with initial condition $C_{-1} = 0, C_0 = 1$ up from the right hand-sides, we get

Theorem 10.10. *The solution $f^{(0,0)}(A; x)$ of $Z(A)$ holomorphic at $x = 0$ (with normalization $f^{(0,0)}(A; 0) = C_0 = 1$) can be expressed as*

$$f^{(0,0)}(A; x)$$

$$(18) \quad = \sum_{n=0}^{\infty} x^n \frac{(A_{----})_n (A_{-++})_n}{n! \left(\frac{3}{2} - A_0\right)_n} {}_4F_3 \left(\begin{matrix} -n, A_0 - n, A_{-++}, A_{-++} \\ A_{++-} - n, A_{+++} - n, 1 - A_0 \end{matrix} ; 1 \right)$$

$$(19) \quad = \sum_{n=0}^{\infty} x^n \frac{(1 - A_0 - A_2)_n (A_{--+})_n (A_{-++})_n}{n! (1 - A_0)_n \left(\frac{3}{2} - A_0\right)_n} {}_4F_3 \left(\begin{matrix} -n, A_0 - A_2 - n, A_{-++}, A_{-++} \\ A_{++-} - n, A_{+++} - n, 1 - A_0 - A_2 \end{matrix} ; 1 \right).$$

It is invariant under the symmetry of $Rc_0(A)$ in Remark 10.3.

REMARK 10.11 (ANOTHER WAY OF DERIVING (19) FROM (18)). By the way, the following identity is known:

$$(20) \quad {}_4F_3 \left(\begin{matrix} -n, a, b, c \\ d, e, f \end{matrix} ; 1 \right) = \frac{(e - a)_n (f - a)_n}{(e)_n (f)_n} {}_4F_3 \left(\begin{matrix} -n, a, d - b, d - c \\ d, a + 1 - n - e, a + 1 - n - f \end{matrix} ; 1 \right),$$

where $a + b + c - n + 1 = d + e + f$ (see Theorem 3.3.3 in [1]). Applying this transformation formula to the expression (18) by putting

$$\left(\begin{matrix} a, b, c \\ d, e, f \end{matrix} \right) = \left(\begin{matrix} A_{-++}, A_0 - n, A_{-++} \\ A_{++-} - n, A_{+++} - n, 1 - A_0 \end{matrix} \right)$$

we get the expression (19).

REMARK 10.12. It is easily seen that

$$(21) \quad {}_2F_1\left(\begin{matrix} \alpha, \beta \\ \gamma \end{matrix}; x\right) {}_2F_1\left(\begin{matrix} a, b \\ c \end{matrix}; x\right) = \sum_{n=0}^{\infty} x^n \frac{(a)_n (b)_n}{(c)_n n!} {}_4F_3\left(\begin{matrix} -n, 1-c-n, \alpha, \beta \\ 1-a-n, 1-b-n, \gamma \end{matrix}; 1\right),$$

which implies

$$\begin{aligned} & {}_2F_1\left(\begin{matrix} A_{-++}, A_{-++} \\ 1-A_0 \end{matrix}; x\right) {}_2F_1\left(\begin{matrix} A_{---}, A_{---} \\ 1-A_0 \end{matrix}; x\right) \\ &= \sum_{n=0}^{\infty} x^n \frac{(A_{---})_n (A_{-++})_n}{n! (1-A_0)_n} {}_4F_3\left(\begin{matrix} -n, A_0-n, A_{-++}, A_{-++} \\ A_{-++}-n, A_{+++}-n, 1-A_0 \end{matrix}; 1\right). \end{aligned}$$

Compare this with (18). It leads the authors to the discovery of the relationship between $\tilde{Z}(A)$ and $L(A)$: $\tilde{Z}(A) = mc_{1/2}(L(A))$ in Theorem 5.2 (see also §10.7).

Let us define the following operation to state the above discovery impressively.

DEFINITION 10.13. The operation $[X^n]$ is defined to pick up the coefficient of X^n from a series $\sum_{n=0}^{\infty} C_n X^n$, that is,

$$[X^n] \sum_{n=0}^{\infty} C_n X^n := C_n.$$

Then, the expression (18) can be rephrased as

$$(22) \quad f^{(0,0)}(A; x) = \sum_{n=0}^{\infty} x^n \frac{(1-A_0)_n}{\left(\frac{3}{2}-A_0\right)_n} [X^n] {}_2F_1\left(\begin{matrix} A_{-++}, A_{-++} \\ 1-A_0 \end{matrix}; X\right) {}_2F_1\left(\begin{matrix} A_{---}, A_{---} \\ 1-A_0 \end{matrix}; X\right).$$

REMARK 10.14. By combining (19) and (21), we can also obtain another expression of $f^{(0,0)}(A; x)$:

$$\begin{aligned} & f^{(0,0)}(A; x) \\ &= \sum_{n=0}^{\infty} x^n \frac{(1-A_0-A_2)_n (1-A_0+A_2)_n}{(1-A_0)_n \left(\frac{3}{2}-A_0\right)_n} [X^n] {}_2F_1\left(\begin{matrix} A_{-++}, A_{-++} \\ 1-A_0-A_2 \end{matrix}; X\right) {}_2F_1\left(\begin{matrix} A_{-++}, A_{-++} \\ 1-A_0+A_2 \end{matrix}; X\right). \end{aligned}$$

10.5. Other local solutions expressed in terms of $f^{(0,0)}$. Recall

$$\tilde{Z}(A) = \text{Ad}\left(x^{-A_0+\frac{1}{2}}\right) Z(A) = x^{-A_0+\frac{1}{2}} \circ Z(A) \circ x^{A_0-\frac{1}{2}}$$

and that $\tilde{Z}(A)$ has symmetries

$$(23) \quad A_j \rightarrow -A_j \quad (j = 0, 1, 2, 3); \quad A_2 \longleftrightarrow A_3;$$

$$(24) \quad (x, A_0, A_1) \longleftrightarrow (1-x, A_1, A_0).$$

In particular, the symmetry $A_0 \rightarrow -A_0$ for $\tilde{Z}(A)$ implies that if $f(A; x)$ is a solution of $Z(A)$, then $x^{-A_0+1/2} f(A; x)$ and $x^{A_0+1/2} f(-A_0, A_1, A_2, A_3; x)$ satisfy $\tilde{Z}(A)$, that is, $x^{2A_0} f(-A_0, A_1, A_2, A_3; x)$ also solves $Z(A)$. Therefore, we get the following:

Proposition 10.15. *Let $f^{(0,2A_0)}(A; x)$ be the normalized local solution of $Z(A)$ at $x = 0$ with exponent $2A_0$. Then, $f^{(0,2A_0)}(A; x)$ is expressible as*

$$f^{(0,2A_0)}(A; x) = x^{2A_0} f^{(0,0)}(-A_0, A_1, A_2, A_3; x).$$

Similarly, the symmetry (24) leads to the following:

Proposition 10.16. *Let $f^{(1,\pm A_1)}(A; x)$ be the normalized local solutions of $Z(A)$ at $x = 1$ with exponents $\frac{1}{2} \pm A_1$, respectively. Then, $f^{(1,\pm A_1)}(A; x)$ are expressed as*

$$f^{(1,\pm A_1)}(A; x) = x^{A_0 - \frac{1}{2}} (1-x)^{\frac{1}{2} \pm A_1} f^{(0,0)}(\mp A_1, A_0, A_2, A_3; 1-x),$$

respectively.

Thus, we have obtained the normalized local solutions at $x = 0$ with exponents $0, 2A_0$, and at $x = 1$ with exponents $1/2 \pm A_1$.

10.6. Local solutions at infinity I: using invariants of the difference equations. In this section, we start to find the normalized local solutions of $Z(A)$ at $x = \infty$. Recall that the local exponents at $x = \infty$ are

$$1 - A_0 \pm A_2, \quad 1 - A_0 \pm A_3.$$

We find the normalized local solution with exponent $1 - A_0 + A_2$. By substituting the expression

$$f^{(\infty, +A_2)}(A; x) := \left(\frac{1}{x}\right)^{1-A_0+A_2} \sum_{n=0}^{\infty} C_n \left(\frac{1}{x}\right)^n, \quad \text{where } C_0 = 1.$$

into $Z(A)$, we see that coefficients C_n satisfy the following 3-term recurrence relation

$$Rc_{\infty}(A) : \quad C_n = \frac{\{2(n+A_2) - 1\}^2 \{2n^2 + 4A_2n - A_0^2 + A_1^2 + A_2^2 - A_3^2 + 1 - 2(n+A_2)\}}{4n(n+2A_2)(n+A_2+A_3)(n+A_2-A_3)} C_{n-1} \\ - \frac{(2n+2A_2-1)(2n+2A_2-3)(n+A_0+A_2-1)(n-A_0+A_2-1)}{4n(n+2A_2)(n+A_2+A_3)(n+A_2-A_3)} C_{n-2}.$$

Thus the invariant $H_{\infty}(n; A)$ of $Rc_{\infty}(A)$ is given by

$$H_{\infty}(n; A) = H_0(n; -A_2, A_1, A_0, A_3).$$

REMARK 10.17 (SYMMETRY). $Rc_{\infty}(A)$ is invariant under $A_j \rightarrow -A_j$ ($j = 0, 1, 3$), not under $A_2 \leftrightarrow A_3$.

Hence, Proposition 10.8 yields the following proposition;

Proposition 10.18. *The two invariants $H^{(0)}(n; \hat{\alpha})$ and $H_{\infty}(n; A)$ of the difference equations $Rc^{(0)}(\hat{\alpha})$ and $Rc_{\infty}(A)$, respectively, agree if and only if*

$$(25) \quad \hat{\alpha} = \begin{pmatrix} 0, \alpha_1, \alpha_2, \alpha_3 \\ \beta_1, \beta_2, \beta_3 \end{pmatrix} = \begin{pmatrix} 0, -A_2, A_{-+++}, A_{+++} \\ A_{-+--}, A_{++-+}, 1 + A_2 \end{pmatrix},$$

$$(26) \quad \text{or} \quad = \begin{pmatrix} 0, -A_0 - A_2, A_{-+++}, A_{-++-} \\ A_{-+--}, A_{++-+}, 1 - A_0 + A_2 \end{pmatrix},$$

$$(27) \quad \text{or} \quad = \begin{pmatrix} 0, -A_2 - A_3, A_{+++-}, A_{-++-} \\ A_{-+--}, A_{++--}, 1 + A_2 - A_3 \end{pmatrix},$$

up to the symmetries of the difference equations.

In a similar way to §10.4, by computing the ratio of the coefficients of the two difference equations $Rc_\infty(A)$ and $Rc^{(0)}(\hat{\alpha})$, we get

Proposition 10.19.

(Solutions of $Rc_\infty(A)$)

$$\begin{aligned} &= \frac{\left(\frac{1}{2} + A_2\right)_n (A_{--+-})_n (A_{+++-})_n}{n! (1 + A_2 - A_3)_n (1 + A_2 + A_3)_n} (\text{Solutions of } Rc^{(0)}(\hat{\alpha}) \text{ with } \hat{\alpha} \text{ as (25)}) \\ &= \frac{\left(\frac{1}{2} + A_2\right)_n (1 - A_0 + A_2)_n (A_{+++-})_n (A_{-++-})_n}{n! (1 + A_2)_n (1 + A_2 - A_3)_n (1 + A_2 + A_3)_n} (\text{Solutions of } Rc^{(0)}(\hat{\alpha}) \text{ with } \hat{\alpha} \text{ as (26)}) \\ &= \frac{\left(\frac{1}{2} + A_2\right)_n (A_{--+-})_n (A_{+++-})_n}{n! (1 + A_2)_n (1 + A_2 + A_3)_n} (\text{Solutions of } Rc^{(0)}(\hat{\alpha}) \text{ with } \hat{\alpha} \text{ as (27)}). \end{aligned}$$

Picking the solutions with initial condition $C_{-1} = 0, C_0 = 1$ up from the right hand-sides, we get

Theorem 10.20. $f^{(\infty, +A_2)}(A; x)$ is given as

$$\begin{aligned} f^{(\infty, +A_2)}(A; x) &= \left(\frac{1}{x}\right)^{1-A_0+A_2} \sum_{n=0}^{\infty} \left(\frac{1}{x}\right)^n \frac{\left(\frac{1}{2} + A_2\right)_n (A_{--+-})_n (A_{+++-})_n}{n! (1 + A_2 - A_3)_n (1 + A_2 + A_3)_n} \\ &\quad \times {}_4F_3\left(\begin{matrix} -n, -A_2 - n, A_{-++}, A_{++-} \\ A_{-+--} - n, A_{++--} - n, 1 + A_2 \end{matrix}; 1\right) \\ &= \left(\frac{1}{x}\right)^{1-A_0+A_2} \sum_{n=0}^{\infty} \left(\frac{1}{x}\right)^n \frac{\left(\frac{1}{2} + A_2\right)_n (1 - A_0 + A_2)_n (A_{+++-})_n (A_{-++-})_n}{n! (1 + A_2)_n (1 + A_2 - A_3)_n (1 + A_2 + A_3)_n} \\ &\quad \times {}_4F_3\left(\begin{matrix} -n, -A_0 - A_2 - n, A_{-++}, A_{-+-} \\ A_{-+--} - n, A_{-++} - n, 1 - A_0 + A_2 \end{matrix}; 1\right) \\ &= \left(\frac{1}{x}\right)^{1-A_0+A_2} \sum_{n=0}^{\infty} \left(\frac{1}{x}\right)^n \frac{\left(\frac{1}{2} + A_2\right)_n (A_{--+-})_n (A_{+++-})_n}{n! (1 + A_2)_n (1 + A_2 + A_3)_n} \\ &\quad \times {}_4F_3\left(\begin{matrix} -n, -A_2 - A_3 - n, A_{++-}, A_{-+-} \\ A_{-+--} - n, A_{++--} - n, 1 + A_2 - A_3 \end{matrix}; 1\right). \end{aligned}$$

The function $x^{1-A_0+A_2} f^{(\infty, +A_2)}(A; x)$ is invariant under the change of parameters $A_j \rightarrow -A_j$ ($j = 0, 1, 3$).

Note that one of these three expressions yield the other two by the help of the transformation formula (20) as stated before. Note also that these expressions are rephrased in terms of products of ${}_2F_1$ using (21):

Theorem 10.21. $f^{(\infty, +A_2)}(A; x)$ is expressible as

$$\begin{aligned}
f^{(\infty, +A_2)}(A; x) &= \left(\frac{1}{x}\right)^{1-A_0+A_2} \sum_{n=0}^{\infty} \left(\frac{1}{x}\right)^n \frac{\left(\frac{1}{2} + A_2\right)_n (1 + A_2)_n}{(1 + A_2 - A_3)_n (1 + A_2 + A_3)_n} \\
&\quad \times [X^n] {}_2F_1\left(\begin{matrix} A_{-+++}, A_{++++} \\ 1 + A_2 \end{matrix}; X\right) {}_2F_1\left(\begin{matrix} A_{+---}, A_{----} \\ 1 + A_2 \end{matrix}; X\right) \\
&= \left(\frac{1}{x}\right)^{1-A_0+A_2} \sum_{n=0}^{\infty} \left(\frac{1}{x}\right)^n \frac{\left(\frac{1}{2} + A_2\right)_n (1 - A_0 + A_2)_n (1 + A_0 + A_2)_n}{(1 + A_2)_n (1 + A_2 - A_3)_n (1 + A_2 + A_3)_n} \\
&\quad \times [X^n] {}_2F_1\left(\begin{matrix} A_{-+++}, A_{-+++} \\ 1 - A_0 + A_2 \end{matrix}; X\right) {}_2F_1\left(\begin{matrix} A_{+---}, A_{+---} \\ 1 + A_0 + A_2 \end{matrix}; X\right) \\
&= \left(\frac{1}{x}\right)^{1-A_0+A_2} \sum_{n=0}^{\infty} \left(\frac{1}{x}\right)^n \frac{\left(\frac{1}{2} + A_2\right)_n}{(1 + A_2)_n} \\
&\quad \times [X^n] {}_2F_1\left(\begin{matrix} A_{++++}, A_{-+++} \\ 1 + A_2 - A_3 \end{matrix}; X\right) {}_2F_1\left(\begin{matrix} A_{+---}, A_{----} \\ 1 + A_2 + A_3 \end{matrix}; X\right).
\end{aligned}$$

The other three local solutions at $x = \infty$ are easily obtained from Theorem 10.20 by recalling symmetries of $Z(A)$

$$A_j \rightarrow -A_j \quad (j = 1, 2, 3) \quad \text{and} \quad A_2 \leftrightarrow A_3.$$

Theorem 10.22. Let $f^{(\infty, -A_2)}(A; x)$, $f^{(\infty, \pm A_3)}(A; x)$ be normalized local solutions of $Z(A)$ at $x = \infty$ with exponents $1 - A_0 - A_2$, $1 - A_0 \pm A_3$, respectively. Then, these can be expressed as

$$\begin{aligned}
f^{(\infty, -A_2)}(A; x) &= f^{(\infty, +A_2)}(A_0, A_1, -A_2, A_3; x), \\
f^{(\infty, \pm A_3)}(A; x) &= f^{(\infty, +A_2)}(A_0, A_1, \pm A_3, A_2; x),
\end{aligned}$$

where $f^{(\infty, +A_2)}(A; x)$ are given in Theorems 10.20 and 10.21.

10.7. Correspondence of solutions via the Riemann-Liouville transformation. For a linear differential operator P , the middle convolution $mc_\mu P$ of P with parameter μ is defined in §5.1 as the linear differential operator $\partial^{-\mu} \circ P \circ \partial^\mu$.

On the other hand, for a function $u(x)$, the notion of Riemann-Liouville transformation of u with parameter μ is defined as the function in x :

$$(I_\gamma^\mu u)(x) := \frac{1}{\Gamma(\mu)} \int_\gamma u(s)(x-s)^{\mu-1} ds,$$

where γ is a cycle.

It is known ([Hara2]) that if u is a solution of P , then the function given by the integral above is a solution of $mc_\mu P$.

If $u(x)$ is given locally around $x = 0$ as

$$u = x^\alpha \sum_{j=0}^{\infty} c_n x^n \quad (\alpha \notin \mathbf{Z}),$$

and if $\mu \notin \mathbf{Z}$, then we can choose γ as a path from 0 to x :

$$(I_{[0,x]}^\mu u)(x) = \frac{1}{\Gamma(\mu)} \int_0^x u(s)(x-s)^{\mu-1} ds.$$

Since we assume $\alpha, \mu \notin \mathbf{Z}$, we can apply the beta function formula to get

$$(I_{[0,x]}^\mu u)(x) = \frac{\Gamma(1+\alpha)}{\Gamma(1+\alpha+\mu)} x^{\alpha+\mu} \sum_{n=0}^{\infty} \frac{(1+\alpha)_n}{(1+\alpha+\mu)_n} c_n x^n.$$

10.8. Partial correspondence of local solutions at $x = 0, 1$.

Recall Theorem 5.2:

$$\tilde{Z}(A) := x^{-A_0+1/2} Z(A) x^{A_0-1/2} = mc_{\frac{1}{2}} L(A).$$

This suggests us to apply the above transformation formula for

$$\alpha = -A_0, \quad \mu = 1/2$$

and for u the product of the two Gauss hypergeometric series multiplied by x^{-A_0} :

$$u = x^{-A_0} \sum c_n x^n, \quad c_n = [X^n]_2 F_1 \left(\begin{matrix} A_{-++} & A_{-+-} \\ 1 - A_0 & \end{matrix}; x \right) {}_2 F_1 \left(\begin{matrix} A_{---} & A_{--+} \\ 1 - A_0 & \end{matrix}; x \right).$$

We get a solution of $\tilde{Z}(A)$:

$$(I_{[0,x]}^{1/2} u)(x) = \frac{\Gamma(1 - A_0)}{\Gamma(3/2 - A_0)} x^{-A_0+1/2} \sum \frac{(1 - A_0)_n}{(3/2 - A_0)_n} c_n x^n,$$

which is, by the definition of $f^{(0,0)}(A, x)$, equal to

$$\frac{\Gamma(1 - A_0)}{\Gamma(3/2 - A_0)} x^{-A_0+1/2} f^{(0,0)}(A, x).$$

This rediscovers the expression (22), and we have

Proposition 10.23. *Via the Riemann-Liouville transformation $I_{[0,x]}^{1/2}$ above, the local solution at $x = 0$ to $L(A)$, the product of two Gauss equation, of exponent $\pm A_0$ is sent to the local solution at $x = 0$ to $\tilde{Z}(A)$ of exponent $\frac{1}{2} \pm A_0$.*

Since the equation is stable under the change

$$(x, A_0, A_1, A_2, A_3) \mapsto (1 - x, A_1, A_0, A_2, A_3)$$

as in Remark 4.5, the happening at $x = 1$ reduces to that at $x = 0$.

10.9. Local solutions at infinity II: using middle convolution. Since the local exponents

$$1 \pm A_2, \quad 1 \pm A_3$$

of $L(A)$ at ∞ are non integral, the corresponding solutions with exponents

$$1/2 \pm A_2, \quad 1/2 \pm A_3$$

of $\tilde{Z}(A)$ are obtained via the Riemann-Liouville transformation with parameter $1/2$ from those of $L(A)$. Just apply to the local solution at infinity

$$u(x) = x^\alpha \sum_{n=-\infty}^0 c_n x^n = \sum_{n=-\infty}^0 c_n x^{n+\alpha} \quad \alpha \in \{-1 \pm A_2, 1 \pm A_3\}$$

to get

$$\frac{\Gamma(\alpha+1)}{\Gamma(\alpha+\frac{3}{2})} x^{\alpha+\frac{1}{2}} \sum_{n=-\infty}^0 c_n \frac{(-\alpha-\frac{1}{2})_n}{(-\alpha)_n} x^n.$$

11. Local solutions of $Z(A)$ at $x = 0$ with exponent $A_0 \pm 1/2$

In this section, local solutions of $Z(A)$ at $x = 0$ with exponent $A_0 \pm \frac{1}{2}$ are constructed.

11.1. Recurrence relation $Rc_1(A)$. The coefficients of a local solution

$$x^{A_0-\frac{1}{2}} \sum_{n=0}^{\infty} C_n x^n$$

satisfy

$$Rc_1(A) : \begin{aligned} C_n &= \frac{(n-1)^2 (2n^2 - 4n - A_0^2 + A_1^2 - A_2^2 - A_3^2 + \frac{5}{2})}{n(n-1)(n+A_0-\frac{1}{2})(n-A_0-\frac{1}{2})} C_{n-1} \\ &\quad - \frac{(n+A_2-\frac{3}{2})(n-A_2-\frac{3}{2})(n+A_3-\frac{3}{2})(n-A_3-\frac{3}{2})}{n(n-1)(n+A_0-\frac{1}{2})(n-A_0-\frac{1}{2})} C_{n-2}. \end{aligned} \quad (n = 2, 3, \dots)$$

For arbitrary given C_0 and C_1 , remaining coefficients $C_n (n \geq 2)$ are uniquely determined. The invariant of $Rc_1(A)$ is given by

$$H_1(n; A) = \frac{-n(n-1)(2n^2 - 4n - A_0^2 + A_1^2 - A_2^2 - A_3^2 + \frac{5}{2})(2n^2 - A_0^2 + A_1^2 - A_2^2 - A_3^2 + \frac{1}{2})}{(n+A_0-\frac{1}{2})(n-A_0-\frac{1}{2})(n+A_2-\frac{1}{2})(n-A_2-\frac{1}{2})(n+A_3-\frac{1}{2})(n-A_3-\frac{1}{2})}.$$

REMARK 11.1 (SYMMETRY). $Rc_1(A)$ is invariant under

$$(28) \quad A_j \rightarrow -A_j \quad (j = 0, 1, 2, 3) \quad \text{and} \quad A_2 \leftrightarrow A_3.$$

11.2. Special values of non-terminating ${}_4F_3$ at $x = 1$. The difference equation $Rc_0(A)$ for holomorphic solutions at $x = 0$ was solved by special values of the *terminating* series ${}_4F_3(*; 1)$, which satisfy the difference equation $Rc^{(0)}(\hat{\alpha})$. The key was to find the parameters $\hat{\alpha}$ so that the invariants of the two difference equations

$$H_0(n; A) \quad \text{and} \quad H^{(0)}(n; \hat{\alpha})$$

agree.

For the difference equation $Rc_1(A)$ for general parameters $A = (A_0, A_1, A_2, A_3)$, we can not find $\hat{\alpha}$ so that the invariants $H_1(n; A)$ and $H^{(0)}(n; \hat{\alpha})$ agree. In other words, the *terminating* series ${}_4F_3(*; 1)$ can not solve our equation $Rc_1(A)$.

We introduce some special values of non-terminating ${}_4F_3$ at 1: We first introduce

$${}_4\tilde{f}_3\left(\begin{matrix} a_0, a_1, a_2, a_3 \\ b_0, b_1, b_2, b_3 \end{matrix}; x\right) := \sum_{n=0}^{\infty} \frac{\Gamma(a_0+n)\Gamma(a_1+n)\Gamma(a_2+n)\Gamma(a_3+n)}{\Gamma(b_0+n)\Gamma(b_1+n)\Gamma(b_2+n)\Gamma(b_3+n)} x^n,$$

and⁹

$${}_4f_3\left(\begin{matrix} a_0, a_1, a_2, a_3 \\ b_1, b_2, b_3 \end{matrix}; x\right) := {}_4\tilde{f}_3\left(\begin{matrix} a_0, a_1, a_2, a_3 \\ 1, b_1, b_2, b_3 \end{matrix}; x\right) = \frac{\Gamma(a_0)\Gamma(a_1)\Gamma(a_2)\Gamma(a_3)}{\Gamma(b_1)\Gamma(b_2)\Gamma(b_3)} {}_4F_3\left(\begin{matrix} a_0, a_1, a_2, a_3 \\ b_1, b_2, b_3 \end{matrix}; x\right).$$

For

$$\boldsymbol{\alpha} = \begin{pmatrix} \alpha_0, \alpha_1, \alpha_2, \alpha_3 \\ \beta_1, \beta_2, \beta_3 \end{pmatrix},$$

we define

$$\begin{aligned} y_0(\boldsymbol{\alpha}) &:= {}_4f_3(\boldsymbol{\alpha}; 1) := {}_4f_3\left(\begin{matrix} \alpha_0, \alpha_1, \alpha_2, \alpha_3 \\ \beta_1, \beta_2, \beta_3 \end{matrix}; 1\right), \\ y_i(\boldsymbol{\alpha}) &:= {}_4\tilde{f}_3\left(\begin{matrix} \alpha_0 + 1 - \beta_i, \alpha_1 + 1 - \beta_i, \alpha_2 + 1 - \beta_i, \alpha_3 + 1 - \beta_i \\ 2 - \beta_i, \beta_1 + 1 - \beta_i, \beta_2 + 1 - \beta_i, \beta_3 + 1 - \beta_i \end{matrix}; 1\right), \quad (i = 1, 2, 3) \\ y_{i+4}(\boldsymbol{\alpha}) &:= -{}_4\tilde{f}_3\left(\begin{matrix} \alpha_i, \alpha_i + 1 - \beta_1, \alpha_i + 1 - \beta_2, \alpha_i + 1 - \beta_3 \\ \alpha_i + 1 - \alpha_0, \alpha_i + 1 - \alpha_1, \alpha_i + 1 - \alpha_2, \alpha_i + 1 - \alpha_3 \end{matrix}; 1\right). \quad (i = 0, 1, 2, 3) \end{aligned}$$

From now on we always assume that the parameters are balanced:

$$(29) \quad \beta_1 + \beta_2 + \beta_3 - \alpha_0 - \alpha_1 - \alpha_2 - \alpha_3 = 1;$$

so all the infinite series above are convergent.

11.3. Difference equation $Rc^{(1)}(\boldsymbol{\alpha})$: an extension of $Rc^{(0)}(\hat{\boldsymbol{\alpha}})$. Set

$$\mathbf{e}_1 = \begin{pmatrix} 1, 0, 0, 0 \\ 1, 0, 0 \end{pmatrix}, \quad \mathbf{e}_2 = \begin{pmatrix} 0, 1, 0, 0 \\ 0, 1, 0 \end{pmatrix}, \quad \mathbf{e}_{12} = \begin{pmatrix} 1, 1, 0, 0 \\ 1, 1, 0 \end{pmatrix}.$$

as in §10.3. We have

Proposition 11.2 ([4]).

$$\begin{aligned} y_i(\boldsymbol{\alpha}) &= U_1^{(1)}(\boldsymbol{\alpha})y_i(\boldsymbol{\alpha} + \mathbf{e}_1) + V_1^{(1)}(\boldsymbol{\alpha})y_i(\boldsymbol{\alpha} + \mathbf{e}_{12}) + \frac{1}{\alpha_0(\beta_3 - \alpha_0 - 1)}, \\ y_i(\boldsymbol{\alpha}) &= U_2^{(1)}(\boldsymbol{\alpha})y_i(\boldsymbol{\alpha} + \mathbf{e}_2) + V_2^{(1)}(\boldsymbol{\alpha})y_i(\boldsymbol{\alpha} + \mathbf{e}_{12}) + \frac{1}{\alpha_1(\beta_3 - \alpha_1 - 1)} \end{aligned}$$

hold for any i ($i = 0, 1, \dots, 7$). Here,

$$\begin{aligned} U_1^{(1)}(\boldsymbol{\alpha}) &= \frac{-(\beta_1 - \alpha_1)(\beta_1 + \beta_2 - \alpha_2 - \alpha_3)}{\alpha_0(\beta_3 - \alpha_0 - 1)}, & V_1^{(1)}(\boldsymbol{\alpha}) &= \frac{-(\beta_2 - \alpha_2)(\beta_2 - \alpha_3)}{\alpha_0(\beta_3 - \alpha_0 - 1)}, \\ U_2^{(1)}(\boldsymbol{\alpha}) &= U_1^{(1)}(\boldsymbol{\alpha})|_{\alpha_0 \leftrightarrow \alpha_1, \beta_1 \leftrightarrow \beta_2}, & V_2^{(1)}(\boldsymbol{\alpha}) &= V_1^{(1)}(\boldsymbol{\alpha})|_{\alpha_0 \leftrightarrow \alpha_1, \beta_1 \leftrightarrow \beta_2}. \end{aligned}$$

As Proposition 10.4 led to Lemma 10.5 and Proposition 10.6, we have

Lemma 11.3. Assume condition (29). Set $(n; \boldsymbol{\alpha}) := -n\mathbf{e}_{12} + \boldsymbol{\alpha}$. $y_i(n; \boldsymbol{\alpha})$ ($i = 0, 1, \dots, 7$) satisfies the non-homogeneous linear difference equation

$$D_n = q_1^{(1)}(n; \boldsymbol{\alpha})D_{n-1} + q_2^{(1)}(n; \boldsymbol{\alpha})D_{n-2} + q_0^{(1)}(n; \boldsymbol{\alpha}),$$

⁹This series is known to be convergent at $x = 1$ if

$$\Re(b_0 + b_1 + b_2 + b_3 - a_0 - a_1 - a_2 - a_3) > 1.$$

where

$$\begin{aligned} q_1^{(1)}(\boldsymbol{\alpha}) &= U_1^{(1)}(\boldsymbol{\alpha})U_2^{(1)}(\boldsymbol{\alpha} + \mathbf{e}_1) + V_1^{(1)}(\boldsymbol{\alpha}) + \frac{U_1^{(1)}(\boldsymbol{\alpha})V_2^{(1)}(\boldsymbol{\alpha} + \mathbf{e}_1)}{U_1^{(1)}(\boldsymbol{\alpha} + \mathbf{e}_{12})}, \\ q_2^{(1)}(\boldsymbol{\alpha}) &= \frac{-U_1^{(1)}(\boldsymbol{\alpha})V_1^{(1)}(\boldsymbol{\alpha} + \mathbf{e}_{12})V_2^{(1)}(\boldsymbol{\alpha} + \mathbf{e}_1)}{U_1^{(1)}(\boldsymbol{\alpha} + \mathbf{e}_{12})}, \\ q_0^{(1)}(\boldsymbol{\alpha}) &= \frac{(\beta_1 - \alpha_2 + 1)(\beta_1 - \alpha_3 + 1)(\beta_1 + \beta_2 - \alpha_2 - \alpha_3)}{\alpha_0\alpha_1(\beta_3 - \alpha_0 - 1)(\beta_3 - \alpha_1 - 1)(\beta_1 + \beta_2 - \alpha_2 - \alpha_3 + 2)} \\ &\quad + \frac{\alpha_0\alpha_1 - \beta_1(\beta_1 + \beta_2 - \alpha_2 - \alpha_3)}{\alpha_0\alpha_1(\beta_3 - \alpha_0 - 1)(\beta_3 - \alpha_1 - 1)}. \end{aligned}$$

Proposition 11.4. *Assume condition (29).*

$$D_{ij}^{(1)}(n; \boldsymbol{\alpha}) := y_i(n; \boldsymbol{\alpha}) - y_j(n; \boldsymbol{\alpha}) \quad i, j \in \{0, 1, 2, \dots, 7\}$$

satisfies the homogeneous linear difference equation

$$Rc^{(1)}(\boldsymbol{\alpha}) : D_n = q_1^{(1)}(n; \boldsymbol{\alpha})D_{n-1} + q_2^{(1)}(n; \boldsymbol{\alpha})D_{n-2}.$$

The invariant of $Rc^{(1)}(\boldsymbol{\alpha})$ is

$$H^{(1)}(n; \boldsymbol{\alpha}) := \frac{q_1^{(1)}(n; \boldsymbol{\alpha})q_1^{(1)}(n+1; \boldsymbol{\alpha})}{q_2^{(1)}(n+1; \boldsymbol{\alpha})}.$$

Note that this invariant is a generalization of the former one:

$$H^{(1)}(n; \boldsymbol{\alpha})|_{\alpha_0=0} = H^{(0)}(n; \hat{\boldsymbol{\alpha}}).$$

REMARK 11.5 (SYMMETRY). The homogeneous linear difference equation $Rc^{(1)}(\boldsymbol{\alpha})$ has symmetries

$$(30) \quad \alpha_0 \leftrightarrow \alpha_1, \quad \alpha_2 \leftrightarrow \alpha_3, \quad \beta_1 \leftrightarrow \beta_2,$$

(31)

$$\boldsymbol{\alpha} \rightarrow (\alpha_0 + 1 - \beta_3, \alpha_1 + 1 - \beta_3, \alpha_2 + 1 - \beta_3, \alpha_3 + 1 - \beta_3, \beta_1 + 1 - \beta_3, \beta_2 + 1 - \beta_3, 2 - \beta_3),$$

(32)

$$\boldsymbol{\alpha} \rightarrow (\alpha_0, \alpha_0 + 1 - \beta_3, \alpha_0 + 1 - \beta_1, \alpha_0 + 1 - \beta_2, \alpha_0 + 1 - \alpha_2, \alpha_0 + 1 - \alpha_3, \alpha_0 + 1 - \alpha_1).$$

11.4. Local solutions at zero II: solving $Rc_1(A)$. We eventually find $\boldsymbol{\alpha}$ to solve $Rc_1(A)$.

Proposition 11.6. *The invariant $H_1(n; A)$ of the equation $Rc_1(A)$ and the invariant $H^{(1)}(n; \boldsymbol{\alpha})$ of the equation $Rc^{(1)}(\boldsymbol{\alpha})$ agree if and only if $\boldsymbol{\alpha} = (\alpha_0, \alpha_1, \alpha_2, \alpha_3; \beta_1, \beta_2, \beta_3)$ is equal to*

$$(33) \quad \boldsymbol{\alpha} = \left(\frac{1}{2}, A_0 + \frac{1}{2}, A_{+---}, A_{++-+}; A_{+---} + \frac{1}{2}, A_{++-+} + \frac{1}{2}, A_0 + 1 \right),$$

$$(34) \quad \text{or} \quad = \left(\frac{1}{2}, A_2 + \frac{1}{2}, A_{--+-}, A_{+-++}; A_{--+-} + \frac{1}{2}, A_{+-++} + \frac{1}{2}, A_2 + 1 \right),$$

$$(35) \quad \text{or} \quad = \left(A_0 + \frac{1}{2}, A_2 + \frac{1}{2}, A_{++-+}, A_{+-+}; A_{++-+} + \frac{1}{2}, A_{++-+} + \frac{1}{2}, A_0 + A_2 + 1 \right),$$

$$(36) \quad \text{or} \quad = \left(A_2 + \frac{1}{2}, A_3 + \frac{1}{2}, A_{+-++}, A_{-++}; A_{-++} + \frac{1}{2}, A_{-++} + \frac{1}{2}, A_2 + A_3 + 1 \right),$$

up to the symmetries (28), (30), (31), (32) of the difference equations.

Proof is parallel to that of Proposition 10.8, and goes as follows: The numerator of $H_1(n; A) - H^{(1)}(n; \alpha)$ is a polynomial in n of degree 14. The coefficient of the top term decomposes as

$$\text{constant} \times (\alpha_0 + \alpha_1 - \beta_1 - \beta_2 + 1 - A_1)(\alpha_0 + \alpha_1 - \beta_1 - \beta_2 + 1 + A_1).$$

If we set $\alpha_0 = -\alpha_1 + \beta_1 + \beta_2 - 1 \pm A_1$, then the second top term decomposes as

$$\text{constant} \times (2A_1 - 1)(2A_1 + 1)(\alpha_2 + \alpha_3 - \beta_1 - \beta_2 + 1).$$

Since we assume A_1 is a free parameter, we have

$$(37) \quad \alpha_2 + \alpha_3 - \beta_1 - \beta_2 + 1 = 0.$$

Eliminating β_2 and β_3 from $H^{(1)}(n; \alpha)$ by making use of the condition (29) and the relation (37), we get an expression

$$H^{(1)}(n; \alpha) = \frac{-\gamma^{(1)} + O\left(\left(n - \frac{1}{2}\right)^2\right)}{\gamma^{(1)} + O\left(\left(n - \frac{1}{2}\right)^2\right)},$$

where

$$\gamma^{(1)} := (2\alpha_0 - 1)^2(2\alpha_1 - 1)^2(2\alpha_2 - 2\beta_1 + 1)^2(2\alpha_3 - 2\beta_1 + 1)^2.$$

Now, assume $\gamma^{(1)} \neq 0$. Then, $H^{(1)}\left(\frac{1}{2}; \alpha\right) = -1$, whereas $H_1\left(\frac{1}{2}; A\right)$ is parameter dependent:

$$H_1\left(\frac{1}{2}; A\right) = \frac{-\left(A_0^2 - A_1^2 + A_2^2 + A_3^2 - 1\right)^2}{4A_0^2 A_2^2 A_3^2}.$$

So it does not happen that $H_1(n; A) = H^{(1)}(n; \alpha)$ in the case $\gamma^{(1)} \neq 0$. Thus $\gamma^{(1)}$ must be 0. Thanks to the symmetry (30) of $Rc^{(1)}(\alpha)$, we have only to consider two cases:

$$\text{Case 1 : } \alpha_0 = \frac{1}{2} \quad \text{and} \quad \text{Case 2 : } \beta_1 = \alpha_2 + \frac{1}{2}.$$

Case 1: Eliminating β_2 and β_3 from $H^{(1)}(n; \alpha)$ by the condition (29) and the relation (37), we get an expression

$$H^{(1)}(n; \alpha) = \frac{-n(n-1)\epsilon_1^{(1)}(n)\epsilon_1^{(1)}(n+1)}{(n-\alpha_1)(n+\alpha_1-1)(n+\alpha_2-\beta_1)(n-\alpha_2+\beta_1-1)(n+\alpha_3-\beta_1)(n-\alpha_3+\beta_1-1)},$$

where $\epsilon_1^{(1)}(n) = 2(n^2 - 2n - \alpha_1\alpha_2 - \alpha_1\alpha_3 + \alpha_2\alpha_3 + \alpha_2\beta_1 + \alpha_3\beta_1 - \beta_1^2 + \alpha_1 - \alpha_2 - \alpha_3 + \beta_1 + 1)$.

Equating this and $H_1(n; A)$, we conclude that α satisfies

$$\bullet \quad \epsilon_1^{(1)}(0) = -A_0^2 + A_1^2 - A_2^2 - A_3^2 + \frac{5}{2},$$

$$\begin{aligned} & \bullet \left\{ \pm \left(\alpha_1 - \frac{1}{2} \right) - \frac{1}{2}, \pm \left(-\alpha_2 + \beta_1 - \frac{1}{2} \right) - \frac{1}{2}, \pm \left(-\alpha_3 + \beta_1 - \frac{1}{2} \right) - \frac{1}{2} \right\} \\ & = \left\{ \pm A_0 - \frac{1}{2}, \pm A_2 - \frac{1}{2}, \pm A_3 - \frac{1}{2} \right\}. \end{aligned}$$

These lead to (33) and (34), up to the symmetries (28) and (30).

Case 2: In the same way we have

$$H^{(1)}(n; \alpha) = \frac{-n(n-1)\epsilon_2^{(1)}(n)\epsilon_2^{(1)}(n+1)}{(n-\alpha_0)(n+\alpha_0-1)(n-\alpha_1)(n+\alpha_1-1)\left(n-\alpha_2+\alpha_3-\frac{1}{2}\right)\left(n+\alpha_2-\alpha_3-\frac{1}{2}\right)},$$

$$\text{where } \epsilon_2^{(1)}(n) = 2\left(n^2 - 2n + \alpha_0\alpha_1 - \alpha_0\alpha_2 - \alpha_0\alpha_3 - \alpha_1\alpha_2 - \alpha_1\alpha_3 + 2\alpha_2\alpha_3 + \frac{\alpha_0}{2} + \frac{\alpha_1}{2} + 1\right),$$

and the conditions

$$\begin{aligned} & \bullet \epsilon_2^{(1)}(0) = -A_0^2 + A_1^2 - A_2^2 - A_3^2 + \frac{5}{2}, \\ & \bullet \left\{ \pm \left(\alpha_0 - \frac{1}{2} \right) - \frac{1}{2}, \pm \left(\alpha_1 - \frac{1}{2} \right) - \frac{1}{2}, \pm (\alpha_2 - \alpha_3) - \frac{1}{2} \right\} \\ & = \left\{ \pm A_0 - \frac{1}{2}, \pm A_2 - \frac{1}{2}, \pm A_3 - \frac{1}{2} \right\}, \end{aligned}$$

which lead to (35) and (36), up to the symmetries (28) and (30). These complete the proof of the proposition.

This proposition together with Corollary 10.2 implies

Proposition 11.7.

(Solutions of $Rc_1(A)$)

$$\begin{aligned} & = \frac{\left(\frac{1}{2}\right)_n}{n!} (\text{Solutions of } Rc^{(1)}(\alpha) \text{ with } \alpha \text{ as (33)}) \\ & = \frac{\left(\frac{1}{2}\right)_n \left(-A_2 + \frac{1}{2}\right)_n \left(A_2 + \frac{1}{2}\right)_n}{n! \left(-A_0 + \frac{1}{2}\right)_n \left(A_0 + \frac{1}{2}\right)_n} (\text{Solutions of } Rc^{(1)}(\alpha) \text{ with } \alpha \text{ as (34)}) \\ & = \frac{\left(-A_2 + \frac{1}{2}\right)_n \left(A_2 + \frac{1}{2}\right)_n}{n! \left(\frac{1}{2}\right)_n} (\text{Solutions of } Rc^{(1)}(\alpha) \text{ with } \alpha \text{ as (35)}) \\ & = \frac{\left(-A_2 + \frac{1}{2}\right)_n \left(A_2 + \frac{1}{2}\right)_n \left(-A_3 + \frac{1}{2}\right)_n \left(A_3 + \frac{1}{2}\right)_n}{n! \left(\frac{1}{2}\right)_n \left(-A_0 + \frac{1}{2}\right)_n \left(A_0 + \frac{1}{2}\right)_n} (\text{Solutions of } Rc^{(1)}(\alpha) \text{ with } \alpha \text{ as (36)}). \end{aligned}$$

So far we got many solutions of the difference equation $Rc_1(A)$, whose solution space is two dimensional. Among these, there are many linearly independent pairs, but it is not so obvious to pick two independent ones. Set

$$W_{1,ij}^{(1)}(n; A) := \left. \frac{\left(\frac{1}{2}\right)_n}{n!} D_{ij}^{(1)}(n; \alpha) \right|_{\alpha=(33)},$$

$$\begin{aligned}
W_{2,ij}^{(1)}(n; A) &:= \frac{\left(\frac{1}{2}\right)_n \left(-A_2 + \frac{1}{2}\right)_n \left(A_2 + \frac{1}{2}\right)_n}{n! \left(-A_0 + \frac{1}{2}\right)_n \left(A_0 + \frac{1}{2}\right)_n} D_{ij}^{(1)}(n; \alpha) \Big|_{\alpha=(34)}, \\
W_{3,ij}^{(1)}(n; A) &:= \frac{\left(-A_2 + \frac{1}{2}\right)_n \left(A_2 + \frac{1}{2}\right)_n}{n! \left(\frac{1}{2}\right)_n} D_{ij}^{(1)}(n; \alpha) \Big|_{\alpha=(35)}, \\
W_{4,ij}^{(1)}(n; A) &:= \frac{\left(-A_2 + \frac{1}{2}\right)_n \left(A_2 + \frac{1}{2}\right)_n \left(-A_3 + \frac{1}{2}\right)_n \left(A_3 + \frac{1}{2}\right)_n}{n! \left(\frac{1}{2}\right)_n \left(-A_0 + \frac{1}{2}\right)_n \left(A_0 + \frac{1}{2}\right)_n} D_{ij}^{(1)}(n; \alpha) \Big|_{\alpha=(36)},
\end{aligned}$$

and

$$\Omega := \{(k, i, j) \mid k = 1, 2, 3, 4; i, j = 0, 1, \dots, 7; i \neq j\}.$$

We have

Theorem 11.8. For $(k, i, j) \in \Omega$,

$$f_{k,ij}^{(0,A_0-1/2)}(A; x) := x^{A_0-\frac{1}{2}} \sum_{n=0}^{\infty} W_{k,ij}^{(1)}(n; A) x^n$$

is a solution of $Z(A)$ at $x = 0$ with exponent $A_0 - \frac{1}{2}$.

Examples of linearly independent pairs:

$$\{f_{1,01}^{(0,A_0-1/2)}, f_{1,02}^{(0,A_0-1/2)}\}, \{f_{1,03}^{(0,A_0-1/2)}, f_{1,54}^{(0,A_0-1/2)}\}, \{f_{2,03}^{(0,A_0-1/2)}, f_{2,54}^{(0,A_0-1/2)}\}.$$

12. Local solutions of $Q(A)$

Let $P_Q(A; x) (= P_Q(A_0, A_1, A_2, A_3; x))$ denote the space of solutions of $Q(A)$ appeared in §6.2. Remark 6.3 implies the following equivalence.

Proposition 12.1.

$$(38) \quad P_Q(A_0, A_1, A_2, A_3; x) = P_Q(\pm A_0, \pm A_1, A_2, \pm A_3; x)$$

$$(39) \quad = P_Q(\pm A_1, \pm A_0, A_2, \pm A_3; 1-x)$$

$$(40) \quad = \left(-\frac{1}{x}\right)^{1+2A_2} P_Q\left(\pm A_3, \pm A_1, A_2, \pm A_0; \frac{1}{x}\right)$$

$$(41) \quad = (1-x)^{-1-2A_2} P_Q\left(\pm A_0, \pm A_3, A_2, \pm A_1; \frac{x}{x-1}\right).$$

To help understand the following propositions, we tabulate several expressions of the Riemann scheme of $Q(A)$:

$$\left\{ \begin{array}{ccc} x=0 & x=1 & x=\infty \\ 0 & 0 & 1+2A_2 \\ -A_0-A_2 & -A_1-A_2 & 1+A_2-A_3 \\ A_0-A_2 & A_1-A_2 & 1+A_2+A_3 \end{array} \right\} = \left(-\frac{1}{x}\right)^{1+2A_2} \left\{ \begin{array}{ccc} \frac{1}{x}=\infty & \frac{1}{x}=1 & \frac{1}{x}=0 \\ 1+2A_2 & 0 & 0 \\ 1+A_2-A_0 & -A_1-A_2 & -A_3-A_2 \\ 1+A_2+A_0 & A_1-A_2 & A_3-A_2 \end{array} \right\}$$

$$= (1-x)^{-(1+2A_2)} \begin{cases} \frac{x}{x-1} = 0 & \frac{x}{x-1} = \infty & \frac{x}{x-1} = 1 \\ 0 & 1+2A_2 & 0 \\ -A_0 - A_2 & 1+A_2 - A_1 & -A_3 - A_2 \\ A_0 - A_2 & 1+A_2 + A_1 & A_3 - A_2 \end{cases}.$$

Thanks to the identities (39) and (40), three linearly independent local solutions at $x = 0$ give those at other singular points. We find a holomorphic solution $f_Q^{(0,0)}(A; x)$ and a solution $f_Q^{(0,\pm)}(A; x)$ of local exponent $\pm A_0 - A_2$ at $x = 0$ as follows.

12.1. Holomorphic solution $f_Q^{(0,0)}(A; x)$ to $Q(A)$ at $x = 0$. Set

$$f_Q^{(0,0)}(A; x) = \sum_{n=0}^{\infty} C_n x^n, \quad C_0 = 1.$$

The coefficients C_n satisfy the recurrence relation

$$RcQ(A) : \quad C_n = \frac{(n+A_2 - \frac{1}{2})(2n^2 + 2(2A_2 - 1)n - A_0^2 + A_1^2 + A_2^2 - A_3^2 - 2A_2 + 1)}{n(n+A_0 + A_2)(n-A_0 + A_2)} C_{n-1} - \frac{(n+2A_2 - 1)(n+A_2 + A_3 - 1)(n+A_2 - A_3 - 1)}{n(n+A_0 + A_2)(n-A_0 + A_2)} C_{n-2},$$

whose invariant will be called $H_Q(n; A)$.

REMARK 12.2 (SYMMETRY). $RcQ(A)$ is invariant under

$$(42) \quad A_j \rightarrow -A_j \quad (j = 0, 1, 3).$$

Proposition 12.3. *Let A be generic. Then, the two invariants $H_Q(n; A)$ and $H^{(0)}(n; \hat{\alpha})$ of the difference equations $RcQ(A)$ and $Rc^{(0)}(\hat{\alpha})$ agree as functions in n if and only if*

$$\begin{aligned} \hat{\alpha} &= \begin{pmatrix} 0, \alpha_1, \alpha_2, \alpha_3 \\ \beta_1, \beta_2, \beta_3 \end{pmatrix} = \begin{pmatrix} 0, A_0 - A_2, A_{+-+}, A_{++-} \\ A_{+--}, A_{-+-}, 1 + A_0 + A_2 \end{pmatrix}, \\ \text{or} \quad &= \begin{pmatrix} 0, -A_2 + A_3, A_{-++}, A_{++-} \\ A_{--+}, A_{-+-}, 1 + A_2 + A_3 \end{pmatrix}, \\ \text{or} \quad &= \begin{pmatrix} 0, -A_2, A_{--+}, A_{-+-} \\ A_{--+}, A_{-+-}, 1 + A_2 \end{pmatrix}, \end{aligned}$$

up to the symmetries of two difference equations, that is, (42) and Remark 10.3.

Doing the same as we got Theorem 10.10 from Proposition 10.8, we can obtain the following theorem:

Theorem 12.4. *The holomorphic solution $f_Q^{(0,0)}(A; x)$ to $Q(A)$ at $x = 0$ has the following expressions:*

$$(43) \quad f_Q^{(0,0)}(A; x) = \sum_{n=0}^{\infty} x^n \frac{(1+2A_2)_n (A_{-++})_n (A_{-+-})_n}{(1+A_2)_n (1-A_0+A_2)_n n!} {}_4F_3 \left(\begin{matrix} -n, A_0 - A_2 - n, A_{-++}, A_{-+-} \\ A_{+--} - n, A_{-+-} - n, 1 + A_0 + A_2 \end{matrix}; 1 \right)$$

$$(44) \quad = \sum_{n=0}^{\infty} x^n \frac{(1+2A_2)_n (1+A_2+A_3)_n (A_{++-})_n (A_{-++})_n}{(1+A_2)_n (1-A_0+A_2)_n (1+A_0+A_2)_n n!}$$

$$(45) \quad \begin{aligned} & \times {}_4F_3 \left(\begin{matrix} -n, -A_2 + A_3 - n, A_{-++}, A_{++-} \\ A_{--+} - n, A_{+--} - n, 1 + A_2 + A_3 \end{matrix}; 1 \right) \\ & = \sum_{n=0}^{\infty} x^n \frac{(1+2A_2)_n (A_{++++})_n (A_{-++})_n}{(1-A_0+A_2)_n (1+A_0+A_2)_n n!} {}_4F_3 \left(\begin{matrix} -n, -A_2 - n, A_{-++}, A_{++-} \\ A_{--+} - n, A_{+--} - n, 1 + A_2 \end{matrix}; 1 \right). \end{aligned}$$

REMARK 12.5. With the help of the transformation formula (20), one of the three (43), (44), (45) implies the other two.

REMARK 12.6. By using the formula (21), the expressions (43), (44), (45) can be written also as

$$(46) \quad \begin{aligned} & f_Q^{(0,0)}(A; x) \\ & = \sum_{n=0}^{\infty} x^n \frac{(1+2A_2)_n}{(1+A_2)_n} [Z^n] {}_2F_1 \left(\begin{matrix} A_{-++}, A_{-++} \\ 1-A_0+A_2 \end{matrix}; Z \right) {}_2F_1 \left(\begin{matrix} A_{-++}, A_{-++} \\ 1+A_0+A_2 \end{matrix}; Z \right) \end{aligned}$$

$$(47) \quad \begin{aligned} & \sum_{n=0}^{\infty} x^n \frac{(1+2A_2)_n (1+A_2+A_3)_n (1+A_2-A_3)_n}{(1+A_2)_n (1-A_0+A_2)_n (1+A_0+A_2)_n} \\ & \times [Z^n] {}_2F_1 \left(\begin{matrix} A_{-++}, A_{-++} \\ 1+A_2-A_3 \end{matrix}; Z \right) {}_2F_1 \left(\begin{matrix} A_{-++}, A_{-++} \\ 1+A_2+A_3 \end{matrix}; Z \right) \end{aligned}$$

$$(48) \quad \begin{aligned} & \sum_{n=0}^{\infty} x^n \frac{(1+A_2)_n (1+2A_2)_n}{(1-A_0+A_2)_n (1+A_0+A_2)_n} [Z^n] {}_2F_1 \left(\begin{matrix} A_{-++}, A_{-++} \\ 1+A_2 \end{matrix}; Z \right) {}_2F_1 \left(\begin{matrix} A_{-++}, A_{-++} \\ 1+A_2 \end{matrix}; Z \right). \end{aligned}$$

REMARK 12.7. Applying the trivial symmetry (38)) to $f_Q^{(0,0)}(A; x)$, we have

$$f_Q^{(0,0)}(A; x) = f_Q^{(0,0)}(\pm A_0, \pm A_1, A_2, \pm A_3; x).$$

12.2. Local solution to $Q(A)$ at $x = 0$ with exponent $A_0 - A_2$. Set

$$f_Q^{(0,+)}(A; x) = x^{A_0 - A_2} \sum_{n=0}^{\infty} C_n x^n, \quad C_0 = 1.$$

The coefficients C_n satisfy the recurrence relation:

$$\begin{aligned} C_n &= \frac{\left(n + A_0 - \frac{1}{2}\right) \left(2n^2 + 2(2A_0 - 1)n + A_0^2 + A_1^2 - A_2^2 - A_3^2 - 2A_0 + 1\right)}{n(n + 2A_0)(n + A_0 - A_2)} C_{n-1} \\ &\quad - \frac{(n + A_0 + A_2 - 1)(n + A_0 + A_3 - 1)(n + A_0 - A_3 - 1)}{n(n + 2A_0)(n + A_0 - A_2)} C_{n-2}. \end{aligned}$$

The same argument as in the previous section leads to

Theorem 12.8. *The local solution $f_Q^{(0,+)}(A; x)$ to $Q(A)$ at $x = 0$ with exponent $A_0 - A_2$ has the following expressions:*

$$(49) \quad \begin{aligned} & f_Q^{(0,+)}(A; x) \\ & = x^{A_0 - A_2} \sum_{n=0}^{\infty} x^n \frac{(1+A_0+A_2)_n}{(1+A_0)_n} [Z^n] {}_2F_1 \left(\begin{matrix} A_{-++}, A_{-++} \\ 1+A_0-A_2 \end{matrix}; Z \right) {}_2F_1 \left(\begin{matrix} A_{++++}, A_{++++} \\ 1+A_0+A_2 \end{matrix}; Z \right) \end{aligned}$$

$$(50) \quad = x^{A_0-A_2} \sum_{n=0}^{\infty} x^n \frac{(1+A_0-A_3)_n (1+A_0+A_3)_n}{(1+A_0)_n (1+A_0-A_2)_n} \\ \times [Z^n] {}_2F_1 \left(\begin{matrix} A_{++-}, A_{++-} \\ 1+A_0+A_3 \end{matrix}; Z \right) {}_2F_1 \left(\begin{matrix} A_{++-}, A_{++-} \\ 1+A_0-A_3 \end{matrix}; Z \right)$$

$$(51) \quad = x^{A_0-A_2} \sum_{n=0}^{\infty} x^n \frac{(1+A_0)_n}{(1+A_0-A_2)_n} [Z^n] {}_2F_1 \left(\begin{matrix} A_{++-}, A_{++-} \\ 1+A_0 \end{matrix}; Z \right) {}_2F_1 \left(\begin{matrix} A_{++-}, A_{++-} \\ 1+A_0 \end{matrix}; Z \right).$$

REMARK 12.9. As in the previous section, the formula (21) changes the expressions (49), (50) and (51) into those in terms of ${}_4F_3(1)$.

REMARK 12.10. Applying the trivial symmetry (38) to $f_Q^{(0,+)}(A; x)$, we have

$$f_Q^{(0,+)}(A; x) = f_Q^{(0,+)}(A_0, \pm A_1, A_2, \pm A_3; x).$$

12.3. Pfaff transforms of the solutions of $Q(A)$. From formula (41) we get the following Pfaff transformations:

Proposition 12.11.

$$f_Q^{(0,0)}(A; x) = (1-x)^{-(1+2A_2)} f_Q^{(0,0)} \left(\pm A_0, \pm A_3, A_2, \pm A_1; \frac{x}{x-1} \right), \\ f_Q^{(0,+)}(A; x) = C \times (1-x)^{-(1+2A_2)} f_Q^{(0,+)} \left(A_0, \pm A_3, A_2, \pm A_1; \frac{x}{x-1} \right),$$

where C is a constant depending on the choice of the branch of $x^{A_0-A_2}$.

12.4. Local solutions of $Q(A)$. Applying the symmetries (38), (39) and (40) to local solutions $f_Q^{(0,0)}(A; x)$ and $f_Q^{(0,+)}(A; x)$ of $Q(A)$, we obtain series expressions of other local solutions.

Proposition 12.12. *Local solutions of $Q(A)$ are tabulated in Table 1.*

13. Local solutions of the Dotsenko-Fateev equation

The Dotsenko-Fateev equation S is obtained (Propositions 6.5) from the equation $Q(A)$ by the change of unknown

$$(52) \quad z = x^{-A_0-A_2} (x-1)^{-A_1-A_2} w,$$

where z : solution of $Q(A)$, w : solution of S , and the parameter change given in Propositions 6.6. So we get expressions of local solutions of Dotsenko-Fateev (D-F for short) equation from those obtained in the previous section.

Table 2 tabulates the local solutions of $Q(A)$ appeared in the previous section, and names $f_{DF}^{(*,*)}(a, b, c, g; x)$ of the corresponding solutions of the Dotsenko-Fateev equation.

Table 1. Names of local solutions of $Q(A)$ and their series expressions

names of solutions	(point, exponent)	series expression
$f_Q^{(0,0)}(A; x)$	(0, 0)	Theorem 12.4, Remark 12.6, 12.7, Proposition 12.11
$f_Q^{(0,+)}(A; x)$	(0, $A_0 - A_2$)	Theorem 12.8, Remark 12.10, Proposition 12.11
$f_Q^{(0,-)}(A; x)$	(0, $-A_0 - A_2$)	$f_Q^{(0,+)}(-A_0, \pm A_1, A_2, \pm A_3; x)$
$f_Q^{(1,0)}(A; x)$	(1, 0)	$f_Q^{(0,0)}(\pm A_1, \pm A_0, A_2, \pm A_3; 1 - x)$
$f_Q^{(1,+)}(A; x)$	(1, $A_1 - A_2$)	$f_Q^{(0,+)}(A_1, \pm A_0, A_2, \pm A_3; 1 - x)$
$f_Q^{(1,-)}(A; x)$	(1, $-A_1 - A_2$)	$f_Q^{(0,+)}(-A_1, \pm A_0, A_2, \pm A_3; 1 - x)$
$f_Q^{(\infty,0)}(A; x)$	$(\infty, 2A_2 + 1)$	$\left(-\frac{1}{x}\right)^{1+2A_2} f_Q^{(0,0)}\left(\pm A_3, \pm A_1, A_2, \pm A_0; \frac{1}{x}\right)$
$f_Q^{(\infty,+)}(A; x)$	$(\infty, A_2 + A_3 + 1)$	$\left(-\frac{1}{x}\right)^{1+2A_2} f_Q^{(0,+)}\left(A_3, \pm A_1, A_2, \pm A_0; \frac{1}{x}\right)$
$f_Q^{(\infty,-)}(A; x)$	$(\infty, A_2 - A_3 + 1)$	$\left(-\frac{1}{x}\right)^{1+2A_2} f_Q^{(0,+)}\left(-A_3, \pm A_1, A_2, \pm A_0; \frac{1}{x}\right)$

Table 2. Names of local solutions of D-F equations corresponding to those of $Q(A)$

solutions of $Q(A)$	(point, exponent)	solutions of D-F equation	(point, exponent)
$f_Q^{(0,0)}(A; x)$	(0, 0)	$f_{DF}^{(0,1)}(a, b, c, g; x)$	(0, $a + c + 1$)
$f_Q^{(0,+)}(A; x)$	(0, $A_0 - A_2$)	$f_{DF}^{(0,2)}(a, b, c, g; x)$	(0, $2a + 2c + g + 2$)
$f_Q^{(0,-)}(A; x)$	(0, $-A_0 - A_2$)	$f_{DF}^{(0,0)}(a, b, c, g; x)$	(0, 0)
$f_Q^{(1,0)}(A; x)$	(1, 0)	$f_{DF}^{(1,1)}(a, b, c, g; x)$	(1, $b + c + 1$)
$f_Q^{(1,+)}(A; x)$	(1, $A_1 - A_2$)	$f_{DF}^{(1,2)}(a, b, c, g; x)$	(1, $2b + 2c + g + 2$)
$f_Q^{(1,-)}(A; x)$	(1, $-A_1 - A_2$)	$f_{DF}^{(1,0)}(a, b, c, g; x)$	(1, 0)
$f_Q^{(\infty,0)}(A; x)$	$(\infty, 2A_2 + 1)$	$f_{DF}^{(\infty,1)}(a, b, c, g; x)$	$(\infty, -a - b - 2c - g - 1)$
$f_Q^{(\infty,+)}(A; x)$	$(\infty, A_2 + A_3 + 1)$	$f_{DF}^{(\infty,0)}(a, b, c, g; x)$	$(\infty, -2c)$
$f_Q^{(\infty,-)}(A; x)$	$(\infty, A_2 - A_3 + 1)$	$f_{DF}^{(\infty,2)}(a, b, c, g; x)$	$(\infty, -2a - 2b - 2c - g - 2)$

13.1. Local solutions of the Dotsenko-Fateev equation at $x = 0$.

Proposition 13.1. (1)

$$\begin{aligned}
 & f_{DF}^{(0,0)}(a, b, c, g; x) / (1 - x)^{b+c+1} \\
 &= \sum_{n=0}^{\infty} x^n \frac{(-a - c - g)_n}{(-a - c - \frac{g}{2})_n} [Z^n] {}_2F_1 \left(\begin{matrix} -c, -a - b - c - \frac{g}{2} - 1 \\ -a - c \end{matrix}; Z \right) {}_2F_1 \left(\begin{matrix} b + 1, -a - \frac{g}{2} \\ -a - c - g \end{matrix}; Z \right) \\
 &= \sum_{n=0}^{\infty} x^n \frac{(-2a - b - c - g - 1)_n (b - c + 1)_n}{(-a - c)_n (-a - c - \frac{g}{2})_n} [Z^n] {}_2F_1 \left(\begin{matrix} -c, -c - \frac{g}{2} \\ b - c + 1 \end{matrix}; Z \right) {}_2F_1 \left(\begin{matrix} -a, -a - \frac{g}{2} \\ -2a - b - c - g - 1 \end{matrix}; Z \right)
 \end{aligned}$$

$$= \sum_{n=0}^{\infty} x^n \frac{(-a-c-\frac{g}{2})_n}{(-a-c)_n} [Z^n]_2 F_1 \left(\begin{matrix} -c, -a-b-c-g-1 \\ -a-c-\frac{g}{2} \end{matrix}; Z \right) {}_2 F_1 \left(\begin{matrix} b+1, -a \\ -a-c-\frac{g}{2} \end{matrix}; Z \right).$$

(2)

$$\begin{aligned} & f_{DF}^{(0,1)}(a, b, c, g; x) / x^{a+c+1} (1-x)^{b+c+1} \\ &= \sum_{n=0}^{\infty} x^n \frac{(1-g)_n}{(1-\frac{g}{2})_n} [Z^n]_2 F_1 \left(\begin{matrix} b+1, -a-\frac{g}{2} \\ -a-c-g \end{matrix}; Z \right) {}_2 F_1 \left(\begin{matrix} a+1, -b-\frac{g}{2} \\ a+c+2 \end{matrix}; Z \right) \\ &= \sum_{n=0}^{\infty} x^n \frac{(1-g)_n (a+b+2)_n (-a-b-g)_n}{(1-\frac{g}{2})_n (-a-c-g)_n (a+c+2)_n} [Z^n]_2 F_1 \left(\begin{matrix} c+1, -a-\frac{g}{2} \\ -a-b-g \end{matrix}; Z \right) {}_2 F_1 \left(\begin{matrix} a+1, -c-\frac{g}{2} \\ a+b+2 \end{matrix}; Z \right) \\ &= \sum_{n=0}^{\infty} x^n \frac{(1-\frac{g}{2})_n (1-g)_n}{(-a-c-g)_n (a+c+2)_n} [Z^n]_2 F_1 \left(\begin{matrix} a+1, -a-b-c-g-1 \\ 1-\frac{g}{2} \end{matrix}; Z \right) {}_2 F_1 \left(\begin{matrix} b+1, c+1 \\ 1-\frac{g}{2} \end{matrix}; Z \right). \end{aligned}$$

(3)

$$f_{DF}^{(0,2)}(a, b, c, g; x) = x^{2a+2c+g+2} f_{DF}^{(0,0)} \left(-c - \frac{g}{2} - 1, a+b+c + \frac{g}{2} + 1, -a - \frac{g}{2} - 1, g; x \right).$$

Proof. By the relation (52) and Proposition 12.12, we get (1) and (3). From the expressions (46), (47), (48) of the holomorphic solution $f_Q^{(0,0)}(A; x)$ of $Q(A)$, we have (2). \square

REMARK 13.2. By using the trivial symmetry (38) of the equation $Q(A)$, we can obtain other expressions of $f_{DF}^{(0,i)}(a, b, c, g; x)$ ($i = 0, 1, 2$), which we omit.

Relation (52) between $Q(A)$ and D-F, and the Pfaff transformation (41) lead to other expressions of $f_{DF}^{(0,0)}$ and $f_{DF}^{(0,1)}$:

Proposition 13.3.

$$(1) f_{DF}^{(0,0)}(a, b, c, g; x) = (1-x)^{2c} f_{DF}^{(0,0)} \left(a, -a-b-c-g-2, c, g; \frac{x}{x-1} \right).$$

$$(2) f_{DF}^{(0,1)}(a, b, c, g; x) = C \times (1-x)^{2c} f_{DF}^{(0,1)} \left(a, -a-b-c-g-2, c, g; \frac{x}{x-1} \right),$$

where C is a constant depending on the choice of a branch around $x=0$; or more precisely,

$$\begin{aligned} & f_{DF}^{(0,1)}(a, b, c, g; x) / x^{a+c+1} (1-x)^{b+c+g} \\ &= \sum_{n=0}^{\infty} \left(\frac{x}{x-1} \right)^n \frac{(1-g)_n}{(1-\frac{g}{2})_n} [Z^n]_2 F_1 \left(\begin{matrix} c+1, -b-\frac{g}{2} \\ a+c+2 \end{matrix}; Z \right) {}_2 F_1 \left(\begin{matrix} b+1, -c-\frac{g}{2} \\ -a-c-g \end{matrix}; Z \right) \\ &= \sum_{n=0}^{\infty} \left(\frac{x}{x-1} \right)^n \frac{(1-g)_n (b+c+2)_n (-b-c-g)_n}{(1-\frac{g}{2})_n (a+c+2)_n (-a-c-g)_n} [Z^n]_2 F_1 \left(\begin{matrix} c+1, -a-\frac{g}{2} \\ b+c+2 \end{matrix}; Z \right) {}_2 F_1 \left(\begin{matrix} a+1, -c-\frac{g}{2} \\ -b-c-g \end{matrix}; Z \right) \\ &= \sum_{n=0}^{\infty} \left(\frac{x}{x-1} \right)^n \frac{(1-\frac{g}{2})_n (1-g)_n}{(a+c+2)_n (-a-c-g)_n} [Z^n]_2 F_1 \left(\begin{matrix} a+1, b+1 \\ 1-\frac{g}{2} \end{matrix}; Z \right) {}_2 F_1 \left(\begin{matrix} c+1, -a-b-c-g-1 \\ 1-\frac{g}{2} \end{matrix}; Z \right). \end{aligned}$$

13.2. Local solutions of the Dotsenko-Fateev equation at $x=1$. By the symmetry of (39) of $Q(A)$ and the relation (52) between $Q(A)$ and D-F, we have

Proposition 13.4.

- (1) $f_{DF}^{(1,0)}(a, b, c, g; x) = f_{DF}^{(0,0)}(b, a, c, g; 1-x).$
- (2) $f_{DF}^{(1,1)}(a, b, c, g; x) = f_{DF}^{(0,1)}(b, a, c, g; 1-x).$
- (3) $f_{DF}^{(1,2)}(a, b, c, g; x) = f_{DF}^{(0,2)}(b, a, c, g; 1-x)$

$$= (1-x)^{2b+2c+g+2} f_{DF}^{(0,0)}\left(-c - \frac{g}{2} - 1, a+b+c + \frac{g}{2} + 1, -b - \frac{g}{2} - 1, g; 1-x\right).$$

13.3. Solutions of the Dotsenko-Fateev equation at $x = \infty$. The relation (40) and the relation (52) between $Q(A)$ and D-F lead to

Proposition 13.5.

- (1) $f_{DF}^{(\infty,0)}(a, b, c, g; x) = \left(-\frac{1}{x}\right)^{-2c} f_{DF}^{(0,0)}\left(-a - b - c - g - 2, b, c, g; \frac{1}{x}\right).$
- (2) $f_{DF}^{(\infty,1)}(a, b, c, g; x) = \left(-\frac{1}{x}\right)^{-2a-2b-2c-g-2} f_{DF}^{(0,1)}\left(a, c, b, g; \frac{1}{x}\right).$
- (3) $f_{DF}^{(\infty,2)}(a, b, c, g; x) = \left(-\frac{1}{x}\right)^{-2a-2b-2c-g-2} f_{DF}^{(0,0)}\left(a, c, b, g; \frac{1}{x}\right).$

ACKNOWLEDGEMENTS. The authors thank D. Zagier for presenting them with the challenging system $Z_3(A)$, $a_1 = a_2 = a_3 = 0$. They also thank N. Takayama for instructing them about computer systems as well as various computational skills, and K. Mimachi for telling them his results on the Dotsenko-Fateev equation.

References

- [1] G.E. Andrews, R. Askey and R. Roy: Special functions, Encyclopedia of Mathematics and its Applications **71**, Cambridge University Press, Cambridge, 1999.
- [2] V.S. Dotsenko and V.A. Fateev: *Conformal algebra and multipoint correlation functions in 2D statistical models*, Nuclear Phys. B **240** (1984), 312–348.
- [3] A. Ebisu: *Invariants of difference equations and transformation formulae for hypergeometric functions*, in preparation.
- [4] A. Ebisu and H. Ochiai: *Difference equations for ${}_4F_3(1)$* , in preparation.
- [5] Y. Haraoka: *Prolongability of ordinary differential equations*, J. Nonlinear. Math. Physics **20**, Supplement 1, 70–84.
- [6] Y. Haraoka: *On Oshima’s middle convolution*, Josai Math. Monographs **12** (2020), 19–51.
- [7] E.J. Wilczynski: Projective Differential Geometry of Curves and Ruled Surfaces, reprinted by Chelsea Pub. Co., Teubner, 1906.
- [8] J.A. Wilson: *Three-term contiguous relations and some new orthogonal polynomials*: in Padé and rational approximation (Proc. Internat. Sympos., Univ. South Florida, Tampa, Fla., 1976), 227–232, Academic Press, New York, 1977.

Akihito Ebisu
Faculty of Information and Computer Science
Chiba Institute of Technology
Chiba 275-0023
Japan
e-mail: akihito.ebisu@p.chibakoudai.jp

Yoshishige Haraoka
Department of Mathematics, Kumamoto University
Kumamoto 860-8555
Japan
e-mail: haraoka@kumamoto-u.ac.jp

Masanobu Kaneko
Department of Mathematics, Kyushu University
Fukuoka 819-0395
Japan
e-mail: mkaneko@math.kyushu-u.ac.jp

Hiroyuki Ochiai
Department of Mathematics, Kyushu University
Fukuoka 819-0395
Japan
e-mail: ochiai@imi.kyushu-u.ac.jp

Takeshi Sasaki
Kobe University
Kobe 657-8501
Japan
e-mail: sasaki@math.kobe-u.ac.jp

Masaaki Yoshida
Kyushu University
Fukuoka 819-0395
Japan
e-mail: myoshida@math.kyushu-u.ac.jp