

Title	On almost M-projectives
Author(s)	Harada, Manabu; Mabuchi, Tuguo
Citation	Osaka Journal of Mathematics. 1989, 26(4), p. 837-848
Version Type	VoR
URL	<a href="https://doi.org/10.18910/9145">https://doi.org/10.18910/9145</a>
rights	
Note	

*Osaka University Knowledge Archive : OUKA*

<https://ir.library.osaka-u.ac.jp/>

Osaka University

## ON ALMOST $M$ -PROJECTIVES

Dedicated to Professor Teruo Kanzaki on his 60th birthday

MANABU HARADA AND TUGUO MABUCHI

(Received February 15, 1989)

We have defined a new concept of almost  $M$ -projectives [7] and given several properties of them [4]. This paper is a continuous work of [4] and [7]. If a module  $M_0$  is  $M_i$ -projective for a finite set of modules  $M_i$ , then  $M_0$  is  $\Sigma \oplus M_i$ -projective [2]. However this fact is not true for almost relative projectives [7]. As far as we know this is one of great differences between relative projectives and almost relative projectives. The main purpose of this paper is to fill this gap. Let  $R$  be a semiperfect ring whose Jacobson radical is nil. When  $M_0$  is a local  $R$ -module and the  $M_i$  are  $R$ -modules whose endomorphism rings are local, we shall give a necessary and sufficient condition for  $M_0$  to be almost  $\Sigma \oplus M_i$ -projective (Theorem 2), which is dual to [3], Theorem. We shall study this problem in [6] when  $R$  is right artinian.

First we take any ring  $R$ . Let  $M_0$  be an  $R$ -module and  $M_1$  an indecomposable and non-hollow  $R$ -module. Then we shall show, in §1, that  $M_0$  is  $M_1$ -projective if  $M_0$  is almost  $M_1$ -projective (Theorem 1). Next we shall assume that  $R$  is semiperfect. In §2 we study almost relative projectivity among local modules. From the results in this section we can understand differences between relative projectives and almost relative projectives. Using those results, we shall give the main theorem above in §3.

### 1. Non cyclic modules

Throughout this paper we always assume that a ring  $R$  is a *semiperfect ring* with identity except in Theorem 1 and every module  $M$  is a unitary right  $R$ -module. We denote the *Jacobson radical*, the *length* of  $M$  by  $J(M)$  and  $|M|$ , respectively.  $e_i$  means always a primitive idempotent in  $R$ . We shall use the same terminologies in [4].

Let  $M$  and  $N$  be  $R$ -modules. For any exact sequence with  $K$  a submodule of  $M$ :

$$(1) \quad \begin{array}{c} M \xrightarrow{v} M/K \rightarrow 0 \\ \quad \quad \quad \uparrow h \\ \quad \quad \quad N \end{array}$$

if either there exists  $\tilde{h}: N \rightarrow M$  with  $\nu\tilde{h}=h$  or there exist a non-zero direct summand  $M_1$  of  $M$  and  $\tilde{h}: M_1 \rightarrow N$  with  $h\tilde{h}=\nu|_{M_1}$ ,  $N$  is called *almost  $M$ -projective* [7]. (If we obtain only the first case, we call  $N$   *$M$ -projective* [2].)

We note the following fact: when  $N$  is almost  $M$ -projective and  $M$  is indecomposable,

(#) if  $h$  in the diegram (1) is not an epimorphism, there exists an  $\tilde{h}: N \rightarrow M$  with  $\nu\tilde{h} = h$ .

The concept of almost relative projectives was introduced in [4] and [7] to study the structure of lifting module [8] and extending module [9]. We refer [4] and [7] for the details.

If every proper submodule of an  $R$ -module  $T$  is small in  $T$ ,  $T$  is called a *hollow module*. In particular if  $T$  is a cyclic hollow, we call  $T$  a *local module*.

First we shall give the following theorem for any ring  $R$ .

**Theorem 1.** *Let  $R$  be any ring. Let  $M$  be a non-hollow and indecomposable  $R$ -module and  $M_0$  an  $R$ -module. If  $M_0$  is almost  $M$ -projective, then  $M_0$  is  $M$ -projective.*

*Proof.* Take any diagram with row exact:

$$(2) \quad \begin{array}{ccccccc} 0 & \rightarrow & K & \rightarrow & M & \xrightarrow{\nu} & M/K & \rightarrow & 0 \\ & & & & & & \uparrow h & & \\ & & & & & & M_0 & & \end{array}$$

First assume that  $K$  is not small in  $M$ . Then there exists a submodule  $K_1$  in  $M$  such that  $K_1 \neq M$  and  $M=K_1+K$ . Now we obtain a derived diagram from the above:

$$\begin{array}{ccccccc} 0 & \rightarrow & K_1 \cap K & \rightarrow & M & \xrightarrow{\nu'} & M/(K_1 \cap K) & \approx & M/K \oplus M/K_1 & \rightarrow & 0 \\ & & & & & & \uparrow h+0 & & & & \\ & & & & & & M_0 & & & & \end{array}$$

Since  $M/K_1 \neq 0$ , by assumption there exists  $\tilde{h}: M_0 \rightarrow M$  such that  $h=\pi\nu'\tilde{h}=\nu\tilde{h}'$ , where  $\pi: M/(K_1 \cap K) \rightarrow M/K$  is the projection, and hence  $\pi\nu'=\nu$  from the construction. Therefore we may assume that  $K$  is small in  $M$ . Since  $M$  is not hollow, there exists two proper submodules  $K_1, K_2$  of  $M$  with  $M=K_1+K_2$ . We may assume  $K_1 \supset K$  for  $i=1, 2$ , since  $K$  is small in  $M$ . Then we obtain as above a derived diagram:

$$\begin{array}{ccccccc} M & \xrightarrow{\nu} & M/K & \xrightarrow{\nu'} & M/(K_1 \cap K_2) & \approx & M/K_1 \oplus M/K_2 & \rightarrow & 0 \\ & & & & \uparrow \nu'h & & & & \\ & & & & M_0 & & & & \end{array}$$

Let  $\pi_1$  be the projection of  $M/(K_1 \cap K_2)$  onto  $M/K_1$  and  $i_1$  the inclusion of  $M/K_1$

into  $M/(K_1 \cap K_2)$ . Then  $i_1 \pi_1 \nu' h: M_0 \rightarrow M/(K_1 \cap K_2)$  is not an epimorphism. Hence there exists  $\tilde{h}: M_0 \rightarrow M$  with  $\nu' \nu \tilde{h} = i_1 \pi_1 \nu' h$ , and so  $(h - \nu \tilde{h})(M_0) \subset \ker(\nu' - i_1 \pi_1 \nu') = K_1/K$ . Accordingly we have a diagram

$$\begin{array}{ccc} M & \xrightarrow{\nu} & M/K \rightarrow 0 \\ & & \uparrow h - \nu \tilde{h} \\ & & M_0 \end{array}$$

and  $(h - \nu \tilde{h})(M_0) \subset K_1/K \subsetneq M/K$ . Hence there exists again  $\tilde{h}_1: M_0 \rightarrow M$  with  $\nu \tilde{h}_1 = h - \nu \tilde{h}$ , and so  $h = \nu(\tilde{h} + \tilde{h}_1)$ . Therefore  $M_0$  is  $M$ -projective.

**Corollary 1.** *Let  $R$  be semiperfect, and let  $M$  be an indecomposable module not isomorphic to any local modules  $eR/A$  and  $M_0$  an  $R$ -module such that  $M_0 J$  is small in  $M_0$ . Then  $M_0$  is  $M$ -projective if  $M_0$  is almost  $M$ -projective, where  $e$  is a primitive idempotent in  $R$  and  $J$  is the Jacobson radical of  $R$ .*

Proof. If  $M$  is not hollow,  $M_0$  is  $M$ -projective by Theorem 1. Hence we assume that  $M$  is hollow. If the  $h$  in the proof of Theorem 1 is an epimorphism,  $M/K \cong (M/K)J$  by assumption. Hence since  $M$  is hollow,  $M/K$  is local and so  $M$  is also local, which is a contradiction. Accordingly  $h$  is always not an epimorphism, and hence  $M_0$  is  $M$ -projective.

Let  $Z$  be the ring of integers and  $p$  a prime. Then  $E(Z/p)$ , the injective hull of  $Z/p$ , is a uniserial  $Z_p$ -module and hence  $E(Z/p)$  is almost  $E(Z/p)$ -projective. However  $E(Z/p)$  is not  $E(Z/p)$ -projective. Hence we need the assumption on  $M_0$  in Corollary 1.

From Corollary 1 if  $M_0$  is almost  $M$ -projective, but not  $M$ -projective for an indecomposable module  $M$ ,  $M$  must be a local module  $eR/B$  whenever  $M_0$  is finitely generated, where  $e$  is a primitive idempotent.

**Proposition 1.** *Let  $M_0$  be as in Corollary 1 and  $M_1$  an indecomposable module. Assume that  $M_0$  is almost  $M_1$ -projective. Then  $M_0$  is  $M_1$ -projective if and only if either  $M_1$  is not of a form  $eR/A$  or  $M_1 \cong eR/A$  and any homomorphism:  $M_0/J(M_0) \rightarrow M_1/J(M_1)$  is liftable to an element  $f: M_0 \rightarrow M_1$ .*

Proof. We assume "if" part. If  $M_1 \cong eR/A$ ,  $M_0$  is  $M_1$ -projective by Corollary 1. Hence suppose  $M_1 = eR/A$  and put  $\bar{M}_0 = M_0/M_0 J = \sum_i \oplus e_i \bar{R}$ . If  $e \not\cong e_i$  for all  $i$ , there are no epimorphisms  $h': M_0 \rightarrow M_1/K_1$ , where  $K_1 \subsetneq M_1$ . Hence  $M_0$  is  $M_1$ -projective. Assume that  $e \cong e_1$ . Take a diagram:

$$\begin{array}{ccc} M_1 & \xrightarrow{\nu} & M_1/K_1 \rightarrow 0 \\ & & \uparrow h \\ & & M_0 \end{array}$$

If  $h$  is an epimorphism, then  $h$  induces an epimorphism  $\bar{h}: M_0/J(M_0) \rightarrow M_1/J(M_1)$ . By assumption there exists  $h_1: M_0 \rightarrow M_1$  such that  $\bar{h}_1 = \bar{h}$ , i.e.,  $(\nu h_1 - h)(M_0) \subset$

$(M_1/K_1)J$ . Hence there exists  $\tilde{h}: M_0 \rightarrow M_1$  with  $v\tilde{h}=h$  from (#). If  $h$  is not an epimorphism, by (#) we obtain always  $\tilde{h}'$  similar to the above  $\tilde{h}$ . Hence  $M_0$  is  $M_1$ -projective. “only if” part is clear.

**Corollary 2.** *Let  $M_0$  and  $M_1$  be as in Proposition 1. If  $M_0$  is almost  $M_1$ -projective but not  $M_1$ -projective, there exists a homomorphism  $\tilde{h}: M_1 \rightarrow M_0$  which induces a monomorphism of  $M_1/J(M_1)$  into  $M_0/J(M_0)$ .*

Proof. Since  $M_0$  is not  $M_1$ -projective, we have an epimorphism  $h: M_0/J(M_0) \rightarrow M_1/J(M_1)$ , which is not liftable, by Proposition 1. Hence there exists the desired homomorphism  $\tilde{h}: M_1 \rightarrow M_0$ .

**2. Local modules**

We shall study almost relative projectives among local modules. We recall here the definition of the lifting property of simple modules modulo radical (briefly 1.p.s.m) [5]. Let  $T_1$  and  $T_2$  be local modules. If for any simple submodule  $U$  in  $T_1/J(T_1) \oplus T_2/J(T_2)$  there exists a direct summand  $T'$  of  $T=T_1 \oplus T_2$  such that  $T' + (J(T_1) \oplus J(T_2)) / (J(T_1) \oplus J(T_2)) = U$ , then we say that  $T$  has the 1.p.s.m.. This is equivalent to the following: for every element  $f$  in  $\text{Hom}_R(T_1/J(T_1), T_2/J(T_2))$  is liftable to an element in  $\text{Hom}_R(T_1, T_2)$  or so is  $f^{-1}$  to an element in  $\text{Hom}_R(T_2, T_1)$ , provided  $|T_1|$  and  $|T_2|$  are finite. Now in this paper we call the latter equivalent property the 1.p.s.m. even if  $|T_i|$  is infinite.

Let  $A, B$  be right ideals in  $eR$ . If  $eR/B$  is epimorphic to  $eR/A$ , there exists a unit  $v$  in  $eRe$  such that  $vB \subset A$  and  $eR/B \approx eR/vB$ . We denote this situation by  $B \lesssim A$ .

**Proposition 2.** *Let  $R$  be a semi-perfect ring and  $A, B$  right ideals in  $eR$  such that either  $eR/A$  or  $eR/B$  is noetherian. Then  $eR/A$  is almost  $eR/B$ -projective if and only if  $eJeA \subset B$  and  $eR/A \oplus eR/B$  has the 1.p.s.m.. In this case if  $eR/A$  is not  $eR/B$ -projective, then  $eR/B$  is  $eR/A$ -projective.*

Proof. If  $eR/A$  is almost  $eR/B$ -projective,  $eJeA \subset B$  by [4], Proposition 2 and  $eR/A \oplus eR/B$  has the 1.p.s.m. by definition. Conversely if  $eR/A \oplus eR/B$  has the 1.p.s.m., then 1)  $eR/B$  is epimorphic to  $eR/A$  or 2)  $eR/A$  is epimorphic to  $eR/B$  by definition. In either case we may assume 1)  $A \supset B$  or 2)  $A \subset B$  by the remark above (note that  $veJe = eJe = eJev$ ).

Case 1) Assume  $eJeA \subset B \subset A$ . Take the diagram (2), where  $M_0 = eR/A$  and  $M = eR/B$ . If  $h$  is not an epimorphism,  $h$  is given by an element  $j$  in  $eJe$ . Since  $jA \subset B$ ,  $h$  is liftable to an  $\tilde{h} = j_i: eR/A \rightarrow eR/B$ , where  $j_i$  is the left-sided multiplication of  $j$ . Next we assume that  $h$  is an epimorphism. Then  $h$  is given by a unit  $u$  in  $eRe$ . Since  $eR/A \oplus eR/B$  has the 1.p.s.m. and either  $eR/A$  or  $eR/B$  is noetherian, there exists a unit  $u'$  in  $eRe$  such that  $u^{-1} \equiv u' \pmod{eJe}$  and  $u'B \subset A$ . Put  $u' = u^{-1} + j'$ ;  $j' \in eJe$ . Then  $A \supset u'B = (u^{-1} + j')B$  and  $j'B \subset j'A \subset B \subset A$ .

Hence  $u^{-1}B \subset A$ . Putting  $\tilde{h}=(u^{-1})_1$ ,  $h\tilde{h}=\nu$ . Hence  $eR/A$  is almost  $eR/B$ -projective.

Case 2) We can show in the same manner that  $eR/A$  is  $eR/B$ -projective. Finally assume that  $eR/A$  is not  $eR/B$ -projective. Then we may assume  $B \subset A$  by Corollary 2. Further  $eJeB (\subset eJeA \subset B) \subset A$ . Hence  $eR/B$  is almost  $eR/A$ -projective by the first statement. While  $B \subset A$  implies that  $eR/B$  is  $eR/A$ -projective by Corollary 2.

We shall apply the above proposition to a particular case, e.g. an algebra over an algebraically closed field.

**Lemma 1.** *Let  $M_0=eR/A$  and  $M_1=eR/B$ . Then  $M_0$  is  $M_1$ -projective if and only if for any generator  $a_0=a_0e$  of  $M_0$  (resp.  $a_1=a_1e$  of  $M_1$ ), a mapping  $a_0 \rightarrow a_1$  gives us an epimorphism of  $M_0$  onto  $M_1$ .*

Proof. Since  $a_i e = a_i$  ( $i=0, 1$ ),  $a_i$  is a unit in  $eRe$ . The last statement of the lemma is equivalent to  $\{x \in eR \mid a_0 x \in A, \text{ i.e. } x \in a_0^{-1}A\} \subset \{x \in eR \mid a_1 x \in B, \text{ i.e. } x \in a_1^{-1}B\}$ . Hence  $A \subset B$  by taking  $a_0 = a_1 = e$  and  $uA \subset B$  for any unit  $u$  in  $eRe$  by taking  $a_0 = u^{-1}$  and  $a_1 = e$ . Let  $j$  be any element in  $eJe$ . Then  $(e+j)A \subset B$  and  $eA \subset B$  from the above. Hence  $jA \subset B$ . Therefore  $eReA \subset B$ , and so  $M_0$  is  $M_1$ -projective by [1], p. 22, Exercise 4. The converse is clear from the above and [1].

**Proposition 3.** *Let  $M$  be an  $R$ -module and  $M_0=eR/A$ . Then  $M_0$  is  $M$ -projective if and only if for any  $m=me$  in  $M$  and any generator  $a_0=a_0e$  of  $M_0$ , a mapping  $a_0 \rightarrow m$  gives us an epimorphism of  $M_0$  onto  $mR$ .*

Proof. If  $M_0$  is  $M$ -projective, then  $M_0$  is  $N$ -projective for any submodule  $N$  of  $M$  by definition. Hence we obtain "only if" part from Lemma 1, since  $mR \approx eR/B$  for some  $B$ . Conversely take  $m=me$  in  $M$  with  $h(e+A)=\nu(m)$  in the diagram (2). Since there exists  $\tilde{h}: M_0 \rightarrow mR (\subset M)$  with  $\tilde{h}(e+A)=m$  by assumption,  $\nu\tilde{h}=h$ .

From the above result we shall define a new concept. Let  $M_0=eR/A$  be a local module. An  $R$ -module  $N$  is called *locally generated* by  $M_0$  if every cyclic submodule  $nR$  of  $N$  with  $ne=n$  is a homomorphic image of  $M_0$ .

Now we assume that  $eJ/B$  is locally generated by  $eR/A$ . For any element  $x$  in  $eJe$  we obtain an epimorphism  $f: eR/A \rightarrow (xR+B)/B \subset eR/B$ . Then  $f(e+A) = xr+B$  and  $r$  is a unit in  $eRe$  and there exists  $y$  in  $eRe$  such that  $yA \subset B$  and  $y \equiv xr \pmod{B}$ . Put  $y = xr + b; b \in B$ . Then  $B \supset yA = (xr+b)A$ . Hence since  $bA \subset B, xrA \subset B$ . Therefore  $eJ/B$  is locally generated by  $eR/A$  if and only if

(3) for any element  $x$  in  $eJe$ , there exists a unit  $u_x$  in  $eRe$  such that  $xu_x A \subset B$ .

If  $eJeA \subset B$ , (3) is trivially satisfied.

**Lemma 2.** *Let  $R$  be a right artinian ring and assume that  $eR/A \oplus eR/A$  has the 1.p.s.m.. Then 1): for  $B \subset eR$   $eR/A$  is almost  $eR/B$ -projective if and only*

if i)  $eJ/B$  is locally generated by  $eR/A$  and ii)  $A \lesssim B$  or  $A \gtrsim B$ . 2): For an  $R$ -module  $M$   $eR/A$  is  $M$ -projective if and only if  $M$  is locally generated by  $eR/A$ .

Proof. 1) We assume that  $eR/A$  is almost  $eR/B$ -projective. Then i) and ii) are clear from Proposition 2 and the remark after (3). Conversely we assume i) and ii). We shall show  $eJ^i eA \subset B$  for each  $i$  by induction on  $i$ . Assume  $eJ^{i+1} eA \subset B$  and take an element  $x$  in  $eJ^i e - eJ^{i+1} e$ . Then from (3) there exists a unit  $r$  in  $eRe$  such that  $xrA \subset B$ . By assumption; 1.p.s.m.

(4)  $r = u + j$ ;  $u$  is a unit in  $eRe$  with  $uA = A$  and  $j \in eJe$ .

Then  $B \subset xrA = (xu + xj)A$  and  $xj \in eJ^{i+1} e$ . Hence  $xA = xuA \subset B$  by induction hypothesis, and so  $eJeA \subset B$  by taking  $i=1$ . From ii) we may assume  $A \subset B$  or  $A \supset B$ . Hence it is clear that  $eR/A \oplus eR/B$  has the 1.p.s.m. for  $eR/A \oplus eR/A$  does. Therefore  $eR/A$  is almost  $eR/B$ -projective by Proposition 2.

2) Assume that  $M$  is locally generated by  $eR/A$ . Let  $m$  be an element in  $M$  with  $me = m$ . Then  $mR \approx eR/B$  for some  $B$ . Now we shall show that  $eR/A$  is  $eR/B$ -projective. Since  $eR/B$  is locally generated by  $eR/A$ , (3) holds for any element in  $eRe$  from the argument given before (3). Hence the observation after (4) shows  $eReA \subset B$  and hence  $eR/A$  is  $eR/B$ -projective by [1]. Accordingly  $eR/A$  is  $M$ -projective by Lemma 1 and Proposition 3. The converse is clear from Proposition 3.

**Proposition 4.** Let  $R$  be a right artinian ring and  $M$  an  $R$ -module. We assume that  $eR/A \oplus eR/A$  has the 1.p.s.m.. Then  $eR/A$  is almost  $M$ -projective if and only if for any element  $m = me$  in  $M$ , we obtain one of the following:

1) If  $mR$  is not a direct summand of  $M$ , then  $mR$  is a homomorphic image of  $eR/A$ .

2) If  $mR$  is a direct summand of  $M$ , then either  $mR$  is a homomorphic image of  $eR/A$  or  $eR/A$  is that of  $mR$ .

Proof. Assume that  $eR/A$  is almost  $M$ -projective. Let  $m = me$  be in  $M$  and  $mR$  not a direct summand of  $M$ . We shall show that  $eR/A$  is  $mR$ -projective. Consider a diagram with  $K$  a submodule of  $mR$ :

$$\begin{array}{ccc} mR & \xrightarrow{v} & mR/K \rightarrow 0 \\ & & \uparrow h \\ & & eR/A \end{array}$$

Then we obtain a derived diagram

$$\begin{array}{ccc} M & \xrightarrow{v_M} & M/K \rightarrow 0 \\ \cup & & \cup \\ mR & \xrightarrow{v} & mR/K \rightarrow 0 \\ & & \uparrow h \\ & & eR/A. \end{array}$$

Since  $eR/A$  is almost  $M$ -projective, a) there exists  $\tilde{h}: eR/A \rightarrow M$  with  $\nu_M \tilde{h} = h$  or b) there exist a direct summand  $M_1$  of  $M$  and  $\tilde{h}: M_1 \rightarrow eR/A$  with  $h\tilde{h} = \nu_M|_{M_1}$ . Assume b). Since  $\nu_M(M_1) \subset h(eR/A) \subset \nu(mR)$  and  $K \subset mR$ ,  $M_1 \subset mR$ . Hence  $M_1 = mR$  for  $mR$  is hollow, which contradicts the initial assumption. Therefore, if  $mR$  is not a direct summand of  $M$ , we always obtain the case a). Then since  $\nu_M(\tilde{h}(eR/A)) \subset h(eR/A) \subset \nu(mR)$ ,  $\tilde{h}(eR/A) \subset mR$ . Hence  $eR/A$  is  $mR$ -projective, whence  $mR$  is a homomorphic image of  $eR/A$  by Proposition 3. Next we assume that  $mR$  is a direct summand of  $M$ . Then  $eR/A$  is almost  $mR$ -projective by definition. Then we obtain 2) from Lemma 2-1)-ii). Conversely assume 1) and 2). Take any diagram with  $K \subset M$ :

$$\begin{array}{ccc} M & \xrightarrow{\nu} & M/K \rightarrow 0 \\ & & \uparrow h \\ & & eR/A \end{array}$$

and put  $h(\tilde{e}) = \nu(m)$  for some  $m = me \in M$ , where  $\tilde{e} = e + A$  in  $eR/A$ . Assume that  $mR$  is not a direct summand of  $M$ . Then since  $mR$  is hollow,  $m'R$  is not a direct summand of  $M$  for any  $m' (=m'e)$  in  $mR$ . Accordingly  $mR$  is locally generated by  $eR/A$  from 1) and so  $eR/A$  is  $mR$ -projective by Lemma 2-2). Hence there exists a homomorphism  $\tilde{h}: eR/A \rightarrow mR \subset M$  with  $\tilde{h}(\tilde{e}) = m$  by Lemma 1. Therefore  $\nu\tilde{h} = h$ . Assume the case 2). Since  $mR$  is a local module, any proper submodule of  $mR$  is not a direct summand of  $M$ . Hence  $eR/A$  is almost  $mR$ -projective by 1) and Lemma 2-1). Take the derived diagram from the above one

$$\begin{array}{ccc} mR & \xrightarrow{\nu} & mR/(K \cap mR) \rightarrow 0 \\ & & \uparrow h \\ & & eR/A \end{array}$$

Since  $eR/A$  is almost  $mR$ -projective, we obtain an  $\tilde{h}: eR/A \rightarrow mR$  (or  $mR \rightarrow eR/A$ ) which makes the above diagram commutative. Noting that  $mR$  is a direct summand of  $M$ , we know that  $eR/A$  is almost  $M$ -projective.

REMARK. We don not need Lemma 2 in the first half of the proof of Proposition 4, and hence it shows the following fact: Let  $R$  be semiperfect and  $eR/A$  almost  $M$ -projective. Then for  $m = me$  in  $M$  such that  $mR$  is not a direct summand of  $M$ ,  $eR/A$  is  $mR$ -projective.

We note that if  $R$  is an algebra over an algebraically closed field of finite dimension,  $eR/A \oplus eR/A$  has always the l.p.s.m.. Further Lemma 2 and Proposition 4 are not true without the assumption: l.p.s.m. of  $eR/A \oplus eR/A$  (see the next examples).

EXAMPLE 1. Let  $L \supset K$  be fields with  $L = aK \oplus bK$ . Put  $R_1 = L \oplus uL$ , a trivial extension with  $J(R_1) = 0 \oplus uL$  and  $V = xL \oplus yL$ , a vector space over  $L$ .



Set

$$R = \begin{pmatrix} R_1 & V \\ 0 & R \end{pmatrix}$$

with  $(ud) x=yd$  and  $uy=0; d \in L$ . Put  $A=(0 x(aK) \oplus yL)$  and  $B=(0 y(aK))$ . Then for  $e=e_{11}$   $eJe=J(R_1)$ ,  $eJeA \subset B$  and for any  $c'=cu(\neq 0)$  in  $eJe c'c^{-1}A \subset B$  ((3)), and hence  $eJ/B$  is locally generated by  $eR/A$ . Further  $eR/A \oplus eR/B$  has the l.p.s.m.. However  $eR/A$  is not almost  $eR/B$ -projective for  $eJeA \subset B$ .

2. Put  $A'=(0 x(aK) \oplus y(aK))$ . Since  $eJeB=0$  and  $A' \supset B$ ,  $eR/A'$  is locally generated by  $eR/B$ . However  $eR/B$  is not  $eR/A'$ -projective.

### 3. Direct sums

Let  $M_0, M_1$  and  $M_2$  be indecomposable modules and let  $M_0$  be almost  $M_i$ -projective for  $i=1, 2$ . In this section we shall study a condition under which  $M_0$  is almost  $M_1 \oplus M_2$ -projective, when  $M_0$  is cyclic. This is dual to [3], Theorem. We note that if  $M_0$  is almost  $M_1 \oplus M_2$ -projective, then  $M_0$  is almost  $M_i$ -projective for  $i=1, 2$  by definition. If  $\text{End}_R(M)$  is a local ring, we say  $M$  is an *l.e. module*.

**Proposition 5.** *Let  $M_0$  be a finitely generated  $R$ -module and let  $M_1$  be a local and l.e. module  $e_1R/A_1$  and  $M_2$  an l.e. module. Assume that i)  $M_0$  is almost  $M_1 \oplus M_2$ -projective, but  $M_0$  is not  $M_1$ -projective, and ii) for any  $m(\neq 0)$  in  $M_2$  with  $me_1=m$  we take any isomorphism  $f: M_1/M_1J \approx mR/mJ$ . Then  $f$  (or  $f^{-1}$  if  $M_2=mR$ ) is liftable to  $f': M_1 \rightarrow M_2$  (or  $f': M_2 \rightarrow M_1$ ).*

Proof. Since  $M_0$  is almost  $M_1$ -projective but not  $M_1$ -projective, there exist a maximal submodule  $B$  of  $M_0$  and an isomorphism  $g: M_0/B \rightarrow M_1/J(M_1)$  which is not liftable to an element:  $M_0 \rightarrow M_1$  (cf. the proof of Proposition 1). Let  $f: M_1/J(M_1) \rightarrow mR/mJ$  be the given isomorphism and take a diagram:

$$\begin{array}{ccc} M_1 \oplus M_2 & \xrightarrow{\nu_1 + \nu_2} & M_1/J(M_1) \oplus M_2/mJ \rightarrow 0 \\ & & \uparrow h \\ & & M_0/B \\ & & \uparrow \nu_0 \\ & & M_0, \end{array}$$

where  $h=g+fg$ . Since  $M_0$  is almost  $M_1 \oplus M_2$ -projective, either there exists  $\tilde{h}: M_0 \rightarrow M_1 \oplus M_2$  with  $(\nu_1 + \nu_2)\tilde{h} = h\nu_0$  or there exist a non-zero direct summand  $N$  of  $M_1 \oplus M_2$  and  $\tilde{h}: N \rightarrow M_0$  with  $h\nu_0\tilde{h} = (\nu_1 + \nu_2)|_N$ . If the former occurs, taking the projection of  $M_1 \oplus M_2$  onto  $M_1$ , we have a contradiction to the choice of  $g$ . Hence we should obtain the latter. We may assume that  $N$  is an indecomposable module. Since  $N$  has the exchange property by assumption

$$M_1 \oplus M_2 = N \oplus M_1 \text{ or } = N \oplus M_2 .$$

The first case: Let  $x_2$  be any element in  $M_2$ . Then

$$x_2 = n + x_1; n \in N, x_1 \in M_1 \text{ and } n = y_1 + y_2, y_i \in M_i .$$

Hence  $x_2 = y_2$  and  $x_1 = -y_1$ . Put  $z = \nu_0 \tilde{h}(n)$ , and  $\nu_1(y_1) = g(z)$ ,  $\nu_2(y_2) = fg(z)$ , i.e.,  $\nu_2(x_2) = f(\nu_1(-x_1))$ . Then  $M_2/mJ = f(M_1/J(M_1)) = mR/mJ$ . Accordingly,  $M_2 = mR$  and  $-\pi|_{M_2}: M_2 \rightarrow M_1$  is a lifted element of  $f^{-1}$ , where  $\pi: N \oplus M_1 \rightarrow M_1$  is the projection. We obtain a similar result for the second case.

**Lemma 3.** *Let  $\{M_i\}_{i=1}^n$  be a set of indecomposable  $R$ -modules and let  $N$  and  $M_0$  be  $R$ -modules. Assume that  $M_0$  is almost  $M_i$ -projective for all  $i$  and  $N$ -projective. Take a diagram with row exact:*

$$0 \rightarrow K \rightarrow (\sum_i \oplus M_i) \oplus N \xrightarrow{\nu} H \rightarrow 0$$

$\uparrow h$   
 $M_0$

If there exists a small submodule  $T$  in  $\sum_i \oplus M_i$  such that  $h(M_0) \subset \nu(T \oplus N)$ , then there exists  $\tilde{h}: M_0 \rightarrow (\sum_i \oplus M_i) \oplus N$  with  $\nu \tilde{h} = h$ .

Proof. Put  $M^* = \sum_i \oplus M_i \oplus N$  and  $\pi_1: M^* \rightarrow \sum_i \oplus M_i$ ,  $\pi_2: M^* \rightarrow N$  the projections. Further put  $K^i = \pi_i(M^*)$  for  $i=1, 2$ . We can derive the following diagram (cf. [4]):

$$\sum_i \oplus M_i \xrightarrow{\nu'} (\sum_i \oplus M_i)/K^1 \rightarrow 0$$

$\uparrow \pi'_1 h$   
 $M_0$

where  $\pi'_1: H \xrightarrow{\nu^*} M^*/(K^1 \oplus K^2) \rightarrow (\sum_i \oplus M_i)/K^1$  is the projection (we note that  $K \subset (K^1 \oplus K^2)$  and  $H = M^*/K$ , and so we obtain the natural epimorphism  $\nu^*$ ). From the assumption  $\pi'_1 h(M_0)$  is small in  $(\sum_i \oplus M_i)/K^1$ . Hence there exists  $\tilde{h}_1: M_0 \rightarrow \sum_i \oplus M_i$  with  $\nu' \tilde{h}_1 = \pi'_1 h$  by [4], Lemma 1. Since  $M_0$  is  $N$ -projective, we obtain the desired homomorphism from the remark before [4], Lemma 1.

The following theorem is dual to [3], Theorem and will be generalized in [6] to a case where  $M_0$  is a finitely generated module, when  $R$  is right artinian.

**Theorem 2.** *Assume that  $R$  is a semiperfect ring and  $J$  is nil. Let  $\{M_i\}_{i=1}^n$  be a set of l.e. modules and  $M_0$  a local module  $e_0R/A_0$ . Then the following are equivalent:*

- 1)  $M_0$  is almost  $\sum_{i=1}^n \oplus M_i$ -projective.
- 2) The following are fulfilled:
  - i)  $M_0$  is almost  $M_i$ -projective for all  $i \geq 1$ .
  - ii) If  $M_0$  is not  $M_k$ -projective for  $k=i$  and  $j$ , then  $M_i \oplus M_j$  has the l.p.s.m. (in this case  $M_i \approx e_0R/A_i$ ,  $M_j \approx e_0R/A_j$ ).

Proof. 2)→1) We may assume that there exists an integer  $m$  such that  $M_0$  is  $M_i$ -projective for all  $i > m$  and  $M_0$  is not  $M_j$ -projective for all  $j \leq m$  and hence all  $M_j$  ( $j \leq m$ ) are local modules  $e_0R/A_j$  by Corollary 1. Take a diagram with row exact:

$$(5) \quad \begin{array}{ccccccc} 0 & \rightarrow & K & \rightarrow & M = \sum \oplus M_i & \xrightarrow{\nu} & M/K \rightarrow 0 \\ & & & & & \uparrow h & \\ & & & & & M_0 = eR/A & \end{array}$$

Let  $h(\tilde{e}_0) = (\sum a_i) + K$ ;  $a_i \in M_i$ , where  $\tilde{e}_0 = e_0 + A$  in  $M_0$ . We may assume  $a_i e_0 = a_i$ . We show that

there exists  $\tilde{h}: M_0 \rightarrow M$  (or there exist a non-zero direct summand  $N$  of  $M$  and a homomorphism  $\tilde{h}: N \rightarrow M_0$ ) such that  $\nu\tilde{h} = h$  (or  $h\tilde{h} = \nu|N$ ).

If  $a_i \in J(M_i)$  for all  $(m \geq i) i \geq 1$ , there exists  $\tilde{h}: M_0 \rightarrow M$  such that  $\nu\tilde{h} = h$  by Lemma 3. Hence we assume that there exists an integer  $k$  such that  $a_j \in J(M_j)$  for  $(m \geq j) j > k$  and  $a_{j'} \notin J(M_{j'})$  for  $1 \leq j' \leq k$ . Then  $a_{j'}$  is a generator of  $M_{j'}$ , since  $M_{j'}$  is local. Now  $M_0$  is not  $M_t$ -projective for  $t=1, s \leq k$ , and so  $M_1 \oplus M_s$  has the l.p.s.m. by assumption. Hence there exists  $f: M_1 \rightarrow M_s$  (or  $M_s \rightarrow M_1$ ) such that  $f(a_1) = a_s + a_s j_s$  (or  $f(a_s) = a_1 + a_1 j_s$ ) for some  $j_s \in J$ . We take a new decomposition  $M = M_1(f) \oplus M_s \oplus \sum_{i \neq 1, s} \oplus M_i$  (or  $M_1 \oplus M_s(f) \oplus \sum_{i \neq 1, s} \oplus M_i$ ), where  $M_1(f) = \{x + f(x) \mid x \in M_1\} \subset M_1 \oplus M_s$ . Then  $a_1 + a_s = (a_1 + f(a_1)) + (a_s - f(a_1)) = (a_1 + f(a_1)) - a_s j_s$  and  $(a_1 + f(a_1)) \in M_1(f)$ ,  $a_s j_s \in J(M_s)$  (similar for another case). Hence iterating this argument, we remain ourselves a case  $k=1$ , i.e.,  $M_0$  is not  $M_1$ -projective and  $a_t \in J(M_t)$  for all  $(m \geq t) t > 1$ . Since  $a_t R \subset J(M_t)$  for  $1 < t \leq m$ , there exists

$$(6) \quad \tilde{h}_t: M_0 \rightarrow M_t \quad \text{such that} \quad \tilde{h}_t(\tilde{e}_0) = a_t, \quad (n \geq t > 1)$$

by Lemma 1 and Remark in §2. On the other hand, consider  $f_1: M_0/J(M_0) \approx M_1/J(M_1)$  ( $f_1(e_0 + J(M_0)) = a_1 + J(M_1)$ ). Since  $M_0 \oplus M_1$  has the l.p.s.m. by assumption i) and Proposition 2, there exists  $\tilde{h}_1: M_1 \rightarrow M_0$  (or  $M_0 \rightarrow M_1$ ) such that  $\tilde{h}_1(a_1) \equiv \tilde{e}_0 \pmod{J(M_0)}$  (or  $\tilde{h}_1(\tilde{e}_0) \equiv a_1 \pmod{J(M_1) = a_1 J}$ ), i.e.,

$$(7) \quad \tilde{h}_1(a_1) = \tilde{e}_0 + \tilde{e}_0 j_0; \quad j_0 \in J, \quad \text{or}$$

$$(7') \quad \tilde{h}_1(\tilde{e}_0) = a_1 + a_1 j_1; \quad j_1 \in J$$

Case (7'): Put  $g = \sum_{t=1}^n \tilde{h}_t: M_0 \rightarrow M$  and  $h' = h - \nu g$ . Then  $h'(\tilde{e}_0) = \nu(a_1 j_1)$  and  $a_1 j_1 \in J(M_1)$ . Hence there exists  $h^*: M_0 \rightarrow M$  such that  $\nu h^* = h'$  by Lemma 3 and so  $h = \nu(g + h^*)$ .

Case (7): Now put  $g = (\sum_{t \geq 2} \tilde{h}_t) \tilde{h}_1: M_1 \rightarrow \sum_{t \geq 2} \oplus M_t$ . Then  $g(a_1) = \sum_{t \geq 2} \tilde{h}_t(\tilde{e}_0) + \sum_{t \geq 2} \tilde{h}_t(\tilde{e}_0) j_0 = \sum_{t \geq 2} a_t + \sum_{t \geq 2} a_t j_0$ . Taking a decomposition  $M = M_1(g) \oplus \sum_{t \geq 2} \oplus M_t$ ,  $\sum_{t=1}^n a_t = a_1 + g(a_1) - \sum_{t \geq 2} a_t j_0$  and  $a_1 + g(a_1) \in M_1(g)$ ,  $a_t j_0 \in M_t$  ( $t > 1$ ). Similarly to (6), we obtain by Lemma 1 and Remark in §2.

$$(8) \quad \tilde{h}'_t: M_0 \rightarrow M_t \quad \text{with} \quad \tilde{h}'_t(\tilde{e}) = a_t j_0 \quad (n \geq t > 1).$$

While since  $M_1 \approx M_1(g)$  ( $a_1 \mapsto a_1 + g(a_1)$ ), from (7) we obtain  $\tilde{h}'_1: M_1(g) \rightarrow M_0$  with

$$(9) \quad \tilde{h}'_1(a_1 + g(a_1)) = \tilde{e}_0 + \tilde{e} j_0.$$

Put  $g' = (\sum_{t \geq 2} \tilde{h}'_t) \tilde{h}'_1$ , and  $\sum_{i=1}^n \oplus M_i = (M_1(g)) (g') \oplus \sum_{t \geq 2} \oplus M_t$ ,  $\sum_{s+1}^n a_t = (a_1 + g(a_1) + g'(a_1 + g(a_1)) - \sum_{t \geq 2} a_t j_0^2)$ . Repeating this procedure we obtain the final decomposition  $M = M'_1 \oplus M_2 \oplus \dots \oplus M_n$  and  $\sum_{i=1}^n a_i \in M'_1 \approx M_1$ , since  $J$  is nil. Thus we have derived the following diagram from (5):

$$\begin{array}{ccc} M'_1 & \xrightarrow{\nu} & \nu(M'_1) \rightarrow 0 \\ & \uparrow h & \\ & & M_0 \end{array}$$

Therefore there exists  $\tilde{h}: M_0 \rightarrow M'_1$  (or  $\tilde{h}: M'_1 \rightarrow M_0$ ) such that  $\nu \tilde{h} = h$  (or  $h \tilde{h} = \nu | M'_1$ ).

1)  $\rightarrow$  2) (cf. [7]).  $M_0$  is almost  $M_t$ -projective for all  $t$  by definition. Let  $M_i$  and  $M_j$  be as in 2)-ii). Since  $M_0$  is almost  $M_1 \oplus M_2$ -projective as above,  $M_1 \oplus M_2$  has the l.p.s.m. by Proposition 5.

We shall show in [6] that Theorem 2 is useful when we characterize right Nakayama rings in terms of almost relative projectives.

---

### References

- [1] T. Albu and C. Nastasescu: Relative finiteness in module theory, Monographs Textbooks Pure and Appl. Math. 84, Marcel Dekker, Inc., New York and Basel
- [2] G. Azumaya, F. Mbuntum and K. Varadarajan: *On  $M$ -projective and  $M$ -injective modules*, Pacific J. Math. **59** (1975), 9–16.
- [3] Y. Baba: *Note on almost  $M$ -injectives*, Osaka J. Math. **26** (1989), 687–698.
- [4] ——— and M. Harada: *On almost  $M$ -projectives and almost  $M$ -injectives*, Tukuba J. Math **14**, in press.
- [5] M. Harada: *On modules with lifting properties*, Osaka J. Math. **19** (1982), 189–221.
- [6] ———: *On almost relative projectives over perfect rings*, to appear.
- [7] ——— and A. Tozaki: *Almost  $M$ -projectives and Nakayama rings*, J. Algebra **122** (1989), 447–474.
- [8] K. Oshiro: *Semiperfect modules and quasi-semiperfect modules*, Osaka J. Math. **20** (1983), 337–373.
- [9] ———: *Continuous modules and quasi-continuous modules*, Osaka J. Math. **20** (1983), 681–694.

Manabu HARADA  
Department of Mathematics

Osaka City University  
Sugimoto-3, Sumiyoshi-ku  
558 Osaka, Japan

Tuguo MABUCHI  
Ohtemae High School  
Higashi-ku  
541 Osaka, Japan