

Title	Does Treatment Stigma among adolescents with autism spectrum disorder and their guardians affect the effectiveness of cognitive behavioral therapy?A secondary of a randomized controlled trial
Author(s)	大下,惠美子
Citation	大阪大学, 2023, 博士論文
Version Type	VoR
URL	https://doi.org/10.18910/92004
rights	
Note	

The University of Osaka Institutional Knowledge Archive : OUKA

https://ir.library.osaka-u.ac.jp/

The University of Osaka

Does Treatment Stigma among adolescents with autism spectrum disorder and their guardians affect the effectiveness of cognitive behavioral therapy? A secondary of a randomized controlled trial

(児童思春期の自閉スペクトラム症児および保護者の治療スティグマは認知行動療法の 効果に影響があるか?ランダム化比較試験の二次解析)

> 大阪大学大学院 大阪大学・金沢大学・浜松医科大学・千葉大学・福井大学 連合小児発達学研究科 小児発達学専攻

大下惠美子

2023年2月 博士学位論文



Does Treatment Stigma among Adolescents with Autistic Spectrum Disorder and Their Guardians Affect the Effectiveness of Cognitive Behavioral Therapy? A Secondary Analysis of a Randomized Controlled Trial

Emiko Oshita^{1), 2), *}, Fumiyo Oshima^{1), 2)}, Minako Hongo^{1), 2)}, Siqing Guan^{2), 3)}, Yusuke Nitta^{2), 3)}, and Eiji Shimizu^{1), 2), 4)}

1) United Graduate School of Child Development, Osaka University, Osaka, Japan

- 2) Research Center for Child Mental Development, Chiba University, Chiba, Japan
- 3) Graduate School of Human Sciences, Waseda University, Chiba, Japan

- University, Chiba, Japan
- * E-mail: u794531b@ecs.osaka-u.ac.jp/ccda0334@chiba-u.jp

Summary

This study explored the effect of treatment stigma on the intervention effects of cognitive behavioral therapy (CBT) on adolescents with autism spectrum disorder (ASD) and their guardians. Secondary analyses were conducted using data from a randomized controlled trial, and only the intervention group (adolescents with ASD [n=23; 16 boys and 7 girls; mean age=12.8±2.2 years] and their guardians [n=23; 22 women and 1 man; mean age=44.8±6.1 years]) was used. We used the Japanese version of the Barriers to Access to Care Evaluation scale version 3 (BACEv3) to assess treatment stigma. Correlations between the BACEv3 scores at pre-intervention and the difference scores for pre-to-post-intervention in each measure were analyzed using Spearman's rank correlation coefficient. Adolescents' BACEv3 scores at pre-intervention were not correlated with difference scores in any measure. However, guardians' BACEv3 scores at pre-intervention demonstrated a significant positive correlation with adolescents' knowledge of ASD and a significant negative correlation with the communication of adaptive behaviors. Interventions through CBT are expected to be somewhat effective, regardless of adolescents' or their guardians' pre-treatment stigma. Further research is required to clarify whether treatment stigma in adolescents with ASD and their guardians affects CBT intervention effectiveness,

Key words: treatment stigma, adolescents with ASD, guardians, cognitive behavioral therapy, autism spectrum disorder

Background

Autism spectrum disorder (ASD) is a group of neurodevelopmental disorders characterized by "impaired interpersonal communication" and "limited and repetitive behavioral patterns."⁽¹⁾ Its heterogeneity ranges across the full range of IQ and language functioning, communication, and social and behavioral deficits.⁽²⁾ Adolescents with ASD may have poor adaptive behaviors even with normal cognitive abilities.⁽³⁾⁻⁽⁷⁾ Therefore, although high cognitive ability cases require diagnosis and support,⁽⁸⁾ stigma can still hinder support and treatment.^{(9), (10)} In addition, the stigma held by the person concerned hinders support-seeking behavior and, therefore, does not allow them to receive adequate support.^{(11), (12)} Stigma refers to prejudice against "undesirable kinds of attributes."⁽¹³⁾ Self-stigma refers to prejudice and discrimination where the party stigmatizes themselves, sees themselves as less worthy, and limits their own behavior.⁽¹⁴⁾ Treatment stigma refers to prejudice and discrimination against receiving treatment and support,⁽¹⁵⁾ and it is a stigma held by the person concerned. It is a factor that inhibits seeking help from treatment and support agencies.^{(16), (17)}

In contrast, regarding adolescents, Calzada et al. suggest that a diagnosis of ASD may have a negative impact on exacerbating stigma, with a

⁴⁾ Department of Cognitive Behavioral Physiology, School of Medicine, Chiba



lack of interest in the diagnosis on the part of adolescents with ASD being shown to exacerbate treatment stigma.⁽¹⁸⁾ Therefore, treatment stigma may be a disincentive for support. Guardians of adolescents with ASD also experience treatment stigma. Studies on parents of children with ASD show that the more the parents internalize stigma, the more their help-seeking behavior decreases.⁽¹⁹⁾ Parents of children with ASD are more likely to experience severe psychological distress than those of children with other developmental disabilities; they are also less likely to receive adequate support.^{(20), (21)}

Through studying the relationship between autism and stigma, Japanese researchers developed and implemented an awareness program about public stigma regarding disabilities; it resulted in reduced public stigma and improved knowledge.⁽²²⁾ Online classes on ASD improved college students' knowledge of ASD and reduced the stigma against it.^{(23), (24)} However, these studies were solely about the public stigma around ASD, and no study has thus far examined treatment stigma in relation to ASD. Therefore, it is valuable to examine the impact of treatment stigma.

Cognitive behavioral therapy (CBT) has been effective for both children and adults with ASD.⁽²⁵⁾⁻⁽²⁷⁾ CBT reduces anxiety symptoms in young people and patients with ASD,⁽²⁸⁾⁻⁽³⁰⁾ decreases depression and stress symptoms,⁽³¹⁾ and is effective in treating comorbid obsessive-compulsive disorder.⁽³²⁾ The Program for Education and Enrichment of Relationship Skills (PEERS) for Young Adults improves the social skills of adolescents and young adults.⁽³³⁾⁻⁽³⁵⁾

Psychoeducation about disability increases knowledge and positive attitudes toward disabilities.⁽³⁶⁾ These programs teach adolescents about their disability, increasing their knowledge and awareness of ASD.⁽³⁷⁾ Public stigma around ASD also undermines the mental health and quality of life of the guardians of adolescents with ASD,⁽³⁸⁾⁻⁽⁴⁰⁾ whereas a decrease in guardianship treatment stigma contributes to improved parental mental health.^{(41), (42)} We developed Aware and Care for my Autistic Traits (ACAT)—a program that integrates CBT and psychoeducation to promote understanding of ASD for diagnosed adolescents and their guardians. We implemented it for two years, beginning in 2018.⁽⁴³⁾ The results showed that the intervention group of children with ASD had significantly improved awareness and mental health and reduced treatment stigma. Awareness of ASD significantly improved in the guardian intervention group as well, as compared to the control group.⁽⁴⁴⁾

Our study purpose was to determine whether the treatment stigma of adolescents with ASD and their guardians, who participated in the ACAT clinical trials as the intervention group, affected the effectiveness of the trial intervention. Two hypotheses were formulated in this study. First, adolescents with ASD who have higher treatment stigma have poorer outcomes from ACAT interventions than those who have lower treatment stigma. Second, when guardians' treatment stigma is high, they are less likely to have positive outcomes from ACAT interventions. High guardian treatment stigma also has a negative effect on the effectiveness of ACAT for adolescents with ASD.

Methods

Our study was a secondary analysis that used data from the ACAT clinical trial⁽⁴⁴⁾—a six-week psychosocial education program by CBT at Chiba and Fukushima University.

Participants

The intervention group constituted groups receiving both outpatient psychotherapy with a psychiatrist and ACAT in the randomization, consisting of adolescents with ASD, n=23 (16 boys and 7 girls; mean age=12.8 years, SD=2.2); and their guardians, n=23 (22 women and 1 man;

mean age=44.8 years, SD=6.1). A control group was created from groups receiving only outpatient psychotherapy with a psychiatrist and observed for six weeks, n=22 (17 boys and 5 girls; mean age=14.1 years, SD=2.0); and their guardians, n=22 (all women; mean age=46.6 years, SD=4.5). An *a priori* power calculation showed that a sample of at least 23 participants was required to determine the α and power levels using a correlation analyses (α =0.05, power=0.75, correlations among repeated measures=0.5).

Ethics approval

This study was approved by the Chiba University Clinical Research Ethics Review Committee (CRB3180015, no. G29027). It was conducted in accordance with the Declaration of Helsinki. This clinical trial has been registered with the University Hospitals Medical Information Network (UMIN 000029851). Informed written consent was obtained from a parent or legally authorized representative.

Relevant standards

For adolescents with ASD, the inclusion criteria were as follows: (1) those diagnosed and notified by a psychiatrist with ASD as their primary diagnosis and those who met the criteria for ASD on either the Autism Diagnostic Observation Schedule (ADOS-2) or Autism Diagnostic Interview-Revised (ADI-R), (2) those with a verbal IQ of 90 or higher on the Wechsler Intelligence Test, (3) those with a score of moderate (some need) or higher on the Strengths and Difficulties Questionnaires (SDQ), and (4) those aged 10–17 years at the time of obtaining consent.

For guardians, the inclusion criteria were as follows: (1) those living with adolescents with ASD as a guardian or grandparent and acting as guardian and (2) those able to attend CBT once a week accompanied by the adolescent with ASD.

For adolescents with ASD, the exclusion crite-

ria were as follows: (1) those with active suicidal intent, (2) those with active repetitive antisocial behavior, and (3) those with an active severe degenerative physical disorder that could disrupt CBT.

For guardians, the exclusion criteria were as follows: (1) those who met the diagnostic criteria for psychiatric disorders measured by the Mini International Neuropsychiatric Interview; (2) those who were unable to accompany the patient to the sessions; and (3) those who were diagnosed with any of the following: organic brain disorder (including dementia), psychotic disorder, bipolar disorder, drug dependence, or other mental severe disorders complicated by psychosis, imminent suicide risk, antisocial behavior, or significant progressive physical illness.

The demographic characteristics of participants are summarized in Tables 1 and 2.

Materials

This study was exploratory and based on the ACAT intervention study; we sought to determine the association between the measures used and treatment stigma.

We used the Autism Knowledge Quiz-Child (AKQ-C) and the Autism Knowledge Quiz-Guardians/Parents (AKQ-P),⁽³⁷⁾ a five-item structured interview that measures understanding of ASD characteristics, and 15 sections of the item knowledge quiz. Both adolescent self-administered and guardian self-administered questions were to be answered. An interviewer asked adolescents about their strengths and weaknesses related to ASD characteristics and scored for ASD-related responses (e.g., they could be scored if they answered, "I am good at rule-based behavior"). Guardians were asked about their child's strengths and weaknesses. The Barriers to Access to Care Evaluation scale version 3 (BACEv3) Japanese Version⁽⁴⁵⁾ is a 30-item guestionnaire that measures treatment stigma related to mental health and has a four-point



Demographic variables	Total N=45	TAU group n=22	COMB group n=23	р
Sex				.5589
Male	33	17 (51.5)	16 (48.5)	
Female	12	5 (41.7)	7 (58.3)	
School attendance				.295
Every day	29	14 (48.3)	15 (51.7)	
Less than one-third of the semester	1	1 (10.0)	0 (0.0)	
No attendance	13	5 (38.5)	8 (61.5)	
Non-student	2	2 (100.0)	0 (0.0)	
Academic background (%)				.139
Primary-high school students	43 (95.6)	20 (46.5)	23 (53.3)	
Junior high school graduates	2(4.4)	2 (100.0)	0 (0.0)	
Total years of education				.0541
Mean (<i>SD</i>)	8.0 (2.2)	8.6 (1.9)	7.3(2.3)	
Support and consideration at school				.117
Support classes	5(11.1)	1 (20.0)	4 (80.0)	
Special classes	2(4.4)	0 (0.0)	2 (100.0)	
Consideration	2(4.4)	2 (100.0)	0 (0.0)	
Non-support consideration	36 (80.0)	19 (52.8)	17 (47.2)	
Age at diagnosis (in years)				.878
Mean (<i>SD</i>)	10.3 (4.1)	10.4 (4.9)	10.2(3.3)	
Complications				.668
Without	26 (57.8)	12 (46.2)	14 (53.8)	
With	19 (42.2)	10 (52.6)	9 (47.4)	
ADHD				.466
Without ADHD	3 (15.8)	1 (33.3)	2(66.7)	
With ADHD	16 (84.2)	9(56.3)	7 (43.8)	
LD				.213
Without LD	15 (78.9)	9 (60.0)	6 (40.0)	
With LD	4 (21.1)	1 (25.0)	3 (75.0)	

Table 1 Demographics of adolescents with ASD in COMB/TAU groups (from the original study).

COMB, Intervention group, groups receiving both outpatient psychotherapy with a psychiatrist and ACAT; TAU, Control group, groups receiving only outpatient psychotherapy.

Complications are presented verbatim from the original study. ADHD: Attention-Deficit/Hyperactivity Disorder, LD: Specific Learning Disorder. Note that the mean (SD) is not given in the original study.

scale ranging from "*not at all*" to "*quite a bit*." Both adolescents and guardians were asked to respond. The scale consists of two subscales: treatment stigma and non-stigma. The mean value in the previous study was 1.31.⁽⁴⁵⁾

The Depression Self-Rating Scale for Children (DSRS-C)⁽⁴⁶⁾ is an 18-item self-assessment questionnaire for depression in children. The adolescents were asked to rate their level of depression by choosing from "*always*," "*sometimes*," and "*never*."

The Strengths and Difficulties Questionnaire $(SDQ)^{(47)}$ is a self-administered questionnaire designed to comprehensively assess adjustment and mental health status from early childhood through adolescence. It comprises 25 items, each requiring a response of either "*yes*," "somewhat yes," or "no," and responses were rated on three levels: "low need," "some need," and "high need." Guardians were asked to complete the forms. For **the Vineland Adaptive Behavior Scales–Second Edition** (Vineland II),⁽⁴⁸⁾ the adolescents' adaptive behavior was graded across four domains: "communication domain," "daily skill domain," "sociality domain," and "motor skills." Three of the domains were graded; the Does Treatment Stigma among Adolescents with Autistic Spectrum Disorder and Their Guardians Affect the Effectiveness of Cognitive Behavioral Therapy

Demographic variables	Total N=45	TAU group n=22	COMB group n=23	р
Sex				.3226
Father	1(2.2)	0 (0.0)	1 (100.0)	
Mother	44 (97.8)	22 (50.0)	22 (50.0)	
Total years of education				.3075
Mean (<i>SD</i>)	14.3 (1.6)	14.6 (1.4)	14.1 (1.8)	
Total years of work				.1874
Mean (<i>SD</i>)	14.4 (8.2)	12.7 (8.4)	16.0 (7.8)	
Current employment status n (%)				.0981
No (homemaker)	17 (37.0)	11 (64.7)	6(35.3)	
Yes	28 (62.2)	11 (39.3)	17 (60.7)	
Full-time employment	13 (46.4)	5(38.5)	8 (61.5)	
Part-time employment	15 (51.7)	6 (40.0)	9 (60.0)	
Household income n (%)				.5311
4,000,000 yen or more ^a	6 (13.4)	4 (66.7)	2(33.3)	
5,000,000–8,000,000 yen	25(55.6)	13(52.0)	12 (48.0)	
9,000,000 yen or more	14 (31.0)	5 (35.7)	9 (64.3)	
Parental use of support n (%)				.4773
Non-use	37 (82.2)		18 (48.6)	
Use	8 (17.8)		5 (62.5)	
Past medical history				.7655
No	36 (80.0)		18 (50.0)	
Yes	9 (20.0)		5 (55.6)	

Table 2	Domographics of	fguardiane	of COMB/TALL	group (f	rom tho	original	etudy
	Demographics c	n guardians		group (i	romine	onginai	sludy

^a1 US dollar=115 Japanese yen.

COMB, Intervention group, groups receiving both outpatient psychotherapy with a psychiatrist and ACAT; TAU, Control group, groups receiving only outpatient psychotherapy.

"motor skills," for which the evaluation target was up to six-years-old, was excluded. We conducted a semi-structured interview with the guardians, who were asked to respond to **the 12-item General Health Questionnaire** (GHQ-12)⁽⁴⁹⁾, a self-administered questionnaire that measures the presence of mental and physical problems in the child from the first day of the child's life to the present. It comprises 12 items to be answered using a four-point scale: "*less than usual*," "*no more than usual*," "*rather more than usual*," or "*much more than usual*." Guardians were asked to respond to **the Parenting Resilience Elements Questionnaire** (PREQ)⁽⁵⁰⁾ for guardians of children with developmental disabilities. This is a self-administered questionnaire that measures the strength of nurturing resilience, with 16 items to be answered using a seven-point scale ranging from "*not at all true*" to "*very true*."

Analysis method

The hypotheses were as follows: (1) adolescents with ASD who have higher treatment stigma have poorer outcomes from ACAT interventions than those who have lower treatment stigma; and (2) when guardians' treatment stigma is high, they are less likely to have positive out-



comes from ACAT interventions. High guardian treatment stigma also negative effects the effectiveness of ACAT for adolescents with ASD. To clarify this, we used Spearman's rank correlation coefficient to calculate the correlation coefficient between the BACEv3 scores at pre-intervention and the difference scores for pre-to-post-intervention. For the effectiveness measures used (for adolescents, AKQ-C, SDQ, DSRS, and Vineland II; for guardians, AKQ-P, GHQ-12, and PREQ were used), we calculated difference scores (prevalue-post-value) and correlated their differences. We used Spearman's correlation coefficient because it is adaptable to small samples and less affected by outliers. Analyses were performed using SPSS Version 27.0 (IBM Armonk, New York, USA). A p-value <.05 indicated significance.

Results

The result of the hypothesis (1) analysis indicated that the BACEv3 scores of adolescents with ASD in the intervention group were not correlated with the difference scores in any measure. For hypothesis (2), we found that the BACEv3 scores of the guardians in the intervention group were not correlated with the difference scores for the AKQ-P, GHQ-12, and PREQ. However, we found a positive (r=0.681, p<.01) and moderate correlation between the guardians' BACEv3 scores at pre-intervention and their difference scores for AKQ-C knowledge. There was a negative (r=-0.534, p<.05) and moderate correlation between the guardians' BACEv3 scores at pre-intervention and their difference scores got Vineland II "communication" (receptive/expressive language, reading, and writing) (Table 3). For the BACEv3, which measures treatment

Difference scores on the questionnaires before – and after the intervention		Baseline intervention group			
		Adolescents with ASD BACEv3	Guardians of adolescents with ASD BACEv3	n	
	AKQ-C strength	.094	036	21	
	AKQ-C weak	251	103	21	
Adolescents with ASD	AKQ-C knowledge	288	.681**	18	
	SDQ	170	205	21	
	DSRS-C	333	.204	21	
	Vineland II Communication	.102	534*	21	
	Vineland II Daily living	.209	390	21	
	Vineland II Social	.204	.048	21	
Guardians of adolescents with ASD	AKQ-P strength	085	125	21	
	AKQ-P weak	022	186	21	
	AKQ-P knowledge	316	.326	19	
	GHQ-12	412	.158	21	
	PREQ	.253	050	21	

Table 3 Correlation coefficients between the BACEv3 scores at pre-intervention and the difference scores in the efficacy indicators before and after ACAT intervention.

Difference scores reflect the difference between scores pre- and post-intervention.

Significance was evaluated by Spearman's correlation coefficient.

p < .05, p < .01.

AKQ-C, Autism Knowledge Quiz-Child; AKQ-P, Autism Knowledge Quiz-Guardians; SDQ, Strengths and Difficulties Questionnaire; DSRS-C, Depression Self-Rating Scale for Children; Vineland II, Vineland Adaptive Behavior Scale Second Edition; GHQ, the 12-item General Health Questionnaire; PREQ, Parenting Resilience Elements Questionnaire. stigma, the mean pre-intervention values for adolescents with ASD and guardians in this study were 1.09 and 0.84, respectively. In Hongo et al., the mean score was 1.31.⁽⁴⁵⁾ As the participants in the previous study were adults, only parents were used as comparators in this study.

Discussion

The study examined whether the treatment stigma of adolescents with ASD and their guardians affected the effectiveness of the ACAT intervention. However, Spearman's rank correlation coefficient analysis showed that the severity of treatment stigma in adolescents with ASD was not correlated with the difference scores for any of the measures. Moreover, the guardians' severity of treatment stigma was not correlated with their difference scores. However, we found a positive and moderate correlation between the guardians' BACEv3 scores at pre-intervention and their difference scores for AKQ-C knowledge. This study suggested that even with high guardian treatment stigma, interventions improved knowledge in adolescents with ASD.

This study posited two main hypotheses:

- Adolescents with ASD with higher treatment stigma have poorer outcomes from ACAT interventions than those who have lower treatment stigma.
- (2) When guardians' treatment stigma is high, they are less likely to have positive outcomes from ACAT interventions. High guardian treatment stigma also negatively impacts the effectiveness of ACAT for adolescents with ASD.

Hypothesis (1) was rejected, as no differences in outcomes between adolescents with ASD with and those without treatment stigma were observed. Therefore, we suggest that adolescents' stigma did not affect intervention effectiveness. Hypothesis (2) was also rejected as there were no observable differences in outcome between guardians of ASD individuals with and those without treatment stigma.

Notably, the data suggested a correlation between the guardians' treatment stigma and an increase in the child's ASD knowledge. This may be because the adolescent encountered vital information related to the disease during the intervention, which the guardians, reluctant regarding the treatment, did not provide. Consequently, the adolescent was more willing to acquire knowledge to better understand the disease. Another interesting result showed a low score in communication skills in adolescents who had guardians with high stigma. As this score was evaluated from their perspective, guardians' lack of knowledge before the intervention may have interfered with communication with the child, leading to an overestimation. However, after the intervention, the guardian may have corrected their impression.

Limitations

Our data raise the issue of selection bias. The inclusion criteria for the study were individuals diagnosed with ASD and their guardians. Patients who had never received treatment for ASD were omitted. Therefore, it may be necessary to consider the treatment stigma of autistic adolescents and their guardians who have avoided treatment and to support them both in the future. The use of data from the intervention trial meant that the data sample was quite small; therefore, there is a need to test our hypotheses with a larger amount of data. The authors of a similar intervention study⁽⁵¹⁾ stated that they failed to detect small to moderate effects owing to sample size. Future studies are needed to confirm treatment stigma and ACAT intervention effects. Our participants had received treatment and support before the intervention and thus did not completely avoid it. Therefore, our conclusions cannot be generalized. In addition, the reasons people with treatment stigma received psychoeducation (intervention) may include the following.



Some participants were, for example, children/ students who had experienced truancy. In some cases, they and their guardians were referred to the university for advice by their school or family physician, which led to their participation in the study; however, in other cases, their guardians found out about the study by chance through an online search, which led to their participation. The common denominator was that the need was clear, and the intervention was recommended and attended by those around them. Future studies with more rigorous inclusion assessments are necessary to clarify intervention effects when treatment stigma is high.

Conclusions

In adolescents with ASD with high treatment stigma, no clear association between high treatment stigma and intervention effects was demonstrated. Therefore, treatment stigma in adolescents with ASD could not be said to influence intervention effectiveness. However, even when guardians' treatment stigma was high, ACAT interventions for adolescents with ASD increased their knowledge, suggesting that treatment stigma also did not reduce the effectiveness of this intervention for adolescents with ASD.

Conflicts of interest -

The authors declare no conflicts of interest.

Funding

This study was supported by a special research grant (no. 2017 S-1) from the Association of Japanese Clinical Psychology—a non-profit association for clinical psychology in Japan. The funders had no role in the design of the study, the collection, analysis, and interpretation of data or in the writing of the manuscript.

References

(1) American Psychiatric Association: Diagnostic and Statistical Manual of Mental Disorders. 5th ed. Washington, D.C.: Amer Psychiatric Pub Inc. 2013

- (2) Jones W, Klin A: Heterogeneity and homogeneity across the autism spectrum: The role of development. J Am Acad Child Adolesc Psychiatry. 48: 471– 473, 2009
- (3) Klin A, Saulnier CA, Sparrow SS, et al: Social and communication abilities and disabilities in higher functioning individuals with autism spectrum disorders: The Vineland and the ADOS. J Autism Dev Disord. 37: 748–59, 2007
- (4) Kenworthy L, Case L, Harms MB, et al: Adaptive behavior ratings correlate with symptomatology and IQ among individuals with high-functioning autism spectrum disorders. J Autism Dev Disord. 40: 416-423, 2010
- (5) Volker MA, Lopata C, Smerbeck AM, et al: BASC-2 PRS profiles for students with highfunctioning autism spectrum disorders. J Autism Dev Disord. 40: 188–199, 2010
- (6) Kanne SM, Gerber AJ, Quirmbach LM, et al: The role of adaptive behavior in autism spectrum disorders: Implications for functional outcome. J Autism Dev Disord. 41: 1007-1018, 2011
- (7) Pugliese CE, Anthony L, Strang JF, et al: Increasing adaptive behavior skill deficits from childhood to adolescence in autism spectrum disorder: Role of executive function. J Autism Dev Disord. 45: 1579-1587, 2015
- (8) Tillmann J, San José Cáceres A, Chatham CH, et al: Investigating the factors underlying adaptive functioning in autism in the EU-AIMS Longitudinal European Autism Project. Autism Res. 12: 645-657, 2019
- (9) Aubé B, Follenfant A, Goudeau S, et al: Public stigma of autism spectrum disorder at school: Implicit attitudes matter. J Autism Dev Disord. 51: 1584-1597, 2021
- (10) Clement S, Schauman O, Graham T, Maggioni F et al: What is the impact of mental healthrelated stigma on help-seeking? A systematic review of quantitative and qualitative studies. Psychol Med. 45: 11-27, 2015
- (11) Jorm AF, Oh E: Desire for social distance from people with mental disorders: A review. Aust NZ J Psychiatry. 43: 183–200, 2009
- (12) Wright A, Jorm AF, Mackinnon AJ: Labeling of mental disorders and stigma in young people. Soc Sci Med. 73: 498–506, 2011
- (13) Goffman E. Stigma: Notes on the Management

Does Treatment Stigma among Adolescents with Autistic Spectrum Disorder and Their Guardians Affect the Effectiveness of Cognitive Behavioral Therapy

of Spoiled Identity. Reissue edition. New York, Touchstone, 1986

- (14) Corrigan PT, Watson AC: Understanding the impact of stigma on people with mental illness. World Psychiatry. 1: 16–20, 2002
- (15) Clement S, Brohan E, Jeffery D, et al: Development and psychometric properties the Barriers to Access to Care Evaluation Scale (BACE) related to people with mental ill Health. BMC Psychiatry. 12: 36, 2012
- (16) Baker G, Olukoya A, Agglen P: Young people, social support and help-seeking. Int J Adolesc Med Health. 17: 315–336, 2005
- (17) Beukema L, de Winter AF, KorevaarEL et al: Investigating the use of support in secondary school: the role of self-reliance and stigma towards help-seeking. J Ment Health. 2022: 1-9, 2022
- (18) Calzada LR, Pistrang N, Mandy WPL: Highfunctioning autism and Asperger's disorder: Utility and meaning for families. J Autism Dev Disord. 42: 230-243, 2012
- (19) Mohamadi M, Mohtashami J, Vasli P, et al: The correlation between help-seeking behaviour, social comparison and stigma in caregivers of children with autism. Med Sci. 23(96): 146-151, 2019
- (20) Bromley J, Hare DJ, Davison K, et al: Mothers supporting children with autistic spectrum disorders: Social support, mental health status and satisfaction with services. Autism. 8(4): 409-423, 2004
- (21) Shepherd D, Landon J, Goedeke S, et al: Stress and distress in New Zealand parents caring for a child with autism spectrum disorder. Res De Disabil. 111: 103875, 2021
- (22) Torii M, Umeda M, Someki F, et al: Survey of stigma towards neurodevelopmental disabilities in high school students and effectiveness of the online education program of understanding neurodevelopmental disabilities. Bull Grad Sch Hum Dev Envi. 14: 69–77, 2021 (in Japanese).
- (23) Gillespie-Lynch K, Brooks PJ, Someki F, et al: Changing college students' conceptions of autism: An online training to increase knowledge and decrease stigma. J Autism Dev Disord. 45: 2553-2566, 2015
- (24) Someki F, Torii M, Brooks PJ, et al: Stigma associated with autism among college students in Japan and the United States: An online training study. Res Dev Disabil. 76: 88–98, 2018

- (25) Weston L, Hodgekins J, Langdon PE: Effectiveness of cognitive behavioural therapy with people who have autistic spectrum disorders: A systematic review and meta-analysis. Clin Psychol Rev. 49: 41-54, 2016
- (26) Maddox BB, Miyazaki Y, White SW: Long-term effects of CBT on social impairment in adolescents with ASD. J Autism Dev Disord. 47: 3872–3882, 2017
- (27) Spain D, Happé F: How to optimise cognitive behaviour therapy (CBT) for people with autism spectrum disorders (ASD): A Delphi study. J Ration-Emotive Cogn-Behav Ther. 38: 184–208, 2020
- (28) Sung M, Ooi YP, Goh TJ, et al: Effects of cognitive-behavioral therapy on anxiety in children with autism spectrum disorders: A randomized controlled trial. Child Psychiatry Hum Dev. 42: 634-449, 2011
- (29) White SW, Ollendick T, Albano AM, et al: Randomized controlled trial: Multimodal anxiety and social skill intervention for adolescents with autism spectrum disorder. J Autism Dev Disord. 43: 382-394, 2013
- (30) Wood JJ, Ehrenreich-May J, Alessandri M, et al: Cognitive behavioral therapy for early adolescents with autism spectrum disorders and clinical anxiety: A randomized, controlled trial. Behav Ther. 46: 7–19, 2015
- (31) McGillivray JA, Evert HT: Group cognitive behavioural therapy program shows potential in reducing symptoms of depression and stress among young people with ASD. J Autism Dev Disord. 44: 2041-2051, 2014
- (32) Russell AJ, Jassi A, Fullana MA, et al: Cognitive behavior therapy for comorbid obsessivecompulsive disorder in high-functioning autism spectrum disorders: A randomized controlled trial: Research article: CBT for OCD for adults with ASD. Depress Anxiety. 30: 697–708, 2013
- (33) Gantman A, Kapp SK, Orenski K, et al: Social skills training for young adults with highfunctioning autism spectrum disorders: A randomized controlled pilot study. J Autism Dev Disord. 42: 1094-1103, 2012
- (34) Laugeson EA, Frankel F, Gantman A, et al: Evidence-based social skills training for adolescents with autism spectrum disorders: The UCLA PEERS program. J Autism Dev Disord. 42: 1025–1036, 2012
- (35) Schohl KA, Van Hecke AV, Carson AM, et al: A



replication and extension of the PEERS intervention: Examining effects on social skills and social anxiety in adolescents with autism spectrum disorders. J Autism Dev Disord. 44: 532-545, 2014

- (36) Nussey C, Pistrang N, Murphy T: How does psychoeducation help? A review of the effects of providing information about Tourette syndrome and attention-deficit/hyperactivity disorder: A review of psychoeducational approaches in TS and ADHD. Child Care Health Dev. 39: 617– 627, 2013
- (37) Gordon K, Murin M, Baykaner O, et al: A randomised controlled trial of PEGASUS, A psychoeducational programme for young people with high-functioning autism spectrum disorder. J Child Psychol Psychiatry. 56: 468-476, 2015
- (38) Werner S, Shulman C: Subjective well-being among family caregivers of individuals with developmental disabilities: The role of affiliate stigma and psychosocial moderating variables. Res Dev Disabil. 34: 4103–4114, 2013
- (39) Papadopoulos C, Lodder A, Constantinou G, et al: Systematic review of the relationship between autism stigma and informal caregiver mental health. J Autism Dev Disord. 49: 1665– 1685, 2019
- (40) Shepherd D, Landon J, Goedeke S, et al: Stress and distress in New Zealand parents caring for a child with autism spectrum disorder. Res Dev Disabil. 111: 103875, 2021
- (41) Chan KKS, Lam CB: Trait mindfulness attenuates the adverse psychological impact of stigma on parents of children with autism spectrum disorder. Mindfulness. 8: 984-994, 2017
- (42) Lodder A, Papadopoulos C, Randhawa G: Stigma of living as an autism carer: A brief psycho-social support intervention (SOLACE). Study protocol for a randomised controlled feasibility study. Pilot Feasibility Stud. 5: 34, 2019

- (43) Oshima F, William M, Takahashi N, et al: Cognitive-behavioral family therapy as psychoeducation for adolescents with highfunctioning autism spectrum disorders: Aware and care for my autistic traits (ACAT) program study protocol for a pragmatic multisite randomized controlled trial. Trials. 21: 814, 2020
- (44) Oshima et al: In submission.
- (45) Hongo M, Oshima F, Nishinaka H, et al: Reliability and validity of the Japanese version of the Barriers to Access to Care Evaluation Scale Version 3 for people with mental disorders: an online survey study. Front Psychol. 12: 760184, 2021
- (46) Birleson P, Hudson I, Buchanan DG, et al: Clinical evaluation of a self-rating scale for depressive disorder in childhood (Depression Self-Rating Scale). J Child Psychol Psychiatry. 28: 43-60, 1987
- (47) Goodman R: The strengths and difficulties questionnaire: A research note. J Child Psychol Psychiatry. 38: 581–586, 1997
- (48) Sparrow SS, Cicchetti DV, Balla DA: Vineland adaptive behavior scales (Vineland-II). 2nd edition. Minneapolis: NCS Pearson Inc., 2005
- (49) Montazeri A, Harirchi AM, Shariati M, et al: The 12-item General Health Questionnaire (GHQ-12): Translation and validation study of the Iranian version. Health Qual Life Outcomes. 1: 66, 2003
- (50) Suzuki K, Kobayashi T, Moriyama K, et al: Development and evaluation of a Parenting Resilience Elements Questionnaire (PREQ) measuring resiliency in rearing children with developmental disorders. PLoS One. 10: e0143946, 2015
- (51) Takahashi T, Sugiyama F, Kikai T, et al: Changes in depression and anxiety through mindfulness group therapy in Japan: The role of mindfulness and self-compassion as possible mediators. BioPsychoSocial Med. 13(1): 4, 2019