

Title	Characteristic initial boundary value problems for symmetric hyperbolic systems
Author(s)	Nishitani, Tatsuo; Takayama, Masahiro
Citation	Osaka Journal of Mathematics. 1998, 35(3), p. 629-657
Version Type	VoR
URL	<a href="https://doi.org/10.18910/9254">https://doi.org/10.18910/9254</a>
rights	
Note	

***Osaka University Knowledge Archive : OUKA***

<https://ir.library.osaka-u.ac.jp/>

Osaka University

# CHARACTERISTIC INITIAL BOUNDARY VALUE PROBLEMS FOR SYMMETRIC HYPERBOLIC SYSTEMS

TATSUO NISHITANI and MASAHIRO TAKAYAMA

(Received May 2, 1997)

## 1. Introduction

Let  $\Omega$  be a bounded open subset of  $\mathbf{R}^n$  with smooth boundary  $\partial\Omega$ . In  $\mathbf{R} \times \Omega$  we consider a first order symmetric hyperbolic system:

$$Lu = \sum_{j=0}^n A_j(t, x) \partial_j u + B(t, x)u,$$

$$A_j(t, x), B(t, x) \in C^\infty(\mathbf{R} \times \overline{\Omega}), \quad A_j^*(t, x) = A_j(t, x)$$

with  $\partial_0 = \partial/\partial t$ ,  $\partial_j = \partial/\partial x_j$ ,  $j = 1, \dots, n$  and  $u = (u_1, \dots, u_N)$  where  $A_0(t, x)$  is positive definite on  $\mathbf{R} \times \Omega$ . We assume that  $A_j(t, x)$  and  $B(t, x)$  are independent of  $t$  outside a compact subset of  $\mathbf{R} \times \overline{\Omega}$ . Recall that the boundary matrix is given by

$$A_b(t, x) = \sum_{j=1}^n \nu_j(x) A_j(t, x) \quad \text{for } (t, x) \in \mathbf{R} \times \partial\Omega$$

where  $\nu(x) = (\nu_1(x), \dots, \nu_n(x))$  is the unit outward normal to  $\Omega$  at  $x \in \partial\Omega$ . In this paper we study the initial boundary value problems for  $L$  assuming that the boundary space is maximal positive.

A general theory of initial boundary value problem for non singular  $A_b$  with maximal positive boundary space was developed by Friedrichs [2], Lax-Phillips [4], Rauch-Massey III [13] and so on. The case of the characteristic boundary has been studied by Lax-Phillips [4], Majda-Osher [6], Rauch [12] and so on. In particular, when  $\dim \text{Ker} A_b$  is constant on the boundary, in [11] we find a detailed study on the initial boundary value problem where the regularity was measured by conormal Sobolev spaces. In the characteristic case, one can not expect full regularity even if  $f \in H^s(\Omega)$  (see [6], [17]). In [9], [14], in a similar situation, the initial boundary value problems were studied in usual Sobolev space setting aimed to study non linear perturbations. For a concrete problem of this type see [18] which motivated our study.

When  $\dim \text{Ker} A_b$  is not constant it is well known (see [5], [10]) that one does not in general get a well posed boundary value problem by merely taking maximal

positive boundary conditions, while in [11] we can find some positive results. In [7] we proved the existence of regular solutions in the case that  $A_b$  is definite apart from an embedded  $n - 2$  dimensional submanifold of  $\partial\Omega$  on which  $A_b$  vanishes under the same conditions assumed also in this paper. In [15] the same question is studied in a similar situation.

In this paper we continue studying the same problem when  $A_b$  is non singular outside a set, assumed to be an open set with smooth boundary on which  $A_b$  is definite.

Let us set

$$(1.1) \quad O^+(O^-) = \{(t, x) \in \mathbf{R} \times \partial\Omega; A_b(t, x) \text{ is positive (negative) definite}\}$$

and denote by  $\gamma^\pm$  the boundaries of  $O^\pm$  in  $\mathbf{R} \times \partial\Omega$ . Letting  $\gamma = \gamma^+ \cup \gamma^-$  we assume that  $\gamma$  is a smooth embedded  $n - 1$  dimensional submanifold of  $\mathbf{R} \times \partial\Omega$ , the boundary matrix  $A_b(t, x)$  is non singular on  $(\mathbf{R} \times \partial\Omega) \setminus \gamma$  and that  $\text{Ker} A_b(t, x)$  is a smooth vector bundle over  $\gamma$ .

The boundary condition takes the form:

$$u(t, x) \in M(t, x) \quad \text{for } (t, x) \in \mathbf{R} \times \partial\Omega$$

where  $M(t, x)$  is a linear subspace of  $\mathbf{C}^N$ . We assume that the boundary space  $M(t, x)$  is maximal positive in the sense that

$$\langle A_b(t, x)v, v \rangle \geq 0 \quad \text{for all } v \in M(t, x),$$

$$\dim M(t, x) = \#\{\text{non negative eigenvalues of } A_b(t, x) \text{ counting multiplicity}\}.$$

In particular, (1.1) implies that

$$(1.2) \quad M(t, x) = \begin{cases} \mathbf{C}^N & \text{on } O^+ \\ \{0\} & \text{on } O^- \end{cases}$$

We also assume that  $M(t, x)$  is smooth on each component of  $(\mathbf{R} \times \partial\Omega) \setminus \gamma$  up to the boundary and independent of  $t$  outside a compact subset of  $\mathbf{R} \times \partial\Omega$ .

We study the following initial boundary value problem:

$$(IBVP) \quad \begin{cases} Lu = f & \text{in } I \times \Omega \\ u \in M & \text{at } I \times \partial\Omega \\ u(0, \cdot) = u_0 & \text{on } \Omega \end{cases}$$

where  $I = (0, T)$ . In what follows, we introduce the notation  $\mathcal{O} = I \times \Omega$ ,  $\Gamma = I \times \partial\Omega$ ,  $\mathcal{R} = \mathbf{R} \times \Omega$  and  $\Delta = \mathbf{R} \times \partial\Omega$ .

We make our assumptions precise. Let  $(\bar{t}, \bar{x}) \in \gamma$  and we work in a neighborhood  $U$  of  $(\bar{t}, \bar{x})$ . Let  $\{v_1(t, x), \dots, v_p(t, x)\} \subset C^\infty(U)$  be a basis for  $\text{Ker} A_b(t, x)$  on  $\gamma \cap U$

and set  $V(t, x) = (v_1(t, x), \dots, v_p(t, x))$ . Take  $h(t, x) \in C^\infty(U)$  so that  $\gamma \cap U = (\mathbf{R} \times \partial\Omega) \cap \{h(t, x) = 0\}$  where  $dh(t, x)$  and  $\nu(x)$  are linearly independent on  $\gamma \cap U$ . Since  $(V^* A_b V)(t, x)$  vanishes on  $\gamma \cap U$  we can factor out  $h(t, x)$ ;

$$(1.3) \quad (V^* A_b V)(t, x) = h(t, x) A_{b, \gamma}(t, x) \quad \text{on } (\mathbf{R} \times \partial\Omega) \cap U.$$

Moreover we set

$$(1.4) \quad A_{\gamma/b}(t, x) = V^*(t, x) \left( \sum_{j=0}^n (\partial_j h) A_j \right) (t, x) V(t, x).$$

For more intrinsic definitions of  $A_{b, \gamma}$  and  $A_{\gamma/b}$ , see [8]. Our assumption is stated as follows:

$$(1.5) \quad A_{b, \gamma}(t, x) \text{ and } A_{\gamma/b}(t, x) \text{ have the same definiteness on } \gamma \cap U.$$

Clearly this condition does not depend on the choice of  $v_j(t, x)$  and  $h(t, x)$ .

Under the conditions (1.5) we discuss the existence and regularity of solutions to (IBVP). We also study asymptotic behavior of solutions near  $\gamma$ .

## 2. Results for zero initial data

In what follows, if  $u = u(t, x)$  is a function of  $t$  and  $x$  then we denote by  $u(t)$  the function of  $x$  obtained by freezing  $t$ ;  $u(t)(x) = u(t, x)$ .

We denote the formal adjoint of  $L$  by  $L^*$ :

$$L^* u = - \sum_{j=0}^n \partial_j A_j(t, x) u + B^*(t, x) u.$$

For  $u, v \in C^{0,1}(\overline{\mathcal{O}})$ , Green's identity yields

$$\begin{aligned} (Lu, v)_{L^2(\mathcal{O})} &= (u, L^* v)_{L^2(\mathcal{O})} + \iint_{\Gamma} \langle A_b u, v \rangle dt d\sigma \\ &\quad + (A_0(T)u(T), v(T))_{L^2(\Omega)} - (A_0(0)u(0), v(0))_{L^2(\Omega)}. \end{aligned}$$

The adjoint boundary space  $M^*(t, x)$  is defined by

$$M^*(t, x) = [A_b(t, x)M(t, x)]^\perp \quad \text{for } (t, x) \in \Delta.$$

In particular, (1.2) implies that

$$(2.1) \quad M^*(t, x) = \begin{cases} \{0\} & \text{on } O^+ \\ \mathbf{C}^N & \text{on } O^-. \end{cases}$$

We recall the following definition (see [1], [2]).

**DEFINITION.** For  $f \in L^2(\mathcal{O})$  and  $u_0 \in L^2(\Omega)$  we say  $u \in L^2(\mathcal{O})$  is a weak solution to (IBVP) if and only if the identity

$$(u, L^* \psi)_{L^2(\mathcal{O})} = (f, \psi)_{L^2(\mathcal{O})} + (A_0(0)u_0, \psi(0))_{L^2(\Omega)}$$

holds for all  $\psi \in C^{0,1}(\overline{\mathcal{O}})$  with  $\psi \in M^*$  at  $\Gamma$  and  $\psi(T) = 0$ .

Take  $r(x) \in C^\infty(\overline{\Omega})$  with  $dr(x) \neq 0$  on  $\partial\Omega$  so that  $\Omega = \{r(x) > 0\}$  and  $h_\pm(t, x) \in C^\infty(\overline{\mathcal{R}})$  such that  $O^\pm = \Delta \cap \{h_\pm(t, x) > 0\}$  where  $dh_\pm(t, x)$  and  $\nu(x)$  are linearly independent on  $\gamma^\pm$ . Similarly, take  $h(t, x) \in C^\infty(\overline{\mathcal{R}})$  such that  $\gamma = \Delta \cap \{h(t, x) = 0\}$  where  $dh(t, x)$  and  $\nu(x)$  are linearly independent on  $\gamma$ . We assume that  $h_\pm(t, x)$  and  $h(t, x)$  are independent of  $t$  outside a compact subset of  $\overline{\mathcal{R}}$ . Let us set

$$m_\pm(t, x) = \{r(x)^2 + h_\pm(t, x)^2\}^{1/2}, \quad m(t, x) = \{r(x)^2 + h(t, x)^2\}^{1/2}, \\ \phi_\pm(t, x) = m_\pm(t, x) - h_\pm(t, x).$$

Note that  $\phi_\pm(t, x) > 0$  if  $(t, x) \in \overline{\mathcal{R}} \setminus (O^\pm \cup \gamma^\pm)$  and that  $\phi_\pm(t, x) = 0$  if  $(t, x) \in O^\pm \cup \gamma^\pm$ .

We first get the following two propositions.

**Proposition 2.1.** If  $f \in \phi_-^\tau L^2(\mathcal{O})$  and  $u_0 \in \phi_-^\tau(0)L^2(\Omega)$  for some  $\tau \geq 1$  then there exists a weak solution  $u \in \phi_-^\tau L^2(\mathcal{O})$  to (IBVP) satisfying

$$\|\phi_-^{-\tau} u\|_{L^2(\mathcal{O})}^2 \leq C \{ \|\phi_-^{-\tau} f\|_{L^2(\mathcal{O})}^2 + \|\phi_-^{-\tau}(0)u_0\|_{L^2(\Omega)}^2 \}$$

where  $C = C(\tau) > 0$  is independent of  $f$ ,  $u_0$  and  $u$ .

**Proposition 2.2.** If  $f \in L^2(\mathcal{O})$  and  $u_0 \in L^2(\Omega)$  then a weak solution  $u \in m_- L^2(\mathcal{O})$  to (IBVP) is unique.

An immediate corollary to Proposition 2.1 and Proposition 2.2 is

**Corollary 2.3.** If  $f \in \phi_-^\tau L^2(\mathcal{O})$  and  $u_0 \in \phi_-^\tau(0)L^2(\Omega)$  for some  $\tau \geq 1$  and if  $u \in m_- L^2(\mathcal{O})$  is a weak solution to (IBVP) then we have  $u \in \phi_-^\tau L^2(\mathcal{O})$  and it follows that

$$\|\phi_-^{-\tau} u\|_{L^2(\mathcal{O})}^2 \leq C \{ \|\phi_-^{-\tau} f\|_{L^2(\mathcal{O})}^2 + \|\phi_-^{-\tau}(0)u_0\|_{L^2(\Omega)}^2 \}$$

where  $C = C(\tau) > 0$  is independent of  $f$ ,  $u_0$  and  $u$ .

Our main concern is the regularity of solutions  $u$  to (IBVP). Hence we introduce the following spaces: For  $q \in \mathbf{Z}_+$  and  $\sigma, \tau \in \mathbf{R}$  we set

$$(2.2) \quad X_{(\sigma, \tau)}^q(\mathcal{O}) = \bigcap_{j=0}^q \phi_+^{\sigma+q-j} \phi_-^{\tau+q-j} H^j(\mathcal{O}),$$

$$(2.3) \quad X_{0(\sigma, \tau)}^q(\Omega) = \bigcap_{j=0}^q (\phi_+^{\sigma+q-j} \phi_-^{\tau+q-j})(0) H^j(\Omega)$$

where  $H^j(\mathcal{O})$  and  $H^j(\Omega)$  are usual Sobolev spaces of order  $j$ . We define  $X_{(\sigma, \tau)}^q(\mathcal{O}; \Gamma)$  by (2.2) with  $H^j(\mathcal{O}; \Gamma)$ , the conormal Sobolev space of order  $j$  with respect to  $\Gamma$ , instead of  $H^j(\mathcal{O})$ . The space  $X_{0(\sigma, \tau)}^q(\Omega; \partial\Omega)$  is defined similarly (see also [8]). Note that if  $f \in X_{(\sigma, \tau)}^q(\mathcal{O})$  (resp.  $X_{(\sigma, \tau)}^q(\mathcal{O}; \Gamma)$ ) for some  $q \in \mathbf{Z}_+$ ,  $q \geq 1$  and  $\sigma, \tau \in \mathbf{R}$  then  $(\partial_0^k f)(0) \in X_{0(\sigma, \tau)}^{q-1-k}(\Omega)$  (resp.  $X_{0(\sigma, \tau)}^{q-1-k}(\Omega; \partial\Omega)$ ) for  $k = 0, \dots, q-1$ .

We can now obtain regular solutions to (IBVP) with zero initial data (results for the general case is described in Theorem 5.4 and Theorem 5.5 in Section 5).

**Theorem 2.4.** *For  $q \in \mathbf{Z}_+$ ,  $q \geq 1$  there is a  $\Sigma(q) > 0$  such that if  $f \in X_{(-\sigma, \tau)}^q(\mathcal{O}; \Gamma) \cap \phi_- L^2(\mathcal{O})$ , for some  $\sigma, \tau > \Sigma(q)$ , satisfies  $(\partial_0^k f)(0) = 0$  for  $k = 0, \dots, q-1$  then there exists a weak solution  $u \in X_{(-\sigma, \tau)}^q(\mathcal{O}; \Gamma) \cap \phi_- L^2(\mathcal{O})$  to (IBVP) with zero initial data which satisfies*

$$\|u\|_{X_{(-\sigma, \tau)}^q(\mathcal{O}; \Gamma)}^2 + \|\phi_-^{-1} u\|_{L^2(\mathcal{O})}^2 \leq C \{ \|f\|_{X_{(-\sigma, \tau)}^q(\mathcal{O}; \Gamma)}^2 + \|\phi_-^{-1} f\|_{L^2(\mathcal{O})}^2 \}$$

where  $C = C(q, \sigma, \tau) > 0$  is independent of  $f$  and  $u$ .

We can get a rough estimate of asymptotic behavior of weak solutions near  $\gamma$ .

**Theorem 2.5.** *For  $q \in \mathbf{Z}_+$  there is a  $\Sigma(q) > 0$  such that if  $f \in X_{(-\sigma, \tau)}^{q+[n/2]+1}(\mathcal{O}) \cap \phi_- L^2(\mathcal{O})$ , for some  $\sigma, \tau > \Sigma(q)$ , satisfies  $(\partial_0^k f)(0) = 0$  for  $k = 0, \dots, q + [n/2]$  and if  $u \in m_- L^2(\mathcal{O})$  is a weak solution to (IBVP) with zero initial data then we have  $u \in m^{-(q+[n/2]+1)} \phi_+^{-\sigma} \phi_-^{\tau} C^q(\overline{\mathcal{O}})$ .*

### 3. Existence and uniqueness of solutions to (IBVP)

Let us set

$$m_{\pm}(t, x; \kappa, \mu) = \{ \kappa r(x)^2 + (\mu r(x) - h_{\pm}(t, x))^2 \}^{1/2},$$

$$\phi_{\pm}(t, x; \kappa, \mu) = m_{\pm}(t, x; \kappa, \mu) + \mu r(x) - h_{\pm}(t, x)$$

for  $\kappa > 0$  and  $\mu \in \mathbf{R}$ . Then we can choose a  $C = C(\kappa, \mu) > 0$  satisfying

$$C^{-1} m_{\pm}(t, x) \leq m_{\pm}(t, x; \kappa, \mu) \leq C m_{\pm}(t, x),$$

$$C^{-1}\phi_{\pm}(t, x) \leq \phi_{\pm}(t, x; \kappa, \mu) \leq C\phi_{\pm}(t, x) \quad \text{for } (t, x) \in \mathcal{R}.$$

Thus it suffices to prove the results in Section 2 with  $m_{\pm}(t, x; \kappa, \mu)$  and  $\phi_{\pm}(t, x; \kappa, \mu)$  instead of  $m_{\pm}(t, x)$  and  $\phi_{\pm}(t, x)$ . In what follows, we simply write  $m_{\pm}(t, x)$  and  $\phi_{\pm}(t, x)$  for  $m_{\pm}(t, x; \kappa, \mu)$  and  $\phi_{\pm}(t, x; \kappa, \mu)$  respectively.

We denote by  $\|\cdot\|_{\mathcal{O}}$ ,  $\|\cdot\|_{\mathcal{R}}$  and  $\|\cdot\|_{\Omega}$  the norm in  $L^2(\mathcal{O})$ ,  $L^2(\mathcal{R})$  and in  $L^2(\Omega)$  respectively. The following a priori estimate is obtained by much the same way as in [8]. (for details see Lemma 5.4 in [8]).

**Lemma 3.1.** *There are  $c_0, C_1 > 0$  such that for  $\tau \geq 0$  we can take a  $\Lambda(\tau) \in \mathbf{R}$  having the following properties: If  $\operatorname{Re} \lambda > \Lambda(\tau)$ ,  $-\infty < T_1 < T_2 < \infty$  and if  $u \in C_0^{0,1}(\overline{\mathcal{R}})$  with  $u \in M^*$  at  $(T_1, T_2) \times \partial\Omega$  then it follows that*

$$\begin{aligned} & (\operatorname{Re} \lambda - \Lambda(\tau)) \|u\|_{(T_1, T_2) \times \Omega}^2 + c_0 \|u(T_1)\|_{\Omega}^2 \\ & \leq C_1 \{ \|\phi_-^{\tau}(L^* + \bar{\lambda}A_0)\phi_-^{-\tau}u\|_{(T_1, T_2) \times \Omega}^2 + \|u(T_2)\|_{\Omega}^2 \}. \end{aligned}$$

Applying this we can prove Proposition 2.1.

**Proof of Proposition 2.1.** Let us set

$$E = \{ \phi_-^{\tau}(L^* + \bar{\lambda}A_0)e^{\bar{\lambda}t}\psi; \psi \in C_0^{0,1}(\overline{\mathcal{O}}) \text{ with } \psi \in M^* \text{ at } \Gamma \text{ and } \psi(T) = 0 \}$$

and we study the map

$$T : E \ni \phi_-^{\tau}(L^* + \bar{\lambda}A_0)e^{\bar{\lambda}t}\psi \mapsto (f, \psi)_{\mathcal{O}} + (A_0(0)u_0, \psi(0))_{\Omega} \in \mathbf{C}.$$

From Lemma 3.1 with  $u = e^{\bar{\lambda}t}\phi_-^{\tau}\psi$  and  $T_1 = 0$ ,  $T_2 = T$  we obtain that

$$\begin{aligned} & |T\phi_-^{\tau}(L^* + \bar{\lambda}A_0)e^{\bar{\lambda}t}\psi|^2 \\ & \leq C \{ \|e^{\bar{\lambda}t}\phi_-^{\tau}\psi\|_{\mathcal{O}}^2 \|e^{-\bar{\lambda}t}\phi_-^{-\tau}f\|_{\mathcal{O}}^2 + \|(\phi_-^{\tau}\psi)(0)\|_{\Omega}^2 \|\phi_-^{-\tau}(0)u_0\|_{\Omega}^2 \} \\ & \leq C'(\lambda) \|\phi_-^{\tau}(L^* + \bar{\lambda}A_0)e^{\bar{\lambda}t}\psi\|_{\mathcal{O}}^2 \{ \|e^{-\bar{\lambda}t}\phi_-^{-\tau}f\|_{\mathcal{O}}^2 + \|\phi_-^{-\tau}(0)u_0\|_{\Omega}^2 \}. \end{aligned}$$

By Hahn-Banach theorem there is a  $w \in L^2(\mathcal{O})$  such that

$$\begin{aligned} & \|w\|_{\mathcal{O}}^2 \leq C(\lambda) \{ \|e^{-\bar{\lambda}t}\phi_-^{-\tau}f\|_{\mathcal{O}}^2 + \|\phi_-^{-\tau}(0)u_0\|_{\Omega}^2 \}, \\ & (w, \phi_-^{\tau}(L^* + \bar{\lambda}A_0)e^{\bar{\lambda}t}\psi)_{\mathcal{O}} = (f, \psi)_{\mathcal{O}} + (A_0(0)u_0, \psi(0))_{\Omega} \end{aligned}$$

for every  $\psi \in C_0^{0,1}(\overline{\mathcal{O}})$  with  $\psi \in M^*$  at  $\Gamma$  and  $\psi(T) = 0$ . Then  $u = e^{\lambda t}\phi_-^{\tau}w$  is a desired weak solution.  $\square$

For the proof of Proposition 2.2 and Theorem 2.4 we study the following boundary value problem:

$$(BVP) \quad \begin{cases} (L + \lambda A_0)u = f & \text{in } \mathbf{R} \times \Omega = \mathcal{R} \\ u \in M & \text{at } \mathbf{R} \times \partial\Omega = \Delta \end{cases}$$

where  $\lambda \in \mathbf{C}$  is a parameter.

**DEFINITION.** For  $f \in L^2(\mathcal{R})$  we say  $u \in L^2(\mathcal{R})$  is a weak solution to (BVP) if and only if the identity

$$(u, (L^* + \bar{\lambda}A_0)\psi)_{L^2(\mathcal{R})} = (f, \psi)_{L^2(\mathcal{R})}$$

holds for all  $\psi \in C_0^{0,1}(\bar{\mathcal{R}})$  with  $\psi \in M^*$  at  $\Delta$ .

We now set  $\phi_{\pm, \eta}(t, x) = \phi_{\pm}(t, x) - \eta$  and  $\mathcal{R}_{\pm, \eta} = \mathcal{R} \cap \{\phi_{\pm, \eta} > 0\}$  for  $\eta \geq 0$ . The following proposition is a key result to prove Proposition 2.2 and Theorem 2.4.

**Proposition 3.2.** *There is a  $\Lambda \in \mathbf{R}$  such that if  $\operatorname{Re} \lambda > \Lambda$  and if  $f \in L^2(\mathcal{R})$  with  $\operatorname{supp} f \subset \bar{\mathcal{R}}_{-, \eta} \cap \{t \geq T_0\}$  for some  $\eta > 0$  and  $T_0 \in \mathbf{R}$  then there exists a weak solution  $u \in L^2(\mathcal{R})$  to (BVP) with  $\operatorname{supp} u \subset \bar{\mathcal{R}}_{-, \eta} \cap \{t \geq T_0\}$ .*

To prove this we shall need a few lemmas which are proved by repeating the same arguments as in [8] (see Lemma 5.6, Proposition 5.2 and Corollary 7.8 in [8]).

**Lemma 3.3.** *There is a  $\eta_0 > 0$  such that for  $\tau \geq 0$  we can take a  $\Lambda(\tau) \in \mathbf{R}$  verifying the following properties: If  $0 < \eta < \eta_0$ ,  $\operatorname{Re} \lambda > \Lambda(\tau)$  and if  $u \in C_0^{0,1}(\bar{\mathcal{R}})$  with  $\operatorname{supp} u \cap \{\phi_{-, \eta} = 0\} = \emptyset$  and  $u \in M^*$  at  $\Delta \cap \{\phi_{-, \eta} > 0\}$  then it follows that*

$$\begin{aligned} & (\operatorname{Re} \lambda - \Lambda(\tau)) \|u\|_{\mathcal{R}_{-, \eta}}^2 + c_0(\tau - 1/4) \|\phi_{-, \eta}^{-1/2} u\|_{\mathcal{R}_{-, \eta}}^2 \\ & \leq C_1 \|\phi_{-, \eta}^{\tau+1/2} (L^* + \bar{\lambda}A_0) \phi_{-, \eta}^{-\tau} u\|_{\mathcal{R}_{-, \eta}}^2 \end{aligned}$$

where  $c_0, C_1 > 0$  depend only on  $\eta$ .

**Lemma 3.4.** *There are  $c_0, C_1, \Sigma_0 > 0$  such that for  $\sigma, \tau > \Sigma_0$  we can take a  $\Lambda(\sigma, \tau) \in \mathbf{R}$  verifying the following properties: If  $\operatorname{Re} \lambda > \Lambda(\sigma, \tau)$ ,  $-\infty < T_1 < T_2 < \infty$  and if  $u \in C_0^{0,1}(\bar{\mathcal{R}})$  with  $u \in M$  at  $(T_1, T_2) \times \partial\Omega$  then it follows that*

$$\begin{aligned} & (\operatorname{Re} \lambda - \Lambda(\sigma, \tau)) \|m^{1/2} u\|_{(T_1, T_2) \times \Omega}^2 \\ & + c_0(\min(\sigma, \tau) - \Sigma_0) \|u\|_{(T_1, T_2) \times \Omega}^2 + c_0 \|(m^{1/2} u)(T_2)\|_{\Omega}^2 \\ & \leq C_1 \{ \|m \phi_+^{\sigma} \phi_-^{-\tau} (L + \lambda A_0) \phi_+^{-\sigma} \phi_-^{\tau} u\|_{(T_1, T_2) \times \Omega}^2 + \|(m^{1/2} u)(T_1)\|_{\Omega}^2 \}. \end{aligned}$$



**Lemma 3.5.** *Let  $u \in L^2(\mathcal{R})$ , with  $\text{supp} u \subset \overline{\mathcal{R}}_{-, \eta}$  for some  $\eta > 0$ , be a weak solution to (BVP). Then there is a  $\{u_\epsilon\} \subset C_0^\infty(\overline{\mathcal{R}})$  with  $\text{supp} u_\epsilon \subset \overline{\mathcal{R}}_{-, \eta_0}$  and  $u_\epsilon \in M$  at  $\Delta$  such that if  $\sigma \geq 4$  and  $\tau \in \mathbf{R}$  then  $\phi_+^\sigma \phi_-^\tau u_\epsilon$  is also a weak solution to (BVP). Moreover we have*

$$u_\epsilon \rightarrow u, \quad \phi_+^\sigma \phi_-^\tau (L + \lambda A_0) u_\epsilon \rightarrow \phi_+^\sigma \phi_-^\tau (L + \lambda A_0) u \quad \text{in } L^2(\mathcal{R}) \quad \text{as } \epsilon \rightarrow 0$$

where  $\eta_0 > 0$  depends only on  $\eta$ .

**Proof of Proposition 3.2.** Using Lemma 3.3 and repeating arguments similar to those in Proposition 3.2 in [8] we can find a  $u \in L^2(\mathcal{R})$  with  $\text{supp} u \subset \overline{\mathcal{R}}_{-, \eta}$  which is a weak solution to (BVP). We choose a  $\{u_\epsilon\}$  as in Lemma 3.5. Then Lemma 3.4 shows that

$$\begin{aligned} & (\min(\sigma, \tau) - \Sigma_0) \|\phi_+^\sigma \phi_-^\tau u_\epsilon\|_{(S_0, T_0) \times \Omega}^2 \\ & \leq C_1 \{ \|m \phi_+^\sigma \phi_-^\tau (L + \lambda A_0) u_\epsilon\|_{(S_0, T_0) \times \Omega}^2 + \|(m^{1/2} \phi_+^\sigma \phi_-^\tau u_\epsilon)(S_0)\|_\Omega^2 \}. \end{aligned}$$

Letting  $S_0 \rightarrow -\infty$  and  $\epsilon \rightarrow 0$  we have

$$\begin{aligned} & (\min(\sigma, \tau) - \Sigma_0) \|\phi_+^\sigma \phi_-^\tau u\|_{(-\infty, T_0) \times \Omega}^2 \\ & \leq C_1 \|m \phi_+^\sigma \phi_-^\tau (L + \lambda A_0) u\|_{(-\infty, T_0) \times \Omega}^2 = \|m \phi_+^\sigma \phi_-^\tau f\|_{(-\infty, T_0) \times \Omega}^2 = 0. \end{aligned}$$

This implies  $\text{supp} u \subset \{t \geq T_0\}$  which proves the assertion.  $\square$

We now give the proof of Proposition 2.2.

**Proof of Proposition 2.2.** Assuming that  $u \in m_- L^2(\mathcal{O})$  is a weak solution to (BVP) with  $f = 0$  and  $u_0 = 0$  we wish to show  $u = 0$ . Let  $g \in C_0^\infty(\mathcal{O})$ . Repeating the same arguments as in Proposition 3.2 we can find a  $v \in L^2(\mathcal{R})$ , with  $\text{supp} v \subset \overline{\mathcal{R}}_{+, \eta} \cap \{t \leq T - \eta\}$  for some  $\eta > 0$ , which is a weak solution to the following adjoint boundary value problem

$$(BVP^*) \quad \begin{cases} (L + \bar{\lambda} A_0) v = g & \text{in } \mathbf{R} \times \Omega = \mathcal{R} \\ v \in M^* & \text{at } \mathbf{R} \times \partial\Omega = \Delta. \end{cases}$$

Let us choose  $\chi \in C_0^\infty(\mathbf{R})$  so that  $\chi = 1$  near 0 and set

$$v_k = \chi(k^{-1}t)(1 - \chi(km_-))v, \quad g_k = (L^* + \bar{\lambda} A_0)v_k$$

for  $k > 0$  large enough. Then  $v_k$  is also a weak solution to (BVP\*) replaced  $g$  by  $g_k$ . Since  $\text{supp} v_k$  is compact and  $\text{supp} v_k \cap \gamma = \emptyset$  then Theorem 4 in [12] gives a  $\{v_{k, \epsilon}\} \subset C_0^1(\overline{\mathcal{R}})$  with  $v_{k, \epsilon} \in M^*$  at  $\Delta$  such that

$$v_{k, \epsilon} \rightarrow v_k, \quad (L^* + \bar{\lambda} A_0)v_{k, \epsilon} \rightarrow g_k \quad \text{in } L^2(\mathcal{R}) \quad \text{as } \epsilon \rightarrow 0.$$

Noticing  $e^{-\bar{\lambda}t}v_{k,\epsilon} \in C^1(\bar{\mathcal{O}})$  with  $e^{-\bar{\lambda}t}v_{k,\epsilon} \in M^*$  at  $\Gamma$  and  $(e^{-\bar{\lambda}t}v_{k,\epsilon})(T) = 0$  and recalling that  $u$  is a weak solution to (BVP) with  $f = 0$  and  $u_0 = 0$  we obtain  $(u, L^*e^{-\bar{\lambda}t}v_{k,\epsilon})_{\mathcal{O}} = 0$ . Letting  $\epsilon \rightarrow 0$  we have  $(e^{-\lambda t}u, g_k)_{\mathcal{O}} = 0$ . We note that  $(e^{-\lambda t}u, g_k)_{\mathcal{O}} \rightarrow (e^{-\lambda t}u, g)_{\mathcal{O}}$  as  $k \rightarrow \infty$ . Indeed we can write

$$\begin{aligned} (e^{-\lambda t}u, g_k)_{\mathcal{O}} &= (e^{-\lambda t}u, \chi(k^{-1}t)(1 - \chi(km_-))g)_{\mathcal{O}} \\ &\quad - (e^{-\lambda t}u, k^{-1}\chi'(k^{-1}t)\chi(km_-)A_0v)_{\mathcal{O}} \\ &\quad + (e^{-\lambda t}u, \chi(k^{-1}t)k\chi'(km_-)M_-v)_{\mathcal{O}} \\ &= I_1 + I_2 + I_3 \end{aligned}$$

with  $M_- = \sum_{j=0}^n (\partial_j m_-) A_j$ . The dominated convergence theorem shows that  $I_1 \rightarrow (e^{-\lambda t}u, g)_{\mathcal{O}}$ ,  $I_2 \rightarrow 0$  as  $\epsilon \rightarrow 0$ . We turn to  $I_3$ . Since  $u = m_-w$  for some  $w \in L^2(\mathcal{O})$  and  $|\theta\chi'(\theta)| \leq C$  for some  $C > 0$  the dominated convergence theorem again proves that  $I_3 = (e^{-\lambda t}w, \chi(k^{-1}t)km_- \chi'(km_-)M_-v)_{\Omega} \rightarrow 0$  as  $\epsilon \rightarrow 0$ . Thus we have  $(e^{-\lambda t}u, g)_{\Omega} = 0$ . Noticing that  $C_0^\infty(\mathcal{O})$  is dense in  $L^2(\mathcal{O})$  we conclude  $u = 0$ .  $\square$

#### 4. Proofs of results for zero initial data

We start with the proof of Theorem 2.4. Recalling that  $A_j(t, x)$ ,  $B(t, x)$  and  $h_{\pm}(t, x)$  are independent of  $t$  outside a compact subset of  $\bar{\mathcal{R}}$  and repeating the same arguments as in [8] we can prove the following two propositions (see Proposition 3.1 and Proposition 11.3 in [8]).

**Proposition 4.1.** *For  $q \in \mathbf{Z}_+$  there is a  $\Sigma(q) > 0$  such that for  $\sigma, \tau > \Sigma(q)$  we can take a  $\Lambda(q, \sigma, \tau) \in \mathbf{R}$  verifying the following properties: If  $\operatorname{Re} \lambda > \Lambda(q, \sigma, \tau)$  and  $f \in X_{(-\sigma, \tau)}^q(\mathcal{R}; \Delta) \cap L^2(\mathcal{R})$  and if  $u \in L^2(\mathcal{R})$ , with  $\operatorname{supp} u \subset \bar{\mathcal{R}}_{-, \eta}$  for some  $\eta > 0$ , is a weak solution to (BVP) then it follows that  $u \in X_{(-\sigma, \tau)}^q(\mathcal{R}; \Delta)$ .*

**Proposition 4.2.** *For  $q \in \mathbf{Z}_+$  there is a  $\Sigma(q) > 0$  such that for  $\sigma, \tau > \Sigma(q)$  we can take a  $\Lambda(q, \sigma, \tau) \in \mathbf{R}$  verifying the following properties: If  $\operatorname{Re} \lambda > \Lambda(q, \sigma, \tau)$  and if  $u \in X_{(-\sigma, \tau)}^{q+[n/2]+2}(\mathcal{R}; \Delta) \cap L^2(\mathcal{R})$  is a weak solution to (BVP) with  $f \in C_0^\infty(\mathcal{R})$  then it follows that*

$$\|u\|_{X_{(-\sigma, \tau)}^q(\mathcal{R}; \Delta)}^2 \leq C_1 \|(L + \lambda A_0)u\|_{X_{(-\sigma, \tau)}^q(\mathcal{R}; \Delta)}^2$$

where  $C_1 = C_1(q, \sigma, \tau, \lambda) > 0$  is independent of  $u$  and  $f$ .

An immediate corollary to these propositions is

**Corollary 4.3.** *For  $q \in \mathbf{Z}_+$  there is a  $\Sigma(q) > 0$  such that for  $\sigma, \tau > \Sigma(q)$  we can take a  $\Lambda(q, \sigma, \tau) \in \mathbf{R}$  verifying the following properties: If  $\operatorname{Re} \lambda > \Lambda(q, \sigma, \tau)$  and*

if  $u \in L^2(\mathcal{R})$ , with  $\text{supp } u \subset \overline{\mathcal{R}}_{-\eta}$  for some  $\eta > 0$ , is a weak solution to (BVP) with  $f \in C_0^\infty(\mathcal{R})$  then it follows that  $u \in X_{(-\sigma, \tau)}^q(\mathcal{R}; \Delta)$  and

$$\|u\|_{X_{(-\sigma, \tau)}^q(\mathcal{R}; \Delta)}^2 \leq C_1 \|(L + \lambda A_0)u\|_{X_{(-\sigma, \tau)}^q(\mathcal{R}; \Delta)}^2$$

where  $C_1 = C_1(q, \sigma, \tau, \lambda) > 0$  is independent of  $u$  and  $f$ .

Proof of Theorem 2.4. We can take a  $\tilde{f} \in X_{(-\sigma, \tau)}^q(\mathcal{R}; \Delta)$  with  $\text{supp } \tilde{f} \subset \{0 \leq t \leq \tilde{T}\}$  such that  $\tilde{f} = f$  on  $\mathcal{O}$  and

$$(4.1) \quad \|\tilde{f}\|_{X_{(-\sigma, \tau)}^q(\mathcal{R}; \Delta)} \leq C \|f\|_{X_{(-\sigma, \tau)}^q(\mathcal{O}; \Gamma)}$$

where  $C > 0$  and  $\tilde{T} > T$  are independent of  $f$  and  $\tilde{f}$  (we give the proof of this fact in Corollary 7.11 below). Let us choose  $\chi \in C_0^\infty(\mathbf{R})$  so that  $\chi = 1$  near 0 and  $\rho \in C_0^\infty(\mathbf{R}^{n+1})$  with  $\text{supp } \rho \subset \{(t, x); 0 < t < 1, |x| < 1\}$  such that  $\rho \geq 0$  and  $\iint \rho dt dx = 1$  and set

$$f_{k, \epsilon}(t, x) = (((1 - \chi(kr))\tilde{f}) * \rho_\epsilon)(t, x), \quad \rho_\epsilon(t, x) = \epsilon^{-(n+1)} \rho(\epsilon^{-1}t, \epsilon^{-1}x)$$

for  $k > 0$  large enough and  $0 < \epsilon < 1$  small enough where  $r = r(x)$  is a defining function of  $\Omega$ . Then we have  $f_{k, \epsilon} \in C_0^\infty(\mathcal{R})$  with  $\text{supp } f_{k, \epsilon} \subset \{0 \leq t \leq \tilde{T} + 1\}$  for  $\epsilon > 0$  small enough. Moreover it follows from the proof of Lemma 6.4 in [8] that

$$\phi_-^{-1} f_{k, \epsilon} \rightarrow \phi_-^{-1} f \quad \text{in } L^2(\mathcal{O}), \quad f_{k, \epsilon} \rightarrow \tilde{f} \quad \text{in } X_{(-\sigma, \tau)}^q(\mathcal{R}; \Delta)$$

as  $\epsilon \rightarrow 0$  and  $k \rightarrow \infty$ . Let  $\lambda \in \mathbf{C}$  be  $\text{Re } \lambda > 0$  large enough and set  $F_{k, \epsilon} = e^{-\lambda t} f_{k, \epsilon}$ . Then Proposition 3.2 gives a weak solution  $U_{k, \epsilon} \in L^2(\mathcal{R})$  to (BVP) with  $f$  replaced by  $F_{k, \epsilon}$  with  $\text{supp } U_{k, \epsilon} \subset \overline{\mathcal{R}}_{-\eta} \cap \{t \geq 0\}$  for some  $\eta > 0$ . From Corollary 4.3 it follows that  $U_{k, \epsilon} \in X_{(-\sigma, \tau)}^q(\mathcal{R}; \Delta)$  and

$$\|U_{k, \epsilon}\|_{X_{(-\sigma, \tau)}^q(\mathcal{R}; \Delta)}^2 \leq C_1 \|F_{k, \epsilon}\|_{X_{(-\sigma, \tau)}^q(\mathcal{R}; \Delta)}^2.$$

Now if we write  $u_{k, \epsilon} = e^{\lambda t} U_{k, \epsilon}$  then we have  $u_{k, \epsilon} \in X_{(-\sigma, \tau)}^q(\mathcal{O}; \Gamma) \cap \phi_- L^2(\mathcal{O})$  and

$$(4.2) \quad \|u_{k, \epsilon}\|_{X_{(-\sigma, \tau)}^q(\mathcal{O}; \Gamma)}^2 \leq C_1 \|f_{k, \epsilon}\|_{X_{(-\sigma, \tau)}^q(\mathcal{R}; \Delta)}^2.$$

We first show that  $u_{k, \epsilon}$  is a weak solution to (IBVP) replaced  $f$  and  $u_0$  by  $f_{k, \epsilon}$  and 0. Let  $\psi \in C^{0,1}(\overline{\mathcal{O}})$  with  $\psi \in M^*$  at  $\Gamma$  and  $\psi(T) = 0$ . We choose a  $\tilde{\psi} \in C_0^{0,1}(\overline{\mathcal{R}})$  with  $\text{supp } \tilde{\psi} \subset \{t \leq T\}$  such that  $\tilde{\psi} \in M^*$  at  $\Delta$  and  $\tilde{\psi}(T) = 0$ . Since  $U_{k, \epsilon}$  is a weak solution to (BVP) it follows that

$$(U_{k, \epsilon}, (L^* + \bar{\lambda} A_0) e^{-\bar{\lambda} t} \tilde{\psi})_{\mathcal{R}} = (F_{k, \epsilon}, e^{-\bar{\lambda} t} \tilde{\psi})_{\mathcal{R}}.$$

Noticing  $\text{supp } U_{k,\epsilon} \cap \text{supp } \tilde{\psi} \subset \overline{\mathcal{O}}$  we get  $(u_{k,\epsilon}, L^* \psi)_{\mathcal{O}} = (f_{k,\epsilon}, \psi)_{\mathcal{O}}$ , and hence  $u_{k,\epsilon}$  is a weak solution to (IBVP). Therefore it follows from Corollary 2.3 that

$$(4.3) \quad \|\phi^{-1} u_{k,\epsilon}\|_{\mathcal{O}}^2 \leq C \|\phi^{-1} f_{k,\epsilon}\|_{\mathcal{O}}^2.$$

Combining (4.2) and (4.3) we obtain

$$(4.4) \quad \|u_{k,\epsilon}\|_{X_{(-\sigma,\tau)}^q(\mathcal{O};\Gamma)}^2 + \|\phi^{-1} u_{k,\epsilon}\|_{\mathcal{O}}^2 \leq C_1 \{\|f_{k,\epsilon}\|_{X_{(-\sigma,\tau)}^q(\mathcal{R};\Delta)}^2 + \|\phi^{-1} f_{k,\epsilon}\|_{\mathcal{O}}^2\}.$$

Since  $\{f_{k,\epsilon}\}$  is a Cauchy sequence in  $X_{(-\sigma,\tau)}^q(\mathcal{R};\Delta) \cap \phi_{-} L^2(\mathcal{O})$  then  $\{u_{k,\epsilon}\}$  has a limit point  $u$  in  $X_{(-\sigma,\tau)}^q(\mathcal{O};\Gamma) \cap \phi_{-} L^2(\mathcal{O})$ . Then  $u$  is a desired weak solution to (IBVP) with zero initial data and the desired estimate follows from (4.4) and (4.1).  $\square$

We turn to the proof of Theorem 2.5. The proof easily follows from

**Proposition 4.4.** *Let  $u \in X_{(\sigma,\tau)}^q(\mathcal{O};\Gamma)$  and  $Lu \in X_{(\sigma,\tau)}^q(\mathcal{O})$  for some  $q \in \mathbf{Z}_+$  and  $\sigma, \tau \in \mathbf{R}$ . Then it follows that  $u \in m^{-q} X_{(\sigma,\tau)}^q(\mathcal{O})$  and*

$$\|m^q u\|_{X_{(\sigma,\tau)}^q(\mathcal{O})} \leq C \{\|u\|_{X_{(\sigma,\tau)}^q(\mathcal{O};\Gamma)} + \|Lu\|_{X_{(\sigma,\tau)}^q(\mathcal{O})}\}$$

where  $C = C(q, \sigma, \tau) > 0$  is independent of  $u$ .

Admitting for the moment that Proposition 4.4 holds we shall prove Theorem 2.5.

**Proof of Theorem 2.5.** Let  $q' = q + [n/2] + 1$ . Theorem 2.4 gives a weak solution  $v \in X_{(-\sigma,\tau)}^{q'}(\mathcal{O};\Gamma) \cap \phi_{-} L^2(\mathcal{O})$  to (IBVP) with zero initial data, and hence it follows from Proposition 2.2 that  $u = v \in X_{(-\sigma,\tau)}^{q'}(\mathcal{O};\Gamma)$ . Therefore Proposition 4.4 implies that

$$u \in m^{-q'} X_{(-\sigma,\tau)}^{q'}(\mathcal{O}) \hookrightarrow m^{-q'} \phi_{+}^{-\sigma} \phi_{-}^{\tau} H^{q'}(\mathcal{O}) \hookrightarrow m^{-q'} \phi_{+}^{-\sigma} \phi_{-}^{\tau} C^q(\overline{\mathcal{O}})$$

which shows the assertion.  $\square$

To prove Proposition 4.4 we localize the problem. Let us take a covering  $\{U_i\}_{i=0}^l$  of  $\mathcal{O}$  as follows: First we cover  $\Gamma$  by coordinate patches  $U_i$ ,  $i = 1, \dots, l$ , with coordinate systems  $\chi_i : U_i \cap \mathcal{O} \rightarrow \{(\tau, \xi); a_i < \tau < b_i, |\xi| < 1, \xi_1 > 0\}$  such that  $\tau = t \circ \chi_i^{-1}$  and  $\xi_1 = r \circ \chi_i^{-1}$  where  $0 \leq a_i < b_i \leq T$ . Next we cover  $\mathcal{O} \setminus \bigcup_{i=1}^l U_i$  by  $U_0 \subset \subset \mathbf{R} \times \Omega$ . Choose a partition of unity  $\{\psi_i\}_{i=0}^l$  subordinate to this covering  $\{U_i\}_{i=0}^l$  and set  $u_i = \psi_i u$ . If  $U_i \cap \Gamma = \emptyset$  then Proposition 4.4 with  $u_i$  instead of  $u$  is easily checked.

Now we suppose that  $U_i \cap \Gamma \neq \emptyset$ . Performing a change of independent variables we may assume that  $r = x_1$  and  $\text{supp } u_i \subset \overline{I_i} \times \{|x| < 1, x_1 \geq 0\}$  where  $I_i = (a_i, b_i)$ .

In what follows, we write

$$\begin{aligned}\partial &= (\partial_0, \partial_1, \partial_2, \dots, \partial_n), & \partial_x &= (\partial_1, \partial_2, \dots, \partial_n), \\ Z &= (\partial_0, x_1 \partial_1, \partial_2, \dots, \partial_n), & Z_x &= (x_1 \partial_1, \partial_2, \dots, \partial_n).\end{aligned}$$

Proposition 4.4 is an immediate consequence of Lemma 4.5 below.

**Lemma 4.5.** *Let  $p \in \mathbf{Z}_+$ ,  $\alpha \in \mathbf{Z}_+^{n+1}$  and assume that  $p + |\alpha| \leq q$ . Then we have*

$$\begin{aligned}(4.5) \quad & \|m^p \phi_+^{-\sigma-q+p+|\alpha|} \phi_-^{-\tau-q+p+|\alpha|} \partial_1^p Z^\alpha u_i\|_{I_i \times \mathbf{R}_+^n} \\ & \leq C \{ \|u\|_{X_{(\sigma,\tau)}^q(\mathcal{O};\Gamma)} + \|Lu\|_{X_{(\sigma,\tau)}^q(\mathcal{O})} \}\end{aligned}$$

where  $C = C(q, \sigma, \tau, p, \alpha) > 0$ .

**Proof of Proposition 4.4.** Let  $\alpha \in \mathbf{Z}_+^{n+1}$  and assume that  $|\alpha| \leq q$ . If we write  $\alpha' = (\alpha_0, 0, \alpha_2, \dots, \alpha_n)$  then it follows that

$$\begin{aligned}& \|m^q \phi_+^{-\sigma-q+|\alpha|} \phi_-^{-\tau-q+|\alpha|} \partial^\alpha u_i\|_{I_i \times \mathbf{R}_+^n} \\ & \leq C \|m^{\alpha_1} \phi_+^{-\sigma-q+\alpha_1+|\alpha'|} \phi_-^{-\tau-q+\alpha_1+|\alpha'|} \partial_1^{\alpha_1} Z^{\alpha'} u_i\|_{I_i \times \mathbf{R}_+^n} \\ & \leq C' \{ \|u\|_{X_{(\sigma,\tau)}^q(\mathcal{O};\Gamma)} + \|Lu\|_{X_{(\sigma,\tau)}^q(\mathcal{O})} \}.\end{aligned}$$

Arguments similar to those in Lemma 6.1 in [8] imply that

$$\|m^q u_i\|_{X_{(\sigma,\tau)}^q(\mathcal{O})} \leq C \sum_{|\alpha| \leq q} \|m^q \phi_+^{-\sigma-q+|\alpha|} \phi_-^{-\tau-q+|\alpha|} \partial^\alpha u_i\|_{I_i \times \mathbf{R}_+^n}$$

which shows that

$$(4.6) \quad \|m^q u_i\|_{X_{(\sigma,\tau)}^q(\mathcal{O})} \leq C \{ \|u\|_{X_{(\sigma,\tau)}^q(\mathcal{O};\Gamma)} + \|Lu\|_{X_{(\sigma,\tau)}^q(\mathcal{O})} \}.$$

Summing (4.6) from  $i = 0$  to  $i = l$  we get the desired estimate.  $\square$

We shall prove Lemma 4.5. The interesting patches are at  $\gamma$ . Note that  $h_\pm \partial_1$  is written as a sum of  $a(t, x) Z^\beta$ ,  $|\beta| \leq 1$  and  $a(t, x) L$  where  $a(t, x) \in \mathcal{B}^\infty(\mathbf{R}^{n+1})$ , the set of all smooth functions on  $\mathbf{R}^{n+1}$  with bounded derivatives of all order, which may differ from line to line.

**Lemma 4.6.**  *$(h_\pm \partial_1)^p Z^\alpha$ ,  $p \geq 1$  and  $|\alpha| \leq q$ , is written as a sum of the following terms:*

$$a(t, x) m^{-k} x_1^i h_\pm^j (h_\pm \partial_1)^l Z^\beta, \quad 0 \leq l \leq p-1, \quad 0 \leq k \leq 2(q+p-1-l),$$

$$\begin{aligned}
0 \leq k - i - j \leq q, \quad |\beta| \leq q - k + i + j + 1, \\
a(t, x)m^{-k}x_1^i h_{\pm}^j (h_{\pm}\partial_1)^l Z^{\beta} L, \quad 0 \leq l \leq p-1, \quad 0 \leq k \leq 2(q+p-1-l), \\
0 \leq k - i - j \leq q, \quad |\beta| \leq q - k + i + j.
\end{aligned}$$

**Proof.** We first consider the case  $p = 1$ . Note that  $(h_{\pm}\partial_1)Z^{\alpha}$  is written as a sum of  $a(t, x)Z^{\beta}Z^{\alpha}$ ,  $|\beta| \leq 1$  and  $a(t, x)LZ^{\alpha}$ . Here  $a(t, x)Z^{\beta}Z^{\alpha}$  can be written as a desired sum. We turn to  $a(t, x)LZ^{\alpha}$ . Since  $LZ^{\alpha} = Z^{\alpha}L + [L, Z^{\alpha}]$ , Lemma 10.5 in [7] shows that  $LZ^{\alpha}$  can be also written as a desired sum.

We next consider the case  $p \geq 2$ . Using that  $(h_{\pm}\partial_1)^p Z^{\alpha} = (h_{\pm}\partial_1)^{p-1}(h_{\pm}\partial_1)Z^{\alpha}$  and the results for the case  $p = 1$  we conclude the assertion.  $\square$

**Lemma 4.7.**  $\|m^p \phi_+^{-\sigma-q+p+|\alpha|} \phi_-^{-\tau-q+p+|\alpha|} \partial_1^p Z^{\alpha} u_i\|_{I_i \times \mathbf{R}_+^n}$ ,  $p+|\alpha| \leq q$  and  $p \geq 1$ , is bounded from above by a sum of the following terms:

$$\begin{aligned}
\|m^l \phi_+^{-\sigma-q+l+|\beta|} \phi_-^{-\tau-q+l+|\beta|} \partial_1^l Z^{\beta} u_i\|_{I_i \times \mathbf{R}_+^n}, \quad 0 \leq l \leq p-1, \quad l+|\beta| \leq q, \\
\|m^l \phi_+^{-\sigma-q+l+|\beta|} \phi_-^{-\tau-q+l+|\beta|} \partial_1^l Z^{\beta} L u_i\|_{I_i \times \mathbf{R}_+^n}, \quad 0 \leq l \leq p-1, \quad l+|\beta| \leq q.
\end{aligned}$$

**Proof.** Since  $m \leq C(|x_1| + |h_{\pm}|)$  for some  $C > 0$  it follows that

$$\begin{aligned}
& \|m^p \phi_+^{-\sigma-q+p+|\alpha|} \phi_-^{-\tau-q+p+|\alpha|} \partial_1^p Z^{\alpha} u_i\|_{I_i \times \mathbf{R}_+^n} \\
& \leq C \{ \|x_1^p \phi_+^{-\sigma-q+p+|\alpha|} \phi_-^{-\tau-q+p+|\alpha|} \partial_1^p Z^{\alpha} u_i\|_{I_i \times \mathbf{R}_+^n} \\
& \quad + \|h_{\pm}^p \phi_+^{-\sigma-q+p+|\alpha|} \phi_-^{-\tau-q+p+|\alpha|} \partial_1^p Z^{\alpha} u_i\|_{I_i \times \mathbf{R}_+^n} \} \\
& = C \{ I_1 + I_2 \}.
\end{aligned}$$

Noticing that  $x_1^p \partial_1^p$  can be written as a sum of  $Z_1^l$ ,  $0 \leq l \leq p$ , we have

$$I_1 \leq C \sum_{l=0}^p \|\phi_+^{-\sigma-q+(l+|\alpha|)} \phi_-^{-\tau-q+(l+|\alpha|)} Z_1^l Z^{\alpha} u_i\|_{I_i \times \mathbf{R}_+^n}.$$

Thus  $I_1$  is bounded from above by a desired sum. Moreover Lemma 4.6 implies that  $I_2$  can be also bounded from above by a desired sum.  $\square$

**Proof of Lemma 4.5.** We proceed by induction on  $p$ . From Lemma 6.1 in [8] the case  $p = 0$  is trivial. Inductively assume that the statement is true up to  $p-1$ . Lemma 4.7 shows that  $\|m^p \phi_+^{-\sigma-q+p+|\alpha|} \phi_-^{-\tau-q+p+|\alpha|} \partial_1^p Z^{\alpha} u_i\|_{I_i \times \mathbf{R}_+^n}$  is bounded from above by a sum of the following terms:

$$\begin{aligned}
\|m^l \phi_+^{-\sigma-q+l+|\beta|} \phi_-^{-\tau-q+l+|\beta|} \partial_1^l Z^{\beta} u_i\|_{I_i \times \mathbf{R}_+^n} &= I_1, \quad 0 \leq l \leq p-1, \quad l+|\beta| \leq q, \\
\|m^l \phi_+^{-\sigma-q+l+|\beta|} \phi_-^{-\tau-q+l+|\beta|} \partial_1^l Z^{\beta} L u_i\|_{I_i \times \mathbf{R}_+^n} &= I_2, \quad 0 \leq l \leq p-1, \quad l+|\beta| \leq q.
\end{aligned}$$

By the inductive hypothesis  $I_1$  can be bounded from above by the right-hand side of (4.5). We turn to  $I_2$ . Since  $Lu_i = \psi_i Lu + \sum_{j,k} (\partial_j \psi_i) A_j u_k$  the inductive hypothesis implies that  $I_2$  can also be bounded from above by the right-hand side of (4.5). This proves the assertion for  $p$ .  $\square$

## 5. Results for general initial data

Now we shall extend the definition of a weak solution to (IBVP). Let us set

$$\mathcal{L}^2(\mathcal{O}) = \bigcup_{\sigma, \tau \geq 0} \phi_+^{-\sigma} \phi_-^{\tau} L^2(\mathcal{O}), \quad \mathcal{L}_0^2(\Omega) = \bigcup_{\sigma, \tau \geq 0} (\phi_+^{-\sigma} \phi_-^{\tau})(0) L^2(\Omega).$$

Noticing (2.1) we introduce the following definition.

**DEFINITION.** For  $f \in \mathcal{L}^2(\mathcal{O})$  and  $u_0 \in \mathcal{L}_0^2(\Omega)$  we say  $u \in \mathcal{L}^2(\mathcal{O})$  is a weak solution to (IBVP) if and only if the identity

$$(u, L^* \psi)_{L^2(\mathcal{O})} = (f, \psi)_{L^2(\mathcal{O})} + (A_0(0)u_0, \psi(0))_{L^2(\Omega)}$$

holds for all  $\psi \in C^{0,1}(\overline{\mathcal{O}})$  with  $\psi \in M^*$  at  $\Gamma$ ,  $\psi(T) = 0$  and  $\psi = 0$  on a neighborhood of  $O^+$ .

Then by using arguments similar to those in the proof of Proposition 2.1, Proposition 2.2 and Corollary 2.3 we obtain

**Proposition 5.1.** *If  $f \in \phi_+^{-\sigma} \phi_-^{\tau} L^2(\mathcal{O})$  and  $u_0 \in (\phi_+^{-\sigma} \phi_-^{\tau})(0) L^2(\Omega)$  for some  $\sigma \geq 0$  and  $\tau \geq 1$  then there exists a weak solution  $u \in \phi_+^{-\sigma} \phi_-^{\tau} L^2(\mathcal{O})$  to (IBVP) satisfying*

$$\|\phi_+^{\sigma} \phi_-^{-\tau} u\|_{L^2(\mathcal{O})}^2 \leq C \{ \|\phi_+^{\sigma} \phi_-^{-\tau} f\|_{L^2(\mathcal{O})}^2 + \|(\phi_+^{\sigma} \phi_-^{-\tau})(0) u_0\|_{L^2(\Omega)}^2 \}$$

where  $C = C(\sigma, \tau) > 0$  is independent of  $f$ ,  $u_0$  and  $u$ .

**Proposition 5.2.** *If  $f \in \mathcal{L}^2(\mathcal{O})$  and  $u_0 \in \mathcal{L}_0^2(\Omega)$  then a weak solution  $u \in m_- L^2(\mathcal{O})$  to (IBVP) is unique.*

**Corollary 5.3.** *If  $f \in \phi_+^{-\sigma} \phi_-^{\tau} L^2(\mathcal{O})$  and  $u_0 \in (\phi_+^{-\sigma} \phi_-^{\tau})(0) L^2(\Omega)$  for some  $\sigma \geq 0$  and  $\tau \geq 1$  and if  $u \in m_- L^2(\mathcal{O})$  is a weak solution to (IBVP) then we have  $u \in \phi_+^{-\sigma} \phi_-^{\tau} L^2(\mathcal{O})$  and it follows that*

$$\|\phi_+^{\sigma} \phi_-^{-\tau} u\|_{L^2(\mathcal{O})}^2 \leq C \{ \|\phi_+^{\sigma} \phi_-^{-\tau} f\|_{L^2(\mathcal{O})}^2 + \|(\phi_+^{\sigma} \phi_-^{-\tau})(0) u_0\|_{L^2(\Omega)}^2 \}$$

where  $C = C(\sigma, \tau) > 0$  is independent of  $f$ ,  $u_0$  and  $u$ .

In order to get regularity results we introduce "compatibility conditions". Let  $f \in X_{(-\sigma, \tau)}^q(I \times \Omega)$  and  $u_0 \in X_{0(-\sigma, \tau)}^q(\Omega)$  for  $q \in \mathbf{Z}_+$ ,  $q \geq 1$  and  $\sigma, \tau \geq 0$ . Then we define  $u^{(k)}$ ,  $k = 0, \dots, q-1$ , as follows:

$$u^{(0)} = u_0, \quad u^{(k)} = (\partial_0^{k-1} A_0^{-1} f)(0) - \sum_{i=0}^{k-1} \binom{k-1}{i} K_i u^{(k-1-i)} \quad \text{for } k \geq 1$$

where  $K_i = \sum_{j=1}^n (\partial_0^i A_0^{-1} A_j)(0) \partial_j + (\partial_0^i A_0^{-1} B)(0)$ . Note that

$$u^{(k)} \in X_{0(-\sigma, \tau)}^{q-k}(\Omega) \hookrightarrow X_{0(-\sigma, \tau)}^1(\Omega) \hookrightarrow (\phi_+^{-\sigma} \phi_-^{\tau})(0) H^1(\Omega),$$

and hence  $(\phi_+^{\sigma} \phi_-^{\tau})(0) u^{(k)} \in L^2(\partial\Omega)$ . We write  $T_k(f, u_0) = u^{(k)}$  for  $k = 0, \dots, q-1$ .

Let  $\delta > 0$  be small enough and choose  $P(t, x) \in C^\infty((-\delta, \delta) \times \partial\Omega; M_N(\mathbf{C}))$  such that  $v \in M(t, x)$  if and only if  $P(t, x)v = 0$  for every  $(t, x) \in (-\delta, \delta) \times \partial\Omega$ .

**DEFINITION.** For  $q \in \mathbf{Z}_+$ ,  $q \geq 1$  and  $\sigma, \tau \geq 0$  we say  $f \in X_{(-\sigma, \tau)}^q(\mathcal{O})$  and  $u_0 \in X_{0(-\sigma, \tau)}^q(\Omega)$  satisfy the compatibility conditions up to order  $q-1$  if and only if the following identities hold:

$$\sum_{i=0}^k \binom{k}{i} (\partial_0^i P)(0) (\phi_+^{\sigma} \phi_-^{\tau})(0) u^{(k-i)} = 0 \quad \text{on } \partial\Omega \setminus \gamma_0 \quad \text{for } k = 0, \dots, q-1.$$

Here  $u^{(k)} = T_k(f, u_0)$ ,  $k = 0, \dots, q-1$ , and  $\gamma_0 = \{x \in \partial\Omega; (0, x) \in \gamma\}$ .

**Theorem 5.4.** For  $q \in \mathbf{Z}_+$ ,  $q \geq 1$  there is a  $\Sigma(q) > 0$  such that if  $f \in X_{(-\sigma, \tau)}^q(\mathcal{O})$  and  $u_0 \in X_{0(-\sigma, \tau)}^q(\Omega)$ , for some  $\sigma, \tau > \Sigma(q)$ , satisfy the compatibility conditions up to order  $q-1$  then there exists a weak solution  $u \in X_{(-\sigma, \tau)}^q(\mathcal{O}; \Gamma)$  to (IBVP) which satisfies

$$\|u\|_{X_{(-\sigma, \tau)}^q(\mathcal{O}; \Gamma)}^2 \leq C \{ \|f\|_{X_{(-\sigma, \tau)}^q(\mathcal{O})}^2 + \|u_0\|_{X_{0(-\sigma, \tau)}^q(\Omega)}^2 \}$$

where  $C = C(q, \sigma, \tau) > 0$  is independent of  $f$ ,  $u_0$  and  $u$ .

From Theorem 5.4 and Proposition 4.4 we can derive a rough estimate of asymptotic behavior of weak solutions near  $\gamma$ .

**Theorem 5.5.** For  $q \in \mathbf{Z}_+$  there is a  $\Sigma(q) > 0$  such that if  $f \in X_{(-\sigma, \tau)}^{q+[n/2]+1}(\mathcal{O})$  and  $u_0 \in X_{0(-\sigma, \tau)}^{q+[n/2]+1}(\Omega)$ , for some  $\sigma, \tau > \Sigma(q)$ , satisfy the compatibility conditions up to order  $q+[n/2]$  and if  $u \in m_- L^2(\mathcal{O})$  is a weak solution to (IBVP) then we have  $u \in m^{-(q+[n/2]+1)} \phi_+^{-\sigma} \phi_-^{\tau} C^q(\overline{\mathcal{O}})$ .



## 6. Proofs of results for general initial data

In this section we give the proof of Theorem 5.4. From Lemma 3.4 we recall that

**Lemma 6.1.** *There are  $C, \Sigma_0 > 0$  such that for  $\sigma, \tau > \Sigma_0$  we can take a  $\Lambda(\sigma, \tau) \in \mathbf{R}$  verifying the following properties: If  $\operatorname{Re} \lambda > \Lambda(\sigma, \tau)$  and if  $u \in C_0^{0,1}(\overline{\mathcal{R}})$ , with  $\operatorname{supp} u \cap (O^- \cup \gamma^-) = \emptyset$ , is a weak solution to (IBVP) then it follows that*

$$\begin{aligned} & (\min(\sigma, \tau) - \Sigma_0) \|e^{-\lambda t} \phi_+^\sigma \phi_-^\tau u\|_{\mathcal{O}}^2 \\ & \leq C \{ \|me^{-\lambda t} \phi_+^\sigma \phi_-^\tau Lu\|_{\mathcal{O}}^2 + \|(\phi_+^\sigma \phi_-^\tau u)(0)\|_{\Omega}^2 \}. \end{aligned}$$

This implies the following a priori estimate.

**Proposition 6.2.** *For  $q \in \mathbf{Z}_+$ ,  $q \geq 1$  there is a  $\Sigma(q) > 0$  such that if  $\sigma, \tau > \Sigma(q)$  and if  $u \in C^{q+1}(\overline{\mathcal{O}})$ , with  $\operatorname{supp} u \cap (O^- \cup \gamma^-) = \emptyset$ , is a weak solution to (IBVP) then it follows that*

$$\begin{aligned} & \|u\|_{X_{(-\sigma, \tau)}^q(\mathcal{O}; \Gamma)}^2 \\ & \leq C \left\{ \|mLu\|_{X_{(-\sigma, \tau)}^q(\mathcal{O})}^2 + \sum_{k=0}^{q-1} \|(\partial_0^k Lu)(0)\|_{X_{0(-\sigma, \tau)}^{q-1-k}(\Omega)}^2 + \|u_0\|_{X_{0(-\sigma, \tau)}^q(\Omega)}^2 \right\} \end{aligned}$$

where  $C = C(q, \sigma, \tau) > 0$  is independent of  $u$ .

**Proof.** Localizing the problem as in Proposition 4.4 and repeating the same arguments as in Proposition 10.1 in [8] we can obtain that

$$\|u\|_{X_{(-\sigma, \tau)}^q(\mathcal{O}; \Gamma)}^2 \leq C \left\{ \|mLu\|_{X_{(-\sigma, \tau)}^q(\mathcal{O}; \Gamma)}^2 + \sum_{k=0}^q \|(\partial_0^k u)(0)\|_{X_{0(-\sigma, \tau)}^{q-k}(\Omega)}^2 \right\}.$$

Since  $\partial_0 = A_0^{-1}(L - \sum_{j=1}^n A_j \partial_j - B)$  we note that  $(\partial_0^k u)(0)$ ,  $k \geq 1$ , is written as a sum of  $(\partial_x^\alpha u)(0)$ ,  $|\alpha| \leq k$  and  $(\partial_x^\beta \partial_0^l Lu)(0)$ ,  $l + |\beta| \leq k - 1$ . This completes the proof.  $\square$

Let us set

$$\begin{aligned} \gamma_0^\pm &= \{x \in \partial\Omega; (0, x) \in \gamma^\pm\}, \quad \gamma_0 = \{x \in \partial\Omega; (0, x) \in \gamma\}, \\ O_0^\pm &= \{x \in \partial\Omega; (0, x) \in O^\pm\}. \end{aligned}$$

For the proof of Theorem 5.4 we shall extract from technical details and sum up in the following two lemmas.

**Lemma 6.3.** Let  $q \in \mathbb{Z}_+$  and set  $\tilde{q} = q + [n/2] + 2$ . Suppose that  $f \in H^{\tilde{q}}(\mathcal{O})$  and  $u_0 \in H^{\tilde{q}}(\Omega)$  with  $\text{supp} f \cap (O^+ \cup O^- \cup \gamma) = \emptyset$  and  $\text{supp} u_0 \cap (O_0^+ \cup O_0^- \cup \gamma_0) = \emptyset$  and that  $u \in X_{(-\sigma+\tilde{q}, \tau+\tilde{q})}^{\tilde{q}}(\mathcal{O}; \Gamma)$ , for some  $\sigma, \tau \geq \tilde{q}$ , is a weak solution to (IBVP) (we remark that  $X_{(-\sigma+\tilde{q}, \tau+\tilde{q})}^{\tilde{q}}(\mathcal{O}; \Gamma) \hookrightarrow X_{(-\sigma, \tau)}^q(\mathcal{O}; \Gamma)$ ). Then there exists a  $\{u_l\} \subset C^{q+1}(\overline{\mathcal{O}})$  with  $\text{supp} u_l \cap (O^+ \cup O^- \cup \gamma) = \emptyset$  and  $u_l \in M$  at  $\Gamma$  which satisfies that

$$\begin{aligned} mLu_l &\rightarrow mf && \text{in } X_{(-\sigma, \tau)}^q(\mathcal{O}), \\ (\partial_0^k Lu_l)(0) &\rightarrow (\partial_0^k f)(0) && \text{in } X_{0(-\sigma, \tau)}^{q-1-k}(\Omega), \quad k = 0, \dots, q-1, \\ u_l(0) &\rightarrow u_0 && \text{in } X_{0(-\sigma, \tau)}^q(\Omega), \\ u_l &\rightarrow u && \text{in } X_{(-\sigma, \tau)}^q(\mathcal{O}; \Gamma), \end{aligned}$$

as  $l \rightarrow \infty$ .

**Lemma 6.4.** Let  $f \in X_{(-\sigma, \tau)}^q(\mathcal{O})$  and  $u_0 \in X_{0(-\sigma, \tau)}^q(\Omega)$ , for  $q \in \mathbb{Z}_+$  and  $q \geq 1$ , satisfy the compatibility conditions up to order  $q-1$ . Then for  $q' \in \mathbb{Z}_+$ ,  $q' \geq q$  there exist  $\{f_l\} \subset H^{q'}(\mathcal{O})$  and  $\{u_{0l}\} \subset H^{q'}(\Omega)$  with  $\text{supp} f_l \cap (O^+ \cup O^- \cup \gamma) = \emptyset$  and  $\text{supp} u_{0l} \cap (O_0^+ \cup O_0^- \cup \gamma_0) = \emptyset$  such that  $f_l$  and  $u_{0l}$  satisfy the compatibility conditions up to order  $q'-1$  and moreover

$$(6.1) \quad mf_l \rightarrow mf \quad \text{in } X_{(-\sigma, \tau)}^q(\mathcal{O}),$$

$$(6.2) \quad (\partial_0^k f_l)(0) \rightarrow (\partial_0^k f)(0) \quad \text{in } X_{0(-\sigma, \tau)}^{q-1-k}(\Omega), \quad k = 0, \dots, q-1,$$

$$(6.3) \quad u_{0l} \rightarrow u_0 \quad \text{in } X_{0(-\sigma, \tau)}^q(\Omega),$$

as  $l \rightarrow \infty$ .

Admitting these lemmas we give the proof of Theorem 5.4.

**Proof of Theorem 5.4.** Let us set  $\tilde{q} = q + [n/2] + 2$  and  $q' = 2\tilde{q} + 1$ . First we suppose that  $f \in H^{q'}(\mathcal{O})$  and  $u_0 \in H^{q'}(\Omega)$ , with  $\text{supp} f \cap (O^+ \cup O^- \cup \gamma) = \emptyset$  and  $\text{supp} u_0 \cap (O_0^+ \cup O_0^- \cup \gamma_0) = \emptyset$ , satisfy the compatibility conditions up to order  $q'-1$ . Noticing that the rank of  $M(t, x)$  is constant on each component of  $\Gamma \setminus \text{supp} f$  and repeating the same arguments as in Lemma 3.1 in [13] we can find a  $w \in H^{\tilde{q}+1}(\mathcal{O})$  with  $\text{supp} w \cap (O^+ \cup O^- \cup \gamma) = \emptyset$  such that  $w(t, x) \in M(t, x)$  for  $(t, x) \in \Gamma$ ,  $w(0) = u_0$  and  $(\partial_0^k (Lw - f))(0) = 0$  for  $k = 0, \dots, \tilde{q}-1$ .

Now we set  $g = Lw$  and consider the following initial boundary value problem:

$$(IBVP') \quad \begin{cases} Lv = f - g & \text{in } I \times \Omega = \mathcal{O} \\ v \in M & \text{at } I \times \partial\Omega = \Gamma \\ v(0) = 0 & \text{on } \Omega. \end{cases}$$

Since  $f - g \in X_{(-\sigma+\bar{q}, \tau+\bar{q})}^{\bar{q}}(\mathcal{O}; \Gamma) \cap \phi_- L^2(\mathcal{O})$  and  $(\partial_0^k(f - g))(0) = 0$  for  $k = 0, \dots, \bar{q} - 1$  Theorem 2.4 gives a weak solution  $v \in X_{(-\sigma+\bar{q}, \tau+\bar{q})}^{\bar{q}}(\mathcal{O}; \Gamma)$  to (IBVP'). If we set  $u = v + w$  then it follows that  $u \in X_{(-\sigma+\bar{q}, \tau+\bar{q})}^{\bar{q}}(\mathcal{O}; \Gamma)$  and  $u$  is a weak solution to (IBVP). Moreover combining Proposition 6.2 and Lemma 6.3 we have

$$\begin{aligned} & \|u\|_{X_{(-\sigma, \tau)}^q(\mathcal{O}; \Gamma)}^2 \\ & \leq C \left\{ \|mLu\|_{X_{(-\sigma, \tau)}^q(\mathcal{O})}^2 + \sum_{k=0}^{q-1} \|(\partial_0^k Lu)(0)\|_{X_{0(-\sigma, \tau)}^{q-1-k}(\Omega)}^2 + \|u_0\|_{X_{0(-\sigma, \tau)}^q(\Omega)}^2 \right\} \\ & \leq C' \{ \|f\|_{X_{(-\sigma, \tau)}^q(\mathcal{O})}^2 + \|u_0\|_{X_{0(-\sigma, \tau)}^q(\Omega)}^2 \}. \end{aligned}$$

Next we suppose that  $f \in X_{(-\sigma, \tau)}^q(\mathcal{O})$  and  $u_0 \in X_{0(-\sigma, \tau)}^q(\Omega)$  satisfy the compatibility conditions up to order  $q - 1$ . Then by using Lemma 6.4 and standard limiting argument we conclude the assertion.  $\square$

**Proof of Lemma 6.3.** Proposition 4.4 implies that  $u \in m^{-\bar{q}} X_{(-\sigma+\bar{q}, \tau+\bar{q})}^{\bar{q}}(\mathcal{O}) \hookrightarrow X_{(-\sigma, \tau)}^{\bar{q}}(\mathcal{O})$ . Let us choose  $\chi \in C_0^\infty(\mathbf{R})$  so that  $\chi = 1$  near 0 and set

$$\alpha_l(t, x) = 1 - \chi(l(\phi_+ \phi_-)(t, x)), \quad u_l(t, x) = (\alpha_l u)(t, x)$$

for  $l > 0$  large enough. Then  $\{u_l\}$  is a desired sequence. Indeed since  $\alpha_l \in C_0^\infty(\overline{\mathcal{R}})$  and  $\alpha_l = 0$  on a neighborhood of  $O^+ \cup O^- \cup \gamma$  it follows that  $u_l \in H^{\bar{q}}(\mathcal{O}) \hookrightarrow C^{q+1}(\overline{\mathcal{O}})$ . Thus it is easily checked that  $u_l \in M$  at  $\Gamma$ . For the proof of the desired convergences it suffices to show that

$$(6.4) \quad Lu_l \rightarrow f \quad \text{in } X_{(-\sigma, \tau)}^q(\mathcal{O}), \quad u_l \rightarrow u \quad \text{in } X_{(-\sigma, \tau)}^{q+1}(\mathcal{O}) \quad \text{as } l \rightarrow \infty.$$

Note that  $f \in X_{(-\sigma, \tau)}^q(\mathcal{O})$ ,  $\phi_+^{-1} \phi_-^{-1} u \in X_{(-\sigma, \tau)}^q(\mathcal{O})$ ,  $u \in X_{(-\sigma, \tau)}^{q+1}(\mathcal{O})$  and

$$\begin{aligned} Lu_l - f &= -\chi(l\phi_+ \phi_-)f - \tilde{\chi}(l\phi_+ \phi_-) \sum_{j=0}^n (\partial_j \phi_+ \phi_-) A_j \phi_+^{-1} \phi_-^{-1} u, \\ u_l - u &= -\chi(l\phi_+ \phi_-)u \end{aligned}$$

where  $\tilde{\chi}(\theta) = \theta \chi'(\theta)$ . Thus using arguments similar to those in Lemma 6.5 in [8] we can prove (6.4).  $\square$

**Proof of Lemma 6.4.** The proof of Lemma 6.4 proceeds in three steps.

**FIRST STEP:** If we write  $u^{(k)} = T_k(f, u_0)$  for  $k = 0, \dots, q - 1$  then we can find a  $u \in X_{(-\sigma, \tau)}^q(\mathcal{O})$  such that  $(\partial_0^k u)(0) = u^{(k)}$  for  $k = 0, \dots, q - 1$  (we give the proof

of this fact in Proposition 7.1 below). Now let us choose  $\chi \in C_0^\infty(\mathbf{R})$  so that  $\chi = 1$  near 0 and set with  $\alpha_l(t, x) = 1 - \chi(lm(t, x))$

$$(6.5) \quad f_l(t, x) = (\alpha_l f)(t, x) - \sum_{j=0}^n ((\partial_j \alpha_l) A_j u)(t, x), \quad u_{0l}(x) = (\alpha_l(0) u_0)(x)$$

for  $l > 0$  large enough. Then we remark that  $f_l \in X_{(-\sigma, \tau)}^q(\mathcal{O})$  and  $u_{0l} \in X_{0(-\sigma, \tau)}^q(\Omega)$  with  $\text{supp} f_l \cap \gamma = \emptyset$  and  $\text{supp} u_{0l} \cap \gamma_0 = \emptyset$ .

**Lemma 6.5.** *Let  $f_l$  and  $u_{0l}$  be given by (6.5). Then*

- (i)  $f_l$  and  $u_{0l}$  satisfy the compatibility conditions up to order  $q - 1$ .
- (ii)  $f_l$  and  $u_{0l}$  satisfy (6.1), (6.2) and (6.3).

*Proof.* We first consider the assertion (i). We note that

$$(\phi_+^\sigma \phi_-^{-\tau})(0)(\partial_0^k P u)(0) = \sum_{i=0}^k \binom{k}{i} (\partial_0^i P)(0)(\phi_+^\sigma \phi_-^{-\tau})(0) u^{(k-i)} = 0 \quad \text{on } \partial\Omega \setminus \gamma_0$$

for  $k = 0, \dots, q - 1$ . If we write  $u_l^{(k)} = T_k(f_l, u_{0l})$  for  $k = 0, \dots, q - 1$  then we have  $u_l^{(k)} = (\partial_0^k \alpha_l u)(0)$ , and hence it follows that

$$\begin{aligned} \sum_{i=0}^k \binom{k}{i} (\partial_0^i P)(0)(\phi_+^\sigma \phi_-^{-\tau})(0) u_l^{(k-i)} &= (\phi_+^\sigma \phi_-^{-\tau})(0)(\partial_0^k P \alpha_l u)(0) \\ &= \sum_{i=0}^k \binom{k}{i} (\partial_0^i \alpha_l)(0)(\phi_+^\sigma \phi_-^{-\tau})(0)(\partial_0^{k-i} P u)(0) = 0 \quad \text{on } \partial\Omega \setminus \gamma_0. \end{aligned}$$

We turn to the assertion (ii). With  $\tilde{\chi}(\theta) = \theta \chi'(\theta)$  we can write

$$f_l - f = -\chi(lm)f - m^{-1} \tilde{\chi}(lm) \sum_{j=0}^n (\partial_j m) A_j u, \quad u_{0l} - u_0 = -\chi(lm(0)) u_0.$$

Therefore (6.1) and (6.3) are easily checked and since  $(\partial_0^k f)(0) \in X_{0(-\sigma, \tau)}^{q-1-k}(\Omega)$  and  $(\partial_0^k u)(0) = u^{(k)} \in X_{0(-\sigma, \tau)}^{q-k}(\Omega)$  for  $k = 0, \dots, q - 1$  we can prove (6.2).  $\square$

In what follows, we may assume that  $\text{supp} f \cap \gamma = \emptyset$  and  $\text{supp} u_0 \cap \gamma_0 = \emptyset$ .

**SECOND STEP:** Let  $\chi \in C_0^\infty(\mathbf{R})$  be as in First step and set with  $\alpha_l(t, x) = 1 - \chi(l(\phi_+ \phi_-)(t, x))$

$$(6.6) \quad f_l(t, x) = (\alpha_l f)(t, x), \quad u_{0l}(x) = (\alpha_l(0) u_0)(x)$$

for  $l > 0$  large enough. Then we remark that  $f_l \in X_{(-\sigma, \tau)}^q(\mathcal{O})$  and  $u_{0l} \in X_{0(-\sigma, \tau)}^q(\Omega)$  with  $\text{supp} f_l \cap (O^+ \cup O^- \cup \gamma) = \emptyset$  and  $\text{supp} u_{0l} \cap (O_0^+ \cup O_0^- \cup \gamma_0) = \emptyset$ . In particular,

this implies that  $f_l \in H^q(\mathcal{O})$  and  $u_{0l} \in H^q(\Omega)$ .

**Lemma 6.6.** *Let  $f_l$  and  $u_{0l}$  be given by (6.6). Then the same conclusion as in Lemma 6.5 holds.*

*Proof.* We first consider the assertion (i). Noticing that  $\alpha_l = 1$  near  $\Gamma \setminus (O^+ \cup O^- \cup \text{supp} f)$  and  $\alpha_l = 0$  near  $(O^+ \cup O^-) \setminus \text{supp} f$  and that  $f_l = f = 0$  on  $\text{supp} f$  we obtain that

$$f_l = f \quad \text{near } \Gamma \setminus (O^+ \cup O^-), \quad f_l = 0 \quad \text{near } O^+ \cup O^- \cup \gamma_0.$$

Similarly we have

$$u_{0l} = u_0 \quad \text{near } \partial\Omega \setminus (O_0^+ \cup O_0^-), \quad u_{0l} = 0 \quad \text{near } O_0^+ \cup O_0^- \cup \gamma_0.$$

Therefore if we write  $u^{(k)} = T_k(f, u_0)$  and  $u_l^{(k)} = T_k(f_l, u_{0l})$  for  $k = 0, \dots, q-1$  then it follows that

$$u_l^{(k)} = u^{(k)} \quad \text{near } \partial\Omega \setminus (O_0^+ \cup O_0^-), \quad u_l^{(k)} = 0 \quad \text{near } O_0^+ \cup O_0^- \cup \gamma_0.$$

This proves the assertion (i). The assertion (ii) is easily checked.  $\square$

In what follows, we may assume that  $f \in H^q(\mathcal{O})$  and  $u_0 \in H^q(\Omega)$  with  $\text{supp} f \subset \overline{\mathcal{O}} \cap \{\phi_+ > \eta, \phi_- > \eta\}$  and  $\text{supp} u_0 \subset \overline{\Omega} \cap \{\phi_+(0) > \eta, \phi_-(0) > \eta\}$  for some  $\eta > 0$ .

**THIRD STEP:** Recalling that  $A_b(t, x)$  is non singular on  $\Gamma \cap \{\phi_+ > \eta, \phi_- > \eta\}$  and using the same arguments as in Lemma 3.3 in [13] we can find  $\{f_l\} \subset H^{q'}(\mathcal{O})$  and  $\{u_{0l}\} \subset H^{q'}(\Omega)$ , with  $\text{supp} f_l \subset \overline{\mathcal{O}} \cap \{\phi_+ > \delta, \phi_- > \delta\}$  and  $\text{supp} u_{0l} \subset \overline{\Omega} \cap \{\phi_+(0) > \delta, \phi_-(0) > \delta\}$  for some  $\delta = \delta(\eta) > 0$ , such that  $f_l$  and  $u_{0l}$  satisfy the compatibility conditions up to order  $q' - 1$  and it follows that

$$f_l \rightarrow f \quad \text{in } H^q(\mathcal{O}), \quad u_{0l} \rightarrow u_0 \quad \text{in } H^q(\Omega) \quad \text{as } l \rightarrow \infty.$$

In particular, this implies that

$$f_l \rightarrow f \quad \text{in } X_{(-\sigma, \tau)}^q(\mathcal{O}), \quad u_{0l} \rightarrow u_0 \quad \text{in } X_{0(-\sigma, \tau)}^q(\Omega) \quad \text{as } l \rightarrow \infty$$

which shows (6.1), (6.2) and (6.3). Therefore  $\{f_l\}$  and  $\{u_{0l}\}$  are desired sequences. Thus we conclude the proof of Lemma 6.4.  $\square$

## 7. Auxiliary lemmas

In this section we first show the following proposition which is used in the proof of Lemma 6.4.

**Proposition 7.1.** *Let  $q \in \mathbf{Z}_+$ ,  $q \geq 1$  and  $\sigma, \tau \in \mathbf{R}$ . If  $u^{(k)} \in X_{0(\sigma, \tau)}^{q-k}(\Omega)$  for  $k = 0, \dots, q-1$  then there exists a  $u \in X_{(\sigma, \tau)}^q(\mathbf{R} \times \Omega)$  with  $(\partial_0^k u)(0) = u^{(k)}$ ,  $k = 0, \dots, q-1$ , such that*

$$\|u\|_{X_{(\sigma, \tau)}^q(\mathbf{R}_+ \times \Omega)} \leq C \sum_{k=0}^{q-1} \|u^{(k)}\|_{X_{0(\sigma, \tau)}^{q-k}(\Omega)}$$

where  $C = C(q, \sigma, \tau) > 0$  is independent of  $u^{(k)}$  and  $u$ .

For the proof of Proposition 7.1 it suffices to prove the assertion for  $\sigma = \tau = -q$ . Indeed assume that the statement for  $\sigma = \tau = -q$  is true. We consider the general case. Let  $u^{(k)} \in X_{0(\sigma, \tau)}^{q-k}(\Omega)$  for  $k = 0, \dots, q-1$ . We define  $v^{(k)}$ ,  $k = 0, \dots, q-1$ , as follows:

$$\begin{aligned} v^{(0)} &= (\phi_+^{-\sigma-q} \phi_-^{-\tau-q})(0) u^{(0)}, \\ v^{(k)} &= (\phi_+^{-\sigma-q} \phi_-^{-\tau-q})(0) \left\{ u^{(k)} - \sum_{i=0}^{k-1} \binom{k}{i} (\partial_0^{k-i} \phi_+^{\sigma+q} \phi_-^{\tau+q})(0) v^{(i)} \right\} \quad \text{for } k \geq 1. \end{aligned}$$

Then for each  $v^{(k)}$  it follows that  $v^{(k)} \in X_{0(-q, -q)}^{q-k}(\Omega)$  and

$$\|v^{(k)}\|_{X_{0(-q, -q)}^{q-k}(\Omega)} \leq C \sum_{i=0}^k \|u^{(i)}\|_{X_{0(\sigma, \tau)}^{q-i}(\Omega)}.$$

By the hypothesis we can find a  $v \in X_{(-q, -q)}^q(\mathcal{O})$ , with  $(\partial_0^k v)(0) = v^{(k)}$  for  $k = 0, \dots, q-1$ , such that

$$\|v\|_{X_{(-q, -q)}^q(\mathcal{O})} \leq C \sum_{k=0}^{q-1} \|v^{(k)}\|_{X_{0(-q, -q)}^{q-k}(\Omega)}.$$

If we set  $u = \phi_+^{\sigma+q} \phi_-^{\tau+q} v$  then we have  $u \in X_{(\sigma, \tau)}^q(\mathcal{O})$ . Noticing that  $\partial_0^k u = \sum_{i=0}^k \binom{k}{i} (\partial_0^{k-i} \phi_+^{\sigma+q} \phi_-^{\tau+q}) \partial_0^i v$  we obtain  $(\partial_0^k u)(0) = u^{(k)}$  for  $k = 0, \dots, q-1$ . Moreover it follows that

$$\begin{aligned} \|u\|_{X_{(\sigma, \tau)}^q(\mathcal{O})} &= \|v\|_{X_{(-q, -q)}^q(\mathcal{O})} \\ &\leq C \sum_{k=0}^{q-1} \|v^{(k)}\|_{X_{0(-q, -q)}^{q-k}(\Omega)} \leq C' \sum_{k=0}^{q-1} \|u^{(k)}\|_{X_{0(\sigma, \tau)}^{q-k}(\Omega)}. \end{aligned}$$

Therefore  $u$  is a desired function, and hence we conclude the assertion for  $\sigma, \tau \in \mathbf{R}$ .

To prove Proposition 7.1 for  $\sigma = \tau = -q$  we shall localize the problem. Let us take a covering  $\{U_i\}_{i=0}^l$  of  $\{t = 0\} \times \Omega$  as follows: First we cover  $\{t = 0\} \times \partial\Omega$

by coordinate patches  $U_i$ ,  $i = 1, \dots, l$ , with coordinate systems  $\chi_i : U_i \cap (\mathbf{R} \times \Omega) \rightarrow \{(\tau, \xi); |\tau| < \delta, |\xi| < 1, \xi_1 > 0\}$  such that  $\tau = t \circ \chi_i^{-1}$  and  $\xi_1 = r \circ \chi_i^{-1}$  where  $\delta > 0$  is small enough. Next we cover  $(\{t = 0\} \times \Omega) \setminus \bigcup_{i=1}^l U_i$  by  $U_0 \subset \subset \mathbf{R} \times \Omega$ . Choose a partition of unity  $\{\psi_i\}_{i=0}^l$  subordinate to this covering  $\{U_i\}_{i=0}^l$  and set  $u_i^{(k)} = \psi_i u^{(k)}$ . It suffices to show Proposition 7.1 for  $\sigma = \tau = -q$  with  $u_i^{(k)}$  instead of  $u^{(k)}$ . Performing a change of independent variables we are led to the case where

$$U = \{(t, x); |t| < \delta, |x| < 1\}, \quad \Omega = \mathbf{R}_+^n = \{x; x_1 > 0\}, \quad r = x_1, \\ \text{supp} u^{(k)} \subset \{x; |x| < 1 - \epsilon_0, x_1 \geq 0\} \quad \text{for } k = 0, \dots, q-1$$

with  $\epsilon_0 > 0$  small enough.

Now suppose that  $q \in \mathbf{Z}_+$  ( $q \geq 1$ ) is given. For a fixed  $k \in \mathbf{Z}_+$  ( $0 \leq k \leq q-1$ ) and a fixed  $v \in X_{0(-q, -q)}^{q-k}(\mathbf{R}_+^n)$  we consider the following functions:

$$w(t, x) = \psi(t)t^k \Phi(t, x) V(t, x), \quad \Phi(t, x) = \chi(t(\phi_+^{-1} \phi_-^{-1})(t, x)), \\ V(t, x) = \int_{\mathbf{R}^n} v(x + ty) \rho(y) dy \quad \text{for } (t, x) \in \overline{\mathbf{R}}_+ \times \mathbf{R}_+^n$$

where

$$(7.1) \quad \psi \in C_0^\infty(\mathbf{R}) \quad \text{with} \quad \text{supp} \psi \subset \{t; |t| < \delta\},$$

$$(7.2) \quad \chi \in C_0^\infty(\mathbf{R}) \quad \text{with} \quad \text{supp} \chi \subset \{\theta; |\theta| < 1\},$$

$$(7.3) \quad \rho \in C_0^\infty(\mathbf{R}^n) \quad \text{with} \quad \text{supp} \rho \subset \{y; |y| < \epsilon_0, y_1 > \epsilon_0/2, y_2 < 0\}$$

and they satisfy  $\psi = 1$  near 0,  $\chi(0) = 1$  and  $\int \rho(y) dy = 1$ . Then we obtain the following two results.

**Lemma 7.2.** *It follows that  $w \in X_{(-q, -q)}^q(\mathbf{R}_+ \times \mathbf{R}_+^n)$  and*

$$\|w\|_{X_{(-q, -q)}^q(\mathbf{R}_+ \times \mathbf{R}_+^n)} \leq C \|v\|_{X_{0(-q, -q)}^{q-k}(\mathbf{R}_+^n)}$$

where  $C = C(q, k) > 0$ .

**Lemma 7.3.**  *$(\partial_0^i w)(0, x)$ ,  $i = 0, \dots, q-1$ , has the following properties:*

- (i)  $(\partial_0^i w)(0, x) = 0$  for  $i = 0, \dots, k-1$  and  $(\partial_0^k w)(0, x) = v(x)$ .
- (ii)  $(\partial_0^i w)(0) \in X_{0(-q, -q)}^{q-i}(\mathbf{R}_+^n)$  with  $\text{supp}(\partial_0^i w)(0) \subset \{x; |x| < 1 - \epsilon_0, x_1 \geq 0\}$  and

$$\|(\partial_0^i w)(0)\|_{X_{0(-q, -q)}^{q-i}(\mathbf{R}_+^n)} \leq C \|v\|_{X_{0(-q, -q)}^{q-k}(\mathbf{R}_+^n)}$$

where  $C = C(q, k, i) > 0$ .

Admitting that these results hold we shall complete the proof of Proposition 7.1.

**Proof of Proposition 7.1.** Let  $u^{(k)} \in X_{0(-q,-q)}^{q-k}(\mathbf{R}_+^n)$  with  $\text{supp} u^{(k)} \subset \{x; |x| < 1 - \epsilon_0, x_1 \geq 0\}$  for  $k = 0, \dots, q-1$ . Let us set

$$u(t, x) = \sum_{k=0}^{q-1} w_k(t, x), \quad w_k(t, x) = \psi(t) t^k \Phi(t, x) \int v^{(k)}(x + ty) \rho(y) dy$$

where  $\psi$ ,  $\rho$  and  $\Phi$  are as above. Here we define  $v^{(i)}$ ,  $i = 0, \dots, q-1$ , as follows:

$$v^{(0)} = u^{(0)}, \quad v^{(i)} = u^{(i)} - \sum_{k=0}^{i-1} (\partial_0^i w_k)(0) \quad \text{for } i \geq 1.$$

Then for each  $v^{(i)}$  it follows from Lemma 7.3 that  $v^{(i)} \in X_{0(-q,-q)}^{q-i}(\mathbf{R}_+^n)$  with  $\text{supp} v^{(i)} \subset \{x; |x| < 1 - \epsilon_0, x_1 \geq 0\}$  and

$$\|v^{(i)}\|_{X_{0(-q,-q)}^{q-i}(\mathbf{R}_+^n)} \leq C \|u^{(k)}\|_{X_{0(-q,-q)}^{q-k}(\mathbf{R}_+^n)},$$

and hence Lemma 7.2 shows that  $u \in X_{(-q,-q)}^q(\mathbf{R}_+ \times \mathbf{R}_+^n)$  and

$$\begin{aligned} \|u\|_{X_{(-q,-q)}^q(\mathbf{R}_+ \times \mathbf{R}_+^n)} &\leq \sum_{k=0}^{q-1} \|w_k\|_{X_{(-q,-q)}^q(\mathbf{R}_+ \times \mathbf{R}_+^n)} \\ &\leq C \sum_{k=0}^{q-1} \|v^{(k)}\|_{X_{0(-q,-q)}^{q-k}(\mathbf{R}_+^n)} \leq C' \sum_{k=0}^{q-1} \|u^{(k)}\|_{X_{0(-q,-q)}^{q-k}(\mathbf{R}_+^n)}. \end{aligned}$$

Moreover we have  $(\partial_0^k u)(0) = u^{(k)}$  for  $k = 0, \dots, q-1$ . Therefore  $u$  is a desired function.  $\square$

We shall show Lemma 7.2 and Lemma 7.3. For the proofs we prepare several lemmas.

**Lemma 7.4.**  $t^i(\partial^\alpha V)(t, x)$ ,  $0 \leq i \leq |\alpha|$ , is written as a sum of the following terms:

$$\int (\partial_x^\beta v)(x + ty) \tilde{\rho}(y) dy, \quad |\beta| = |\alpha| - i$$

where  $\tilde{\rho}$  is of type (7.3).

**Proof.** We first consider the case  $i = 0$ . Since  $\partial_0\{v(x + ty)\} = \sum_{j=1}^n y_j(\partial_j v)(x + ty)$  the assertion for  $i = 0$  is clear. We turn to the case  $i \geq 1$ . For  $j = 1, \dots, n$  it



follows from  $t(\partial_j v)(x + ty) = \partial_{y_j} \{v(x + ty)\}$  that

$$t \int (\partial_j v)(x + ty) \tilde{\rho}(y) dy = \int v(x + ty) (-\partial_j \tilde{\rho})(y) dy.$$

Thus the assertion is proved.  $\square$

From this we obtain the following lemma which is easily checked.

**Lemma 7.5.**  $(\partial^\alpha w)(t, x)$ ,  $|\alpha| \leq q$ , is written as a sum of the following terms:

$$\begin{aligned} \tilde{\psi}(t) t^{k-j} (\partial^\beta \Phi)(t, x) \int (\partial_x^\gamma v)(x + ty) \tilde{\rho}(y) dy, \\ j + |\beta| + |\gamma| \leq |\alpha|, \quad 0 \leq j \leq k, \quad |\gamma| \leq q - k \end{aligned}$$

where  $\tilde{\psi}$  and  $\tilde{\rho}$  are of type (7.1) and (7.3) respectively.

To get the estimate for  $w$  the following lemmas will be used.

**Lemma 7.6.** For  $i \in \mathbf{Z}_+$  and  $\alpha \in \mathbf{Z}_+^{n+1}$  there is a  $C > 0$  such that

$$|t^i (\partial^\alpha \Phi)(t, x)| \leq C (\phi_+^{i-|\alpha|} \phi_-^{i-|\alpha|})(t, x) \quad \text{for } 0 \leq t \leq \delta, \quad |x| \leq 1, \quad x_1 \geq 0.$$

**Proof.** We first consider the case  $|\alpha| = 0$ . If we write  $\tilde{\chi}(\theta) = \theta^i \chi(\theta)$  then it follows that  $t^i \Phi(t, x) = (\phi_+^i \phi_-^i)(t, x) \tilde{\chi}(t(\phi_+^{-1} \phi_-^{-1})(t, x))$  which proves the assertion for  $|\alpha| = 0$ . We turn to the case  $|\alpha| \geq 1$ . Note that  $t^i (\partial^\alpha \Phi)(t, x)$  is written as a sum of the following terms:

$$\begin{aligned} t^{i+j} (\partial^{\beta_1} \phi_+^{-1} \phi_-^{-1})(t, x) \cdots (\partial^{\beta_l} \phi_+^{-1} \phi_-^{-1})(t, x) \chi^{(l)}(t(\phi_+^{-1} \phi_-^{-1})(t, x)), \\ 1 \leq l \leq |\alpha|, \quad 0 \leq j \leq l, \quad |\beta_1| + \cdots + |\beta_l| + l = |\alpha| + j \end{aligned}$$

with  $\chi^{(l)}(\theta) = d^l \chi(\theta) / d\theta^l$ . Using  $|\partial^\beta \phi_+^{-1} \phi_-^{-1})(t, x)| \leq C(\phi_+^{-|\beta|-1} \phi_-^{-|\beta|-1})(t, x)$  and repeating the same arguments as above we can conclude the proof.  $\square$

**Lemma 7.7.** Taking  $\mu > 0$  large enough we have

$$\phi_\pm(t, x) \leq \phi_\pm(0, x + ty) \quad \text{for } 0 \leq t \leq \delta, \quad |x| \leq 1, \quad x_1 \geq 0, \quad |y| < \epsilon_0, \quad y_1 > \epsilon_0/2.$$

**Proof.** If we set  $f(\xi, \eta) = (\kappa \xi^2 + \eta^2)^{1/2} + \eta$  for  $(\xi, \eta) \in \mathbf{R}_+ \times \mathbf{R}$  then we can write  $\phi_\pm(t, x) = f(x_1, \mu x_1 - h_\pm(t, x))$ . Since  $(\partial_\xi f)(\xi, \eta) \geq 0$  and  $(\partial_\eta f)(\xi, \eta) \geq 0$  it

suffices to show that  $\mu x_1 - h_{\pm}(t, x) \leq \mu(x_1 + ty_1) - h_{\pm}(0, x + ty)$  because  $x_1 \leq x_1 + ty_1$ . Since  $|h_{\pm}(0, x + ty) - h_{\pm}(t, x)| \leq Ct$  for some  $C > 0$  it follows that

$$\begin{aligned} & \{\mu(x_1 + ty_1) - h_{\pm}(0, x + ty)\} - \{\mu x_1 - h_{\pm}(t, x)\} \\ &= \mu ty_1 - (h_{\pm}(0, x + ty) - h_{\pm}(t, x)) \geq (\mu \epsilon_0/2 - C)t. \end{aligned}$$

Therefore taking  $\mu > 0$  large enough we can prove the assertion.  $\square$

The following lemma is easily checked.

**Lemma 7.8.** *Let  $u \in L^2(\mathbf{R}_+^n)$  with  $\text{supp } u \subset \{|x| \leq 1, x_1 \geq 1\}$  and let  $\tilde{\rho}$  be of type (7.3). Suppose that  $a(t, x, y)$  with  $\text{supp } a \subset \{0 \leq t \leq \delta\}$  satisfies  $|a(t, x, y)| \leq C$  for  $t \in \mathbf{R}_+$ ,  $|x| \leq 1$ ,  $x_1 \geq 1$ ,  $y \in \text{supp } \tilde{\rho}$  where  $C > 0$  is independent of  $t, x$  and  $y$ . If we set*

$$U(t, x) = \int a(t, x, y)u(x + ty)\tilde{\rho}(y)dy$$

then it follows that  $U(t, x) \in L^2(\mathbf{R}_+ \times \mathbf{R}_+^n)$  and

$$\|U\|_{\mathbf{R}_+ \times \mathbf{R}_+^n} \leq C' \|u\|_{\mathbf{R}_+^n}$$

where  $C' > 0$  is independent of  $u$  and  $U$ .

Now we give the proofs of Lemma 7.2 and Lemma 7.3

**Proof of Lemma 7.2.** By using a reasoning similar to that in Lemma 6.1 in [8] it suffices to show that

$$\|\phi_+^{|\alpha|}\phi_-^{|\alpha|}\partial^\alpha w\|_{\mathbf{R}_+ \times \mathbf{R}_+^n} \leq C \sum_{|\beta| \leq q-k} \|(\phi_+^{k+|\beta|}\phi_-^{k+|\beta|})(0)\partial_x^\beta v\|_{\mathbf{R}_+^n}$$

for  $|\alpha| \leq q$ . From Lemma 7.5 we recall that  $\phi_+^{|\alpha|}\phi_-^{|\alpha|}\partial^\alpha w$  is written as a sum of the following terms:

$$\int a(t, x, y)((\phi_+^{k+|\gamma|}\phi_-^{k+|\gamma|})(0)\partial_x^\gamma v)(x + ty)\tilde{\rho}(y)dy$$

where

$$\begin{aligned} a(t, x, y) &= \tilde{\psi}(t)(\phi_+^{|\alpha|}\phi_-^{|\alpha|})(t, x)t^{k-j}(\partial^\beta \Phi)(t, x)(\phi_+^{-k-|\gamma|}\phi_-^{-k-|\gamma|})(0, x + ty), \\ & j + |\beta| + |\gamma| \leq |\alpha|, \quad 0 \leq j \leq k, \quad |\gamma| \leq q - k. \end{aligned}$$

From Lemma 7.6 and Lemma 7.7 it follows that  $|a(t, x, y)| \leq C$  for some  $C > 0$ , and hence using Lemma 7.8 we conclude the proof.  $\square$

Proof of Lemma 7.3. The assertion (i) is clear. We consider the assertion (ii). We may assume  $k \leq i \leq q-1$ . For the proof it suffices to show that

$$(7.4) \quad \|(\phi_+^{i+|\alpha|}\phi_-^{i+|\alpha|})(0)\partial_x^\alpha(\partial_0^i w)(0)\|_{\mathbf{R}_+^n} \leq C \sum_{|\gamma| \leq q-k} \|(\phi_+^{k+|\gamma|}\phi_-^{k+|\gamma|})(0)\partial_x^\gamma v\|_{\mathbf{R}_+^n}$$

for  $|\alpha| \leq q-i$ . Lemma 7.5 shows that  $\partial_x^\alpha(\partial_0^i w)(0, x)$  is written as a sum of  $(\partial^\beta \Phi)(0, x)(\partial_x^\gamma v)(x)$ ,  $|\beta|+|\gamma| \leq i+|\alpha|-k$ . Thus the left-hand side of (7.4) is bounded from above by a sum of the following terms:

$$\|a_{\beta, \gamma}(\cdot)(\phi_+^{k+|\gamma|}\phi_-^{k+|\gamma|})(0)\partial_x^\gamma v\|_{\mathbf{R}_+^n}$$

where  $a_{\beta, \gamma}(x) = (\phi_+^{i+|\alpha|-k-|\gamma|}\phi_-^{i+|\alpha|-k-|\gamma|}\partial^\beta \Phi)(0, x)$ . From Lemma 7.6 it follows that  $|a_{\beta, \gamma}(x)| \leq C$  for some  $C > 0$ , and hence we can prove the assertion (ii).  $\square$

An immediate consequence of Lemma 7.1 is

**Corollary 7.9.** *Let  $q \in \mathbf{Z}_+$  and  $\sigma, \tau \in \mathbf{R}$ . For  $f \in X_{(\sigma, \tau)}^q(\mathbf{R}_- \times \Omega)$  there exists a  $\tilde{f} \in X_{(\sigma, \tau)}^q(\mathbf{R} \times \Omega)$  with  $\tilde{f} = f$  on  $\mathbf{R}_- \times \Omega$  such that*

$$\|\tilde{f}\|_{X_{(\sigma, \tau)}^q(\mathbf{R} \times \Omega)} \leq C\|f\|_{X_{(\sigma, \tau)}^q(\mathbf{R}_- \times \Omega)}$$

where  $C = C(q, \sigma, \tau) > 0$  is independent of  $\tilde{f}$  and  $f$ .

We can also obtain the following proposition.

**Proposition 7.10.** *Let  $q \in \mathbf{Z}_+$ ,  $q \geq 1$  and  $\sigma, \tau \in \mathbf{R}$ . If  $u^{(k)} \in X_{0(\sigma, \tau)}^{q-k}(\Omega; \partial\Omega)$  for  $k = 0, \dots, q-1$  there exists a  $u \in X_{(\sigma, \tau)}^q(\mathbf{R} \times \Omega; \mathbf{R} \times \partial\Omega)$  with  $(\partial_0^k u)(0) = u^{(k)}$ ,  $k = 0, \dots, q-1$ , such that*

$$\|u\|_{X_{(\sigma, \tau)}^q(\mathcal{O}; \Gamma)} \leq C \sum_{k=0}^{q-1} \|u^{(k)}\|_{X_{0(\sigma, \tau)}^{q-k}(\Omega; \partial\Omega)}$$

where  $C = C(q, \sigma, \tau) > 0$  is independent of  $u^{(k)}$  and  $u$ .

An immediate corollary to this proposition is

**Corollary 7.11.** *Let  $q \in \mathbf{Z}_+$  and  $\sigma, \tau \in \mathbf{R}$ . For  $f \in X_{(\sigma, \tau)}^q(\mathbf{R}_- \times \Omega; \mathbf{R}_- \times \partial\Omega)$  there exists a  $\tilde{f} \in X_{(\sigma, \tau)}^q(\mathbf{R} \times \Omega; \mathbf{R} \times \partial\Omega)$  with  $\tilde{f} = f$  on  $\mathbf{R}_- \times \Omega$  such that*

$$\|\tilde{f}\|_{X_{(\sigma, \tau)}^q(\mathbf{R} \times \Omega; \mathbf{R} \times \partial\Omega)} \leq C\|f\|_{X_{(\sigma, \tau)}^q(\mathbf{R}_- \times \Omega; \mathbf{R}_- \times \partial\Omega)}$$

where  $C = C(q, \sigma, \tau) > 0$  is independent of  $\tilde{f}$  and  $f$ .

**Proof of Proposition 7.10.** By the same arguments as in Proposition 7.1 it suffices to prove Proposition 7.10 for  $\sigma = -q$  and  $\tau = q$ . By localization we may assume that  $u^{(k)} \in X_{0(-q,q)}^{q-k}(\mathbf{R}_+^n)$  with  $\text{supp} u^{(k)} \subset \{x; |x| < 1 - \epsilon_0, x_1 \geq 0\}$  for  $k = 0, \dots, q-1$ . Now let us set

$$u(t, x) = \sum_{k=0}^{q-1} w_k(t, x), \quad w_k(t, x) = \psi(t) t^k \Phi(t, x) \int v^{(k)}(x_1 e^{ty_1}, x' + ty') \rho(y) dy$$

with  $x = (x_1, x') = (x_1, x_2, \dots, x_n)$  where  $\psi, \rho, \Phi$  and  $v^{(i)}, i = 0, \dots, q-1$ , are as in the proof of Proposition 7.1. Then  $u$  is shown to be a desired function using the following lemma instead of Lemma 7.7.  $\square$

**Lemma 7.12.** *Taking coordinate patches  $U$  small enough and coordinate systems  $\chi$  appropriate, if necessary, we obtain that*

$$(7.5) \quad \phi_+(t, x) \leq C \phi_+(0, x_1 e^{ty_1}, x' + ty'),$$

$$(7.6) \quad \phi_-^{-1}(t, x) \leq C \phi_-^{-1}(0, x_1 e^{ty_1}, x' + ty')$$

for  $0 \leq t \leq \delta, |x| \leq 1, x_1 \geq 0, |y| < \epsilon_0, y_2 < 0$  where  $C > 0$  is independent of  $t, x$  and  $y$ .

**Proof of Lemma 7.12.** Let us set  $U_0 = \{x; (0, x) \in U\}$ . We shall prove the case  $U_0 \cap \gamma_0^\pm \neq \emptyset$ . Otherwise the proof is easier. There are two cases as follows:

- (I)  $(\partial_2 h_\pm, \dots, \partial_n h_\pm)(0, x) \neq (0, \dots, 0)$  for any  $x \in U_0 \cap \gamma_0^\pm$ .
- (II)  $(\partial_2 h_\pm, \dots, \partial_n h_\pm)(0, \bar{x}) = (0, \dots, 0)$  for some  $\bar{x} \in U_0 \cap \gamma_0^\pm$ .

We first consider the case (I). Then we may assume that  $\chi$  satisfies not only  $\tau = t \circ \chi^{-1}$  and  $\xi_1 = r \circ \chi^{-1}$  but also  $\xi_2 = \pm h_\pm \circ \chi^{-1}$ . Performing a change of independent variables we can write

$$\phi_\pm(t, x) = \{\kappa x_1^2 + (\mu x_1 \mp x_2)^2\}^{1/2} + \mu x_1 \mp x_2.$$

Since  $(\partial_1 \phi_-)(t, x) \geq 0$  and  $(\partial_2 \phi_-)(t, x) \geq 0$  it follows from  $x_1 e^{ty_1} \leq e^{\delta \epsilon_0} x_1$  and  $x_2 + ty_2 \leq x_2$  that

$$\begin{aligned} & \phi_-(0, x_1 e^{ty_1}, x' + ty') \\ & \leq \phi_-(0, e^{\delta \epsilon_0} x_1, x') = \phi_-(t, e^{\delta \epsilon_0} x_1, x') = \phi_-(t, x; \kappa e^{2\delta \epsilon_0}, \mu e^{\delta \epsilon_0}) \\ & \leq C \phi_-(t, x; \kappa, \mu) = C \phi_-(t, x) \end{aligned}$$

which shows (7.6). Similarly we can prove (7.5).

We turn to the case (II). We note that  $\pm(\partial_0 h_{\pm})(0, \bar{x}) > 0$ . Indeed if the identity  $(\partial_0 h_-)(0, \bar{x}) = 0$  holds then  $A_{\gamma/b}(0, \bar{x}) = 0$  would follow from (1.4). This is incompatible with (1.5). Suppose that  $(\partial_0 h_-)(0, \bar{x}) > 0$  holds. Then we would have  $h_-(t, \bar{x}) > 0$  if  $0 < t < \delta$  and  $h_-(t, \bar{x}) < 0$  if  $-\delta < t < 0$ . In particular, we obtain  $(t, \bar{x}) \in O^-$  if  $0 < t < \delta$ , and hence  $A_b(t, \bar{x})$  is negative definite there. On the other hand, since it follows from (1.4) that  $A_{\gamma/b}(0, \bar{x})$  is positive definite then  $A_{b,\gamma}(0, \bar{x})$  is also positive definite. This is incompatible with (1.1) and (1.3). Therefore we have  $(\partial_0 h_-)(0, \bar{x}) < 0$ . Similarly we can prove  $(\partial_0 h_+)(0, \bar{x}) > 0$ . Thus taking  $U$  small enough we may assume that  $\pm(\partial_0 h_{\pm})(t, x) \geq c_0$  on  $U$  for some  $c_0 > 0$ .

Now we shall show (7.6). If we set  $f(\xi, \eta) = \{\kappa\xi^2 + (\mu\xi - \eta)^2\}^{1/2} + \mu\xi - \eta$  for  $(\xi, \eta) \in \mathbf{R}_+ \times \mathbf{R}$  then we can write  $\phi_-(t, x) = f(x_1, h_-(t, x))$ . Since  $(\partial_{\xi} f)(\xi, \eta) \geq 0$  and  $(\partial_{\eta} f)(\xi, \eta) \leq 0$  it suffices to show that  $h_-(0, x_1 e^{ty_1}, x' + ty') \geq h_-(t, x)$  because  $x_1 e^{ty_1} \leq e^{\delta\epsilon_0} x_1$ . Indeed admitting this assertion we have

$$\phi_-(0, x_1 e^{ty_1}, x' + ty') \leq \phi_-(t, x; \kappa e^{2\delta\epsilon_0}, \mu e^{\delta\epsilon_0}) \leq C\phi_-(t, x)$$

which concludes (7.6). Note that

$$\begin{aligned} & h_-(0, x_1 e^{ty_1}, x' + ty') - h_-(t, x) \\ &= -t \int_0^1 (\partial_0 h_-)(t - \theta t, x_1 + \theta x_1(e^{ty_1} - 1), x' + \theta ty') d\theta \\ &\quad + x_1(e^{ty_1} - 1) \int_0^1 (\partial_1 h_-)(\cdots) d\theta + \sum_{j=2}^n ty_j \int_0^1 (\partial_j h_-)(\cdots) d\theta \\ &\geq c_0 t - C\epsilon_0 t. \end{aligned}$$

Therefore taking  $\epsilon_0 > 0$  small enough we can prove the assertion. Similarly we can obtain (7.5).  $\square$

## References

- [1] K.O. Friedrichs: *The identity of weak and strong extensions of differential operators*, Trans. Amer. Math. Soc. **55** (1944), 132–151.
- [2] K.O. Friedrichs: *Symmetric positive linear differential operators*, Comm. Pure Appl. Math. **11** (1958), 333–418.
- [3] L. Hörmander: *Linear Partial Differential Operators*, Springer-Verlag, Berlin-Göttingen-Heidelberg, 1963.
- [4] P.D. Lax and R.S. Phillips: *Local boundary conditions for dissipative symmetric linear differential operators*, Comm. Pure Appl. Math. **13** (1960), 427–455.
- [5] R. Moyer: *On the nonidentity of weak and strong extensions of differential operators*, Proc. Amer. Math. Soc. **19** (1968), 487–488.
- [6] A. Majda and S. Osher: *Initial-boundary value problems for hyperbolic equations with uniformly characteristic boundary*, Comm. Pure Appl. Math. **28** (1975), 607–675.

- [7] T. Nishitani and M. Takayama: *A characteristic initial boundary value problem for a symmetric positive system*, Hokkaido Math. J. **25** (1996), 167–182.
- [8] T. Nishitani and M. Takayama: *Regularity of solutions to characteristic boundary value problem for symmetric systems*, in “Geometrical optics and related topics”(F. Colombini and N. Lerner), Birkhäuser, 1997.
- [9] M. Ohno, Y. Shizuta and T. Yanagisawa: *The initial boundary value problems for linear symmetric hyperbolic systems with characteristic boundary*, Proc. Japan Acad. **67** (1991), 191–196.
- [10] S. Osher: *An ill-posed problem for a hyperbolic equation near a corner*, Bull. Amer. Math. Soc. **79** (1973), 1043–1044.
- [11] R.S. Phillips and L. Sarason: *Singular symmetric positive first order differential operators*, J. Math. Mech. **15** (1966), 235–272.
- [12] J. Rauch: *Symmetric positive systems with boundary characteristic of constant multiplicity*, Trans. Amer. Math. Soc. **291** (1985), 167–187.
- [13] J. Rauch and F. Massey III: *Differentiability of solutions to hyperbolic initial-boundary value problems*, Trans. Amer. Math. Soc. **189** (1974), 303–318.
- [14] P. Secchi: *Linear symmetric hyperbolic systems with characteristic boundary*, Math. Methods Appl. Sci. **18** (1995), 855–870.
- [15] P. Secchi: *A symmetric positive system with non uniformly characteristic boundary*, preprint, (1996).
- [16] D. Tartakoff: *Regularity of solutions to boundary value problems for first order systems*, Indiana Univ. Math. J. **21** (1972), 1113–1129.
- [17] M. Tsuji: *Regularity of solutions of hyperbolic mixed problems with characteristic boundary*, Proc. Japan Acad. **48** (1972), 719–724.
- [18] T. Yanagisawa and A. Matsumura: *The fixed boundary value problems for the equations of ideal Magneto-Hydrodynamics with a perfectly conducting wall condition*, Comm. Math. Phys. **136** (1991), 119–140.

T. Nishitani  
Department of Mathematics  
Graduate School of Science  
Osaka University  
1-16 Machikaneyama Toyonaka  
Osaka 560-0043, Japan  
e-mail: tatsuo@math.wani.osaka-u.ac.jp

M. Takayama  
Department of Mathematics  
Graduate School of Science  
Osaka University  
1-16 Machikaneyama Toyonaka  
Osaka 560-0043, Japan  
e-mail: takayama@math.sci.osaka-u.ac.jp

