

Title	Stable homotopy types of stunted lens spaces mod 4
Author(s)	Kôno, Susumu
Citation	Osaka Journal of Mathematics. 1992, 29(4), p. 697-717
Version Type	VoR
URL	<a href="https://doi.org/10.18910/9267">https://doi.org/10.18910/9267</a>
rights	
Note	

***Osaka University Knowledge Archive : OUKA***

<https://ir.library.osaka-u.ac.jp/>

Osaka University

## STABLE HOMOTOPY TYPES OF STUNTED LENS SPACES MOD 4

Dedicated to Professor Hideki Ozeki on his 60th birthday

SUSUMU KÔNO

(Received December 26, 1991)

### 1. Introduction

Let  $L^n(q) = S^{2n+1}/(\mathbb{Z}/q)$  be the  $(2n+1)$ -dimensional standard lens space mod  $q$ . As defined in [8], we set

$$(1.1) \quad \begin{aligned} L_q^{2n+1} &= L^n(q), \\ L_q^{2n} &= \{[z_0, \dots, z_n] \in L^n(q) \mid z_n \text{ is real} \geq 0\}. \end{aligned}$$

The stable homotopy types ( $S$ -types) of stunted lens spaces  $L_q^m/L_q^n$  have been studied by several authors (e.g. [7], [8], [9], [10], [11] and [12]). For the case  $q=2$ , D.M. Davis and M. Mahowald have completed the classification of the stable homotopy types of stunted real projective spaces in [7]. Their result shows that we can use structures of  $J$ -groups of suspensions of stunted real projective spaces to obtain the necessary conditions for stunted real projective spaces  $RP(m)/RP(n)$  and  $RP(m+t)/RP(n+t)$  to have the same stable homotopy type as follows: if  $RP(m)/RP(n)$  and  $RP(m+t)/RP(n+t)$  have the same stable homotopy type, then there exists a non-negative integer  $N$  such that

$$\tilde{J}(S^j(RP(m)/RP(n))) \cong \tilde{J}(S^{j-t}(RP(m+t)/RP(n+t)))$$

for each integer  $j$  with  $j \geq N$  (see [13]). For the case where  $q$  is an odd prime, T. Kobayashi has obtained some necessary conditions for stunted lens spaces  $L_q^m/L_q^n$  and  $L_q^{m+t}/L_q^{n+t}$  to have the same stable homotopy type (cf. [10]). The conditions are also sufficient if  $k = [m/2] - [(n+1)/2] \not\equiv 0 \pmod{(q-1)}$  or  $n+1 \equiv 0 \pmod{2q^{[k/(q-1)]}}$ . We can use structures of  $J$ -groups of suspensions of stunted lens spaces mod  $q$  to obtain the conditions (see [14]). The object of this paper is to study the stable homotopy types of stunted lens spaces  $L_q^m/L_q^n$  for  $q=4$  or  $8$ .

In order to state our results, we prepare functions  $h_1, h_2, \alpha, \beta_1, \beta_2$  and  $\gamma_1$  defined by

$$(1.2) \quad h_1(k) = \begin{cases} [k/4] + [(k+7)/8] + [(k+4)/8] & (k \geq 2) \\ 0 & (1 \geq k \geq 0). \end{cases}$$

$$(1.3) \quad h_2(k) = \begin{cases} [k/4] + [(k+7)/8] + [k/8] + 1 & (k \geq 4) \\ h_1(k) & (3 \geq k \geq 0) . \end{cases}$$

$$(1.4) \quad \alpha(k, n) = \begin{cases} 1 & (n \equiv 0 \pmod{2} \text{ and } k \equiv 1 \pmod{8} , \\ & \text{or } k = 2([n/2] - [(n-1)/2])) \\ 0 & (\text{otherwise}) . \end{cases}$$

(1.5)  $\beta_1(k, n)$  is equal to the corresponding integer in the following table:

$k \pmod{8} \backslash n \pmod{4}$	0	1	2	3	4	5	6	7
0	0	0	0	0	0	0	0	0
1	0	0	1	1	0	0	1	1
2	0	0	0	1	0	0	1	1
3	0	0	0	0	0	0	0	0

(1.6)  $\beta_2(k, n)$  is equal to the corresponding integer in the following table:

$k \pmod{8} \backslash n \pmod{8}$	0	1	2	3	4	5	6	7
0	0	0	0	0	0	0	0	0
1	0	0	1	1	0	0	1	1
2	0	0	1	1	0	0	1	1
3	0	0	0	0	1	1	0	0
4	0	0	0	0	1	1	0	0
5	0	0	2	2	1	1	1	1
6	0	0	1	2	1	1	1	1
7	0	0	0	0	0	0	0	0

(1.7)  $\gamma_1(m, n)$  is equal to the corresponding integer in the following table:

$m-n$ $n \pmod{8}$	1	2	3	4	5	6
0	0	0	1	0	1	0
1	0	1	1	1	1	2
2	0	0	1	0	2	1
3	0	1	1	1	1	1
4	0	0	1	0	1	0
5	0	1	1	1	1	1
6	0	0	1	0	1	1
7	0	1	1	1	2	2

Let  $\nu_p(s)$  denote the exponent of the prime  $p$  in the prime power decomposition of  $s$ .

**Theorem 1.** If  $\nu_2(t) \geq h_1(m-2[(n+1)/2]) + 1 - \alpha(m-2[(n+1)/2], n)$ , then  $L_4^m/L_4^n$  and  $L_4^{m+t}/L_4^{n+t}$  have the same stable homotopy type.

**Theorem 2.** (1) If  $L_4^m/L_4^n$  and  $L_4^{m+t}/L_4^{n+t}$  have the same stable homotopy type, then

$$\nu_2(t) \geq [m/2] - [(n+1)/2] + \beta_1(m-2[(n+1)/2], n).$$

(2) Suppose  $h_1(m-2[(n+1)/2]) - \alpha(m-2[(n+1)/2], n) \geq i \geq 3$  and  $\max\{\nu_2(n+1), \nu_2(m+1)\} \geq i$ . If  $L_4^m/L_4^n$  and  $L_4^{m+t}/L_4^{n+t}$  have the same stable homotopy type, then  $\nu_2(t) \geq i+1$ .

(3) Suppose  $n < m \leq n+6$ . If  $L_4^m/L_4^n$  and  $L_4^{m+t}/L_4^{n+t}$  have the same stable homotopy type, then

$$\nu_2(t) \geq [m/2] - [(n+1)/2] + \gamma_1(m, n).$$

**REMARK.** It follows from Theorems 1 and 2, that we have obtained necessary and sufficient conditions for spaces  $L_4^m/L_4^n$  and  $L_4^{m+t}/L_4^{n+t}$  to have the same stable homotopy type if one of the following conditions is satisfied:

- (1)  $n < m \leq 2[(n+1)/2] + 3$ ,
- (2)  $\beta_1(m-2[(n+1)/2], n) = 1$ ,
- (3)  $\max\{\nu_2(n+1), \nu_2(m+1)\} \geq h_1(m-2[(n+1)/2]) - \alpha(m-2[(n+1)/2], n)$  and  $m \geq n+5$ .

**Theorem 3.** If  $\nu_2(t) \geq h_2(m-2[(n+1)/2]) + 1 - \alpha(m-2[(n+1)/2], n)$ , then

$L_8^m/L_8^n$  and  $L_8^{m+t}/L_8^{n+t}$  have the same stable homotopy type.

**Theorem 4.** (1) Suppose  $m \geq n+5$ . If  $L_8^m/L_8^n$  and  $L_8^{m+t}/L_8^{n+t}$  have the same stable homotopy type, then

$$\nu_2(t) \geq [m/2] - [(n+1)/2] + \beta_2(m-2[(n+1)/2], n).$$

(2) Suppose  $h_2(m-2[(n+1)/2]) - \alpha(m-2[(n+1)/2], n) \geq i \geq 3$  and  $\max\{\nu_2(n+1), \nu_2(m+1)\} \geq i$ . If  $L_8^m/L_8^n$  and  $L_8^{m+t}/L_8^{n+t}$  have the same stable homotopy type, then  $\nu_2(t) \geq i+1$ .

(3) Suppose  $n < m \leq n+6$ . If  $L_8^m/L_8^n$  and  $L_8^{m+t}/L_8^{n+t}$  have the same stable homotopy type, then

$$\nu_2(t) \geq [m/2] - [(n+1)/2] + \gamma_2(m, n),$$

where  $\gamma_2(m, n)$  is the integer defined by

$$\gamma_1(m, n) = \begin{cases} 1 & (n \equiv 0 \pmod{8} \text{ and } m = n+6) \\ \gamma_1(m, n) & (\text{otherwise}). \end{cases}$$

REMARK. It follows from Theorems 3 and 4, that we have obtained necessary and sufficient conditions for spaces  $L_8^m/L_8^n$  and  $L_8^{m+t}/L_8^{n+t}$  to have the same stable homotopy type if one of the following conditions is satisfied:

- (1)  $n < m \leq 2[(n+1)/2] + 3$ ,
- (2)  $n \equiv 0 \pmod{8}$  and  $m = n+6$ ,
- (3)  $\beta_2(m-2[(n+1)/2], n) = 2$  or  $2[(n+1)/2] \equiv [m/2] - [(n+3)/2] \equiv 2 \pmod{4}$ ,
- (4)  $\max\{\nu_2(n+1), \nu_2(m+1)\} \geq h_2(m-2[(n+1)/2]) - \alpha(m-2[(n+1)/2], n)$  and  $m \geq n+5$ .

This paper is organized as follows. In section 2 we prepare some lemmas and recall known results. We prove Theorems 1, 2, 3 and 4 in the final section.

The author would like to express his gratitude to Professor Akie Tamamura and Professor Kensô Fujii for helpful suggestions.

## 2. Preliminaries

In this section we prepare some lemmas and recall known results which are needed to prove Theorems 1, 2, 3 and 4.

**Lemma 2.1.** Let  $m, n, k$  and  $s$  be non-negative integers with  $2^s \leq k < 2^{s+1}$ . Assume that  $\binom{m}{i} \equiv \binom{n}{i} \pmod{2}$  for  $1 \leq i \leq k$ . Then  $m \equiv n \pmod{2^{s+1}}$ .

Proof. Suppose that  $m = \sum_{j=0}^N a_j 2^j$ ,  $n = \sum_{j=0}^N b_j 2^j$  and  $i = \sum_{j=0}^N c_{i,j} 2^j$  ( $1 \leq i \leq k$ ), where  $a_j, b_j$  and  $c_{i,j}$  are non-negative integers with  $a_j \leq 1, b_j \leq 1$  and  $c_{i,j} \leq 1$  ( $1 \leq i \leq k, 1 \leq j \leq N$ ). Then we have  $\binom{m}{i} \equiv \prod_{j=0}^N \binom{a_j}{c_{i,j}} \pmod{2}$  and  $\binom{n}{i} \equiv \prod_{j=0}^N$

$\binom{b_j}{c_{i,j}} \pmod{2}$  for  $1 \leq i \leq k$ . It follows from the hypothesis that we have  $a_j = b_j$  for  $0 \leq j \leq s$ ; that is,  $m \equiv n \pmod{2^{s+1}}$ . q.e.d.

Let  $q \geq 2$  be an integer and  $q_0, q_1, \dots, q_n$  be integers relatively prime to  $q$ . Consider the  $(\mathbf{Z}/q)$ -action on the unit sphere  $S^{2n+1} \subset \mathbf{C}^{n+1}$  given by

$$\exp(2\pi\sqrt{-1}/q)(z_0, \dots, z_n) = (z_0 \cdot \exp(2q_0\pi\sqrt{-1}/q), \dots, z_n \cdot \exp(2q_n\pi\sqrt{-1}/q)).$$

Then  $S^{2n+1}(q; q_0, \dots, q_n)$  denotes the space  $S^{2n+1}$  with this action,

$$L_q^{2n+1}(q_0, \dots, q_n) = S^{2n+1}(q; q_0, \dots, q_n)/(\mathbf{Z}/q)$$

and  $L_q^{2n}(q_0, \dots, q_n)$  is the subspace of  $L_q^{2n+1}(q_0, \dots, q_n)$  defined by

$$L_q^{2n}(q_0, \dots, q_n) = \{[z_0, \dots, z_n] \in L_q^{2n+1}(q_0, \dots, q_n) \mid z_n \text{ is real} \geq 0\}.$$

For  $0 \leq n < m \leq 2l+1$ , we set

$$L_q^m/L_q^n(q_0, \dots, q_l) = L_q^m(q_0, \dots, q_l)/L_q^n(q_0, \dots, q_l),$$

which is called a stunted lens space mod  $q$ . Then we have

$$(2.2) \quad L_q^m/L_q^n(1, \dots, 1) = L_q^m/L_q^n.$$

Considering the  $(\mathbf{Z}/q)$ -action on  $S^{2l+1}(q; q_0, \dots, q_l) \times \mathbf{C}^k$  given by

$$\begin{aligned} & \exp(2\pi\sqrt{-1}/q)(z, w_1, \dots, w_k) \\ &= (\exp(2\pi\sqrt{-1}/q) \cdot z, w_1 \cdot \exp(2a_1\pi\sqrt{-1}/q), \dots, w_k \cdot \exp(2a_k\pi\sqrt{-1}/q)) \end{aligned}$$

for  $(z, w_1, \dots, w_k) \in S^{2l+1}(q; q_0, \dots, q_l) \times \mathbf{C}^k$ , we have a complex  $k$ -dimensional vector bundle

$$\eta(a_1, \dots, a_k): (S^{2l+1}(q; q_0, \dots, q_l) \times \mathbf{C}^k)/(\mathbf{Z}/q) \rightarrow L_q^{2l+1}(q_0, \dots, q_l).$$

We use same symbol for the restriction of  $\eta(a_1, \dots, a_k)$  to  $L_q^n(q_0, \dots, q_l)$  ( $n \leq 2l+1$ ) and denote the complex line bundle  $\eta(1)$  by  $\eta$ . Then we have

$$(2.3) \quad \eta(a_1, \dots, a_k) \cong \eta^{a_1} \oplus \dots \oplus \eta^{a_k}.$$

Let  $\xi: E(\xi) \rightarrow X$  be a real vector bundle over a finite  $CW$ -complex  $X$  with disk bundle  $B(\xi)$  and sphere bundle  $S(\xi)$ . Then the Thom complex  $X^\xi$  of  $\xi$  is defined as the quotient space  $B(\xi)/S(\xi)$ . Define  $f: S^{2n+1} \times D^{2k} \rightarrow S^{2k+2n+1}$  by

$$f(z_0, \dots, z_n, w) = (w, (1 - \|w\|^2)^{1/2} z_0, \dots, (1 - \|w\|^2)^{1/2} z_n)$$

for  $(z_0, \dots, z_n, w) \in S^{2n+1} \times D^{2k}$ , where  $D^{2k} \subset \mathbf{C}^k$  is the unit disk. Then we have

(2.4) *Let  $a_1, \dots, a_k$  be integers relatively prime to  $q$ . Then  $f$  induces the following homeomorphisms.*

- (1)  $(L_q^{2n+1}(q_0, \dots, q_n))^{r(\eta(a_1, \dots, a_k))} \approx L_q^{2k+2n+1}/L_q^{2k-1}(a_1, \dots, a_k, q_0, \dots, q_n).$
- (2)  $(L_q^{2n}(q_0, \dots, q_n))^{r(\eta(a_1, \dots, a_k))} \approx L_q^{2k+2n}/L_q^{2k-1}(a_1, \dots, a_k, q_0, \dots, q_n).$
- (3)  $(L_q^{2n+1}(q_0, \dots, q_n))^{r(\eta(a_1, \dots, a_k))}/S^{2k} \approx L_q^{2k+2n+1}/L_q^{2k}(a_1, \dots, a_k, q_0, \dots, q_n).$
- (4)  $(L_q^{2n}(q_0, \dots, q_n))^{r(\eta(a_1, \dots, a_k))}/S^{2k} \approx L_q^{2k+2n}/L_q^{2k}(a_1, \dots, a_k, q_0, \dots, q_n).$

We define the function  $h(q, k)$  by setting

$$(2.5) \quad h(q, k) = \text{ord} \langle J(r(\eta) - 2) \rangle,$$

where  $J(r(\eta) - 2)$  is the image of  $r(\eta) - 2 \in \widetilde{KO}(L_q^k)$  by the  $J$ -homomorphism  $J: \widetilde{KO}(L_q^k) \rightarrow \widetilde{J}(L_q^k)$ .

REMARK. The function  $h(q, k)$  have been determined completely (cf. [8]).

Spaces  $X$  and  $Y$  are said to have the same stable homotopy type ( $X \underset{s}{\approx} Y$ ) if there exist non-negative integers  $a$  and  $b$  such that  $S^a X$  and  $S^b Y$  have the same homotopy type. For stunted generalized lens spaces, we have

(2.6) (1) If  $mn \equiv 0 \pmod{2}$ , then  $L_q^m/L_q^n(q_0, \dots, q_{[m/2]})$  and  $L_q^m/L_q^n$  have the same stable homotopy type. In particular,  $L_q^m(q_0, \dots, q_{[m/2]})$  and  $L_q^m$  have the same stable homotopy type.

(2) Let  $a_1, \dots, a_k, b_1, \dots, b_k, q_0, \dots, q_n, r_0, \dots, r_n$  and  $a$  be integers relatively prime to  $q$ .

i) Assume  $q_0 \cdots q_n \equiv \pm a^{n+1} r_0 \cdots r_n \pmod{q}$ . Then

$$L_q^{2n+2k+1}/L_q^{2k-1}(a_1, \dots, a_k, q_0, \dots, q_n)$$

and  $L_q^{2n+2k+1}/L_q^{2k-1}(b_1, \dots, b_k, r_0, \dots, r_n)$  have the same stable homotopy type.

ii) If  $k \equiv 0 \pmod{h(q, 2n+1)}$ , then the spaces

$$L_q^{2n+2k+1}/L_q^{2k-1}(a_1, \dots, a_k, q_0, \dots, q_n)$$

and  $L_q^{2n+2k+1}/L_q^{2k-1}$  have the same stable homotopy type.

Proof. Set  $X = L_q^{2n+1}(q_0, \dots, q_n)$ . According to [17], we have

$$K(X) \cong \mathbf{Z}[\eta]/(\eta^q - 1, (\eta - 1)^{n+1})$$

and  $\text{ord } \tilde{K}(X) = q^n$ . If  $a$  is an integer relatively prime to  $q$ , then

$$\begin{aligned} \bigcap_j (\sum_j j^{f(j)} (\psi^j - 1) \tilde{K}(X)) &= \sum_j (\bigcap_e j^e (\psi^j - 1) \tilde{K}(X)) \\ &\supset \bigcap_e a^e (\psi^a - 1) \tilde{K}(X) = (\psi^a - 1) \tilde{K}(X) \\ &\ni (\psi^a - 1)(\eta - 1) = \eta^a - \eta. \end{aligned}$$

Since the Adams operations are compatible with the real restriction  $r: K \rightarrow KO$  [4],

$$r(\eta^a - \eta) = r(\eta^a) - r(\eta) \in \bigcap_f (\sum_j j^{f(j)} (\psi^j - 1) \widetilde{KO}(X)).$$

By [2], [3] and [19], this implies that

$$J(r(\eta^a) - 2) = J(r(\eta) - 2)$$

in  $\mathcal{J}(X)$ . If  $a_1, \dots, a_k$  are integers relatively prime to  $q$ ,

$$J(r(\eta(a_1, \dots, a_k)) - 2k) = J(k(r(\eta)) - 2k)$$

in  $\mathcal{J}(X)$ . Suppose that  $a_1, \dots, a_k, b_1, \dots, b_k, q_0, \dots, q_n, r_0, \dots, r_n$  and  $a$  be integers relatively prime to  $q$ . According to [5, Proposition (2.6)], we have

$$L_q^{2n+2k+1}/L_q^{2k-1}(a_1, \dots, a_k, q_0, \dots, q_n) \cong_s L_q^{2n+2k+1}/L_q^{2k-1}(a, \dots, a, q_0, \dots, q_n)$$

and

$$L_q^{2n+2k+1}/L_q^{2k-1}(b_1, \dots, b_k, r_0, \dots, r_n) \cong_s L_q^{2n+2k+1}/L_q^{2k-1}(1, \dots, 1, r_0, \dots, r_n).$$

Suppose that  $q_0 \cdots q_n \equiv \pm a^{n+1} r_0 \cdots r_n \pmod{q}$ . Identify  $S^{2n+2k+1}$  with the iterated join

$$S^1 * \cdots * S^1 = \{\lambda_0 z_0 + \cdots + \lambda_{n+k} z_{n+k} \mid \sum_j \lambda_j = 1, \lambda_j \geq 0, z_j \in S^1\}.$$

Choose integers  $\bar{q}_i$  ( $0 \leq i \leq n$ ) with  $q_i \bar{q}_i \equiv 1 \pmod{q}$ . Denote the generator  $\exp(2\pi\sqrt{-1}/q)$  of  $\mathbb{Z}/q$  by  $g$ . Then, the map

$$f: S^{2n+2k+1}(q; a, \dots, a, q_0, \dots, q_n) \rightarrow S^{2n+2k+1}(q; 1, \dots, 1, r_0, \dots, r_n)$$

defined by

$$f = 1 * \cdots * 1 * (ar_0 \bar{q}_0) * \cdots * (ar_n \bar{q}_n),$$

is a map of degree  $a^{n+1} r_0 \cdots r_n \bar{q}_0 \cdots \bar{q}_n \equiv \pm 1 \pmod{q}$  with  $f \circ g = g^a \circ f$ . Modify  $f$  to get a map  $h$  of degree  $\pm 1$  with  $h \circ g = g^a \circ h$  and

$$h|S^{2k-1}(q; a, \dots, a) = f|S^{2k-1}(q; a, \dots, a)$$

(see the proof of [6, (29.4)] for the detail). Then  $h$  induces a homotopy equivalence

$$\bar{h}: L_q^{2n+2k+1}/L_q^{2k-1}(a, \dots, a, q_0, \dots, q_n) \rightarrow L_q^{2n+2k+1}/L_q^{2k-1}(1, \dots, 1, r_0, \dots, r_n).$$

This completes the proof of (2) i).

Now we turn to the proof of (1). Since  $\mathcal{J}(X)$  has a finite order, we may assume that  $k > 1$ . Let  $a$  be an integer with  $aq_0 \cdots q_n \equiv 1 \pmod{q}$ . Then



$$L_q^{2n+2k+1}/L_q^{2k}(a_1, \dots, a_k, q_0, \dots, q_n) \underset{\cong}{\simeq} L_q^{2n+2k+1}/L_q^{2k}(1, \dots, 1, a, q_0, \dots, q_n).$$

By the proof above, there exists an equivariant map

$$h: S^{2n+2k+1}(q; 1, \dots, 1) \rightarrow S^{2n+2k+1}(q; 1, \dots, 1, a, q_0, \dots, q_n)$$

of degree 1, which induces homotopy equivalences

$$\bar{h}: L_q^{2n+2k+1}/L_q^{2k} \rightarrow L_q^{2n+2k+1}/L_q^{2k}(1, \dots, 1, a, q_0, \dots, q_n)$$

and  $\bar{h}|L_q^{2n+2k}/L_q^{2k}: L_q^{2n+2k}/L_q^{2k} \rightarrow L_q^{2n+2k}/L_q^{2k}(1, \dots, 1, a, q_0, \dots, q_n)$ . Thus  $L_q^{2n+2k+1}/L_q^{2k}(a_1, \dots, a_k, q_0, \dots, q_n)$  is stably homotopically equivalent to  $L_q^{2n+2k+1}/L_q^{2k}$  and  $L_q^{2n+2k}/L_q^{2k}(a_1, \dots, a_k, q_0, \dots, q_n)$  is stably homotopically equivalent to  $L_q^{2n+2k}/L_q^{2k}$ . The equivariant map

$$\begin{aligned} f &= 1 * \dots * 1 * q_0 * \dots * q_n: S^{2n+2k+1}(q; 1, \dots, 1) \\ &\rightarrow S^{2n+2k+1}(q; 1, \dots, 1, q_0, \dots, q_n) \end{aligned}$$

induces a homotopy equivalence

$$f|L_q^{2n+2k}/L_q^{2k-1}: L_q^{2n+2k}/L_q^{2k-1} \rightarrow L_q^{2n+2k}/L_q^{2k-1}(1, \dots, 1, q_0, \dots, q_n).$$

Thus  $L_q^{2n+2k}/L_q^{2k-1}(a_1, \dots, a_k, q_0, \dots, q_n)$  is stably homotopically equivalent to  $L_q^{2n+2k}/L_q^{2k-1}$ . This completes the proof of (1).

Finally, we prove the part ii) of (2). Suppose that  $k \equiv 0 \pmod{h(q, 2n+1)}$ . Since the order of  $J(r(\eta)-2) \in \tilde{J}(X)$  coincides with  $h(q, 2n+1)$ , we have

$$\begin{aligned} &L_q^{2n+2k+1}/L_q^{2k-1}(a_1, \dots, a_k, q_0, \dots, q_n) \\ &\underset{\cong}{\simeq} L_q^{2n+2k+1}/L_q^{2k-1}(1, \dots, 1, q_0, \dots, q_n) \\ &\approx (L_q^{2n+1}(q_0, \dots, q_n))^{r(k\eta)} \\ &\underset{\cong}{\simeq} (L_q^{2n+1}(q_0, \dots, q_n))^{2k} \\ &\simeq S^{2k} L_q^{2n+1}(q_0, \dots, q_n) \vee S^{2k} \\ &\underset{\cong}{\simeq} S^{2k} L_q^{2n+1} \vee S^{2k} \\ &\simeq (L_q^{2n+1})^{2k} \underset{\cong}{\simeq} (L_q^{2n+1})^{r(k\eta)} \approx L_q^{2n+2k+1}/L_q^{2k-1}, \end{aligned}$$

by [5, Proposition (2.6)], (2.4) and (2.6) (1). This completes the proof of the part ii) of (2). q.e.d.

Let  $X$  and  $Y$  be pointed  $CW$ -complexes. The stable homotopy group from  $X$  to  $Y$  is defined by

$$(2.7) \quad \{X, Y\} = \lim_{n \rightarrow \infty} [S^n X, S^n Y]_0.$$

The following assertion is proved by making use of Puppe exact sequences.

$$(2.8) \quad (1) \quad \text{If } k > m, \text{ then } \{L_q^m/L_q^n(q_0, \dots, q_{[m/2]}), S^k\} \cong 0.$$

(2) If  $k \leq n$ , then  $\{S^k, L_q^m/L_q^n(q_0, \dots, q_{[m/2]})\} \cong 0$ .

(3) Suppose that  $[m/2] > [(n+1)/2]$ . Then we have

$$i) \quad \{S^{n+1}, L_q^m/L_q^n(q_0, \dots, q_{[m/2]})\} = \langle \{i_{n+1}\} \rangle \cong \begin{cases} \mathbf{Z} & (n: \text{odd}) \\ \mathbf{Z}/q & (n: \text{even}), \end{cases}$$

where  $i_{n+1}$  is the composition

$$S^{n+1} \approx L_q^{n+1}/L_q^n(q_0, \dots, q_{[m/2]}) \subset L_q^m/L_q^n(q_0, \dots, q_{[m/2]}) .$$

$$ii) \quad \{L_q^m/L_q^n(q_0, \dots, q_{[m/2]}), S^m\} = \langle \{p_m\} \rangle \cong \begin{cases} \mathbf{Z} & (m: \text{odd}) \\ \mathbf{Z}/q & (m: \text{even}), \end{cases}$$

where  $p_m$  is the composition

$$L_q^m/L_q^n(q_0, \dots, q_{[m/2]}) \rightarrow L_q^m/L_q^{m-1}(q_0, \dots, q_{[m/2]}) \approx S^m .$$

The following lemma implies that the family of stunted lens spaces is closed under  $S$ -duality.

**Lemma 2.9.** Suppose that  $k=2[m/2]+1-2[(n+1)/2] \geq 3$ ,  $N \equiv 0 \pmod{2h(q, k)}$  and  $N > m+1$ . Then the  $S$ -dual of

$$L_q^m/L_q^n(a_1, \dots, a_{[(n+1)/2]}, q_0, \dots, q_{[k/2]})$$

$$\text{is } L_q^{N-n-2}/L_q^{N-m-2}(b_1, \dots, b_{[(N-m-1)/2]}, q_0, \dots, q_{[k/2]}) .$$

Proof. If  $nm \equiv 1 \pmod{2}$ , then the  $S$ -dual of

$$X = L_q^m/L_q^n(a_1, \dots, a_{[(n+1)/2]}, q_0, \dots, q_{[k/2]}) \\ \cong (L_q^k(q_0, \dots, q_{(k-1)/2}))^{((n+1)/2)r(\eta)}$$

is  $(L_q^k(q_0, \dots, q_{(k-1)/2}))^{-\tau - ((n+1)/2)r(\eta)}$ , where  $\tau$  denotes the tangent bundle of  $L_q^k(q_0, \dots, q_{(k-1)/2})$  (cf. [5, Theorem (3.5)]). Since

$$\tau \oplus 1 \cong r(\eta(q_0, \dots, q_{(k-1)/2})),$$

it follows from [5, Proposition (2.6)] and (2.6) that

$$\begin{aligned} & (L_q^k(q_0, \dots, q_{(k-1)/2}))^{-\tau - ((n+1)/2)r(\eta)} \\ & \cong (L_q^k(q_0, \dots, q_{(k-1)/2}))^{(N/2)r(\eta) - ((k+1)/2)r(\eta) - ((n+1)/2)r(\eta)} \\ & = (L_q^k(q_0, \dots, q_{(k-1)/2}))^{((N-m-1)/2)r(\eta)} \\ & \cong L_q^{N-n-2}/L_q^{N-m-2}(b_1, \dots, b_{[(N-m-1)/2]}, q_0, \dots, q_{[k/2]}) = Y . \end{aligned}$$

This implies that there are positive integers  $a, b$  and  $M$  such that the  $M$ -dual of  $S^a X$  is  $S^b Y$ . Let

$$u: S^a X \wedge S^b Y \rightarrow S^M$$

be a  $M$ -duality pairing. Then the homomorphism

$$\Gamma_u^i: \{S^i, S^a X\} \rightarrow \{S^{i+b} Y, S^M\}$$

defined by  $\Gamma_u^i(\{f\}) = \{S^i u \circ (f \wedge 1)\}$  for  $f: S^{l+i} \rightarrow S^{l+a} X$ , is an isomorphism. It follows from (2.8) that  $M = a + b + N - 1$  and

$$\{S^{n+1}, X\} = \langle \{i_{n+1}\} \rangle \cong \{Y, S^{N-n-2}\} = \langle \{p_{N-n-2}\} \rangle \cong \mathbb{Z}.$$

Hence, by the isomorphism

$$\{S^{n+1}, X\} \cong \{S^{a+n+1}, S^a X\} \xrightarrow{\Gamma_u^{a+n+1}} \{S^{a+b+n+1} Y, S^{a+b+N-1}\} \cong \{Y, S^{N-n-2}\},$$

$\{i_{n+1}\}$  corresponds to either  $\{p_{N-n-2}\}$  or  $-\{p_{N-n-2}\}$ . This implies that there exists a homotopy commutative diagram

$$\begin{array}{ccc} S^{l+a+n+1} \wedge S^b Y & \xrightarrow{1 \wedge S^b(p_{N-n-2})} & S^{l+a+n+1} \wedge S^{b+N-n-2} \\ \downarrow S^{l+a}(i_{n+1}) \wedge 1 & & \approx \downarrow v \\ S^{l+a} X \wedge S^b Y & \xrightarrow{S^l u} & S^{l+a+b+N-1}, \end{array}$$

where  $v$  and  $S^l u$  are  $(l+a+b+N-1)$ -duality pairings. It follows that  $C_{S^{l+a}(i_{n+1})}$  is the  $(l+a+b+N)$ -dual of  $C_{S^b(p_{N-n-2})}$ . Since

$$C_{S^{l+a}(i_{n+1})} \cong S^{l+a}(L_q^m/L_q^{n+1}(a_1, \dots, a_{(n+1)/2}, q_0, \dots, q_{[k/2]}))$$

and

$$C_{S^b(p_{N-n-2})} \cong S^{b+1}(L^{N-n-3}/L^{N-m-2}(b_1, \dots, b_{(N-m-1)/2}, q_0, \dots, q_{[k/2]})),$$

this implies that  $L_q^m/L_q^{n+1}$  is the  $S$ -dual of  $L^{N-n-3}/L^{N-m-2}$ . In the similar way it is shown that  $L_q^{N-n-2}/L_q^{N-m-1}$  is the  $S$ -dual of  $L_q^{m-1}/L_q^n$ . Using this fact, in the similar way it is shown that  $L_q^{m-1}/L_q^{n+1}$  is the  $S$ -dual of  $L_q^{N-n-3}/L_q^{N-m-1}$ . q.e.d.

REMARK. The partial results for the case where  $q$  is a prime of this lemma have been obtained in [18].

It follows from (2.6) and Lemma 2.9 that, in the following cases,

$$L_q^m/L_q^n(a_1, \dots, a_{[(n+1)/2]}, q_0, \dots, q_{[m/2]-[(n+1)/2]})$$

and  $L_q^m/L_q^n$  have the same stable homotopy type:

- (1)  $q=2, 3, 4$  or  $6$ ,
- (2)  $mn \equiv 0 \pmod{2}$ ,
- (3)  $q_0 \cdots q_{[m/2]-[(n+1)/2]} \equiv \pm a^{[m/2]-[(n-1)/2]} \pmod{q}$ ,

- (4)  $n+1 \equiv 0 \pmod{2h(q, m-n-1)}$ ,  
 (5)  $m+1 \equiv 0 \pmod{2h(q, m-n-1)}$ .

QUESTION. Is it true that  $L_q^m/L_q^n(q_0, \dots, q_{[m/2]})$  and  $L_q^m/L_q^n$  have the same stable homotopy type for any case? If it is not, then how many stable homotopy types are there for fixed  $m, n$  and  $q$ ?

From now on, we restrict ourselves to standard stunted lens spaces. According to [5, Propositions (2.6) and (2.9)], (2.4) and Lemma 2.9, we obtain the following.

(2.10) Set  $k=m-2[(n+1)/2]$  and  $l=2[m/2]-n$ .

- (1) If  $t \equiv 0 \pmod{2h(q, k)}$ , then  $L_q^m/L_q^n$  and  $L_q^{m+t}/L_q^{n+t}$  have the same stable homotopy type.  
 (2) If  $k \geq 2$  and  $n+1 \equiv 0 \pmod{2h(q, k)}$ , then  $t \equiv 0 \pmod{2h(q, k)}$  if and only if  $L_q^m/L_q^n$  and  $L_q^{m+t}/L_q^{n+t}$  have the same stable homotopy type.  
 (3) If  $t \equiv 0 \pmod{2h(q, l)}$ , then  $L_q^m/L_q^n$  and  $L_q^{m+t}/L_q^{n+t}$  have the same stable homotopy type.  
 (4) If  $l \geq 2$  and  $m+1 \equiv 0 \pmod{2h(q, l)}$ , then  $t \equiv 0 \pmod{2h(q, l)}$  if and only if  $L_q^m/L_q^n$  and  $L_q^{m+t}/L_q^{n+t}$  have the same stable homotopy type.

(2.11) (1) ([12, I; Theorem 1.1]) Let  $p$  be a prime and  $r$  a positive integer with  $p^r > 2$ . Suppose that  $k=m-2[(n+1)/2] \geq 2$ . Then  $t \equiv 0 \pmod{2p^{[(k-2)/2(p-1)]}}$  if  $L_{p^r}^m/L_{p^r}^n$  and  $L_{p^r}^{m+t}/L_{p^r}^{n+t}$  have the same stable homotopy type.

(2) Let  $r \geq 2$  be a positive integer and set  $k=m-2[(n+1)/2]$ . Then  $v_2(t) \geq [k/2] + \beta_1(k, n)$  if  $L_{2^r}^m/L_{2^r}^n$  and  $L_{2^r}^{m+t}/L_{2^r}^{n+t}$  have the same stable homotopy type, where  $\beta_1$  is the function defined by (1.5).

(3) Suppose that  $q \equiv 0 \pmod{2}$  and  $m \geq n+2$ . Then  $v_2(t) \geq [\log_2 2(m-n-1)]$  if  $L_q^m/L_q^n$  and  $L_q^{m+t}/L_q^{n+t}$  have the same stable homotopy type.

Proof. Suppose that  $q \equiv 0 \pmod{2}$  and  $m \geq n+2$ . It is well known that

$$H^*(L_q^m; \mathbf{Z}/2) \cong \begin{cases} (\mathbf{Z}/2)[u]/(u^{m+1}) & (q \equiv 2 \pmod{4}) \\ (\mathbf{Z}/2)[u, v]/(u^2, v^{[m/2]+1}, uv^{m-[m/2]}) & (q \equiv 0 \pmod{4}), \end{cases}$$

where  $\deg u = 1$  and  $\deg v = 2$ . The action of the Steenrod squares is given by

$$\begin{cases} Sq^i(u^j) = \binom{j}{i} u^{j+i} & (q \equiv 2 \pmod{4}) \\ Sq^{2i}(v^j) = \binom{j}{i} v^{j+i} & (q \equiv 0 \pmod{4}) \\ Sq^{2i}(uv^j) = \binom{j}{i} uv^{j+i} & (q \equiv 0 \pmod{4}) \\ Sq^{2i+1} = 0 & (q \equiv 0 \pmod{4}). \end{cases}$$

Assume that  $L_q^m/L_q^n$  and  $L_q^{m+t}/L_q^{n+t}$  have the same stabel homotopy type. Then  $t \equiv 0 \pmod{2}$ . It follows from the naturality of the Steenrod squares that we have

$$(2.12) \quad \begin{cases} \binom{n+1}{i} \equiv \binom{n+t+1}{i} \pmod{2} & (1 \leq i \leq m-n-1, q \equiv 2 \pmod{4}) \\ \binom{[(n+1)/2]}{i} \equiv \binom{[(n+t+1)/2]}{i} \pmod{2} & (2 \leq 2i \leq m-n-1, q \equiv 0 \pmod{4}). \end{cases}$$

Let  $s$  be the integer with  $2^s \leq m-n-1 < 2^{s+1}$ . By Lemma 2.1, (2.12) implies that  $\nu_2(t) \geq s+1 = [\log_2 2(m-n-1)]$ . This completes the proof of (3).

(2) It follows from [11, Theorem 1.1] that

$$\nu_2(t) \geq [m/2] - [(n+1)/2] + C(m, n)$$

if  $L_{2^r}^m/L_{2^r}^n$  and  $L_{2^r}^{m+t}/L_{2^r}^{n+t}$  have the same stable homotopy type, where  $C(m, n)$  is the function defined by

$$C(m, n) = \begin{cases} 1 & (n \equiv 1, 5 \text{ or } 6 \pmod{8} \text{ and } m \equiv 1, 4 \text{ or } 5 \pmod{8}) \\ 0 & (\text{otherwise}). \end{cases}$$

Then (2) is obtained by making use of the  $S$ -duality (Lemma 2.9). q.e.d.

In order to state the next proposition, we set

$$(2.13) \quad \begin{cases} a_0(m, n) = [m/4] + [(m+7)/8] + [(m+4)/8] \\ \quad - [(n+1)/4] - [(n+1)/8] - [(n+6)/8] \\ b_0(m, n) = [m/8] + [(m+6)/8] - [(n+7)/8] - [(n+5)/8]. \end{cases}$$

$$(2.14) \quad \begin{cases} a(j, m, n) = \min \{ \nu+1, a_0(m+j, n+j) \} \\ b(j, m, n) = \min \{ \nu+1, b_0(m+j, n+j) \} \end{cases},$$

where  $\nu$  is the integer defined by

$$\nu = \begin{cases} \nu_2(j) & (j \neq 0) \\ m & (j = 0). \end{cases}$$

Let  $\mathfrak{m}(s)$  denote the function defined on positive integers as follows (cf. [3]):

$$\nu_p(\mathfrak{m}(s)) = \begin{cases} 0 & (p \neq 2 \text{ and } s \not\equiv 0 \pmod{p-1}) \\ 1 + \nu_p(s) & (p \neq 2 \text{ and } s \equiv 0 \pmod{p-1}) \\ 1 & (p = 2 \text{ and } s \not\equiv 0 \pmod{2}) \\ 2 + \nu_2(s) & (p = 2 \text{ and } s \equiv 0 \pmod{2}). \end{cases}$$

**Proposition 2.15** ([15, Theorem 3]). *Let  $j$ ,  $m$  and  $n$  be non-negative in-*

tegers with  $m > n$  and  $j \equiv n+1 \equiv 0 \pmod{4}$ . Then we have

$$\tilde{J}(S^j(L_4^m/L_4^n)) \cong \begin{cases} \mathbf{Z}/m((n+j+1)/2) \cdot 2^c \oplus \mathbf{Z}/2^{d+i} \oplus \mathbf{Z}/2^k & (b(j, m, n) \geq 0) \\ \mathbf{Z}/m((n+j+1)/2) & (b(j, m, n) < 0), \end{cases}$$

where  $i, k, c$  and  $d$  are integers defined by

$$(2.16) \quad \begin{cases} i = \begin{cases} \min \{v_2(n+1)-1, a(j, m, n)\} & (n+j \equiv 7 \pmod{8}) \\ \min \{v_2(n+1), a(j, m, n)\} & (n+j \equiv 3 \pmod{8}) \end{cases} \\ k = \min \{v_2(n+1)-1, b(j, m, n)\} \\ c = \max \{a(j, m, n)-i, b(j, m, n)-k\} \\ d = \min \{a(j, m, n)-i, b(j, m, n)-k\}. \end{cases}$$

In order to state the next proposition, we set

$$(2.17) \quad \begin{cases} a_3(m, n) = [(m-2)/8] - [(n+5)/8] \\ a_4(m, n) = [m/8] - [(n+7)/8] \\ a_5(m, n) = a_0(m, n) + [m/8] - [(m-4)/8] \\ a_6(m, n) = [(m+4)/8] + [(m-2)/8] - [(n+1)/4]. \end{cases}$$

$$(2.18) \quad \begin{cases} b_1(j, m, n) = \min \{v - a_3(n, n+4), a_5(m, n)\} \\ b_2(j, m, n) = \min \{v - a_3(n-5, n), a_6(m, n)\} \\ b_3(j, m, n) = \min \{v+1, a_4(m, n)\}, \end{cases}$$

where  $v$  is the integer defined in (2.14).

**Proposition 2.19** ([16, Theorem 2]). *Let  $j, m$  and  $n$  be non-negative integers with  $m > n$  and  $j \equiv 0 \pmod{8}$ .*

(1) *If  $n \not\equiv 3 \pmod{4}$  and  $m \geq 2[(n+6)/8] + 2[n/8] + 4[(n+15)/8]$ , then we have*

$$\tilde{J}(S^j(L_8^m/L_8^n)) \cong \begin{cases} (\oplus_{i=1}^3 \mathbf{Z}/2^{b_i(j, m, n)}) \oplus \mathbf{Z}/2 & (n \equiv 2 \pmod{8}) \\ \oplus_{i=1}^3 \mathbf{Z}/2^{b_i(j, m, n)} & (\text{otherwise}). \end{cases}$$

(2) *If  $n \equiv 3 \pmod{4}$  and  $2[(n+6)/8] + 2[n/8] + 4[(n+15)/8] > m > n$ , then we have*

$$\tilde{J}(S^j(L_8^m/L_8^n)) \cong \begin{cases} \mathbf{Z}/2^{b_1(j, m, n)} \oplus \mathbf{Z}/4 & (n \equiv 2 \pmod{8} \text{ and } m \geq n+6) \\ \mathbf{Z}/8 & (n \equiv 1 \pmod{8} \text{ and } m \geq n+3) \\ K\tilde{O}(S^j(L_8^m/L_8^n)) & (\text{otherwise}). \end{cases}$$

(3) *If  $n \equiv 3 \pmod{4}$  and  $m \geq n+5$ , then we have*

$$\tilde{J}(S^j(L_8^m/L_8^n)) \cong \mathbf{Z}/m((n+j+1)/2) \cdot 2^{c_1} \oplus \mathbf{Z}/2^{c_2+i_1} \oplus \mathbf{Z}/2^{i_2} \oplus \mathbf{Z}/2^{b_3(j, m, n)},$$

where  $i_1, i_2, c_1$  and  $c_2$  are integers defined by

$$(2.20) \quad \begin{cases} i_1 = \begin{cases} \min \{b_1(j, m, n), v_2(n+1)-1\} & (n \equiv 7 \pmod{8}) \\ \min \{b_1(j, m, n), v_2(n+1)\} & (n \equiv 3 \pmod{8}) \end{cases} \\ i_2 = \begin{cases} \min \{b_2(j, m, n), v_2(n+1)-2\} & (n \equiv 7 \pmod{8}) \\ \min \{b_2(j, m, n), v_2(n+1)-1\} & (n \equiv 3 \pmod{8}) \end{cases} \\ c_1 = \max \{b_1(j, m, n)-i_1, b_2(j, m, n)-i_2\} \\ c_2 = \min \{b_1(j, m, n)-i_1, b_2(j, m, n)-i_2\}. \end{cases}$$

(4) If  $n \equiv 3 \pmod{4}$  and  $n+5 > m > n$ , then we have

$$J(S^j(L_8^m/L_8^n)) \cong \begin{cases} \mathbf{Z}/m((n+j+1)/2) \oplus \widetilde{KO}(S^j(L_8^m/L_8^{n+1})) & (m \geq n+2) \\ \mathbf{Z}/m((n+j+1)/2) & (m = n+1). \end{cases}$$

In order to state the next proposition, we set

$$(2.21) \quad (1) \quad \begin{cases} \sigma_{2^i} = \eta^{2^i} - 1 & (0 \leq i \leq 2) \\ \sigma_6 = \sigma_4 \sigma_2 \\ \sigma_{2i+1} = \sigma_{2i} \sigma_1 & (1 \leq i \leq 3). \end{cases}$$

(2) Let  $F(x)$  denote the free abelian group generated by  $x_1, x_2, x_3, x_4, x_5, x_6$  and  $x_7$ . Then  $X_i$  and  $X_i(n)$  ( $1 \leq i \leq 7, n \geq 0$ ) denote the elements of  $F(x)$  defined by  $X_1 = 4x_1 + 2x_3 + 2x_5 + x_7$ ,  $X_2 = 2x_2 + x_6$ ,  $X_3 = 2x_3 + x_7$ ,  $X_6 = x_6 + x_7$ ,  $X_i = x_i$  ( $i = 4, 5$  or  $7$ ),  $X_1(n) = 2^{\lfloor n/2 \rfloor} X_1$ ,

$$\begin{aligned} X_2(n) &= 2^{\lfloor n/4 \rfloor} X_2 - 2^{2\lfloor n/4 \rfloor} X_1, \\ X_3(n) &= 2^{\lfloor (n-2)/4 \rfloor} X_3 + (2^{\lfloor n/2 \rfloor} - 2^{2\lfloor (n-2)/4 \rfloor + 1}) X_1, \\ X_4(n) &= 2^{\lfloor n/8 \rfloor} X_4 + (2^{\lfloor n/4 \rfloor} - 2^{2\lfloor n/8 \rfloor}) X_2 + 2^{\lfloor n/4 \rfloor + 2\lfloor n/8 \rfloor} X_1, \\ X_5(n) &= 2^{\lfloor (n-2)/8 \rfloor} X_5 + (2^{\lfloor (n-2)/4 \rfloor} - 2^{2\lfloor (n-2)/8 \rfloor}) X_3 \\ &\quad - 2^{\lfloor (n+2)/4 \rfloor + 2\lfloor (n-2)/8 \rfloor} X_1, \\ X_6(n) &= 2^{\lfloor (n-4)/8 \rfloor} X_6 + (2^{\lfloor n/4 \rfloor} - 2^{2\lfloor (n-4)/8 \rfloor + 1}) X_2 \\ &\quad - 2^{\lfloor n/4 \rfloor + 2\lfloor (n-4)/8 \rfloor + 1} X_1 \end{aligned}$$

and

$$\begin{aligned} X_7(n) &= 2^{\lfloor (n-6)/8 \rfloor} X_7 - (2^{\lfloor (n-2)/4 \rfloor} - 2^{2\lfloor (n-6)/8 \rfloor + 1}) (X_3 - 2X_2) \\ &\quad + 2^{2\lfloor (n+2)/8 \rfloor + \lfloor (n-2)/4 \rfloor} X_1. \end{aligned}$$

(3) Let  $\varphi: F(x) \rightarrow \tilde{K}(L_8^m)$  be the homomorphism defined by setting  $\varphi(x_i) = \sigma_i$  ( $1 \leq i \leq 7$ ).

**Proposition 2.22** (Kobayashi and Sugawara [12]). *The homomorphism  $\varphi$  is an epimorphism, and the kernel of  $\varphi$  coincides with the subgroup of  $F(x)$  generated by  $\{X_i(m) \mid 1 \leq i \leq 7\}$ .*

For each integer  $n$  with  $0 \leq n < m$ , we denote the inclusion map of  $L_8^n$  into

$L_8^m$  by  $i_n^m$ , and denote the kernel of the homomorphism

$$(i_n^m)^!: \tilde{K}(L_8^m) \rightarrow \tilde{K}(L_8^n)$$

by  $V_n$ . Set  $u = [(n+1)/2]$  and  $S_i = \varphi(X_i(2u))$  ( $1 \leq i \leq 7$ ). Then  $V_{2u}$  is the subgroup of  $\tilde{K}(L_8^n)$  generated by  $S_i$  ( $1 \leq i \leq 7$ ), and we have

$$\tilde{K}(L_8^m/L_8^n) \cong \begin{cases} V_{2u} & (n \equiv 0 \pmod{2}) \\ \mathbf{Z} \oplus V_{2u} & (n \equiv 1 \pmod{2}) \end{cases}.$$

According to [1], we have the following lemma.

**Lemma 2.23.** *The Adams operations are given by the following formulae, where  $s_i = \varphi(X_i)$  ( $1 \leq i \leq 7$ ) and  $k \equiv 1 \pmod{2}$ .*

- (1)  $\psi^k(s_i) = s_i$  ( $i = 1, 2$  or 4).
- (2)  $\psi^k(s_3) = \begin{cases} s_3 & (k \equiv 1 \pmod{4}) \\ -s_3 - 2s_2 & (k \equiv 3 \pmod{4}) \end{cases}.$
- (3)  $\psi^k(s_5) = \begin{cases} s_5 & (k \equiv 1 \pmod{8}) \\ s_5 + s_6 & (k \equiv 3 \pmod{8}) \\ -s_5 - 2s_4 & (k \equiv 5 \pmod{8}) \\ -s_5 - 2s_4 - s_6 & (k \equiv 7 \pmod{8}) \end{cases}.$
- (4)  $\psi^k(s_6) = \begin{cases} s_6 & (k \equiv \pm 1 \pmod{8}) \\ -s_6 & (k \equiv \pm 3 \pmod{8}) \end{cases}.$
- (5)  $\psi^k(s_7) = \begin{cases} s_7 & (k \equiv 1 \pmod{8}) \\ -s_7 + 2s_4 & (k \equiv 3 \pmod{8}) \\ s_7 - 2s_6 & (k \equiv 5 \pmod{8}) \\ -s_7 + 2s_4 + 2s_6 & (k \equiv 7 \pmod{8}) \end{cases}.$

### 3. Proofs of Theorems

In this section we prove Theorems 1, 2, 3 and 4.

**3.1. Proof of Theorems 1 and 3.** According to [8], we have

$$h(q, k) = \begin{cases} 2^{h_1(k)} & (q = 4) \\ 2^{h_2(k)} & (q = 8) \end{cases}.$$

Then Theorems 1 and 3 follows from (2.10) for the case  $m \geq n+3$ . Note that we have  $L_q^{n+1}/L_q^n \approx S^{n+1}$  and

$$L_q^{n+2}/L_q^n \cong \begin{cases} S^{n+1} \vee S^{n+2} & (n \equiv 1 \pmod{2}) \\ S^n L_q^2 & (n \equiv 0 \pmod{2}) \end{cases}.$$



This completes the proof of Theorems 1 and 3.

**3.2. Proof of Theorem 2.** The part (1) is obtained by (2.11) (2).

Suppose that the spaces  $L_4^m/L_4^n$  and  $L_4^{m+t}/L_4^{n+t}$  have the same stable homotopy type, where  $\nu_2(n+1) \geq i$  and  $h_1(m-2[(n+1)/2]) \geq i \geq 3$ . Then  $m \geq n+5$  and there exists a homotopy equivalence

$$f: S^j(L_4^m/L_4^n) \rightarrow S^{j-t}(L_4^{m+t}/L_4^{n+t}),$$

which induces an isomorphism

$$(3.1) \quad J(f^!): \tilde{J}(S^{j-t}(L_4^{m+t}/L_4^{n+t})) \rightarrow \tilde{J}(S^j(L_4^m/L_4^n)).$$

We can assume that  $\nu_2(j) \geq h_1(m-2[(n+1)/2]) = h_1(m-n-1)$ . By (2.11) (3),  $t \equiv 0 \pmod{8}$ . It follows from (2.11) (1) and Proposition 2.15 that we have  $\nu_2(t) \geq [(m-n-1)/2]$ ,

$$\tilde{J}(S^j(L_4^m/L_4^n)) \cong \mathbf{Z}/m((n+j+1)/2) \cdot 2^{c_1} \oplus \mathbf{Z}/2^{d_1+i_1} \oplus \mathbf{Z}/2^{k_1}$$

and

$$\tilde{J}(S^{j-t}(L_4^{m+t}/L_4^{n+t})) \cong \mathbf{Z}/m((n+j+1)/2) \cdot 2^{c_2} \oplus \mathbf{Z}/2^{d_2+i_2} \oplus \mathbf{Z}/2^{k_2},$$

where  $i_l = \min \{\nu_2(n+1)-1, h_1(m-n-1)\}$ ,

$$\begin{aligned} i_2 &= \min \{\nu_2(n+t+1)-1, h_1(m-n-1)\}, \\ k_1 &= \min \{\nu_2(n+1)-1, [(m-n-1)/8] + [(m-n+5)/8]\}, \\ k_2 &= \min \{\nu_2(n+t+1)-1, [(m-n-1)/8] + [(m-n+5)/8]\}, \\ c_l &= h_1(m-n-1) - i_l \quad (l=1 \text{ or } 2) \end{aligned}$$

and  $d_l = [(m-n-1)/8] + [(m-n+5)/8] - k_l$  ( $l=1$  or  $2$ ). Since  $c_l \geq d_l \geq 0$  and  $\nu_2(m((n+j+1)/2)) \geq i_l \geq k_l$  ( $l=1$  or  $2$ ), the isomorphism (3.1) implies that  $c_1 = c_2$ , and hence

$$\min \{\nu_2(n+1)-1, h_1(m-n-1)\} = \min \{\nu_2(n+t+1)-1, h_1(m-n-1)\}.$$

Since  $\nu_2(n+1) \geq i$ , this implies that we have  $\nu_2(n+t+1) > i$  if  $\nu_2(n+1) > i$  and  $\nu_2(n+t+1) = i$  if  $\nu_2(n+1) = i$ . Thus we have  $\nu_2(t) \geq i+1$ . The proof of the part (2) of Theorem 2 is completed by making use of the  $S$ -duality (Lemma 2.9).

The part (3) is obtained by (2.11) (3) and the parts (1) and (2) of Theorem 2. This completes the proof of Theorem 2.

**3.3. Proof of Theorem 4.** Suppose that the spaces  $L_8^m/L_8^n$  and  $L_8^{m+t}/L_8^{n+t}$  have the same stable homotopy type. Then there exists a homotopy equivalence

$$f: S^j(L_8^m/L_8^n) \rightarrow S^{j-t}(L_8^{m+t}/L_8^{n+t}),$$

which induces isomorphisms

$$(3.2) \quad f^*: \tilde{K}(S^{j-t}(L_8^{m+t}/L_8^{n+t})) \rightarrow \tilde{K}(S^j(L_8^m/L_8^n))$$

and

$$(3.3) \quad J(f^!): \tilde{J}(S^{j-t}(L_8^{m+t}/L_8^{n+t})) \rightarrow \tilde{J}(S^j(L_8^m/L_8^n)).$$

We can assume that  $v_2(j) \geq a_0(m, n) + 1$ . Suppose  $m \geq n + 5$ . Then, by (2.11) (3),  $t \equiv 0 \pmod{8}$ . If  $n \not\equiv 3 \pmod{4}$ , then Proposition 2.19 asserts that the exponent of the group  $\tilde{J}(S^j(L_8^m/L_8^n))$  is equal to  $2^{b_1(j, m, n)}$  and the exponent of the group  $\tilde{J}(S^{j-t}(L_8^{m+t}/L_8^{n+t}))$  is equal to  $2^{b_1(j-t, m+t, n+t)}$ , and the isomorphism (3.3) implies that  $b_1(j, m, n) = b_1(j-t, m+t, n+t)$ . In the case  $n \equiv 3 \pmod{4}$ ,  $f$  induces a homotopy equivalence

$$\tilde{f}: S^j(L_8^m/L_8^{n+1}) \rightarrow S^{j-t}(L_8^{m+t}/L_8^{n+t+1}),$$

which induces an isomorphism

$$J(\tilde{f}^!): \tilde{J}(S^{j-t}(L_8^{m+t}/L_8^{n+t+1})) \rightarrow \tilde{J}(S^j(L_8^m/L_8^{n+1})).$$

Since  $b_1(j, m, n) = b_1(j, m, n+1)$  and  $b_1(j-t, m+t, n+t) = b_1(j-t, m+t, n+t+1)$ , the isomorphism  $J(\tilde{f}^!)$  implies that  $b_1(j, m, n) = b_1(j-t, m+t, n+t)$ . In either case, we have  $b_1(j, m, n) = b_1(j-t, m+t, n+t)$ . Hence  $v_2(j-t) - a_3(n, n+4) \geq a_5(m, n)$ , and

$$(3.4) \quad v_2(t) \geq \begin{cases} a_5(m, n) - 2 & (n \equiv 0, 1 \text{ or } 7 \pmod{8}) \\ a_5(m, n) - 1 & (\text{otherwise}). \end{cases}$$

By Lemma 2.9, (2.11) (2) and (3.4), the part (1) of Theorem 4 is obtained except for the case  $n \equiv m - n \equiv 2 \pmod{8}$ . So, assume that  $n \equiv m - 2 \equiv 2 \pmod{8}$ . Let  $Y_i$  be the element of  $\tilde{K}(S^j(L_8^m/L_8^n))$ , which corresponds to  $I^{j/2}(S_i)$  by the isomorphism

$$(p_n^m)^!: \tilde{K}(S^j(L_8^m/L_8^n)) \xrightarrow{\cong} I^{j/2}(V_n),$$

where  $I$  denotes the Bott periodicity isomorphism ( $1 \leq i \leq 7$ ). Set  $u_1 = Y_6$ ,

$$\begin{aligned} u_2 &= Y_4 + Y_2 + 2^{(m-n-10)/8}(2Y_2 + Y_1) + 2^{(3m-3n-6)/8} Y_1, \\ u_3 &= Y_7 - Y_2 - Y_1 + 2^{(m-n-10)/8}(2Y_2 + Y_1 - Y_3) + 2^{(3m-3n-14)/8} Y_1 \\ &\quad - 2^{(2m-2n-12)/8} Y_1, \\ u_4 &= 2Y_4 - Y_1 + 2^{(m-n-2)/8}(2Y_2 + Y_1) + 2^{(3m-3n+2)/8} Y_1, \\ u_5 &= Y_5 + Y_1 - 2^{(3m-3n-6)/8} Y_1, \\ u_6 &= 2Y_6 - Y_2 \end{aligned}$$

and  $u_7 = 2Y_7 + Y_3 - 2Y_2 - 2Y_1 + 2^{(m-n-2)/8}(2Y_2 + Y_1 - Y_3) + 2^{(3m-3n-6)/8} Y_1$ . Then, by Proposition 2.22, we have

$$\tilde{K}(S^j(L_8^m/L_8^n)) \simeq \langle \{u_i \mid 1 \leq i \leq 7\} \rangle / \langle \{2^{a(i)} u_i \mid 1 \leq i \leq 7\} \rangle,$$

where

$$\begin{aligned} a(1) &= (m-n+4)/2, \\ a(2) &= a(3) = (m-n+2)/4, \\ a(5) &= a(6) = (m-n-2)/8 \end{aligned}$$

and  $a(4) = a(7) = (m-n-10)/8$ . According to Lemma 2.23, the Adams operation  $\psi^5$  is given by the following formulae.

$$(3.5) \quad \begin{cases} (1) & \psi^5(u_1) = 5^{j/2}(u_1 - u_6). \\ (2) & \psi^5(u_2) = 5^{j/2} u_2. \\ (3) & \psi^5(u_3) = 5^{j/2}(u_3 - u_6). \\ (4) & \psi^5(u_4) = 5^{j/2} u_4. \\ (5) & \psi^5(u_5) = 5^{j/2}(2^{(m-n-2)/8}(2u_2 - u_4) - u_4 - u_5). \\ (6) & \psi^5(u_6) = 5^{j/2}(-u_6). \\ (7) & \psi^5(u_7) = 5^{j/2}(u_7 - 2u_6). \end{cases}$$

Choose  $v_i \in \tilde{K}(S^{j-t}(L_8^{m+t}/L_8^{n+t}))$  similarly as  $u_i \in \tilde{K}(S^j(L_8^m/L_8^n))$  ( $1 \leq i \leq 7$ ), and set

$$f^*(v_i) = \sum_{k=1}^7 a_{ik} u_k \quad (1 \leq i \leq 7).$$

By the equality  $\psi^5 \circ f^* = f^* \circ \psi^5$ , we have

$$(3.6) \quad 5^{j/2} a_{11} \equiv 5^{(j-t)/2}(a_{11} - a_{61}) \pmod{2^{a(1)}}.$$

$$(3.7) \quad 5^{j/2} a_{31} \equiv 5^{(j-t)/2}(a_{31} - a_{61}) \pmod{2^{a(1)}}.$$

Since  $v_2(a_{31}) \geq a(1) - a(3) = (m-n+6)/4$ , (3.7) implies that  $a_{61} \equiv 0 \pmod{2^{a(1)}}$ . It follows from (3.6) that we have  $(5^{t/2} - 1)a_{11} \equiv 0 \pmod{2^{a(1)}}$ . Note that  $a_{11} \equiv 1 \pmod{2}$ . According to [15, Lemma 3.1], we see that  $v_2(t) + 1 \geq a(1) = (m-n+4)/2$ . Hence

$$v_2(t) \geq (m-n+2)/2 = [m/2] - [(n+1)/2] + 1.$$

This completes the proof of the part (1) of Theorem 4.

Suppose that  $v_2(n+1) \geq i$  and  $h_2(m-2[(n+1)/2]) \geq i \geq 3$ . Then  $m \geq n+5$  and  $t \equiv 0 \pmod{8}$  by (2.11) (3). It follows from (2.11) (1) and Proposition 2.19 that we have  $v_2(t) \geq [(m-n-1)/2]$ ,

$$\tilde{J}(S^j(L_8^m/L_8^n)) \simeq \mathbf{Z}/m((n+j+1)/2) \cdot 2^{c_1} \oplus \mathbf{Z}/2^{c_2+i_1} \oplus \mathbf{Z}/2^{i_2} \oplus \mathbf{Z}/2^{i_3}$$

and

$$\tilde{J}(S^{j-t}(L_8^{m+t}/L_8^{n+t})) \simeq \mathbf{Z}/m((n+j+1)/2) \cdot 2^{d_1} \oplus \mathbf{Z}/2^{d_2+k_1} \oplus \mathbf{Z}/2^{k_2} \oplus \mathbf{Z}/2^{k_3},$$

where  $i_1 = \min \{v_2(n+1) - 1, h_2(m-m-1)\}$ ,

$$k_1 = \min \{v_2(n+t+1) - 1, h_2(m-n-1)\},$$

$$\begin{aligned}
 i_2 &= \min \{ \nu_2(n+1) - 2, [(m-n+3)/8] + [(m-n-3)/8] \}, \\
 k_2 &= \min \{ \nu_2(n+t+1) - 2, [(m-n+3)/8] + [(m-n-3)/8] \}, \\
 i_3 &= k_3 = [(m-n-1)/8], \\
 c_1 &= h_2(m-n-1) - i_1, \\
 d_1 &= h_2(m-n-1) - k_1, \\
 c_2 &= [(m-n+3)/8] + [(m-n-3)/8] - i_2,
 \end{aligned}$$

and  $d_2 = [(m-n+3)/8] + [(m-n-3)/8] - k_2$ . Since  $c_1 \geq c_2 \geq 0$ ,  $c_2 + i_1 \geq i_3$ ,

$$\begin{aligned}
 \nu_2(\mathfrak{m}((n+j+1)/2)) &\geq i_1 \geq i_2, \\
 \nu_2(\mathfrak{m}((n+j+1)/2)) + c_1 &\geq h_2(m-n-1) + 1
 \end{aligned}$$

and  $\max \{d_2 + k_1, k_2, k_3\} \leq h_2(m-n-1)$ , the isomorphism (3.3) implies that  $c_1 = d_1$ , and hence

$$\min \{ \nu_2(n+1) - 1, h_2(m-n-1) \} = \min \{ \nu_2(n+t+1) - 1, h_2(m-n-1) \}.$$

Since  $\nu_2(n+1) \geq i$ , this implies that we have  $\nu_2(n+t+1) > i$  if  $\nu_2(n+1) > i$  and  $\nu_2(n+t+1) = i$  if  $\nu_2(n+1) = i$ . Thus we have  $\nu_2(t) \geq i+1$ . The proof of the part (2) of Theorem 4 is completed by making use of the  $S$ -duality (Lemma 2.9).

By (2.11) (3) and the parts (1) and (2) of Theorem 4, the part (3) of Theorem 4 is obtained except for the case  $n \equiv 0 \pmod{8}$  and  $m = n+6$ . So, assume that  $n \equiv 0 \pmod{8}$  and  $m = n+6$ . Let  $Y_i$  be the element of  $\tilde{K}(S^j(L_8^m/L_8^n))$ , which corresponds to  $I^{j/2}(S_i)$  by the isomorphism

$$(p_n^m)^!: \tilde{K}(S^j(L_8^m/L_8^n)) \xrightarrow{\cong} I^{j/2}(V_n)$$

( $1 \leq i \leq 7$ ). Set  $u_1 = Y_5$ ,  $u_2 = Y_6 + 2Y_5$  and  $u_3 = Y_7 + 2Y_5$ . Then, by Proposition 2.22, we have

$$\tilde{K}(S^j(L_8^m/L_8^n)) \cong \langle \{u_i \mid 1 \leq i \leq 3\} \rangle / \langle \{2^{a(i)} u_i \mid 1 \leq i \leq 3\} \rangle,$$

where  $a(1) = 32$  and  $a(2) = a(3) = 4$ . According to Lemma 2.23, we have the following formulae.

$$(3.8) \quad \begin{cases} (1) & \psi^{-1}(u_1) = 5u_1 + u_2. \\ (2) & \psi^5(u_1) = 5^{j/2}(17u_1 + 2u_2). \\ (3) & \psi^{-1}(u_2) = 8u_1 - u_2. \\ (4) & \psi^5(u_2) = 5^{j/2}(16u_1 + u_2). \\ (5) & \psi^{-1}(u_3) = -8u_1 - u_3. \\ (6) & \psi^5(u_3) = 5^{j/2}(16u_1 + u_3). \end{cases}$$

Choose  $v_i \in \tilde{K}(S^{j-t}(L_8^{m+t}/L_8^{n+t}))$  similarly as  $u_i \in \tilde{K}(S^j(L_8^m/L_8^n))$  ( $1 \leq i \leq 3$ ), and set

$$f^*(v_i) = \sum_{k=1}^3 a_{ik} u_k \quad (1 \leq i \leq 3).$$

By the equality  $\psi^{-1} \circ f^* = f^* \circ \psi^{-1}$ , we have

$$(3.9) \quad 8a_{12} - 8a_{13} \equiv a_{21} \pmod{32}.$$

By the equality  $\psi^5 \circ f^* = f^* \circ \psi^5$ , we have

$$(3.10) \quad 5^{j/2}(17a_{11} + 16a_{12} + 16a_{13}) \equiv 5^{(j-t)/2}(17a_{11} + 2a_{21}) \pmod{32}.$$

By (3.9), (3.10) and the fact  $5^{t/2} \equiv 1 \pmod{2^{v_2(t)+1}}$ , we have

$$(5^{t/2} - 1)a_{11} \equiv 0 \pmod{32}.$$

Note that  $a_{11} \equiv 1 \pmod{2}$ . According to [15, Lemma 3.1], we see that  $v_2(t) + 1 \geq 5$ . Hence

$$v_2(t) \geq 4 = [m/2] - [(n+1)/2] + 1.$$

This completes the proof of Theorem 4.

### References

- [1] J.F. Adams: *Vector fields on spheres*, Ann. of Math. **75** (1962), 603–632.
- [2] J.F. Adams: *On the groups  $J(X)$ -I*, Topology **2** (1963), 181–195.
- [3] J.F. Adams: *On the groups  $J(X)$ -II, -III*, Topology **3** (1965), 137–171, 193–222.
- [4] J.F. Adams and G. Walker: *On complex Stiefel manifolds*, Proc. Camb. Phil. Soc. **61** (1965), 81–103.
- [5] M.F. Atiyah: *Thom complexes*, Proc. London Math. Soc. **11** (1961), 291–310.
- [6] M.M. Cohen: *A course in simple-homotopy theory*, Graduate Texts in Mathematics 10, Springer-Verlag, 1973.
- [7] D.M. Davis and M. Mahowald: *Classification of the stable homotopy types of stunted real projective spaces*, Pacific J. Math. **125** (1986), 335–345.
- [8] K. Fujii, T. Kobayashi and M. Sugawara: *Stable homotopy types of stunted lens spaces*, Mem. Fac. Sci. Kochi Univ. (Math.) **3** (1982), 21–27.
- [9] T. Kambe, H. Matsunaga and H. Toda: *A note on stunted lens space*, J. Math. Kyoto Univ. **5** (1966), 143–149.
- [10] T. Kobayashi: *Stable homotopy types of stunted lens spaces mod  $p$* , Mem. Fac. Sci. Kochi Univ. (Math.) **5** (1984), 7–14.
- [11] T. Kobayashi: *Stable homotopy types of stunted lens spaces mod  $2^r$* , Mem. Fac. Sci. Kochi Univ. (Math.) **11** (1990), 17–22.
- [12] T. Kobayashi and M. Sugawara: *On stable homotopy types of stunted lens spaces*, Hiroshima Math. J. **1** (1971), 287–304; **II**, **7** (1977), 689–705.
- [13] S. Kôno and A. Tamamura: *On  $J$ -groups of  $S^l(RP(t-l)/RP(n-l))$* , Math. J. Okayama Univ. **24** (1982), 45–51.
- [14] S. Kôno and A. Tamamura:  *$J$ -groups of the suspensions of the stunted lens spaces mod  $p$* , Osaka J. Math. **24** (1987), 481–498.

- [15] S. Kôno and A. Tamamura: *J-groups of suspensions of stunted lens spaces mod 4*, Osaka J. Math. **26** (1989), 319–345.
- [16] S. Kôno and A. Tamamura: *J-groups of suspensions of stunted lens spaces mod 8*, to appear.
- [17] N. Mahammed: *A propos de la K-théorie des espaces lenticulaires*, C.R. Acad. Sc. Paris **271** (1970), 639–642.
- [18] M. Mimura, J. Mukai and G. Nishida: *Representing elements of stable homotopy groups by symmetric maps*, Osaka J. Math. **11** (1974), 105–111.
- [19] D. Quillen: *The Adams conjecture*, Topology **10** (1971), 67–80.

Department of Mathematics  
Osaka University  
Toyonaka, Osaka 560, Japan

