



Title	One-Step Synthesis of Benzophosphole Derivatives from Arylalkynes by Phosphenium-Dication-Mediated Sequential C-P/C-C Bond Forming Reaction
Author(s)	Nishimura, Kazutoshi; Xu, Shibo; Nishii, Yuji et al.
Citation	Organic Letters. 2023, 25(9), p. 1503-1508
Version Type	AM
URL	https://hdl.handle.net/11094/92684
rights	© 2023 American Chemical Society.
Note	

The University of Osaka Institutional Knowledge Archive : OUKA

<https://ir.library.osaka-u.ac.jp/>

The University of Osaka


One-Step Synthesis of Benzophosphole Derivatives from Arylalkynes by Phosphenium-Dication-Mediated Sequential C–P/C–C Bond Forming Reaction

Kazutoshi Nishimura,^{†,§} Shibo Xu,^{‡,§} Yuji Nishii,[†] and Koji Hirano^{*,†,‡}

[†]Department of Applied Chemistry, Graduate School of Engineering, Osaka University, Suita, Osaka 565-0871, Japan

[‡]Innovative Catalysis Science Division, Institute for Open and Transdisciplinary Research Initiatives (ICS-OTRI), Osaka University, Suita, Osaka 565-0871, Japan

Supporting Information Placeholder




ABSTRACT: A metal-free, phosphenium-dication-mediated sequential C–P and C–C bond forming reaction has been developed. This protocol can provide concise access to the (di)benzophosphole derivatives in one synthetic operation from the readily available and simple arylalkynes and phosphinic acids. Application to the multiple cyclization reaction and the fully intermolecular three-component-coupling-type reaction are also described.

Phosphole derivatives have recently received significant attention in fields of phosphorus-based organic functional materials because of their unique optoelectronic and physical properties.¹ Accordingly, numerous strategies for the preparation and decoration of phosphole nuclei have been developed so far. While most traditional protocols largely relied on the transition-metal-mediated multistep sequence with some toxic and sensitive reagents,^{1,2} P-radical,³ -cation,⁴ and -dication⁵-promoted methods recently appeared to streamline the synthesis of targeted phosphole cores via efficient C–P bond formation. In particular, our previous work using the phosphenium dication equivalent^{5a} enabled the rapid construction of the dibenzophospholes from the corresponding simple biaryls and phosphinic acids (Scheme 1a). However, electron-rich (hetero)aromatics such as indole, thiophene, and phenoxy moieties were generally essential to trigger the reaction, which apparently limited the versatility of the phosphole product. Herein, we report a more flexible approach to the benzophospholes from the arylalkynes and phosphinic acids: the phosphenium-dication-mediated sequential C–P/C–C bond forming reaction is developed (Scheme 1b). By using the alkyne moiety as the reaction trigger, this strategy easily accesses the non-heteroaromatic benzene-fused phosphole derivatives


with high efficiency. The starting arylalkynes are also readily available by the Sonogashira coupling, same as in the preparation of biaryls via the Suzuki-Miyaura coupling. Thus, this protocol significantly expands the applicability of the P-cation-based strategy in the phosphole synthesis. Additionally, the multiple phosphole-constructing cyclization and application to the fully intermolecular three-component-coupling reaction of alkynes, phosphinic acids, and external nucleophiles are achieved. We note that during the course of this study, Huang and Xiao reported the related reaction of *ortho*-MeO- and MeS-substituted diarylalkynes,^{5b} but the products were restricted to the furan- and thiophene-fused dibenzophospholes, which are more readily synthesized by our previous double C–P bond forming reaction of heterobiaryls.^{5a} In addition, the intermolecular process was not mentioned at all.

Scheme 1. Phosphenium-Dication-Mediated Approaches to (Di)Benzophospholes

a) Double C–P bond formation from (hetero)biaryls




b) Sequential C–P/C–C bond formation from arylalkynes (**this work**)



Our reaction design is shown in Scheme 2. The active phosphonium dication equivalent **A** is initially generated from the phosphinic acid **1** by the action of 2 equiv of dehydrating reagent Ti_2O and Lewis base (**L**) via P(V)/P(III) tautomerization.⁶ Subsequent [1+2] cycloaddition with the alkyne moiety of **2** generates the corresponding three-membered phosphirenium species **B**.⁷ The ring-opening process is induced by the intramolecular attack of the tethered Ar group to form **C**. The intermediate **C** has still one positive charge on the phosphorus and thus undergoes the intramolecular phospho-Friedel-Crafts-type reaction⁸ to furnish the benzophosphole derivative **3**. Overall, two C–P bonds and one C–C bond are constructed in one synthetic operation.

Scheme 2. Working Hypothesis




We selected phenylphosphinic acid (**1a**) and the arylalkyne **2a** as model substrates and started optimization studies to identify the suitable conditions including the dehydrating reagent, Lewis base, and solvent. After exten-

sive screening, the targeted benzophosphole oxide derivative **3aa** was found to be isolated in 90% yield by using the combination of Ti_2O and 2,6-lutidine in heated DCE (Scheme 3). Some observations are to be noted: the corresponding P(III) phosphole derivative was not detected at all, and **3aa** was only formed even without any additional oxidative treatment.⁹ Ti_2O was the critical dehydrating reagent as far as we tested. No reaction occurred under Lewis-base-free conditions. Some other pyridine-type Lewis bases also promoted the reaction, whereas sluggish conversion of **1a** was observed in the presence of alkylamines and inorganic bases (see the Supporting Information for details).

With optimal conditions in hand, we first investigated the scope of the arylalkynes **2**. In addition to the model substrate **2a**, electron-donating and -withdrawing substituents both were well compatible under the reaction conditions (**3ab** and **3ac**). The sterically congested *ortho*-Me group (**3ad**) and more condensed naphthalene (**3ae**) and phenanthrene (**3af**) substitutions were also accommodated. The alkyne containing the heteroaromatic thiophene moiety underwent the reaction without any difficulty (**3ag**). The ethylene tether could also be replaced with the oxygen-, nitrogen-, and ester-based linkers, and the corresponding heteroatom-incorporated benzophosphole derivatives **3ah–ak** were isolated in synthetically acceptable yields. In the case of **3ai**, the regioisomeric **3ai'** was also formed probably because of the competitive steric and electronic effects. Notably, in the reaction of the NTs-type substrate **2j**, the additional treatment with KOH in MeOH promoted the elimination of TsH to selectively form the aromatized pyridine-fused phosphole **3aj**. The benzene ring also worked well as the two-carbon tether, and the targeted highly condensed **3al**, **3am**, and **3an** were obtained in good yields. The electronically diverse arylphosphinic acids **1b** and **1c** could also be coupled with **2a** to deliver the corresponding **3ba** and **3ca** in 69% and 72% yields, respectively. Moreover, the aliphatic phosphinic acid was employed with good efficiency (**3da**). On the other hand, the alkene substrate instead of the alkyne did not provide the targeted cyclization product (see the Supporting Information for details).

Scheme 3. Phosphenium-Dication-Mediated Sequential C–C/C–P Bond Formation of Arylalkynes **2** with Phosphinic Acids **1**^a




^a Reaction conditions: **1** (0.20 mmol), **2** (0.10 mmol), Tf_2O (0.48 mmol), 2,6-lutidine (0.48 mmol), DCE (1.5 mL), 60 or 90 °C, 16 h. Isolated yields are shown. ^b On a 1.0 mmol scale. ^c Obtained as a 7:3 mixture of regioisomers. ^d With additional treatment with KOH in MeOH. See the Supporting Information for details. ^e On a 0.20 mmol scale in toluene.

Additionally notable is that the conjugated 1,3-diyne **2o** was readily converted to the phosphole derivative **3ao** by the sequential two C–P and two C–C bond formations in one synthetic operation (Scheme 4a). The phosphonium-dication-promoted strategy was also applied to the double cyclization reaction of the dialkynyl naphthalenes **2p** and **2q** (Scheme 4b and c). In these cases, four C–P and two C–C bonds were successively formed to deliver the corresponding octa- and decacyclic systems **3cp** and **3cq** with two phosphole rings. The structure of *syn*-**3cq** was confirmed by X-ray analysis (CCDC 2210187). Moreover, the dialkynylterphenyl **2r** also participated in the double cyclization reaction to afford benzophosphole-fused dibenzanthracene **3cr** in an acceptable yield (Schemes 4d). Its crystal structure was also successfully obtained (CCDC 2210186). On the other hand, the regioisomeric dialkynylterphenyl **2s** gave the helical, unexpected double

cyclization product **3cs** (Scheme 4e). This is probably because of the acid-mediated aryl ring 1,2-migration after the first cyclization event, a similar process of which is frequently found in the acid-promoted Scholl reaction of the condensed aromatic compounds.¹⁰ The structure was finally determined by the crystallographic analysis after the conversion of P=O to P=S (*anti*-**3cs-S**; CCDC 2231651). Notably, in Schemes 4b–e, the use of CF₃-containing phosphinic acid **1c** was critical to success: with simple phenyl-substituted **1a**, the reaction stopped after the single cyclization. The outcome indicates that the electron-withdrawing group on the phosphorus increases the electrophilicity of the phosphonium dication species and enables the otherwise difficult second cyclization process.

The preliminary optoelectronic properties of highly condensed *syn*-**3cq**, *syn*-**3cr**, *anti*-**3cr**, and **3cs** were investigated.


Scheme 4. Double Cyclizations from 1,3-Diynes and Dialkynylarenes



Their UV/vis absorption, fluorescence spectra, and images in CHCl_3 are shown in Figures S6 and S7, and the absorption/emission properties ($\lambda_{\text{abs}}/\lambda_{\text{em}}$) and fluorescence quantum yields (Φ_{f}) are summarized in Table S5. All compounds were strongly fluorescent in solution and gave a relatively broad range of their longest wavelength absorption maxima in the visible light region. In particular, the decacyclic system *syn*-**3cq** showed a large bathochromic shift of its λ_{abs} value (470 nm) with a higher molar extinction coefficient (ϵ) and fluorescence in the green color region (492–522 nm). Whereas the quantum efficiencies of *syn*-**3cq**, *syn*-**3cr**, and *anti*-**3cr** were good (Φ_{f} = 62–69%), the helical **3cs** showed a relatively poor value of Φ_{f} = 25%, thus indicating that the planarity is critical for the fluorescence event. The generally observed smaller Stokes shifts reflect their rigid structures associated with the highly condensed acene-type cores. The electrochemical properties of the aforementioned compounds

were examined by cyclic voltammetry (CV) and differential pulse voltammetry (DPV) in MeCN with tetrabutylammonium hexafluorophosphate (Bu_4NPF_6) as an electrolyte *versus* ferrocene/ferrocenium ion (Fc/Fc^+) (Figures S8–10), and their HOMO and LUMO levels were estimated according to the first oxidation potentials and the optical band gaps ($E_{\text{g}}^{\text{opt}}$). Due to its poor solubility, any clear spectra of *anti*-**3cr** were not obtained, and the HOMO/LUMO levels were thus investigated by density functional theory (DFT) calculations at the PBE0/6-31+G(d) level of theory (Figure S11). The data are summarized in Table S6. The CV of *syn*-**3cq**, *syn*-**3cr**, and **3cs** all showed irreversible oxidation waves, and oxidation potential values $E^{1/2}_{\text{ox}}$ are thus determined by the DPV. The HOMO levels were almost identical but an even lower LUMO level was estimated for *syn*-**3cq**, which may suggest the larger intramolecular charge transfer ability.

Scheme 5. Fully Intermolecular Three Component Coupling of Phosphinic Acids, Diarylacetylenes, and Electron-Rich (Hetero)aromatics



We finally attempted to apply the phosphonium dication strategy to the fully intermolecular three component coupling reaction with the external nucleophiles. After the optimization studies (see the Supporting Information for details), we were pleased to find that the reaction of phosphinic acid **1a**, diphenylacetylene (**4a**), and *N*-Me indole (**5a**) proceeded in the presence of TiF_2O and pyridine to afford the C3-indolylbenzophosphole **6aaa** in 60% yield after the oxidative workup with aq. H_2O_2 (Scheme 5).¹¹ The C3-indolyl **6aaa** was isolable but gradually underwent the dehydrogenative cyclization under visible light irradiation to finally deliver the *P,N*-containing heteroacene **7aaa**. The structure of the Me-substituted analogue **7aba** was unambiguously confirmed by X-ray analysis (CCDC 2202706). In addition to the indole **5a**, electron-rich (hetero)aromatics, including 3-methoxythiophene and 1,3-dimethoxybenzene, could also be used as the external nucleophiles, and the corresponding 2,3-diarylbenzophospholes **6aab** and **6aac** were successfully formed. Similar to the indole derivatives **6aaa** and **6aba**, the thiophene-substituted **6aab** could be converted to **7aab** by the visible-light-promoted cyclization with concomitant elimination of MeOH, while the benzene-derived **6aac** was stable and unchanged even under photoirradiation.

In conclusion, we have developed a phosphonium-dication-mediated sequential C–P/C–C bond forming reaction of the relatively simple arylalkynes and phosphinic acids, giving the corresponding benzophosphole derivatives in good to high yields under TiF_2O -promoted, metal-free conditions. The alkyne-triggered reaction enables the construction of arene-fused phosphole rings without any assistance of electron-rich heteroaromatics, which are inevitable in the previous phospho-Friedel-Crafts reaction.^{5a} Additionally, the reaction is successfully applicable to the double cyclization of dialkynyl arenes with 2 equiv of phosphinic acids, thus delivering the highly condensed P-containing acene-type molecules of potent interest in material chemistry. Moreover, the fully intermolecular three-component-coupling reaction of arylalkynes, phosphinic

acids, and electron-rich arenes is possible to provide potentially modular access to the C2,C3-diarylated benzo-phospholes. Detailed mechanistic studies and further applications to the design and synthesis of phosphole-based organic functional materials are ongoing in our laboratory.

ASSOCIATED CONTENT

Data Availability Statement

The data underlying this study are openly available in the published article and its online Supporting Information.

Supporting Information

The Supporting Information is available free of charge on the ACS Publications website at DOI: 10.1021/acs.orglett.xxxx.

^1H , $^{13}\text{C}\{^1\text{H}\}$, $^{19}\text{F}\{^1\text{H}\}$, and $^{31}\text{P}\{^1\text{H}\}$ NMR spectra, ORTEP drawing, detailed optimization studies, optoelectronic properties, and DFT calculations (PDF)

Accession Codes

CCDC 2202706, 2210186, 2210187, and 2202706 contain the supplementary crystallographic data for this paper. These data can be obtained free of charge via www.ccdc.cam.ac.uk/data_request/cif, or by emailing data_request@ccdc.cam.ac.uk, or by contacting The Cambridge Crystallographic Data Centre, 12 Union Road, Cambridge CB2 1EZ, UK; fax: +44 1223 336033.

AUTHOR INFORMATION

Corresponding Author

Koji Hirano – Department of Applied Chemistry, Graduate School of Engineering and Innovative Catalysis Science Division, Institute for Open and Transdisciplinary Research Initiatives (ICS-OTRI), Osaka University, Suita, Osaka 565-0871, Japan; orcid.org/0000-0001-9752-1985; Email: k_hirano@chem.eng.osaka-u.ac.jp.

Authors

Kazutoshi Nishimura – Department of Applied Chemistry, Graduate School of Engineering, Osaka University, Suita, Osaka 565-0871, Japan

Shibo Xu – Innovative Catalysis Science Division, Institute for Open and Transdisciplinary Research Initiatives (ICS-OTRI), Osaka University, Suita, Osaka 565-0871, Japan

Yuji Nishii – Department of Applied Chemistry, Graduate School of Engineering, Osaka University, Suita, Osaka 565-0871, Japan; orcid.org/0000-0002-6824-0639.

Complete contact information is available at: <https://pubs.acs.org/10.1021/xxxx>.

Author Contributions

[§]K.N. and S.X. contributed equally to this work.

Notes

The authors declare no competing financial interest.

ACKNOWLEDGMENT

This work was supported by JSPS KAKENHI Grant Nos. JP 21J10947 (Grant-in-Aid for JSPS Research Fellow, to K.N.) and JP 22H02077 (Grant-in-Aid for Scientific Research (B), to K.H.) as well as by JST FOREST Program, Grant Number JPMJFR211X to K.H. We also thank Mr. Shotaro Nakamura (Osaka University) for his assistance with X-ray analysis.

REFERENCES

(1) Reviews and books: (a) Baumgartner, T.; Réau, R. Organophosphorus π -Conjugated Materials. *Chem. Rev.* **2006**, *106*, 4681–4727. (b) Matano, Y.; Imahori, H. *Org. Biomol. Chem.* Design and synthesis of phosphole-based π systems for novel organic materials. **2009**, *7*, 1258–1271. (c) Duffy, M. P.; Delaunay, W.; Bouit, P.-A.; Hissler, M. π -Conjugated phospholes and their incorporation into devices: components with a great deal of potential. *Chem. Soc. Rev.* **2016**, *45*, 5296–5310. (d) Baumgartner, T.; Jäkle, F. *Main Group Strategies towards Functional Hybrid Materials*, Wiley, 2018. Selected examples: (e) Tsuji, H.; Sato, K.; Ilies, L.; Itoh, Y.; Sato, Y.; Nakamura, E. Modular Synthesis of Benzo[b]phosphole Derivatives via BuLi-Mediated Cyclization of (o-Alkynylphenyl)phosphine. *Org. Lett.* **2008**, *10*, 2263–2265. (f) Fukazawa, A.; Yamada, H.; Yamaguchi, S. Phosphonium- and Borate-Bridged Zwitterionic Ladder Stilbene and Its Extended Analogues. *Angew. Chem., Int. Ed.* **2008**, *47*, 5582–5585. (g) Tsuji, H.; Sato, K.; Sato, Y.; Nakamura, E. Benzo[b]phosphole sulfides. Highly electron-transporting and thermally stable molecular materials for organic semiconductor devices. *J. Mater. Chem.* **2009**, *19*, 3364–3366. (h) Tsui, H.; Sato, K.; Sato, Y.; Nakamura, E. Benzophosphole Oxide and Sulfide for Thermally Stable Cathode Buffer Layers in Organic Thin-Film Photovoltaic Devices. *Chem.-Asian J.* **2010**, *5*, 1294–1297. (i) Wang, C.; Taki, M.; Sato, Y.; Fukazawa, A.; Higashiyama, T.; Yamaguchi, S. Super-Photostable Phosphole-Based Dye for Multiple-Acquisition Stimulated Emission Depletion Imaging. *J. Am. Chem. Soc.* **2017**, *139*, 10374–10381.

(2) For a recent review: (a) Wu, B.; Yoshikai, N. Recent developments in synthetic methods for benzo[b]heteroles. *Org. Biomol. Chem.* **2016**, *14*, 5402–5416. Selected examples: (b) Balthazor, T. M. Phosphindolin-3-one. A useful intermediate for phosphindole synthesis. *J. Org. Chem.* **1980**, *45*, 2519–2522. (c) Cordaro, J. G.; Stein, D.; Grützmacher, H. A Synthetic Cycle for the Ruthenium-Promoted Formation of 1H-Phosphindoles from Phosphaalkynes. *J. Am. Chem. Soc.* **2006**, *128*, 14962–14971. (d) Xu, Y.; Wang, Z.; Gan, Z.; Xi, Q.; Duan, Z.; Mathey, F. Versatile Synthesis of Phospholides from Open-Chain Precursors. Application to Annulated Pyrrole- and Silole-Phosphole Rings. *Org. Lett.* **2015**, *17*, 1732–1734. (e) Mei, Y.; Yan, Z.; Liu, L. L. Facile Synthesis of the Dicyanophosphide Anion via Electrochemical Activation of White Phosphorus: An Avenue to Organophosphorus Compounds. *J. Am. Chem. Soc.* **2022**, *144*, 1517–1522. (f) Niecke, E.; Nieger, M.; Wenderoth, P. Phosphindolyl Anions by Elimination from 1-Phosphoallyllithium Complexes— η^5 and η^3 Coordination of a Phospholyl Fragment. *Angew. Chem., Int. Ed. Engl.* **1994**, *33*, 353–354. (g) Xu, L.; Chi, Y.; Du, S.; Zhang, W.-X.; Xi, Z. Direct Synthesis of Phospholyl Lithium from White Phosphorus. *Angew. Chem., Int. Ed.* **2016**, *55*, 9187–9190.

(3) (a) Furukawa, S.; Haga, S.; Kobayashi, J.; Kawashima, T. Synthesis of π -Extended Dibenzophospholes by Intramolecular Radical Cyclization and Their Properties. *Org. Lett.* **2014**, *16*, 3228–3231. (b) Unoh, Y.; Hirano, K.; Satoh, T.; Miura, M. An Approach to Benzophosphole Oxides through Silver- or Manganese-Mediated Dehydrogenative Annulation Involving C–C and C–P Bond Formation. *Angew. Chem., Int. Ed.* **2013**, *52*, 12975–12979. (c) Chen, Y.-R.; Duan, W.-L. Silver-Mediated Oxidative C–H/P–H Functionalization: An Efficient Route for the Synthesis of Benzo[b]phosphole Oxides. *J. Am. Chem. Soc.* **2013**, *135*, 16754–16757. (d) Ma, W.; Ackermann, L. Silver-Mediated Alkyne Annulations by C–H/P–H Functionalizations: Step-Economical Access to Benzophospholes. *Synthesis* **2014**, *46*, 2297–2304. (e) Zhang, P.; Gao, Y.; Zhang, L.; Li, Z.; Liu, Y.; Tang, G.; Zhao, Y. Copper-Catalyzed Cycloaddition between Secondary Phosphine Oxides and Alkynes: Synthesis of Benzophosphole Oxides. *Adv. Synth. Catal.* **2016**, *358*, 138–142. (f) Ma, D.; Chen, W.; Hu, G.; Zhang, Y.; Gao, Y.; Yin, Y.; Zhao, Y. $K_2S_2O_8$ -mediated metal-free direct P–H/C–H functionalization: a convenient route to benzo[b]phosphole oxides from unactivated alkynes. *Green Chem.* **2016**, *18*, 3522–3526. (g) Quint, V.; Morlet-Savary, F.; Lohier, J.-F.; Lalevée, J.; Gaumont, A.-C.; Lakhdar, S. Metal-

Free, Visible Light-Photocatalyzed Synthesis of Benzo[b]phosphole Oxides: Synthetic and Mechanistic Investigations. *J. Am. Chem. Soc.* **2016**, *138*, 7436–7441. (h) Kurimoto, Y.; Yamashita, J.; Mitsudo, K.; Sato, E.; Suga, S. Electrosynthesis of Phosphacycles via Dehydrogenative C–P Bond Formation Using DABCO as a Mediator. *Org. Lett.* **2021**, *23*, 3120–3124.

(4) (a) Nishimura, K.; Unoh, Y.; Hirano, K.; Miura, M. Phosphonium Cation Mediated Formal Cycloaddition Approach to Benzophospholes. *Chem.-Eur. J.* **2018**, *24*, 13089–13092. (b) Nishimura, K.; Hirano, K.; Miura, M. Synthesis of Dibenzophospholes by Tf_2O -Mediated Intramolecular Phospha-Friedel-Crafts-Type Reaction. *Org. Lett.* **2019**, *21*, 1467–1470. (c) Ye, W.; Li, X.; Ding, B.; Wang, C.; Shrestha, M.; Ma, X.; Chen, Y.; Tian, H. Facile Synthesis of Nitrogen-Containing Six-Membered Benzofused Phenophosphazine Oxides and Studies of the Photophysical Properties. *J. Org. Chem.* **2020**, *85*, 3879–3886. (d) Ishida, K.; Higashino, T.; Wada, Y.; Kaji, H.; Saeki, A.; Imahori, H. Thiophene-Fused Naphthodiphospholes: Modulation of the Structural and Electronic Properties of Polycyclic Aromatics by Precise Fusion of Heteroles. *ChemPlusChem* **2021**, *86*, 130–136.

(5) (a) Nishimura, K.; Hirano, K.; Miura, M. Direct Synthesis of Dibenzophospholes from Biaryls by Double C–P Bond Formation via Phosphonium Dication Equivalents. *Org. Lett.* **2020**, *22*, 3185–3189. (b) Li, C.; Huang, H.; Liu, F.; Yuan, C.; Chen, S.; Hua, Y.; Ding, H.; Song, X.-R.; Xiao, Q. Synthesis of (Thio)Furan-Fused Phospholes via Phosphonation Cyclization and a Base-Promoted Phospha-Friedel–Crafts Reaction. *J. Org. Chem.* **2022**, *87*, 2632–2639.

(6) Janesko, B. G.; Fisher, H. C.; Bridle, M. J.; Montchamp, J.-L. P(=O)H to P–OH Tautomerism: A Theoretical and Experimental Study. *J. Org. Chem.* **2015**, *80*, 10025–10032.

(7) (a) Breslow, R.; Deuring, L. Concerning phosphirenium cations. *Tetrahedron Lett.* **1984**, *25*, 1345–1348. (b) Marinetti, A.; Mathey, F. The chemistry of free and complexed phosphirenes: reactivity toward electrophiles, nucleophiles, and conjugated dienes. *J. Am. Chem. Soc.* **1985**, *107*, 4700–4706.

(8) (a) Olah, G. A.; Hehemann, D. Organometallic compounds. 14. Friedel–Crafts type preparation of triphenylphosphine. *J. Org. Chem.* **1977**, *42*, 2190. (b) Wang, Z.-W.; Wang, L.-S. Preparation of dichlorophenylphosphine via Friedel–Crafts reaction in ionic liquids. *Green Chem.* **2003**, *5*, 737–739. (c) Diaz, A. A.; Young, J. D.; Khan, M. A.; Wehmschulte, R. J. Facile Synthesis of Unsymmetrical 9-Phospha- and 9-Arsafluorenes. *Inorg. Chem.* **2006**, *45*, 5568–5575. (d) Diaz, A. A.; Buster, B.; Schomisch, D.; Khan, M. A.; Baum, J. C.; Wehmschulte, R. J. Size Matters: Room Temperature P–C Bond Formation Through C–H Activation in *m*-Terphenyldiiodophosphines. *Inorg. Chem.* **2008**, *47*, 2858–2863. (e) Hashimoto, S.; Nakatsuka, S.; Nakamura, M.; Hatakeyama, T. Construction of a Highly Distorted Benzene Ring in a Double Helicene. *Angew. Chem., Int. Ed.* **2014**, *53*, 14074–14076. (f) Hatakeyama, T.; Hashimoto, S.; Nakamura, M. Tandem Phospha-Friedel–Crafts Reaction toward Curved π -Conjugated Frameworks with a Phosphorus Ring Junction. *Org. Lett.* **2011**, *13*, 2130–2133. (g) Romero-Nieto, C.; López-Andarias, A.; Egler-Lucas, C.; Gebert, F.; Neus, J.-P.; Pilgram, O. Paving the Way to Novel Phosphorus-Based Architectures: A Non-catalyzed Protocol to Access Six-Membered Heterocycles. *Angew. Chem., Int. Ed.* **2015**, *54*, 15872–15875.

(9) A similar trend was observed in our previous work on the Tf_2O -mediated phosphole synthesis. See refs 4a, 4b, and 5a. The initially formed P(III) phosphole could be spontaneously oxidized with residual Tf_2O and/or its derivatives.

(10) For example, see: Skrabal-Joiner, S. L.; McLaughlin, E. C.; Ajaz, A.; Thamatam R.; Johnson, R. P. Scholl Cyclizations of Aryl Naphthalenes: Rearrangement Precedes Cyclization. *J. Org. Chem.* **2015**, *80*, 9578–9583.

(11) In the three component coupling reaction, the phosphole product was somewhat resistant to the in-situ oxidation, and a mixture of P(III) and P=O benzophospholes was detected after

the reaction. Thus, for ease of the purification, additional treatment with aq. H_2O_2 was performed for full conversion into the P=O benzophosphole.