

Title	Rhodium-Catalyzed C4-Selective C-H Alkenylation of 2-Pyridones by Traceless Directing Group Strategy
Author(s)	Hazra, Sunit; Hirano, Koji; Miura, Masahiro
Citation	Organic Letters. 2021, 23(4), p. 1388-1393
Version Type	AM
URL	https://hdl.handle.net/11094/92698
rights	© 2021 American Chemical Society.
Note	

Osaka University Knowledge Archive : OUKA

<https://ir.library.osaka-u.ac.jp/>


Osaka University

Rhodium-Catalyzed C4-Selective C–H Alkenylation of 2-Pyridones by Traceless Directing Group Strategy

Sunit Hazra, Koji Hirano,* and Masahiro Miura*

Department of Applied Chemistry, Graduate School of Engineering, Osaka University, Suita, Osaka 565-0871, Japan

Supporting Information Placeholder



ABSTRACT: A rhodium-catalyzed C4-selective C–H alkenylation of 3-carboxy-2-pyridones with styrenes has been developed. The carboxylic group at the C3 position works as the traceless directing group, and the corresponding C4-alkenylated 2-pyridones are obtained exclusively with concomitant decarboxylation. Unlike the reported procedures, the exclusive C4 selectivity is uniformly observed even in the presence of potentially more reactive C–H bonds at the C5 and C6 positions. By using this strategy, the multiply substituted 2-pyridone can be prepared via sequential C–H functionalization reactions.

A 2-pyridone that has the unique unsaturated system in the *N*-containing six-membered ring is one of the most widely occurring heterocyclic cores in natural products, biologically active molecules, and pharmaceutical agents.¹ Such ubiquity has promoted the development of protocols for the preparation of 2-pyridones, particularly, the multiply substituted ones in synthetic communities. Strategically, the substituted 2-pyridone can be obtained either by functionalization of the pyridone ring² or by constructing the ring from suitable acyclic precursors.^{1a} The former can provide a more convergent and modular approach to the target structure, but the reported methodologies largely relied on the stoichiometric halogenation and metalation.





Figure 1. Reactivity profile of C–H bonds on 2-pyridone in metal-mediated C–H activation.

In the last two decades, the metal-mediated C–H activation has been utilized for a wide range of transformations of C–H bonds to C–C or C–hetero atom bonds with better atom efficiency compared to the traditional cross-coupling methodologies.³ In this context, synthetic chemists have been greatly prompted to adopt the 2-pyr-


idone in the C–H activation.⁴ However, there are four possibly reactive C–H bonds on the 2-pyridone ring, and the control of site selectivity is thus the most important and challenging issue. While the C3-, C5-, and C6-selective C–H functionalizations have greatly progressed to date, the selective access to the C–H bond at the C4 position still remains largely elusive (Figure 1). To the best of our knowledge, only a few successful examples include the in-situ protection/lithiation strategy using CO₂ and BuLi/*t*-BuLi (Scheme 1a),⁵ Ni/Al-cooperative alkylation with alkenes (Scheme 1b),⁶ and sterically controlled Ir-catalyzed borylation with pinB–Bpin (Scheme 1c).⁷ The former is the stoichiometric reaction and suffers from the harsh conditions associated with the strongly basic organolithium reagents. The latter two cases are more attractive catalytic reactions, but the high C4-selectivity is obtained only when the competitively reactive C–H at the C6 position is blocked by substituents. Thus, there still remains a large demand for the C4 site selectivity that is independent of the substituent. Herein, we report a Cp*Rh(III)-catalyzed, carboxylic acid-directed highly C4-selective C–H alkenylation of 2-pyridones with styrenes (Scheme 1d): the key to success is the introduction of the carboxylic acid group at the C3 position, which works as the traceless directing group,⁸ and the corresponding C4-alkenylated 2-pyridones are obtained with the concomitant decarboxylation. The introduction of the carboxyl group at the C3 position sometimes requires some synthetic steps (see the Supporting Information for details), but unlike the aforementioned precedents, the exclusive C4 selectivity is uniformly observed even in the presence of potentially more

reactive C–H bond at the C6 position. Additionally, the installation of the vinyl group is possible, which is difficult to achieve by other C4-selective C–H functionalization protocols and thus complementary and useful from the synthetic point of view.

Scheme 1. C4-Selective C–H Functionalizations on 2-Pyridones




Scheme 2. Working Hypothesis of Carboxylic Acid-Directed C4-Selective C–H Alkenylation of 2-Pyridone **1a** with Styrene **2a** with Concomitant Decarboxylation



The blue print for the catalytic C4-selective C–H functionalization of 2-pyridones is based on the recent progress of carboxylic acid directed formal *meta*-selective C–H functionalization of benzenes, which was developed by our group,⁹ Larrosa,¹⁰ and others.¹¹ Our working hypothesis is shown in Scheme 2. The initial ligand exchange between the catalyst **A** and 3-carboxy-2-pyridone **1a** is followed by the directed C–H cleavage to form the corre-

sponding metallacycle **C**. Subsequent insertion of styrene **2a** into the pyridone C(4)–M bond and β -H elimination afford the intermediate **E**. The C4-alkenylated product **3aa** then follows from decarboxylation and reductive elimination. The catalytic cycle is closed by the reoxidation of reduced **G** with the external oxidant.

Table 1. Optimization Studies for Decarboxylative C4-Selective C–H Alkenylation of **1a** with **2a**^a



entry	catalyst	oxidant	additives	yield (%) ^b
1	[Cp*RhCl ₂] ₂	Cu(OAc) ₂ •H ₂ O	K ₂ HPO ₄	43
2	[Cp*IrCl ₂] ₂	Cu(OAc) ₂ •H ₂ O	K ₂ HPO ₄	0
3	Cp*Co(CO) ₂	Cu(OAc) ₂ •H ₂ O	K ₂ HPO ₄	0
4	[RuCl ₂ (<i>p</i> -cymene)] ₂	Cu(OAc) ₂ •H ₂ O	K ₂ HPO ₄	0
5	Pd(OAc) ₂	Cu(OAc) ₂ •H ₂ O	K ₂ HPO ₄	0
6	[Cp*RhCl ₂] ₂	AgOAc	K ₂ HPO ₄	11
7	[Cp*RhCl ₂] ₂	Ag ₂ CO ₃	K ₂ HPO ₄	17
8 ^c	[Cp*RhCl ₂] ₂	Cu(OAc) ₂ •H ₂ O	K ₂ HPO ₄ MS 4A	60
9 ^c	[Cp*RhCl ₂] ₂	Cu(OAc) ₂ •H ₂ O	Na ₂ CO ₃ MS 4A	26
10 ^c	[Cp*RhCl ₂] ₂	Cu(OAc) ₂ •H ₂ O	K ₂ CO ₃ MS 4A	11
11 ^c	[Cp*RhCl ₂] ₂	Cu(OAc) ₂ •H ₂ O	CS ₂ CO ₃ MS 4A	0
12 ^c	[Cp*RhCl ₂] ₂	Cu(OAc) ₂ •H ₂ O	NaHCO ₃ MS 4A	23
13 ^c	[Cp*RhCl ₂] ₂	Cu(OAc) ₂ •H ₂ O	KHCO ₃ MS 4A	25
14 ^c	[Cp*RhCl ₂] ₂	Cu(OAc) ₂ •H ₂ O	K ₄ P ₂ O ₇ MS 4A	82 (78)


^a Conditions: **1a** (0.20 mmol), **2a** (0.40 mmol), catalyst (0.010 mmol on metal), oxidant (0.20 mmol), additives (0.40 mmol), toluene (1.0 mL), 160 °C, 16–24 h, N₂. ^b Estimated by ¹H NMR with CH₂Br₂ as the internal standard. Isolated yield in parentheses. ^c With 0.60 mmol of **2a** and 100 mg of MS 4A.

On the basis of the above assumption, we started the optimization studies with **1a** (0.20 mmol) and **2a** (0.40 mmol; Table 1). After the initial screening of catalyst, oxidant, additive, and solvent, we were pleased to find that the reaction proceeded in the presence of [Cp*RhCl₂]₂ catalyst, Cu(OAc)₂•H₂O oxidant, and K₂HPO₄ base in heated toluene (160 °C) to deliver the targeted **3aa** in 43% ¹H NMR yield (entry 1). Notably, the C–C bond formation occurred exclusively at the C4 position, and the COOH-remaining byproduct **3aa-COOH** was not detected at all. Additionally, the structure of **3aa** was unambiguously confirmed by X-ray analysis (CCDC 2032339). No alkenylated product **3aa** was observed with other catalysts including [Cp*IrCl₂]₂, Cp*Co(CO)₂, [RuCl₂(*p*-cymene)]₂, and Pd(OAc)₂ (entries 2–5), some of which are known to promote the related decarboxylative C–H functionalization of benzene derivatives.^{8–11} The Cu(OAc)₂•H₂O was


found to be better than the Ag-based oxidants (entries 6 and 7). The ^1H NMR yield further increased to 60% by using 3.0 equivalent of **2a** and addition of MS 4A (entry 8). Final investigation of base (entries 9–14) revealed that $\text{K}_4\text{P}_2\text{O}_7$ showed the best performance,¹² and **3aa** was finally isolated in 78% yield (82% ^1H MR yield; entry 14). Some additional observations are to be noted: the carboxylic acid group was indispensable for the alkenylation, and no conversion of the corresponding ester and simple 2-pyridone substrate was observed. Other ethereal, halogenated, and polar solvents were also tested, but toluene proved to be best.¹³

With the optimal conditions in hand (Table 1, entry 14), we examined the scope and limitation of this strategy (Scheme 3). The larger alkyl groups on the pyridone nitrogen were compatible to form the corresponding C4-alkenylated 2-pyridones **3ba–da** in good yields. The substituents at the C6 position were also accommodated (**3ea–ha**). Particularly notable is the successful reaction of 2-pyridones that bear the amino (**3ia**) and alkoxy (**3ja**) functionalities. On the other hand, the C5-substituted pyridone was the reluctant substrate probably due to the steric factors (**3ka**). The reaction was scalable, and **3aa** was obtained in 69% on a 1.0 mmol scale.


Scheme 3. Rhodium-Catalyzed C4-Selective C–H Alkenylation of 3-Carboxy-2-pyridones **1** with Styrenes **2** by Traceless Directing Group Strategy^a



• scope of 2-pyridones **1**



• scope of styrenes **2**




^a Reaction conditions: **1** (0.20 mmol), **2** (0.60 mmol), $[\text{Cp}^*\text{RhCl}_2]_2$ (0.0050 mmol), $\text{Cu}(\text{OAc})_2 \cdot \text{H}_2\text{O}$ (0.20 mmol), $\text{K}_4\text{P}_2\text{O}_7$ (0.40 mmol), MS 4A (100 mg), toluene (1.0 mL), 160°C , 24 h, N_2 . Isolated yields are shown. ^b On a 1.0 mmol scale. ^c NMR yield.

Several *para*-substituted styrenes were smoothly coupled with **1a**: electron-donating and -withdrawing as well as halogenated substituents all were tolerated under the reaction conditions (**3ab–3ah**).¹⁴ The *meta*-substituted substrates were also viable substrates (**3ai** and **3aj**), but the sterically demanding *ortho*-substitution was somewhat detrimental to the reaction (**3ak**). Unfortunately, the acrylate showed lower reactivity than the styrene derivatives (**3al**), but the reactivity trend was highly dependent on the position of carboxy group (*vide infra*).


The obtained product **3da** underwent some derivatizations (Scheme 4). The vinylenes selective reduction was possible under the hydrogenation conditions using $\text{Pd}(\text{OH})_2/\text{H}_2$ in AcOH solvent to furnish the C4-alkylated 2-pyridone **4** in 81% yield. On the other hand, the saturated piperidin-2-one structure **5** was selectively obtained under the hydrogenation in EtOH. Additionally, the 2-pyridone has an electron-rich vinylogous enamide character, and **3da** thus underwent the Diels-Alder reaction with the triazolidione. After additional treatment with PPTS,¹⁵ the corresponding tricyclic system **6** was isolated in 61% overall yield.

Scheme 4. Derivatizations of **3da**




We next implemented some control experiments to gain mechanistic insight (Scheme 5). First, to check the role of $\text{Cu}(\text{OAc})_2$, the stoichiometric reaction to Rh was performed in the absence of $\text{Cu}(\text{OAc})_2$: the targeted **3aa** was formed in 70% yield without any difficulties. Thus, $\text{Cu}(\text{OAc})_2$ can work as just an oxidant in the catalyst regeneration step (**G** to **A** in Scheme 2). Additionally, the reaction with styrene- d_8 (**2a-d₈**) successfully incorporated a partial but significant amount of deuterium selectively at the C3 position.¹⁶ These outcomes suggest that as shown in Scheme 2 the decarboxylative C–C bond forming process is operated by the Rh alone rather than the stepwise C–C bond formation and Cu-promoted protodecarboxylation, which is consistent with no observation of COOH-remaining **3aa-COOH** (Table 1) and the lower yield of **3aa** in the presence of Ag-based oxidants (entries 6 and 7 in Table 1) that more strongly accelerate the nonproductive simple protodecarboxylation.¹⁷ A similar reaction mechanism was proposed in the related Ru-catalyzed decarboxylative C–H alkenylation with alkynes.^{11a–c}

Scheme 5. Control Experiments



We also investigated the applicability and site selectivity of this strategy in the reaction of regioisomeric carboxy-2-pyridones (Scheme 6). As a general trend, K_2HPO_4 instead of $K_4P_2O_7$ showed better performance in these cases. The 4-carboxy-2-pyridone **7a** afforded the C3-alkenylated product **8aa** exclusively, the site selectivity of which can be controlled by the additional coordination of the pyridone carbonyl.¹⁸ Additionally notable is the successful coupling of acrylate in this case (**8al**). The 5-carboxy-2-pyridone **9a** specifically reacted only with the acrylate **2l** to form the C4- and C6-doubly alkenylated **10al** as the major product. A mixture of the corresponding five-membered lactones¹⁹ **11al** and **12al** was also isolated, thus suggesting the competitive reaction order at the C3 and C6 positions. In the reaction of the 6-carboxy-2-pyridone **13a**, as in the case of the 3-carboxy-2-pyridone **1a**, only the styrene **2a** was reactive, and the expected C5-alkenylated product **14aa** was obtained in 60% yield.

Scheme 6. Attempts To Apply Regioisomeric Carboxy-2-pyridones



Finally, we attempted to synthesize the multiply substituted 2-pyridone by the sequential site-selective C–H functionalization reactions (Scheme 7). The starting platform is the *N*-(2-pyridyl)-3-methoxycarbonyl-2-pyridone

(**15**), and the rhodium-catalyzed, pyridine-directed C6-selective C–H arylation with the aryltrifluoroborate²⁰ was first conducted to afford **16** in 78% yield. The subsequent ester hydrolysis and *N*-substituent switch from Py to Me formed **1h** in 54% (4 steps). The styryl group was introduced at the C4 position by the present traceless directing group strategy (**3ha**). The C–H at the C3 position was then selectively arylated with the hydrazine derivative under radical conditions²¹ to form the trisubstituted 2-pyridone **17** with two different aryl groups and one alkenyl group at the C3, C6, and C4 positions. As demonstrated in Scheme 7, the combination of the presently developed protocol and literature procedures can provide a modular C–H functionalization approach to the multiply substituted 2-pyridones, which will find wide applications in the development of more complex, pyridone-based bioactive molecules.¹

In conclusion, we have developed a rhodium-catalyzed decarboxylative C4-selective C–H alkenylation of 3-carboxy-2-pyridones with styrenes. Taking advantage of the carboxy group as the traceless directing group, we can access the otherwise challenging C–H bond at the C4 position, irrespective of the substituents at the competitively reactive C6 position. Additionally, by using the newly developed strategy, the multiply substituted 2-pyridone can be prepared via sequential C–H functionalization reactions.

ASSOCIATED CONTENT

Supporting Information

The Supporting Information is available free of charge on the ACS Publications website at DOI: 10.1021/acs.orglett.xxxx.

¹H, ¹³C{¹H}, and ¹⁹F{¹H} NMR spectra, ORTEP drawing, detailed optimization studies, detailed D-labeling experiments, and protodecarboxylation test (PDF)

Accession Codes

CCDC 2032339 contains the supplementary crystallographic data for this paper. These data can be obtained free of charge via www.ccdc.cam.ac.uk/data_request/cif, or by emailing data_request@ccdc.cam.ac.uk, or by contacting The Cambridge Crystallographic Data Centre, 12 Union Road, Cambridge CB2 1EZ, UK; fax: +44 1223 336033.

AUTHOR INFORMATION

Corresponding Authors

Koji Hirano – Department of Applied Chemistry, Graduate School of Engineering, Osaka University, Suita, Osaka 565-0871, Japan; orcid.org/0000-0001-9752-1985; Email: k_hirano@chem.eng.osaka-u.ac.jp.


Masahiro Miura – Department of Applied Chemistry, Graduate School of Engineering, Osaka University, Suita, Osaka 565-0871, Japan; orcid.org/0000-0001-8288-6439; Email: miura@chem.eng.osaka-u.ac.jp.

Author

Sunit Hazra – Department of Applied Chemistry, Graduate School of Engineering, Osaka University, Suita, Osaka 565-0871, Japan

Complete contact information is available at: <https://pubs.acs.org/10.1021/xxxx>.

Scheme 7. Synthesis of Multiply Substituted 2-Pyridone 17 by Sequential Site-Selective C–H Functionalization Reactions



Notes

The authors declare no competing financial interest.

ACKNOWLEDGMENT

This work was supported by JSPS KAKENHI Grant Nos. JP 18K19078 (Grant-in-Aid for Challenging Research (Exploratory)) to K.H. and JP 17H06092 (Grant-in-Aid for Specially Promoted Research) to M.M. We thank Dr. Yuji Nishii (Osaka University) for his assistance with the X-ray analysis.

REFERENCES

(1) (a) Torres, M.; Gil, S.; Parra, M. New Synthetic Methods to 2-Pyridone Rings. *Curr. Org. Chem.* **2005**, *9*, 1757-1779. (b) Hibi, S.; Ueno, K.; Nagato, S.; Kawano, K.; Ito, K.; Norimine, Y.; Takenaka, O.; Hanada, T.; Yonaga, M. Discovery of 2-(2-Oxo-1-phenyl-5-pyridin-2-yl)-1,2-dihydropyridin-3-yl)benzotriazole (Perampanel): A Novel, Noncompetitive α -Amino-3-hydroxy-5-methyl-4-isoxazolepropanoic Acid (AMPA) Receptor Antagonist. *J. Med. Chem.* **2012**, *55*, 10584-10600. (c) Hajek, P.; McRobbie, H.; Myers, K. Efficacy of Cytisine in Helping Smokers Quit: Systematic Review and meta-Analysis *Thorax* **2013**, *68*, 1037-1042. (d) Campello, H. R.; Del Villar, S. G.; Honraedt, A.; Minguez, T.; Oliveira, A. S. F.; Ranaghan, K. E.; Shoemark, D. K.; Bermudez, L.; Gotti, C.; Sessions, R. B.; Mulholland, A. J.; Wonnacott, S.; Gallagher, T. Unlocking Nicotinic Selectivity via Direct C–H Functionalization of (-)-Cytisine. *Chem.* **2018**, *4*, 1710-1725. (e) Fioravanti, R.; Stazi, G.; Zwergel, C.; Valente, S.; Mai, A. Six Years (2012–2018) of Researches on Catalytic EZH2 Inhibitors: The Boom of the 2-Pyridone Compounds. *Chem. Rec.* **2018**, *18*, 1818-1832.

(2) (a) Hasvold, L. A.; Wang, W.; Gwaltney, S. L.; Rockway, T. W.; Nelson, L. T. G.; Mantei, R. A.; Fakhoury, S. A.; Sullivan, G. M.; Li, Q.; Lin, N.-H.; Wang, L.; Zhang, H.; Cohen, J.; Gu, W.-Z.; Marsh, K.; Bauch, J.; Rosenberg, S.; Sham, H. L. Pyridone-Containing farnesyltransferase inhibitors: synthesis and biological evaluation. *Bioorg. Med. Chem. Lett.* **2003**, *13*, 4001-4005. (b) Bengtsson, C.; Almqvist, F. Regioselective Halogenations and Subsequent Suzuki–Miyaura Coupling onto Bicyclic 2-Pyridones. *J. Org. Chem.* **2010**, *75*, 972-975. See also: (c) Hill, M. D.; Movassaghi, M. New Strategies for the Synthesis of Pyrimidine Derivatives. *Chem.–Eur. J.* **2008**, *14*, 6836-6844.

(3) For selected reviews, see: (a) Kakiuchi, F.; Kochi, T. Transition-Metal-Catalyzed Carbon–Carbon Bond Formation via Carbon–Hydrogen Bond Cleavage. *Synthesis* **2008**, *2008*, 3013-3039. (b) Ackermann, L.; Vicente, R.; Kapdi, A. R. Transition-Metal-Catalyzed Direct Arylation of (Hetero)Arenes by C–H Bond Cleavage. *Angew. Chem., Int. Ed.* **2009**, *48*, 9792-9826. (c) Lyons, T. W.; Sanford, M. S. Palladium-Catalyzed Ligand-Directed C–H Functionalization Reactions. *Chem. Rev.* **2010**, *110*, 1147-1169. (d) Satoh, T.; Miura, M. Oxidative Coupling of Aromatic

Substrates with Alkynes and Alkenes under Rhodium Catalysis. *Chem. Eur. J.* **2010**, *16*, 11212-11222. (e) Liu, C.; Zhang, H.; Shi, W.; Lei, A. Bond Formations between Two Nucleophiles: Transition Metal Catalyzed Oxidative Cross-Coupling Reactions. *Chem. Rev.* **2011**, *111*, 1780-1824. (f) Yamaguchi, J.; Yamaguchi, A. D.; Itami, K. C–H Bond Functionalization: Emerging Synthetic Tools for Natural Products and Pharmaceuticals. *Angew. Chem., Int. Ed.* **2012**, *51*, 8960-9009. (g) Hirano, K.; Miura, M. Recent Advances in Copper-mediated Direct Biaryl Coupling. *Chem. Lett.* **2015**, *44*, 868-873. (h) Boyarskiy, V. P.; Ryabukhin, D. S.; Bokach, N. A.; Vasilyev, A. V. Alkenylation of Arenes and Heteroarenes with Alkynes. *Chem. Rev.* **2016**, *116*, 5894-5986. (i) Wang, F.; Yu, S.; Li, X. Transition Metal-Catalysed Couplings Between Arenes and Strained or Reactive Rings: Combination of C–H Activation and Ring Scission. *Chem. Soc. Rev.* **2016**, *45*, 6462-6477. (j) Zhu, R.-Y.; Farmer, M. E.; Chen, Y.-Q.; Yu, J.-Q. A Simple and Versatile Amide Directing Group for C–H Functionalizations. *Angew. Chem., Int. Ed.* **2016**, *55*, 10578-10599. (k) Gulías, M.; Mascareñas, J. L. Metal-Catalyzed Annulations through Activation and Cleavage of C–H Bonds. *Angew. Chem., Int. Ed.* **2016**, *55*, 11000-11019. (l) Ping, L.; Chung, D. S.; Bouffard, J.; Lee, S. Transition Metal-Catalyzed Site- and Regio-divergent C–H Bond Functionalization. *Chem. Soc. Rev.* **2017**, *46*, 4299-4328. (m) Mihai, M. T.; Genov, G. R.; Phipps, R. J. Access to the *meta* Position of Arenes Through Transition Metal Catalysed C–H Bond Functionalisation: A Focus on Metals Other than Palladium. *Chem. Soc. Rev.* **2018**, *47*, 149-171. (n) Chu, J. C. K.; Rovis, T. Complementary Strategies for Directed C(sp³)-H Functionalization: A Comparison of Transition-Metal-Catalyzed Activation, Hydrogen Atom Transfer, and Carbene/Nitrene Transfer. *Angew. Chem., Int. Ed.* **2018**, *57*, 62-101. (o) Sambigiato, C.; Schönbauer, D.; Blicke, R.; Dao-Huy, T.; Pototschnig, G.; Schaaf, P.; Wiesinger, T.; Zia, M. F.; Wencel-Delord, J.; Besset, T.; Maes, B. U. W.; Schnürch, M. A Comprehensive Overview of Directing Groups Applied in Metal-catalysed C–H Functionalisation Chemistry. *Chem. Soc. Rev.* **2018**, *47*, 6603-6743. (p) Rej, S.; Ano, Y.; Chatani, N. Bidentate Directing Groups: An Efficient Tool in C–H Bond Functionalization Chemistry for the Expedient Construction of C–C Bonds. *Chem. Rev.* **2020**, *120*, 1788-1887.

(4) Reviews: (a) Hirano, K.; Miura, M. A Lesson for Site-Selective C–H Functionalization on 2-Pyridones: Radical, Organometallic, Directing Group and Steric Controls. *Chem. Sci.* **2018**, *9*, 22-32. (b) Prendergast, A. M.; McGlacken, G. P. Transition Metal Mediated C–H Activation of 2-Pyrones, 2-Pyridones, 2-Coumarins and 2-Quinolones. *Eur. J. Org. Chem.* **2018**, *2018*, 6068-6082. (c) Kumar, S. V.; Banerjee, S.; Punniyamurthy, T. Transition Metal-catalyzed Coupling of Heterocyclic Alkenes via C–H Functionalization: Recent Trends and Applications. *Org. Chem. Front.*, **2020**, *7*, 1527-1569. (d) Biswas, A.; Maiti, S.; Pan, S.; Samanta, R. Transition Metal-Catalysed Direct C–H Bond

Functionalizations of 2-Pyridone Beyond C3-Selectivity. *Chem.–Asian J.* **2020**, *15*, 2092–2109.

(5) Katritzky, A. R.; Fan, W.-Q.; Koziol, A. E.; Palenik, G. J. Carbon Dioxide: A Reagent for the Simultaneous Protection of Nucleophilic Centres and the Activation of Alternative Locations to Electrophilic Attack: Part 8. A Novel Synthetic Route to 4-Substituted-2-Pyridones. *Tetrahedron*, **1987**, *43*, 2343-2348.

(6) Tamura, R.; Yamada, Y.; Nakao, Y.; Hiyama, T. Alkylation of Pyridone Derivatives by Nickel/Lewis Acid Catalysis. *Angew. Chem., Int. Ed.* **2012**, *51*, 5679-5782.

(7) Miura, W.; Hirano, K.; Miura, M. Iridium-Catalyzed Site-Selective C–H Borylation of 2-Pyridones. *Synthesis* **2017**, *49*, 4745-4752.

(8) (a) Font, M.; Quibell, J. M.; Perry, G. J.; Larrosa, I. The use of carboxylic acids as traceless directing groups for regioselective C–H bond functionalization. *Chem. Commun.* **2017**, *53*, 5584-5597. (b) Paul, K. D.; Luxami, V.; Rani, G. Traceless Directing Group: A Novel Strategy in Regiodivergent C-H Functionalization. *Chem. Commun.* **2020**, *56*, 12479-12521.

(9) Mochida, S.; Hirano, K.; Satoh, T.; Miura, M. Synthesis of Stilbene and Distyrylbenzene Derivatives through Rhodium-Catalyzed Ortho-Olefination and Decarboxylation of Benzoic Acids. *Org. Lett.* **2010**, *12*, 5776-5779.

(10) (a) Cornella, J.; Righi, M.; Larrosa, I. Carboxylic Acids as Traceless Directing Groups for Formal *meta*-Selective Direct Arylation. *Angew. Chem., Int. Ed.* **2011**, *50*, 9429-9432. (b) Luo, J.; Preciado, S.; Larrosa, I. Overriding Ortho–Para Selectivity via a Traceless Directing Group Relay Strategy: The *Meta*-Selective Arylation of Phenols. *J. Am. Chem. Soc.* **2014**, *136*, 4109-4112.

(11) (a) Zhang, J.; Shrestha, R.; Hartwig, J. F.; Zhao, P. A decarboxylative approach for regioselective hydroarylation of alkynes. *Nat. Chem.* **2016**, *8*, 1144-1151. (b) Kumar, N. Y. P.; Bechtoldt, A.; Raghuvanshi, K.; Ackermann, L. Ruthenium(II)-Catalyzed Decarboxylative C–H Activation: Versatile Routes to *meta*-Alkenylated Arenes. *Angew. Chem., Int. Ed.* **2016**, *55*, 6929-6932. (c) Huang, L.; Biafora, A.; Zhang, G.; Bragoni, V.; Gooßen, L. J. Regioselective C–H Hydroarylation of Internal Alkynes with Arenecarboxylates: Carboxylates as Deciduous Directing Groups. *Angew. Chem., Int. Ed.* **2016**, *55*, 6933-6937. (d) Biafora, A.; Krause, T.; Hackenberger, D.; Belitz, F.; Gooßen, L. J. *ortho*-C–H Arylation of Benzoic Acids with Aryl Bromides and Chlorides Catalyzed by Ruthenium. *Angew. Chem., Int. Ed.* **2016**, *55*, 14752-14755. (e) Zhang, Y.; Zhao, H.; Zhang, M.; Su, W. *ortho*-C–H Arylation of Benzoic Acids with Aryl Bromides and Chlorides Catalyzed by Ruthenium. *Angew. Chem., Int. Ed.* **2015**, *54*, 3817-3821. (f) Qin, X.; Sun, D.; You, Q.; Cheng, Y.; Lan, J.; You, J. Rh(III)-Catalyzed Decarboxylative *ortho*-Heteroarylation of Aromatic Carboxylic Acids by Using the Carboxylic Acid as a Traceless Directing Group. *Org. Lett.* **2015**, *17*, 1762-1765.

(12) The reason for the unique effect of $K_4P_2O_7$ additive is unclear at this stage, but its suitable basicity (the pK_a value of the

conjugate acid $HP_2O_7^{3-}$ is 9.25) can play an important role in the ligand exchange step on Rh (Scheme 2, **A** to **B**). In addition, it may suppress the nonproductive protodecarboxylation. For an example of the unique effect of $K_4P_2O_7$ base, see: He, Z.; Song, F.; Sun, H.; Huang, Y. Transition-Metal-Free Suzuki-Type Cross-Coupling Reaction of Benzyl Halides and Boronic Acids via 1,2-Metalate Shift. *J. Am. Chem. Soc.* **2018**, *140*, 2693-2699.

(13) See the Supporting Information for more detailed optimization studies.

(14) We have no explanation for the reason why the yields of **3ab**, **3ad**, and **3af** were lower at this stage. We repeated these reaction two times, but almost the same yields were observed.

(15) The initially formed Diels-Alder adduct was somewhat unstable and gradually underwent the olefin isomerization into **6**, which was accelerated by addition of acid such as PPTS.

(16) We confirmed that the deuterium content at the C3 position did not increase in the reaction even with anhydrous $Cu(OAc)_2$ and toluene- d_8 . Additionally, the isolated **3aa** underwent just a partial H/D exchange (9%) at the C3 position with $AcOH-d_4$ under otherwise identical conditions (see the Supporting Information for details). Thus, we believe that the lower deuterium content is mainly attributed to the H/D exchange on Rh-H species (**E** and/or **F** in Scheme 2) with the carboxylic acid H of the starting pyridone.

(17) Ag: (a) Lu, P.; Sanchez, C.; Cornella, J.; Larrosa, I. Silver-Catalyzed Protodecarboxylation of Heteroaromatic Carboxylic Acids. *Org. Lett.* **2009**, *11*, 5710-5713. Cu: (b) Gooßen, L. J.; Thiel, W. R.; Rodríguez, N.; Linder, C.; Melzer, B. Copper-Catalyzed Protodecarboxylation of Aromatic Carboxylic Acids. *Adv. Synth. Catal.* **2007**, *349*, 2241-2246. We also compared the activity of $Cu(OAc)_2 \cdot H_2O$, $AgOAc$, and Ag_2CO_3 in the protodecarboxylation of **1d** and confirmed no protodecarboxylation only with $Cu(OAc)_2 \cdot H_2O$. See the Supporting Information for details.

(18) Chen, Y.; Wang, F.; Jia A.; Li, X. Palladium-catalyzed selective oxidative olefination and arylation of 2-pyridones. *Chem. Sci.* **2012**, *3*, 3231-3236.

(19) The formation of similar lactones is trivial in the literature. For example, see: Ueura, K.; Satoh, T.; Miura, M. An Efficient Waste-Free Oxidative Coupling via Regioselective C–H Bond Cleavage: Rh/Cu-Catalyzed Reaction of Benzoic Acids with Alkynes and Acrylates under Air. *Org. Lett.* **2007**, *9*, 1407-1409.

(20) Peng, P.; Wang, J.; Jiang, H.; Liu, H. Rhodium(III)-Catalyzed Site-Selective C–H Alkylation and Arylation of Pyridones Using Organoboron Reagents. *Org. Lett.* **2016**, *18*, 5376-5379.

(21) Chauhan, P.; Ravi, M.; Singh, S.; Prajapati, P.; Yadav, P. P. Regioselective α -arylation of coumarins and 2-pyridones with phenylhydrazines under transition-metal-free conditions. *RSC Adv.* **2016**, *6*, 109-118.