

Title	Sobolev's inequality for Riesz potentials of functions in non-doubling Morrey spaces
Author(s)	Mizuta, Yoshihiro; Shimomura, Tetsu; Sobukawa, Takuya
Citation	Osaka Journal of Mathematics. 2009, 46(1), p. 255-271
Version Type	VoR
URL	<a href="https://doi.org/10.18910/9294">https://doi.org/10.18910/9294</a>
rights	
Note	

***Osaka University Knowledge Archive : OUKA***

<https://ir.library.osaka-u.ac.jp/>

Osaka University

# SOBOLEV'S INEQUALITY FOR RIESZ POTENTIALS OF FUNCTIONS IN NON-DOUBLING MORREY SPACES

YOSHIHIRO MIZUTA, TETSU SHIMOMURA and TAKUYA SOBUKAWA

(Received July 6, 2007, revised December 20, 2007)

## Abstract

Our aim in this paper is to give Sobolev's inequality and Trudinger exponential integrability for Riesz potentials of functions in non-doubling Morrey spaces.

## 1. Introduction

The space introduced by Morrey [13] in 1938 has become a useful tool of the study for the existence and regularity of solutions of partial differential equations. In the present paper, we aim to establish Sobolev's inequality for the Riesz potentials of functions in generalized Morrey spaces in the non-doubling setting, as extensions of Gogatishvili-Koskela [4], Orobitg-Pérez [14] and Sawano-Sobukawa-Tanaka [19].

Let  $X$  be a separable metric space with a nonnegative Radon measure  $\mu$ . For simplicity, write  $|x - y|$  for the distance of  $x$  and  $y$ . We assume that  $\mu(\{x\}) = 0$  and  $0 < \mu(B(x, r)) < \infty$  for  $x \in X$  and  $r > 0$ , where  $B(x, r)$  denotes the open ball centered at  $x$  of radius  $r > 0$ . In this paper,  $\mu$  may or may not be doubling.

Let  $G$  be an open set in  $X$ . We define the Riesz potential of order  $\alpha$  for a non-negative measurable function  $f$  on  $G$  by

$$U_\alpha f(x) = \int_G \frac{|x - y|^\alpha f(y)}{\mu(B(x, 4|x - y|))} d\mu(y).$$

Here we introduce the family  $L^{p, \nu; k}(G)$  of all measurable functions  $f$  on  $G$  such that

$$\|f\|_{p, \nu, G; k}^p = \sup_{x \in G, 0 < r \leq d_G} \frac{r^\nu}{\mu(B(x, kr))} \int_{G \cap B(x, r)} |f(y)|^p d\mu(y) < \infty,$$

where  $1 < p < \infty$ ,  $\nu > 0$ ,  $k > 1$  and  $d_G$  denotes the diameter of  $G$ . In case  $X = \mathbf{R}^n$  with a nonnegative Radon measure  $\mu$ , we know that

$$L^{p, \nu; k_1}(G) = L^{p, \nu; k_2}(G)$$

when  $k_2 > k_1 > 1$ , but we have an example (see Remark 2.1) in which

$$L^{p,v;1}(G) \neq L^{p,v;2}(G);$$

see also Sawano-Tanaka [20]. The space  $L^{p,v;2}(G)$  is referred to as a generalized Morrey space.

To obtain Sobolev type inequalities for Riesz potentials of functions belonging to generalized Morrey spaces, we consider a generalized maximal function defined by

$$M_k f(x) = \sup_{r>0} \frac{1}{\mu(B(x, kr))} \int_{G \cap B(x,r)} |f(y)| d\mu(y)$$

for  $k \geq 1$  and a locally integrable function  $f$  on  $G$ . In view of Sawano [18, Corollary 2.1]),  $M_2$  is bounded in  $L^p(X)$ . Further it is useful to remark that in a certain metric measure space  $X$ ,  $M_k$  fails to be bounded in  $L^p(X)$  if and only if  $k < 2$  (see Sawano [18, Proposition 1.1]).

By applying the fact that  $M_2$  is a bounded mapping from  $L^{p,v;2}(X)$  to  $L^{p,v;4}(X)$  (see Sawano-Tanaka [20, Theorem 2.3]), we first show that  $U_\alpha f \in L^{p^\sharp, v;4}(X)$  for  $f \in L^{p,v;2}(X)$ , where  $1/p^\sharp = 1/p - \alpha/v > 0$ ; in the borderline case  $v = \alpha p$ , we consider the exponential integrability. For this, we also refer the reader to Sawano-Sobukawa-Tanaka [19, Theorem 3.1].

Finally, in our Morrey space setting, we establish an exponential integrability for functions satisfying a Poincaré inequality, as an extension of Gogatishvili-Koskela [4] and Orobitg-Pérez [14].

For related results, see Adams [1], Chiarenza-Frasca [3, Theorem 2], Nakai [15, Theorem 2.2] and the authors [11, 12] in the doubling case.

## 2. Sobolev's inequality

Throughout this paper, let  $C$  denote various constants independent of the variables in question.

For a nonnegative measurable function  $f$  on  $G$  and  $k > 1$ , define the maximal function

$$\begin{aligned} M_k f(x) &= \sup_{r>0} \frac{1}{\mu(B(x, kr))} \int_{G \cap B(x,r)} |f(y)| d\mu(y) \\ &= \sup_{0 < r < d_G} \frac{1}{\mu(B(x, kr))} \int_{G \cap B(x,r)} |f(y)| d\mu(y) \end{aligned}$$

for  $x \in G$ , where  $d_G$  denotes the diameter of  $G$ . Recall that

$$\|f\|_{p,v,G;k}^p = \sup_{x \in G, 0 < r \leq d_G} \frac{r^v}{\mu(B(x, kr))} \int_{G \cap B(x,r)} |f(y)|^p d\mu(y).$$

When  $X = \mathbf{R}^n$  with a nonnegative Radon measure  $\mu$ , we see that if  $G$  is an open set of  $\mathbf{R}^n$  and  $1 < k < 2$ , then

$$\|f\|_{p,v,G;k} \leq C \|f\|_{p,v,G;2}$$

for all  $f \in L^{p,v;2}(G)$ , where  $C$  is a constant depending only on  $k$  and  $n$ ; for this fact, see e.g. [20, Proposition 1.1].

REMARK 2.1. We set

$$L^{p,v;k}(G) = \left\{ f \mid \sup_{x \in G, 0 < r \leq d_G} \frac{r^v}{\mu(B(x, kr))} \int_{G \cap B(x,r)} |f(y)|^p d\mu(y) < \infty \right\}.$$

When  $G \subset \mathbf{R}^n$ ,  $L^{p,v;k}(G) = L^{p,v;2}(G)$  for all  $k > 1$ . We show by an example that

$$L^{p,v;1}(G) \neq L^{p,v;k}(G)$$

when  $k > 1$ . For this, consider a measure given by

$$d\mu(y) = e^y dy$$

on  $\mathbf{R}^1$ . For  $0 < \beta < 1$ , letting  $f(y) = y^{-\beta/p}$  for  $y > 0$  and  $f(y) = 0$  for  $y \leq 0$ , we note the following:

(i) if  $0 < x \leq r$ , then

$$\begin{aligned} \frac{r^v}{\mu(B(x, 2r))} \int_{B(x,r)} |f(y)|^p d\mu(y) &\leq \frac{r^v}{e^x(e^{2r} - 1)} e^{x+r} \int_0^{2r} y^{-\beta} dy \\ &\leq C \frac{r^{v-\beta+1}}{e^r - 1}; \end{aligned}$$

(ii) if  $x > r > 0$ , then

$$\begin{aligned} \frac{r^v}{\mu(B(x, 2r))} \int_{B(x,r)} |f(y)|^p d\mu(y) &\leq \frac{r^v}{e^x(e^{2r} - 1)} \int_{x-r}^{x+r} y^{-\beta} e^y dy \\ &\leq \frac{r^v}{e^x(e^{2r} - 1)} \frac{(x+r)^{1-\beta} - (x-r)^{1-\beta}}{1-\beta} e^{x+r} \\ &\leq C \frac{r^{v-\beta+1}}{e^r - 1}; \end{aligned}$$

(iii) if  $x > 0$  and  $r > 0$ , then

$$\frac{r^v}{\mu(B(x, r))} \int_{B(x,r)} |f(y)|^p d\mu(y) \geq r^v (x+r)^{-\beta}.$$

If  $0 < \beta < 1$  and  $\beta \leq \nu$ , then (i) and (ii) imply that  $f \in L^{p, \nu; 2}(\mathbf{R}^1)$ , and if  $0 < \beta < 1$  and  $\beta < \nu$ , then (iii) implies that

$$\limsup_{r \rightarrow \infty} \frac{r^\nu}{\mu(B(x, r))} \int_{B(x, r)} |f(y)|^p d\mu(y) = \infty$$

for every fixed  $x > 0$ , so that  $f \notin L^{p, \nu; 1}(\mathbf{R}^1)$ .

In what follows, if  $f$  is a function on  $G$ , then we assume that  $f = 0$  outside  $G$ .

First we present the boundedness of maximal functions in the Morrey space  $L^{p, \nu; 2}(G)$  due to Sawano-Tanaka [20, Theorem 2.3].

**Lemma 2.2.** *If  $\nu > 0$ , then*

$$\|M_2 f\|_{p, \nu, G; 4} \leq C \|f\|_{p, \nu, G; 2}$$

for all  $f \in L^{p, \nu; 2}(G)$ .

Proof. Let  $\|f\|_{p, \nu, G; 2} \leq 1$ , and fix  $x \in G$  and  $0 < r \leq d_G$ . Write  $A_0 = B(x, 2r)$  and  $A_j = B(x, 2^{j+1}r) \setminus B(x, 2^j r)$  for each positive integer  $j$ . We set

$$f_j = f \chi_{A_j},$$

where  $\chi_E$  denotes the characteristic function of  $E$ . Note that

$$\begin{aligned} \int_{B(x, r)} M_2 f(z)^p d\mu &\leq 2^{p-1} \left( \int_{B(x, r)} M_2 f_0(z)^p d\mu + \int_{B(x, r)} M_2 g_0(z)^p d\mu \right) \\ &\equiv 2^{p-1} (I_1 + I_2), \end{aligned}$$

where  $g_0 = \sum_{j=1}^{\infty} |f_j|$ . We have by Sawano [18, Theorem 1.2 and Proposition 1.1]

$$\begin{aligned} I_1 &\leq \int M_2 f_0(z)^p d\mu \leq C \int |f_0(z)|^p d\mu \\ &= C \int_{B(x, 2r)} |f(z)|^p d\mu \leq C r^{-\nu} \mu(B(x, 4r)). \end{aligned}$$

Next we see that for  $z \in B(x, r)$

$$M_2 f_j(z) \leq C \sup_{\{t: (2^j - 1)r < t < (2^{j+1} + 1)r\}} \frac{1}{\mu(B(z, 2t))} \int_{B(z, t)} |f(y)| d\mu$$

$$\begin{aligned}
&\leq C \sup_{\{t: (2^j-1)r < t < (2^{j+1}-1)r\}} \left( \frac{1}{\mu(B(z, 2t))} \int_{B(z, t)} |f(y)|^p d\mu \right)^{1/p} \\
&\leq C(2^j r)^{-v/p},
\end{aligned}$$

so that

$$M_2 g_0(z) \leq \sum_{j=1}^{\infty} M_2 f_j(z) \leq C \sum_{j=1}^{\infty} (2^j r)^{-v/p} \leq C r^{-v/p}.$$

Hence it follows that

$$I_2 \leq C r^{-v} \int_{B(x, r)} d\mu \leq C r^{-v} \mu(B(x, r)).$$

Thus we obtain

$$\frac{r^v}{\mu(B(x, 4r))} \int_{B(x, r)} M_2 f(z)^p d\mu \leq C,$$

which proves the lemma.  $\square$

**Lemma 2.3.** *If  $f$  is a nonnegative measurable function on  $G$  such that  $\|f\|_{p, v, G; 2} \leq 1$ , then*

$$\int_{B(x, \delta)} \frac{|x - y|^\alpha f(y)}{\mu(B(x, 4|x - y|))} d\mu(y) \leq C \delta^\alpha M_2 f(x)$$

for  $x \in G$  and  $\delta > 0$ .

*Proof.* We have

$$\begin{aligned}
\int_{B(x, \delta)} \frac{|x - y|^\alpha f(y)}{\mu(B(x, 4|x - y|))} d\mu(y) &= \sum_{j=1}^{\infty} \int_{B(x, 2^{-j+1}\delta) \setminus B(x, 2^{-j}\delta)} \frac{|x - y|^\alpha f(y)}{\mu(B(x, 4|x - y|))} d\mu(y) \\
&\leq \sum_{j=1}^{\infty} \frac{(2^{-j+1}\delta)^\alpha}{\mu(B(x, 2^{-j+2}\delta))} \int_{B(x, 2^{-j+1}\delta)} f(y) d\mu(y) \\
&\leq \delta^\alpha M_2 f(x) \sum_{j=1}^{\infty} 2^{(-j+1)\alpha} \\
&= C \delta^\alpha M_2 f(x),
\end{aligned}$$

as required.  $\square$

**Lemma 2.4.** *Let  $v/p \geq \alpha$ . Let  $f$  be a nonnegative measurable function on  $G$  such that  $\|f\|_{p,v,G;2} \leq 1$ . In case  $v/p > \alpha$ ,*

$$\int_{G \setminus B(x,\delta)} \frac{|x-y|^\alpha f(y)}{\mu(B(x, 4|x-y|))} d\mu(y) \leq C\delta^{\alpha-v/p}$$

*and in case  $v/p = \alpha$  and  $G$  is bounded,*

$$\int_{G \setminus B(x,\delta)} \frac{|x-y|^\alpha f(y)}{\mu(B(x, 4|x-y|))} d\mu(y) \leq C \log \frac{1}{\delta}$$

*for  $x \in G$  and small  $\delta > 0$ .*

*Proof.* Let  $j_0$  be the smallest integer such that  $2^{j_0}\delta \geq d_G$ , where  $d_G$  is the diameter of  $G$  as before. By using Hölder's inequality, we have

$$\begin{aligned} & \int_{G \setminus B(x,\delta)} \frac{|x-y|^\alpha f(y)}{\mu(B(x, 4|x-y|))} d\mu(y) \\ &= \sum_{j=0}^{j_0} \int_{B(x, 2^{j+1}\delta) \setminus B(x, 2^j\delta)} \frac{|x-y|^\alpha f(y)}{\mu(B(x, 4|x-y|))} d\mu(y) \\ &\leq \sum_{j=0}^{j_0} (2^{j+1}\delta)^\alpha \frac{1}{\mu(B(x, 2^{j+2}\delta))} \int_{B(x, 2^{j+1}\delta)} f(y) d\mu(y) \\ &\leq C\delta^\alpha \sum_{j=0}^{j_0} 2^{\alpha j} \left( \frac{1}{\mu(B(x, 2^{j+2}\delta))} \int_{B(x, 2^{j+1}\delta)} f(y)^p d\mu(y) \right)^{1/p} \\ &\leq C\delta^\alpha \sum_{j=0}^{j_0} 2^{\alpha j} (2^{j+1}\delta)^{-v/p} \\ &= C\delta^{\alpha-v/p} \sum_{j=0}^{j_0} 2^{(\alpha-v/p)j}, \end{aligned}$$

which proves the required inequality.  $\square$

With the aid of Lemmas 2.2, 2.3 and 2.4, we can apply Hedberg's trick (see [6]) to obtain a Sobolev type inequality for Riesz potentials due to Adams [1, Theorem 3.1], Chiarenza and Frasca [3, Theorem 2], Nakai [15, Theorem 2.2] and Sawano-Tanaka [20, Theorem 3.3].

**Theorem 2.5.** *Let  $1/p^\sharp = 1/p - \alpha/\nu > 0$ . Then there exists a positive constant  $c$  such that*

$$\frac{r^\nu}{\mu(B(z, 4r))} \int_{B(z, r)} \{U_\alpha f(x)\}^{p^\sharp} d\mu(x) \leq c$$

for all  $z \in X$  and  $r > 0$ , whenever  $f$  is a nonnegative measurable function on  $X$  satisfying  $\|f\|_{p, \nu, X; 2} \leq 1$ .

*Proof.* We see from Lemmas 2.3 and 2.4 that

$$\begin{aligned} U_\alpha f(x) &= \int_{B(x, \delta)} \frac{|x - y|^\alpha f(y)}{\mu(B(x, 4|x - y|))} d\mu(y) + \int_{X \setminus B(x, \delta)} \frac{|x - y|^\alpha f(y)}{\mu(B(x, 4|x - y|))} d\mu(y) \\ &\leq C\delta^\alpha M_2 f(x) + C\delta^{\alpha - \nu/p} \end{aligned}$$

for all  $\delta > 0$ . Here, letting

$$\delta = \{M_2 f(x)\}^{-p/\nu},$$

we have

$$U_\alpha f(x) \leq C\{M_2 f(x)\}^{1 - \alpha p/\nu} = C\{M_2 f(x)\}^{p/p^\sharp},$$

which yields

$$\int_{B(z, r)} \{U_\alpha f(x)\}^{p^\sharp} d\mu(x) \leq C \int_{B(z, r)} \{M_2 f(x)\}^p d\mu(x)$$

for  $z \in X$  and  $r > 0$ . Hence Lemma 2.2 gives

$$\frac{r^\nu}{\mu(B(z, 4r))} \int_{B(z, r)} \{U_\alpha f(x)\}^{p^\sharp} d\mu(x) \leq C$$

for such  $z$  and  $r$ , as required.  $\square$

**REMARK 2.6.** Theorem 2.5 implies that the mapping  $f \rightarrow U_\alpha f$  is bounded from  $L^{p, \nu; 2}(X)$  to  $L^{p^\sharp, \nu; 4}(X)$ .

**REMARK 2.7.** When  $X = \mathbf{R}^n$ , consider the potential

$$U_{\alpha, k} f(x) = \int \frac{|x - y|^\alpha f(y)}{\mu(B(x, k|x - y|))} d\mu(y)$$

for  $k > 1$ . Then we can show that the mapping  $f \rightarrow U_{\alpha, k} f$  is bounded from  $L^{p, \nu; 2}(\mathbf{R}^n)$  to  $L^{p^\sharp, \nu; 2}(\mathbf{R}^n)$ , when  $1/p^\sharp = 1/p - \alpha/\nu > 0$ .

REMARK 2.8. We show by an example that the mapping  $f \rightarrow U_{\alpha,1}f$  fails to be bounded in  $L^{p,v;2}(\mathbf{R}^n)$ .

For this purpose, consider  $d\mu(y) = e^y dy$  and

$$f(y) = \begin{cases} y^{-\beta/p} & \text{when } y > 0; \\ 0 & \text{when } y \leq 0. \end{cases}$$

In view of Remark 2.1, we see that  $f \in L^{p,v;2}(\mathbf{R}^1)$  when  $0 < \beta < v \leq 1$ . Further we see that

$$U_{\alpha,1}f(x) \geq \int_x^\infty \frac{t^\alpha (x+t)^{-\beta/p}}{e^x(e^t - e^{-t})} e^{x+t} dt = \infty$$

for all  $x > 0$  when  $\alpha - \beta/p + 1 \geq 0$ . This implies that  $U_{\alpha,1}f$  does not belong to  $L^{p,v;2}(\mathbf{R}^1)$  when  $0 < \beta < v \leq 1$ .

### 3. Exponential integrability

Our aim in this section is to discuss the exponential integrability.

**Theorem 3.1.** *Let  $G$  be bounded and  $v = \alpha p$ . Then there exists a positive constant  $c$  such that*

$$\frac{r^v}{\mu(B(z, 4r))} \int_{G \cap B(z, r)} \{\exp(cU_\alpha f(x)) - 1\} d\mu(x) \leq 1$$

for all  $z \in G$  and  $0 < r \leq d_G$ , whenever  $f$  is a nonnegative measurable function on  $G$  satisfying  $\|f\|_{p,v,G;2} \leq 1$ .

Proof. Let  $\|f\|_{p,v,G;2} \leq 1$ . We see from Lemmas 2.3 and 2.4 that

$$\begin{aligned} U_\alpha f(x) &= \int_{G \cap B(x, \delta)} \frac{|x-y|^\alpha f(y)}{\mu(B(x, 4|x-y|))} d\mu(y) + \int_{G \setminus B(x, \delta)} \frac{|x-y|^\alpha f(y)}{\mu(B(x, 4|x-y|))} d\mu(y) \\ &\leq C\delta^\alpha M_2 f(x) + C \log \frac{1}{\delta} \end{aligned}$$

for  $x \in G$  and small  $\delta > 0$ . Here, letting

$$\delta = \{M_2 f(x)\}^{-1/\alpha} \{\log(M_2 f(x))\}^{1/\alpha}$$

when  $M_2 f(x)$  is large enough, we have

$$\exp(U_\alpha f(x)) \leq C + C\{M_2 f(x)\}^p,$$

so that Lemma 2.2 yields

$$\begin{aligned} \int_{G \cap B(z, r)} \exp(U_\alpha f(x)) d\mu(x) &\leq C\mu(G \cap B(z, r)) + C \int_{G \cap B(z, r)} \{M_2 f(x)\}^p d\mu(x) \\ &\leq C\mu(G \cap B(z, r)) + C \frac{\mu(B(z, 4r))}{r^\nu} \end{aligned}$$

for  $z \in G$  and  $r > 0$ . Hence we find  $c > 0$  such that

$$\frac{r^\nu}{\mu(B(z, 4r))} \int_{G \cap B(z, r)} \{\exp(cU_\alpha f(x)) - 1\} d\mu(x) \leq 1$$

for  $z \in G$  and  $0 < r \leq d_G$ , whenever  $\|f\|_{p, \nu, G; 2} \leq 1$ . Thus the required result is obtained.  $\square$

REMARK 3.2. In Theorem 3.1, we can not add an exponent  $q > 1$  such that

$$\frac{r^\nu}{\mu(B(z, 4r))} \int_{G \cap B(z, r)} \{\exp(cU_\alpha f(x)^q) - 1\} d\mu(x) \leq 1.$$

For this, consider the potential

$$U(x) = \int_{\mathbf{B}} |x - y|^{\alpha-n} |y|^{-\alpha} dy,$$

where  $\mathbf{B} = B(0, 1) \subset \mathbf{R}^n$ . If  $\nu = \alpha p < n$  and  $f(y) = |y|^{-\alpha} \chi_{\mathbf{B}}(y)$ , then

$$\begin{aligned} r^{\nu-n} \int_{B(x, r)} |f(y)|^p dy &\leq r^{\nu-n} \int_{B(x, r)} |x - y|^{-\alpha p} dy \\ &\leq C r^{\nu-n} r^{n-\alpha p} = C \end{aligned}$$

for all  $x \in \mathbf{B}$  and  $r > 0$ , so that  $f \in L^{p, \nu; 1}(\mathbf{B})$ . On the other hand, we see that

$$\begin{aligned} U(x) &\geq \int_{\mathbf{B} \setminus B(x, |x|/2)} |x - y|^{\alpha-n} f(y) dy \\ &\geq 3^{-\alpha} \int_{\mathbf{B} \setminus B(x, |x|/2)} |x - y|^{-n} dy \\ &\geq C \log \frac{2}{|x|} \end{aligned}$$

for  $x \in \mathbf{B}$ , and hence

$$\int_{\mathbf{B}} \exp(cU(x)^q) dx = \infty$$

for  $c > 0$  and  $q > 1$ .

Consider the function

$$e_N(t) = e^t - 1 - t - \frac{t^2}{2!} - \cdots - \frac{t^{N-1}}{(N-1)!}.$$

**Theorem 3.3.** *Let  $v = \alpha p$ . For  $\tilde{v} > v$ , take a positive integer  $N$  such that*

$$N > \frac{\tilde{v}p}{\tilde{v} - \alpha p} = \tilde{p}.$$

*Then there exists a positive constant  $c$  such that*

$$\frac{r^v}{\mu(B(z, 4r))} \int_{B(z, r)} e_N(cU_\alpha f) d\mu(x) \leq 1$$

*for all  $z \in X$  and  $r > 0$ , whenever  $f$  is a nonnegative measurable function on  $X$  satisfying  $\|f\|_{p, v, X; 2} + \|f\|_{p, \tilde{v}, X; 2} \leq 1$ .*

**Proof.** We see from Lemmas 2.3 and 2.4 that

$$\begin{aligned} U_\alpha f(x) &= \int_{B(x, \delta)} \frac{|x - y|^\alpha f(y)}{\mu(B(x, 4|x - y|))} d\mu(y) + \int_{X \setminus B(x, \delta)} \frac{|x - y|^\alpha f(y)}{\mu(B(x, 4|x - y|))} d\mu(y) \\ &\leq C\delta^\alpha M_2 f(x) + C \log \frac{1}{\delta} \end{aligned}$$

for small  $\delta > 0$ . Here, letting

$$\delta = \{M_2 f(x)\}^{-1/\alpha} \{\log(M_2 f(x))\}^{1/\alpha}$$

when  $M_2 f(x)$  is large enough, we have

$$U_\alpha f(x) \leq C \log(2 + M_2 f(x)).$$

We write  $G_1 = \{x \in X : M_2 f(x) > 2\}$  and  $G_2 = \{x \in X : M_2 f(x) \leq 2\}$ . Then we find  $c_1 > 0$  such that

$$\int_{G_1 \cap B(z, r)} e_N(c_1 U_\alpha f(x)) d\mu(x) \leq \int_{B(z, r)} \{M_2 f(x)\}^p d\mu(x)$$

and

$$\begin{aligned} \int_{G_2 \cap B(z, r)} e_N(c_1 U_\alpha f(x)) d\mu(x) &\leq \int_{G_2 \cap B(z, r)} \{U_\alpha f(x)\}^{\tilde{p}} d\mu(x) \\ &\leq \int_{B(z, r)} \{M_2 f(x)\}^p d\mu(x) \end{aligned}$$

for  $z \in X$  and  $r > 0$ . Hence Lemma 2.2 gives

$$\frac{r^v}{\mu(B(z, 4r))} \int_{B(z, r)} e_N(c_2 U_\alpha f(x)) d\mu(x) \leq 1$$

for such  $z$  and  $r$ , whenever  $\|f\|_{p, v, X; 2} + \|f\|_{p, \tilde{v}, X; 2} \leq 1$ . This gives the the required result.  $\square$

REMARK 3.4. Let  $v < \alpha p$  and  $f$  be a nonnegative measurable function on  $X$  belonging to  $L^{p, v; 2}(X)$ . Then  $U_\alpha f(x)$  is seen to be continuous at  $x_0 \in X$  where  $\mu(\partial B(x_0, r)) = 0$  for  $r > 0$  and

$$(3.1) \quad \int \frac{|x_0 - y|^\alpha f(y)}{\mu(B(x_0, 2|x_0 - y|))} d\mu(y) < \infty.$$

In fact, for  $\delta > 0$ , we write

$$\begin{aligned} U_\alpha f(x) &= \int_{B(x_0, 3\delta)} \frac{|x - y|^\alpha f(y)}{\mu(B(x, 4|x - y|))} d\mu(y) + \int_{X \setminus B(x_0, 3\delta)} \frac{|x - y|^\alpha f(y)}{\mu(B(x, 4|x - y|))} d\mu(y) \\ &= U_1(x) + U_2(x). \end{aligned}$$

The proof of Lemma 2.3 implies that

$$U_1(x) \leq C\delta^{\alpha - v/p}$$

for  $x \in B(x_0, \delta)$ , when  $\alpha - v/p > 0$ . Note that  $\mu(B(x, 4|x - y|)) \rightarrow \mu(B(x_0, 4|x_0 - y|))$  as  $x \rightarrow x_0$  for fixed  $y \in X \setminus B(x_0, 3\delta)$  by the assumption that  $\mu(\partial B(x_0, r)) = 0$  for  $r > 0$ . Since  $|x - y|^\alpha / \mu(B(x, 4|x - y|)) \leq C|x_0 - y|^\alpha / \mu(B(x_0, 2|x_0 - y|))$  for  $x \in B(x_0, \delta)$  and  $y \in X \setminus B(x_0, 3\delta)$ , we have by Lebesgue's dominated convergence theorem

$$\lim_{x \rightarrow x_0} U_2(x) = U_2(x_0),$$

which shows that  $U_\alpha f(x)$  is continuous at  $x_0$ .

#### 4. Poincaré inequality

Let  $\mu$  be a nonnegative measure on an open set  $G$ . For a measurable function  $u$  on  $G$ , we define the integral mean over a measurable set  $E \subset G$  of positive measure by

$$u_E = \int_E u(x) d\mu = \frac{1}{\mu(E)} \int_E u(x) d\mu(x).$$

In this section, we assume that  $\mu$  satisfies the lower Ahlfors  $s$ -regularity condition

$$(4.1) \quad c_\mu r^s \leq \mu(B) < \infty$$

for all balls  $B = B(x, r) \subset G$ , where  $s > 0$  and  $c_\mu$  is a positive constant.

We say that a couple  $(u, g)$  satisfies a strong  $(1, p_0)$  Poincaré inequality (in  $G$ ) if

$$(4.2) \quad \int_B |u(x) - u_B| d\mu(x) \leq c_P (\text{diam } B)^{1+s} \left( \frac{1}{\mu(2B)} \int_B |g(y)|^{p_0} d\mu(y) \right)^{1/p_0}$$

for each ball  $B = B(x, r)$  with  $2B = B(x, 2r) \subset G$ , where  $1 \leq p_0 < p$  and  $c_P$  is a positive constant.

Set  $1/p^\sharp = 1/p - 1/p_0 > 0$ .

**Theorem 4.1.** *Let  $\mu$  be a nonnegative measure on  $G$  satisfying (4.1), and assume that a couple  $(u, g)$  satisfies a strong  $(1, p_0)$  Poincaré inequality (4.2). If  $\|g\|_{p, \nu, G; 2} \leq 1$ , then*

$$\frac{r^\nu}{\mu(2B)} \int_B |u(x) - u_B|^{p^\sharp} d\mu(x) \leq C$$

for every ball  $B = B(x, r)$  with  $2B \subset G$ .

Orobitg-Pérez [14] gave a version of Sobolev's inequalities in the  $L^p$  space setting.

Proof of Theorem 4.1. Let  $2B = B(x_0, 2r) \subset G$ . For  $x \in B$ , set

$$B_i(x) = B(x, 2^{-i}r).$$

By the Lebesgue differentiation theorem we have

$$\lim_{i \rightarrow \infty} u_{B_i(x)} = u(x)$$

for  $\mu$ -a.e.  $x$  and hence we may assume that our fixed point  $x$  has this property. Let  $N = N(x)$  be a positive integer whose value will be determined later. Letting  $B_i = B_i(x)$  for  $i = 1, 2, \dots$ , we have by (4.1)

$$\begin{aligned} |u(x) - u_B| &\leq |u_{B_1} - u_B| + \sum_{i=1}^{\infty} |u_{B_i} - u_{B_{i+1}}| \\ &\leq \frac{1}{\mu(B \cap B_1)} \int_B |u - u_B| d\mu + \frac{1}{\mu(B \cap B_1)} \int_{B_1} |u - u_{B_1}| d\mu \\ &\quad + \sum_{i=1}^{\infty} \frac{1}{\mu(B_{i+1})} \int_{B_i} |u - u_{B_i}| d\mu \end{aligned}$$

$$\begin{aligned}
&\leq Cr^{-s} \int_B |u - u_B| d\mu + C \sum_{i=1}^N (2^{-i}r)^{-s} \int_{B_i} |u - u_{B_i}| d\mu \\
&\quad + C \sum_{i=N+1}^{\infty} (2^{-i}r)^{-s} \int_{B_i} |u - u_{B_i}| d\mu \\
&= I_0 + I_1 + I_2.
\end{aligned}$$

By using Hölder's inequality and a strong  $(1, p_0)$  Poincaré inequality, we find

$$\begin{aligned}
I_0 &\leq C(\text{diam } B) \left( \frac{1}{\mu(2B)} \int_B |g|^{p_0} d\mu \right)^{1/p_0} \\
&\leq C(\text{diam } B) \left( \frac{1}{\mu(2B)} \int_B |g|^p d\mu \right)^{1/p} \\
&\leq Cr^{1-\nu/p}
\end{aligned}$$

and

$$\begin{aligned}
I_1 &\leq C \sum_{i=0}^N 2^{-i}r \left( \frac{1}{\mu(2B_i)} \int_{B_i} |g|^{p_0} d\mu \right)^{1/p_0} \\
&\leq C \sum_{i=0}^N 2^{-i}r \left( \frac{1}{\mu(2B_i)} \int_{B_i} |g|^p d\mu \right)^{1/p} \\
&\leq C \sum_{i=0}^N (2^{-i}r)^{1-\nu/p} \\
&\leq C(2^{-N}r)^{1-\nu/p}.
\end{aligned}$$

According to the estimation of  $I_1$ , we obtain

$$\begin{aligned}
I_2 &\leq C \sum_{i=N+1}^{\infty} 2^{-i}r \left( \frac{1}{\mu(2B_i)} \int_{B_i} |g|^{p_0} d\mu \right)^{1/p_0} \\
&\leq C \sum_{i=N+1}^{\infty} 2^{-i}r \{M_2 g_0(x)\}^{1/p_0} \\
&\leq C2^{-N}r \{M_2 g_0(x)\}^{1/p_0},
\end{aligned}$$

where  $g_0(y) = |g(y)|^{p_0} \chi_B(y)$  with  $\chi_B$  denoting the characteristic function of  $B$ . Now, considering  $N$  to be the integer part of  $(2^{-N}r)^{-\nu/p} = \{M_2 g_0(x)\}^{1/p_0}$ , we establish

$$(4.3) \quad |u(x) - u_B| \leq C[r^{-\nu/p^\sharp} + \{M_2 g_0(x)\}^{p/(p^\sharp p_0)}].$$

Therefore, it follows from Lemma 2.2 that

$$\begin{aligned}
 \int_B |u(x) - u_B|^{p^*} d\mu(x) &\leq C \int_B [r^{-v} + \{M_2 g_0(x)\}^{p/p_0}] d\mu(x) \\
 &\leq C \left[ r^{-v} \mu(B) + \int_B g_0(y)^{p/p_0} d\mu(y) \right] \\
 &= C \left[ r^{-v} \mu(B) + \int_B |g(y)|^p d\mu(y) \right] \\
 &\leq Cr^{-v} \mu(2B),
 \end{aligned}$$

as required.  $\square$

**Corollary 4.2.** *Let  $\mu$  be a nonnegative measure on  $G$  satisfying (4.1), and assume that a couple  $(u, g)$  satisfies a strong  $(1, p_0)$  Poincaré inequality. Then*

$$\left( \int_B |u(x) - u_B|^{p^*} d\mu(x) \right)^{1/p^*} \leq Cr^{-s/p^*} \mu(B)^{1/p} \left( \int_B |g(y)|^p d\mu(y) \right)^{1/p}$$

for every  $B = B(x, r)$  with  $2B \subset G$ , where  $1/p^* = 1/p - 1/s > 0$ .

To show this, first suppose  $\int_B |g(y)|^p d\mu(y) \leq 1$ . Then the decay condition (4.1) implies that  $\|g\|_{p,s,G;1}$  is bounded. Now we see from the inequalities after (4.3) that

$$\int_B |u(x) - u_B|^{p^*} d\mu(x) \leq Cr^{-s} \mu(B).$$

Hence we obtain

$$\left( \int_B |u(x) - u_B|^{p^*} d\mu(x) \right)^{1/p^*} \leq C(r^{-s} \mu(B))^{1/p^*} \left( \int_B |g(y)|^p d\mu(y) \right)^{1/p}$$

for a general  $g$ , which gives the required result.

**REMARK 4.3.** Let  $G$  be an open set in  $\mathbf{R}^n$ . We assume that a couple  $(u, g)$  satisfies a  $(1, p_0)$  Poincaré inequality in  $G$ , that is,

$$(4.4) \quad \int_B |u(x) - u_B| d\mu(x) \leq c'_p \mu(B)^{1/s} \left( \int_B |g(y)|^{p_0} d\mu(y) \right)^{1/p_0}$$

for all balls  $B \subset G$ , where  $1 < p_0 < p$  and  $c'_p$  is a positive constant independent of  $(u, g)$ . We further assume that  $\mu(\partial B) = 0$  and

$$\mu(B)^{v/s} \int_B |g(y)|^p d\mu(y) \leq 1$$

for each ball  $B \subset G$ , where  $1 < p < \nu$ . Then

$$\sup_{B \subset G} \mu(B)^{\nu/s} \int_B |u(x) - u_B|^{p^*} d\mu(x) \leq C.$$

For this, we also refer to Hajlasz-Koskela [5].

For a proof of this fact, let  $x \in G$  be a Lebesgue point of  $u$ . As in the proof of Theorem 1.1 by Gogatishvili-Koskela [4], we take a sequence of balls  $\{B_j\}$  such that  $x \in B_{j+1} \subset B_j \subset B$  and  $\mu(B_j) = 2^{-j} \mu(B)$ . Then, as in (4.3), we can prove

$$|u(x) - u_B|^{p^*} \leq C[\mu(B)^{-\nu/s} + \{M_1 g_0(x)\}^{p/p_0}],$$

which gives the required inequality by the boundedness of the maximal operator  $M_1$ .

Finally we discuss the exponential integrability in the same manner as in Theorem 4.1.

**Theorem 4.4.** *Let  $G$  be bounded and  $\nu = p$ . Let  $\mu$  be a nonnegative measure on  $G$  satisfying (4.1), and assume that a couple  $(u, g)$  satisfies a strong  $(1, p_0)$  Poincaré inequality. Then there exists a positive constant  $c$  such that*

$$\frac{r^\nu}{\mu(2B)} \int_B \{\exp(c|u(x) - u_B|) - 1\} d\mu(x) \leq 1$$

for every ball  $B = B(z, r)$ , whenever  $2B \subset G$  and  $\|g\|_{p, \nu, G; 2} \leq 1$ .

**REMARK 4.5.** Let  $\mu$  be a nonnegative measure on  $\mathbf{R}^n$  satisfying (4.1), and assume that a couple  $(u, g)$  satisfies a strong  $(1, p_0)$  Poincaré inequality. If  $g \in L^{p, \nu; 2}(\mathbf{R}^n)$  and  $p > \nu$ , then  $u$  can be corrected almost everywhere to be continuous on  $\mathbf{R}^n$ ; for this, see [10].

In fact, the first part of the proof of Theorem 4.1 implies that

$$|u(x) - u_B| \leq Cr^{(p-\nu)/p}$$

for almost every  $x \in B$ , which proves

$$|u(x) - u(y)| \leq C|x - y|^{(p-\nu)/p}$$

for almost every  $x, y \in B$ .

**ACKNOWLEDGMENT.** We would like to thank Dr. Yoshihiro Sawano and the referee for their kind comments and suggestions.

## References

- [1] D.R. Adams: *A note on Riesz potentials*, Duke Math. J. **42** (1975), 765–778.
- [2] D.R. Adams and L.I. Hedberg: *Function Spaces and Potential Theory*, Springer, Berlin, 1996.
- [3] F. Chiarenza and M. Frasca: *Morrey spaces and Hardy-Littlewood maximal function*, Rend. Mat. Appl. (7) **7** (1987), 273–279.
- [4] A. Gogatishvili and P. Koskela: *A non-doubling Trudinger inequality*, Studia Math. **170** (2005), 113–119.
- [5] P. Hajlasz and P. Koskela: *Sobolev met Poincaré*, Mem. Amer. Math. Soc. **145** (2000).
- [6] L.I. Hedberg: *On certain convolution inequalities*, Proc. Amer. Math. Soc. **36** (1972), 505–510.
- [7] J. Heinonen: *Lectures on Analysis on Metric Spaces*, Springer, New York, 2001.
- [8] P. Mattila: *Geometry of Sets and Measures in Euclidean Spaces*, Cambridge Univ. Press, Cambridge, 1995.
- [9] Y. Mizuta: *Potential Theory in Euclidean Spaces*, Gakkōtoshō, Tokyo, 1996.
- [10] Y. Mizuta and T. Shimomura: *Continuity and differentiability for weighted Sobolev spaces*, Proc. Amer. Math. Soc. **130** (2002), 2985–2994.
- [11] Y. Mizuta and T. Shimomura: *Sobolev embeddings for Riesz potentials of functions in Morrey spaces of variable exponent*, J. Math. Soc. Japan **60** (2008), 583–602.
- [12] Y. Mizuta and T. Shimomura: *Continuity properties for Riesz potentials of functions in Morrey spaces of variable exponent*, to appear in Math. Inequal. Appl.
- [13] C.B. Morrey, Jr.: *On the solutions of quasi-linear elliptic partial differential equations*, Trans. Amer. Math. Soc. **43** (1938), 126–166.
- [14] J. Oröbítg and C. Pérez:  *$A_p$  weights for nondoubling measures in  $\mathbf{R}^n$  and applications*, Trans. Amer. Math. Soc. **354** (2002), 2013–2033.
- [15] E. Nakai: *Generalized fractional integrals on Orlicz-Morrey spaces*; in Banach and Function Spaces, Yokohama Publ., Yokohama, 2004, 323–333.
- [16] W. Orlicz: *Über konjugierte Exponentenfolgen*, Studia Math. **3** (1931), 200–211.
- [17] J. Peetre: *On the theory of  $L_{p,\lambda}$  spaces*, J. Funct. Anal. **4** (1969), 71–87.
- [18] Y. Sawano: *Sharp estimates of the modified Hardy-Littlewood maximal operator on the non-homogeneous space via covering lemmas*, Hokkaido Math. J. **34** (2005), 435–458.
- [19] Y. Sawano, T. Sobukawa and H. Tanaka: *Limiting case of the boundedness of fractional integral operators on nonhomogeneous space*, J. Inequal. Appl. (2006), Art. ID 92470, 16 pp.
- [20] Y. Sawano and H. Tanaka: *Morrey spaces for non-doubling measures*, Acta Math. Sin. (Engl. Ser.) **21** (2005), 1535–1544.

Yoshihiro Mizuta  
Department of Mathematics  
Graduate School of Science  
Hiroshima University  
Higashi-Hiroshima 739-8521  
Japan  
e-mail: yomizuta@hiroshima-u.ac.jp

Tetsu Shimomura  
Department of Mathematics  
Graduate School of Education  
Hiroshima University  
Higashi-Hiroshima 739-8524  
Japan  
e-mail: tshimo@hiroshima-u.ac.jp

Takuya Sobukawa  
Department of Mathematics Education  
Faculty of Education  
Okayama University  
Tsushima-naka 700-8530  
Japan  
e-mail: sobu@cc.okayama-u.ac.jp