

Title	OPTIMAL HARDY-TYPE INEQUALITIES FOR SCHRÖDINGER FORMS
Author(s)	Takeda, Masayoshi
Citation	Osaka Journal of Mathematics. 2023, 60(4), p. 761-776
Version Type	VoR
URL	https://doi.org/10.18910/93059
rights	
Note	

Osaka University Knowledge Archive : OUKA

<https://ir.library.osaka-u.ac.jp/>

Osaka University

OPTIMAL HARDY-TYPE INEQUALITIES FOR SCHRÖDINGER FORMS

MASAYOSHI TAKEDA

(Received April 14, 2022, revised August 31, 2022)

Abstract

We give a method to construct a critical Schrödinger form from the subcritical Schrödinger form by subtracting a suitable positive potential. The method enables us to obtain optimal Hardy-type inequalities.

Contents

1. Introduction	761
2. Extended Schrödinger spaces	764
3. Probabilistic representation of Schrödinger semigroups	765
4. Criticality and Hardy-type inequalities	767
References	775

1. Introduction

In [6], Devyver, Fraas and Pinchover give a method for obtaining *optimal* Hardy weights for second-order non-negative elliptic operators on non-compact Riemannian manifolds, in particular, they show that the criticality of Schrödinger forms is related to the *critical* Hardy weights. In [20] we give a method to construct a critical Schrödinger form from a transient Dirichlet form by subtracting a suitable positive potential. In other words, we give a method to construct critical Hardy weights for a transient Dirichlet form by applying the idea in [6]. In this paper, we will consider subcritical Schrödinger forms instead of transient Dirichlet forms, and extend the method for subcritical Schrödinger forms. As an application, we obtain a method to construct critical Hardy weights for Schrödinger forms. Moreover, we discuss the optimality of Hardy weights in the sense of [6], a stronger notion than the criticality, and give a condition for the critical Hardy weights being optimal ones.

Let E be a locally compact separable metric space and m a positive Radon measure on E with full topological support. Let $X = (P_x, X_t, \zeta)$ be an m -symmetric Hunt process. We assume that X is irreducible and resolvent doubly Feller, in addition, that X generates a regular Dirichlet form $(\mathcal{E}, \mathcal{D}(\mathcal{E}))$ on $L^2(E; m)$.

Denote by $\mathcal{K}_{loc}(X)$ the totality of local Kato measures (Definition 3.1 (1)). For a signed local Kato measure such that the positive (resp. negative) part μ^+ (resp. μ^-) of μ belongs to

$\mathcal{K}_{loc}(X)$ ($\mu \in \mathcal{K}_{loc}(X) - \mathcal{K}_{loc}(X)$ in notation), we define a symmetric form by

$$\mathcal{E}^\mu(u, u) = \mathcal{E}(u, u) + \int_E u^2 d\mu, \quad u \in \mathcal{D}(\mathcal{E}) \cap C_0(E).$$

The regularity of $(\mathcal{E}, \mathcal{D}(\mathcal{E}))$ implies that a measure in $\mathcal{K}_{loc}(X)$ is Radon (Remark 3.2) and the form $(\mathcal{E}^\mu, \mathcal{D}(\mathcal{E}) \cap C_0(E))$ is well-defined. In the sequel, for a symmetric bilinear form $(a, \mathcal{D}(a))$ we simply write $a(u)$ for $a(u, u)$.

We suppose that $(\mathcal{E}^\mu, \mathcal{D}(\mathcal{E}) \cap C_0(E))$ is positive semi-definite:

$$(1) \quad \mathcal{E}^\mu(u) \geq 0 \left(\iff \int_E u^2 d\mu^- \leq \mathcal{E}^{\mu^+}(u) \right), \quad u \in \mathcal{D}(\mathcal{E}) \cap C_0(E).$$

Applying results in [1], we prove in [20] that $(\mathcal{E}^\mu, \mathcal{D}(\mathcal{E}) \cap C_0(E))$ is closable in $L^2(E; m)$. We denote the closure $(\mathcal{E}^\mu, \mathcal{D}(\mathcal{E}^\mu))$ and call it *Schrödinger form* with potential μ . By the Radonness of μ^+ , we see that $\mathcal{D}(\mathcal{E}) \cap L^2(E; \mu^+) \subset \mathcal{D}(\mathcal{E}^\mu)$ and

$$\mathcal{E}^\mu(u) = \mathcal{E}(u) + \int_E \tilde{u}^2 d\mu, \quad u \in \mathcal{D}(\mathcal{E}) \cap L^2(E; \mu^+).$$

Here \tilde{u} is a quasi-continuous version of u . In this paper, we always assume that every function u is represented by its quasi-continuous version if it admits.

The L^2 -semigroup T_t^μ generated by $(\mathcal{E}^\mu, \mathcal{D}(\mathcal{E}^\mu))$ is expressed by Feynman-Kac semigroup ([20, Theorem 4.2]): For a bounded Borel function f in $L^2(E; m)$

$$(2) \quad T_t^\mu f(x) = p_t^\mu f(x) \left(:= E_x \left(e^{-A_t^\mu} f(X_t) \right) \right), \quad m\text{-a.e. } x.$$

Here $A_t^\mu = A_t^{\mu^+} - A_t^{\mu^-}$ and $A_t^{\mu^+}$ (resp. $A_t^{\mu^-}$) is the positive continuous additive functional with Revuz measure μ^+ (resp. μ^-). We suppose that $(\mathcal{E}^\mu, \mathcal{D}(\mathcal{E}^\mu))$ is *subcritical*, that is, there exists the Green function $R^\mu(x, y)$ such that for a positive Borel function f

$$\int_0^\infty p_t^\mu f(x) dt = \int_E R^\mu(x, y) f(y) dm(y), \quad \forall x \in E.$$

Let $\mathcal{K}_{loc}^\mu(X)$ be the set of local Kato measures such that for any compact set $K \subset E$

$$(3) \quad R^\mu(1_K \nu)u(x) = \int_E R^\mu(x, y) 1_K(y) d\nu(y) \in L^\infty(E; m).$$

For a non-trivial measure ν in $\mathcal{K}_{loc}^\mu(X)$ define measures ν^μ and μ^ν by

$$(4) \quad \nu^\mu = \frac{\nu}{R^\mu \nu}$$

and

$$(5) \quad \mu^\nu = \mu - \nu^\mu.$$

We will show in Corollary 4.2 and Lemma 4.3 below that μ^ν belongs to $\mathcal{K}_{loc}(X) - \mathcal{K}_{loc}(X)$ and $(\mathcal{E}^{\mu^\nu}, \mathcal{D}(\mathcal{E}) \cap C_0(E))$ is still positive semi-definite

$$(6) \quad \mathcal{E}^{\mu^\nu}(u) = \mathcal{E}^\mu(u) - \int_E u^2 d\nu^\mu \geq 0, \quad u \in \mathcal{D}(\mathcal{E}) \cap C_0(E).$$

In other words, the measure ν^μ is a *Hardy weight* for $(\mathcal{E}^\mu, \mathcal{D}(\mathcal{E}^\mu))$. As remarked above, $(\mathcal{E}^{\mu^\nu}, \mathcal{D}(\mathcal{E}) \cap C_0(E))$ is closable and its closure defines a new Schrödinger form $(\mathcal{E}^{\mu^\nu}, \mathcal{D}(\mathcal{E}^{\mu^\nu}))$.

Let C be the totality of compact sets of E . We then obtain the following main result in this paper: If a non-trivial positive measure ν in $\mathcal{K}_{loc}^\mu(X)$ satisfies that

$$(7) \quad \sup_{K \in \mathcal{C}} \iint_{K \times K^c} R^\mu(x, y) \nu(dx) \nu(dy) < \infty,$$

then $(\mathcal{E}^{\mu^\nu}, \mathcal{D}(\mathcal{E}^{\mu^\nu}))$ turns out to be a *critical* Schrödinger form. Here K^c is the complement of K . More precisely, the function $R^\mu \nu$ is a ground state of $(\mathcal{E}^{\mu^\nu}, \mathcal{D}(\mathcal{E}^{\mu^\nu}))$, that is, $R^\mu \nu$ belongs to the *extended Schrödinger space* $\mathcal{D}_e(\mathcal{E}^{\mu^\nu})$ of $(\mathcal{E}^{\mu^\nu}, \mathcal{D}(\mathcal{E}^{\mu^\nu}))$ (see Section 2 for the definition of the extended Schrödinger space) and $\mathcal{E}^{\mu^\nu}(R^\mu \nu) = 0$. As a corollary, we see that ν^μ is a *critical* Hardy weight for $(\mathcal{E}^\mu, \mathcal{D}(\mathcal{E}^\mu))$ in the sense that there exists no non-trivial positive function ψ such that

$$(8) \quad \int_E u^2 d(\nu^\mu + \psi m) \leq \mathcal{E}^\mu(u), \quad u \in \mathcal{D}(\mathcal{E}) \cap C_0(E).$$

In particular, if $(\mathcal{E}, \mathcal{D}(\mathcal{E}))$ is transient and $\mu \equiv 0$, then every $\nu \in \mathcal{K}_{loc}(X)$ satisfies (3) by replacing $R^\mu(x, y)$ with the 0-resolvent $R(x, y)$ of X . Indeed, since $1_K \nu$ is Green-tight, $1_K \nu \in \mathcal{K}_\infty(X)$ (Definition 3.1 (2)), the condition (3) is derived from [3, Proposition 2.2]. As a result, for any $\nu \in \mathcal{K}_{loc}$ the next Hardy-type inequality follows:

$$(9) \quad \int_E u^2 \frac{d\nu}{R\nu} \leq \mathcal{E}(u), \quad u \in \mathcal{D}(\mathcal{E}) \cap C_0(E).$$

The inequality (9) is proved in Fitzsimmons [7] (see also [2]). Moreover, we see that if the measure $\nu/R\mu$ is a critical Hardy weight for the Dirichlet form $(\mathcal{E}, \mathcal{D}(\mathcal{E}))$ if ν satisfies (7) obtained by replacing $R^\mu(x, y)$ with $R(x, y)$.

As stated above, the function $R^\mu \nu$ belongs to $\mathcal{D}_e(\mathcal{E}^{\mu^\nu})$ under the condition (7). Lemma 4.3 below tells us that $\mathcal{D}_e(\mathcal{E}^\mu)$ is included in $\mathcal{D}_e(\mathcal{E}^{\mu^\nu})$ and $R^\mu \nu$ does not belong to $\mathcal{D}_e(\mathcal{E}^\mu)$ in general. If ν satisfies the stronger condition than (7),

$$\iint_{E \times E} R^\mu(x, y) \nu(dx) \nu(dy) < \infty,$$

i.e., ν is of finite energy with respect to R^μ , then $R^\mu \nu$ belongs to $L^2(E; \nu^\mu)$ because

$$\int_E (R^\mu \nu)^2 d\nu^\mu = \int_E R^\mu \nu d\nu = \iint_{E \times E} R^\mu(x, y) \nu(dx) \nu(dy) < \infty.$$

Moreover, $R^\mu \nu$ belongs to $\mathcal{D}_e(\mathcal{E}^\mu)$ by Lemma 4.8 below. Hence, $\mathcal{E}^\mu(R^\mu \nu)$ is finite and thus

$$\mathcal{E}^{\mu^\nu}(R^\mu \nu) = 0 \iff \frac{\mathcal{E}^\mu(R^\mu \nu)}{\int_E (R^\mu \nu)^2 d\nu^\mu} = 1.$$

Noting that by (6)

$$(10) \quad \inf_{u \in \mathcal{D}_e(\mathcal{E}^\mu)} \frac{\mathcal{E}^\mu(u)}{\int_E u^2 d\nu^\mu} \geq 1,$$

we see $R^\mu \nu$ is a minimizer for the left hand side of (10). In this case, the Schrödinger form $(\mathcal{E}^{\mu^\nu}, \mathcal{D}(\mathcal{E}^{\mu^\nu}))$ is said to be *positive-critical* ([6, Definition 4.8]).

On the other hand, if ν is not of finite energy,

$$(11) \quad \iint_{E \times E} R^\mu(x, y) \nu(dx) \nu(dy) = \infty,$$

then $R^\mu \nu$ does not belong to $L^2(E; \nu^\mu)$ and $(\mathcal{E}^\mu, \mathcal{D}(\mathcal{E}^\mu))$ is *null-critical* in the sense of [6].

The measure ν^μ is called *optimal at infinity* if for any $K \in \mathcal{C}$

$$\lambda \int_E u^2 d\nu^\mu \leq \mathcal{E}^\mu(u), \quad u \in \mathcal{D}(\mathcal{E}) \cap C_0(K^c),$$

then $\lambda \leq 1$. We see from [12, Corollary 3.4] (or [14, Theorem 3]) that if for any $K \in \mathcal{C}$

$$\iint_{K \times E} R^\mu(x, y) \nu(dx) \nu(dy) < \infty,$$

i.e., $R^\mu \nu$ is locally integrable, then the null-criticality implies the optimality at infinity. In generally, if for any $K \in \mathcal{C}$

$$(12) \quad \iint_{K^c \times E} R^\mu(x, y) \nu(dx) \nu(dy) = \infty,$$

then the optimality at infinity holds. Devyver, Fraas and Pinchover [6], where they call a Hardy-type inequality *optimal* if a Hardy weight is critical, null-critical and optimal at infinity. Noting that (12) implies (11), we can conclude that if a measure ν satisfies (3), (7) and (12), then the measure ν^μ is an optimal Hardy-weight for $(\mathcal{E}^\mu, \mathcal{D}(\mathcal{E}^\mu))$ in the sense of [6].

2. Extended Schrödinger spaces

Let E be a locally compact separable metric space and m a positive Radon measure on E with full topological support. Let $(\mathcal{E}, \mathcal{D}(\mathcal{E}))$ be a regular Dirichlet form on $L^2(E; m)$ (c.f. [9, p.6]). We denote by $u \in \mathcal{D}_{loc}(\mathcal{E})$ if for any relatively compact open set D there exists a function $v \in \mathcal{D}(\mathcal{E})$ such that $u = v$ m -a.e. on D . We assume that $(\mathcal{E}, \mathcal{D}(\mathcal{E}))$ is irreducible (c.f. [9, p.40, p.55]).

We call a positive Borel measure μ on E *smooth* if it satisfies

- (i) μ charges no set of zero capacity,
- (ii) there exists an increasing sequence $\{F_n\}$ of closed sets such that
 - a) $\mu(F_n) < \infty, n = 1, 2, \dots,$
 - b) $\lim_{n \rightarrow \infty} \text{Cap}(K \setminus F_n) = 0$ for any compact set K .

We denote by \mathcal{S} the totality of smooth measures.

For a signed smooth Radon measure $\mu = \mu^+ - \mu^- \in \mathcal{S} - \mathcal{S}$ define a symmetric form on $L^2(E; m)$ by

$$(13) \quad \mathcal{E}^\mu(u, v) = \mathcal{E}(u, v) + \int_E u v d\mu, \quad u, v \in \mathcal{D}(\mathcal{E}) \cap C_0(E).$$

We assume that $(\mathcal{E}^\mu, \mathcal{D}(\mathcal{E}) \cap C_0(E))$ is positive semi-definite:

$$(14) \quad \mathcal{E}^\mu(u) \geq 0 \left(\iff \int_E u^2 d\mu^- \leq \mathcal{E}^{\mu^+}(u) \right), \quad u \in \mathcal{D}(\mathcal{E}) \cap C_0(E).$$

When $(\mathcal{E}^\mu, \mathcal{D}(\mathcal{E}) \cap C_0(E))$ is closable, we denote by $(\mathcal{E}^\mu, \mathcal{D}(\mathcal{E}^\mu))$ its closure and call it *Schrödinger form* with potential μ .

A densely defined, closed, positive semi-definite symmetric bilinear form $(a, \mathcal{D}(a))$ is said to be *positive preserving* if for $u \in \mathcal{D}(a)$, $|u|$ belongs to $\mathcal{D}(a)$ and $a(|u|) \leq a(u)$. It follows from [5, Lemma 1.3.4] that the form $(\mathcal{E}^\mu, \mathcal{D}(\mathcal{E}^\mu))$ is positive preserving because $\mathcal{E}^\mu(|u|) \leq \mathcal{E}^\mu(u)$ for $u \in \mathcal{D}(\mathcal{E}) \cap C_0(E)$. As a result, we see from [17, Proposition 2] that $(\mathcal{E}^\mu, \mathcal{D}(\mathcal{E}^\mu))$ has the *Fatou property*, i.e., if $\{u_n\} \subset \mathcal{D}(\mathcal{E}^\mu)$ satisfies $\sup_n \mathcal{E}^\mu(u_n) < \infty$ and $u_n \rightarrow u \in \mathcal{D}(\mathcal{E}^\mu)$ m -a.e., then $\liminf_{n \rightarrow \infty} \mathcal{E}^\mu(u_n) \geq \mathcal{E}^\mu(u)$. Hence, following [16], we can define a space $\mathcal{D}_e(\mathcal{E}^\mu)$ in the way similar to the extended Dirichlet space: An m -measurable function u with $|u| < \infty$ m -a.e. is said to be in $\mathcal{D}_e(\mathcal{E}^\mu)$ if there exists an \mathcal{E}^μ -Cauchy sequence $\{u_n\} \subset \mathcal{D}(\mathcal{E}^\mu)$ such that $\lim_{n \rightarrow \infty} u_n = u$ m -a.e. We call $\mathcal{D}_e(\mathcal{E}^\mu)$ the *extended Schrödinger space* of $(\mathcal{E}^\mu, \mathcal{D}(\mathcal{E}^\mu))$ and the sequence $\{u_n\}$ an *approximating sequence* of u . For $u \in \mathcal{D}_e(\mathcal{E}^\mu)$ and an approximating sequence $\{u_n\}$ of u , we define

$$(15) \quad \mathcal{E}^\mu(u) = \lim_{n \rightarrow \infty} \mathcal{E}^\mu(u_n).$$

We define the criticality and subcriticality of Schrödinger forms in the way similar to the recurrence and transience of Dirichlet forms.

DEFINITION 2.1. Let $(\mathcal{E}^\mu, \mathcal{D}(\mathcal{E}^\mu))$ be a positive semi-definite Schrödinger form.

(1) $(\mathcal{E}^\mu, \mathcal{D}(\mathcal{E}^\mu))$ is said to be *subcritical* if there exists a bounded function g in $L^1(E; m)$ strictly positive m -a.e. such that

$$(16) \quad \int_E |u|g dm \leq \sqrt{\mathcal{E}^\mu(u)}, \quad u \in \mathcal{D}_e(\mathcal{E}^\mu).$$

(2) $(\mathcal{E}^\mu, \mathcal{D}(\mathcal{E}^\mu))$ is said to be *critical* if there exists a function ϕ in $\mathcal{D}_e(\mathcal{E}^\mu)$ strictly positive m -a.e. such that $\mathcal{E}^\mu(\phi) = 0$. The function ϕ is said to be the *ground state*.

Define the operator G^μ by

$$G^\mu f(x) = \int_0^\infty T_t^\mu f(x) dt \quad (\leq +\infty)$$

for a positive function f . Here T_t^μ is the L^2 -semigroup on $L^2(E; m)$ generated by $(\mathcal{E}^\mu, \mathcal{D}(\mathcal{E}^\mu))$.

Lemma 2.2 ([20, Lemma 2.3]). *Let g be the function in Definition 2.1 (1). Then $G^\mu g$ belongs to $\mathcal{D}_e(\mathcal{E}^\mu)$.*

REMARK 2.3. It is recently proved in [15, Theorem A.3] that if the semigroup T_t^μ is expressed using a density $p_t^\mu(x, y)$, $T_t^\mu f(x) = \int_E p_t^\mu(x, y) f(y) dm(y)$, then $(\mathcal{E}^\mu, \mathcal{D}(\mathcal{E}^\mu))$ is subcritical or critical.

REMARK 2.4. We see from the inequality (16) that if $(\mathcal{E}^\mu, \mathcal{D}(\mathcal{E}^\mu))$ is subcritical, then $(\mathcal{D}(\mathcal{E}^\mu), \mathcal{E}^\mu(\cdot, \cdot))$ is a Hilbert space.

3. Probabilistic representation of Schrödinger semigroups

Let $X = (\Omega, \mathcal{F}, \{\mathcal{F}_t\}_{t \geq 0}, \{P_x\}_{x \in E}, \{X_t\}_{t \geq 0}, \zeta)$ be the symmetric Hunt process generated by $(\mathcal{E}, \mathcal{D}(\mathcal{E}))$, where $\{\mathcal{F}_t\}_{t \geq 0}$ is the augmented filtration and ζ is the lifetime of X . Denote by $\{p_t\}_{t \geq 0}$ and $\{R_\alpha\}_{\alpha \geq 0}$ the semigroup and resolvent of X :

$$p_t f(x) = E_x(f(X_t)), \quad R_\alpha f(x) = \int_0^\infty e^{-\alpha t} p_t f(x) dt.$$

Then $p_t f(x) = T_t f(x)$ m -a.e., $R_\alpha f(x) = \int_0^\infty T_t f(x) dt$ m -a.e., where T_t is the L^2 -semigroup on $L^2(E; m)$ generated by $(\mathcal{E}, \mathcal{D}(\mathcal{E}))$. In the sequel, we assume that X satisfies, in addition, the next condition:

Feller Property (F). For each $t > 0$, $p_t(C_\infty(E)) \subset C_\infty(E)$ and for each $f \in C_\infty(E)$ and $x \in E$, $\lim_{t \rightarrow 0} p_t f(x) = f(x)$, where $C_\infty(E)$ is the space of continuous functions on E vanishing at infinity.

Resolvent Strong Feller Property (RSF). For each $\alpha > 0$, $R_\alpha(\mathcal{B}_b(E)) \subset C_b(E)$, where $\mathcal{B}_b(E)$ (resp. $C_b(E)$) is the space of bounded Borel (resp. continuous) functions on E .

Following [11], a Hunt process is said to be *resolvent doubly Feller* if it enjoys both the Feller property and resolvent strong Feller property. We see from **(RSF)** that the resolvent kernel $R_\alpha(x, dy)$ of X has a non-negative jointly measurable density $R_\alpha(x, y)$ with respect to m : For $x \in E$ and $f \in \mathcal{B}_b(E)$

$$R_\alpha f(x) = \int_E R_\alpha(x, y) f(y) m(dy).$$

Moreover, $R_\alpha(x, y)$ is α -excessive in x and in y ([9, Lemma 4.2.4]). We simply write $R(x, y)$ for $R_0(x, y)$ ($:= \lim_{\alpha \rightarrow 0} R_\alpha(x, y)$). For a measure μ , we define the α -potential of μ by

$$R_\alpha \mu(x) = \int_E R_\alpha(x, y) \mu(dy), \quad \alpha \geq 0.$$

Let S_0 be the set of positive Borel measures μ such that $\mu(E) < \infty$ and $R_1 \mu$ is bounded. We call a Borel measure μ on E *smooth measure in the strict sense* if there exists a sequence $\{E_n\}$ of Borel sets increasing to E such that for each n , $1_{E_n} \mu \in S_0$ and for any $x \in E$

$$P_x(\lim_{n \rightarrow \infty} \sigma_{E \setminus E_n} \geq \zeta) = 1,$$

where $\sigma_{E \setminus E_n}$ is the first hitting time of $E \setminus E_n$. We denote by S^1 the set of smooth measures in the strict sense.

DEFINITION 3.1. Let $\mu \in S^1$.

(1) μ is said to be in the *Kato class* of X ($\mathcal{K}(X)$ in abbreviation) if

$$\lim_{\alpha \rightarrow \infty} \|R_\alpha \mu\|_\infty = 0.$$

μ is said to be in the *local Kato class* ($\mathcal{K}_{loc}(X)$ in abbreviation) if for any compact set K , $1_K \cdot \mu$ belongs to $\mathcal{K}(X)$. (2) Suppose that X is transient. A measure μ is said to be in the class $\mathcal{K}_\infty(X)$ if for any $\epsilon > 0$, there exists a compact set $K = K(\epsilon)$

$$\sup_{x \in E} \int_{K^c} R(x, y) \mu(dy) < \epsilon.$$

μ in $\mathcal{K}_\infty(X)$ is called *Green-tight*.

REMARK 3.2. It is known in [19, Theorem 3.1] that for a measure μ in $\mathcal{K}(X)$ and $\alpha > 0$

$$(17) \quad \int_E u^2 d\mu \leq \|R_\alpha \mu\|_\infty \mathcal{E}_\alpha(u), \quad u \in \mathcal{D}(\mathcal{E}).$$

By the regularity of $(\mathcal{E}, \mathcal{D}(\mathcal{E}))$ and the inequality (17), a measure μ in $\mathcal{K}(X)$ is Radon, and so is a measure μ in $\mathcal{K}_{loc}(X)$. As a result, $\mathcal{D}(\mathcal{E}) \cap L^2(E; \mu^+) \subset \mathcal{D}(\mathcal{E}^\mu)$ and

$$\mathcal{E}^\mu(u) = \mathcal{E}(u) + \int_E u^2 d\mu, \quad u \in \mathcal{D}(\mathcal{E}) \cap L^2(E; \mu^+).$$

If $\mu \in \mathcal{K}_\infty(X)$, then $\|R\mu\|_\infty < \infty$ by [3, Proposition 2.2] and [11, Lemma 4.1], and the equation (17) is meaningful for $\alpha = 0$:

$$(18) \quad \int_E u^2 d\mu \leq \|R\mu\|_\infty \mathcal{E}(u), \quad u \in \mathcal{D}_e(\mathcal{E}).$$

We denote by A_t^μ the PCAF corresponding to $\mu \in \mathcal{S}^1$.

Theorem 3.3 ([20, Theorem 4.2]). *Let $\mu = \mu^+ - \mu^- \in \mathcal{K}_{loc}(X) - \mathcal{K}_{loc}(X)$. If $(\mathcal{E}^\mu, \mathcal{D}(\mathcal{E}) \cap C_0(E))$ is positive semi-definite, then it is closable. Moreover, the semigroup T_t^μ generated by the closure $(\mathcal{E}^\mu, \mathcal{D}(\mathcal{E}^\mu))$ is expressed as*

$$T_t^\mu f(x) = p_t^\mu f(x) = E_x(e^{-A_t^\mu} f(X_t)) \quad m\text{-a.e.}$$

REMARK 3.4. By [9, Theorem 4.2.4], the transition semigroup p_t of X is expressed using transition probability density $p_t(x, y)$, as a result, T_t^μ is also expressed by a kernel $p_t^\mu(x, y)$ by Theorem 3.3. Hence, as discussed in Remark 2.3, $(\mathcal{E}^\mu, \mathcal{D}(\mathcal{E}^\mu))$ is either critical or subcritical.

4. Criticality and Hardy-type inequalities

We maintain the setting in Section 3 and fix a measure $\mu \in \mathcal{K}_{loc}(X) - \mathcal{K}_{loc}(X)$. Though this section, we assume that $(\mathcal{E}^\mu, \mathcal{D}(\mathcal{E}^\mu))$ is positive semi-definite and subcritical. By the subcriticality of $(\mathcal{E}^\mu, \mathcal{D}(\mathcal{E}^\mu))$, $(\mathcal{D}_e(\mathcal{E}^\mu), \mathcal{E}^\mu(\cdot, \cdot))$ becomes a Hilbert space. The α -order resolvent kernel $\{R_\alpha^\mu(x, y)\}_{\alpha>0}$ of $(\mathcal{E}^\mu, \mathcal{D}(\mathcal{E}^\mu))$ can be constructed in the same manner as [9, Lemma 4.2.4] and the Green kernel, i.e., 0-order resolvent kernel $R^\mu(x, y)$ is defined by $R^\mu(x, y) = \lim_{\alpha \rightarrow 0} R_\alpha^\mu(x, y)$. The potential of a positive measure ν is defined by

$$R^\mu \nu(x) = \int_E R^\mu(x, y) \nu(dy).$$

Lemma 4.1. *Let ν be a non-trivial positive measure in $\mathcal{K}_{loc}(X)$. Then for any compact set K*

$$\inf_{x \in K} R^\mu \nu(x) > 0.$$

Proof. For any compact set K , take a relatively compact domain G such that $K \subset G$ and $\nu(G) > 0$. Consider the subprocess $X^{\mu^+} = (\{P_x^{\mu^+}\}_{x \in E}, \{X_t\}_{t \geq 0}, \zeta)$ defined by

$$P_x^{\mu^+}(B; t < \zeta) = \int_{B \cap \{t < \zeta\}} e^{-A_t^{\mu^+}} dP_x, \quad B \in \mathcal{F}_t.$$

Then X^{μ^+} has Properties **(F)** and **(RSF)** by [13, Corollary 6.1], and so the part process $X^{\mu^+, G}$ of X^{μ^+} on G has Property **(RSF)** by [13, Theorem 3.1]. Furthermore, $X^{\mu^+, G}$ is irreducible

because G is a domain.

Since the measure ν^G , the restriction of ν to G , is in the Green-tight Kato class of $X^{\mu^+, G}$, $\nu^G \in \mathcal{K}_\infty(X^{\mu^+, G})$, $R^{\mu^+, G}\nu (= R^{\mu^+, G}\nu^G)$ is bounded by [3, Proposition 2.4] on G . Moreover it is continuous on G . Indeed, by Property (RSF) of $X^{\mu^+, G}$, $R_\alpha^{\mu^+, G}(R^{\mu^+, G}\nu) \in C_b(G)$ and $\|R_\alpha^{\mu^+, G}\nu\|_\infty \rightarrow 0$ as $\alpha \rightarrow \infty$ because of $\nu^G \in \mathcal{K}(X^{\mu^+, G})$. Hence, $R^{\mu^+, G}\nu \in C_b(G)$ because the resolvent equation implies

$$\|R^{\mu^+, G}\nu - \alpha R_\alpha^{\mu^+, G}(R^{\mu^+, G}\nu)\|_\infty = \|R_\alpha^{\mu^+, G}\nu\|_\infty \rightarrow 0, \alpha \rightarrow \infty.$$

By the irreducibility and $\nu(G) > 0$, $R^{\mu^+, G}\nu(x) > 0$ for each $x \in E$, and thus $\inf_{x \in K} R^{\mu^+, G}\nu(x) > 0$. On account of $R^\mu\nu(x) \geq R^{\mu^+, G}\nu(x)$, we have this lemma. \square

By Lemma 4.1, we have the next corollary.

Corollary 4.2. *For a non-trivial positive measure $\nu \in \mathcal{K}_{loc}(X)$, the measure $\nu/R^\mu\nu$ belongs to $\mathcal{K}_{loc}(X)$.*

We define the subclass $\mathcal{K}_{loc}^\mu(X)$ of $\mathcal{K}_{loc}(X)$ by

$$\mathcal{K}_{loc}^\mu(X) = \{\nu \in \mathcal{K}_{loc}(X) \mid \text{For any } K \subset C, \|R^\mu(1_K\nu)\|_\infty < \infty.\},$$

where C is the totality of compact sets of E . If $\mu = 0$, then $\mathcal{K}_{loc}^\mu(X)$ equals $\mathcal{K}_{loc}(X)$ because $1_K\nu \in \mathcal{K}_\infty(X)$ and $\|R(1_K\nu)\|_\infty < \infty$.

Lemma 4.3. *Let ν be a non-trivial measure in $\mathcal{K}_{loc}^\mu(X)$. Then*

$$\int_E \phi^2 \frac{d\nu}{R^\mu\nu} \leq \mathcal{E}^\mu(\phi), \quad \phi \in \mathcal{D}(\mathcal{E}) \cap C_0(E).$$

Proof. Let $\{K_n\}$ be a increasing sequence of compact sets such that $K_n \subset \overset{\circ}{K}_{n+1}$ and $K_n \uparrow E$. We fix the sequence $\{K_n\}$. For $0 < \epsilon < 1$, define $\mu_n^\epsilon = \mu^+ - \epsilon\mu_n^-$, where $\mu_n^-(\cdot) := \mu^-(K_n \cap \cdot)$. The positive semi-definiteness of $(\mathcal{E}^\mu, \mathcal{D}(\mathcal{E}) \cap C_0(E))$ implies that

$$\epsilon \int_E \phi^2 d\mu_n^- \leq \epsilon \mathcal{E}^{\mu^+}(\phi),$$

and

$$(19) \quad (1 - \epsilon) \mathcal{E}^{\mu^+}(\phi) \leq \mathcal{E}^{\mu_n^\epsilon}(\phi) - \epsilon \int_E \phi^2 d\mu_n^- = \mathcal{E}^{\mu_n^\epsilon}(\phi) \leq \mathcal{E}^{\mu^+}(\phi),$$

which implies

$$(20) \quad \mathcal{D}_\epsilon(\mathcal{E}^{\mu_n^\epsilon}) = \mathcal{D}_\epsilon(\mathcal{E}^{\mu^+}) \subset \mathcal{D}_\epsilon(\mathcal{E}).$$

Let $\nu_m = \nu(\cdot \cap K_m)$. We may suppose that ν_1 is non-trivial and $R^{\mu_n^\epsilon}\nu_1(x)$ is bounded below by a positive constant on each compact set $K \subset E$. Noting $\nu_m \in \mathcal{K}_\infty(X)$, we see from (18) and (19) that

$$\begin{aligned} \int_E |\phi| d\nu_m &\leq \nu(K_m)^{1/2} \left(\int_E \phi^2 d\nu_m \right)^{1/2} \leq \nu(K_m)^{1/2} \|R\nu_m\|_\infty^{1/2} \cdot \mathcal{E}(\phi)^{1/2} \\ &\leq C \mathcal{E}^{\mu^+}(\phi)^{1/2} \leq C' \mathcal{E}^{\mu_n^\epsilon}(\phi)^{1/2}. \end{aligned}$$

Hence $R^{\mu_n^\epsilon}\nu_m$ belongs to $\mathcal{D}_\epsilon(\mathcal{E}^{\mu_n^\epsilon})$ and

$$\mathcal{E}^{\mu_n^\epsilon}(R^{\mu_n^\epsilon}v_m, \phi) = \int_E \phi d\nu_m = \int_E R^{\mu_n^\epsilon}v_m \cdot \phi \frac{d\nu_m}{R^{\mu_n^\epsilon}v_m},$$

which implies

$$\mathcal{E}^{\mu_n^\epsilon - \nu_m / R^{\mu_n^\epsilon}v_m}(R^{\mu_n^\epsilon}v_m, \phi) = 0, \quad \phi \in \mathcal{D}(E) \cap C_0(E).$$

Note that $R^{\mu_n^\epsilon}v_m$ is in $\mathcal{D}_e(\mathcal{E})$ by (20) and in $L^\infty(E, m)$ by $R^{\mu_n^\epsilon}v_m \leq R^\mu v_m$. Moreover, it is bounded below by a positive constant on each compact set by Lemma 4.1. We then see from Lemma 4.5 and Lemma 4.6 below that

$$\mathcal{E}^{\mu_n^\epsilon - \nu_m / R^{\mu_n^\epsilon}v_m}(\phi) \geq 0, \quad \phi \in \mathcal{D}(\mathcal{E}) \cap C_0(E),$$

and

$$\mathcal{E}^{\mu_n^\epsilon}(\phi) - \int_E \phi^2 \frac{d\nu_m}{R^{\mu_n^\epsilon}v_m} \geq \mathcal{E}^{\mu_n^\epsilon}(\phi) - \int_E \phi^2 \frac{d\nu_m}{R^{\mu_n^\epsilon}v_m} = \mathcal{E}^{\mu_n^\epsilon - \nu_m / R^{\mu_n^\epsilon}v_m}(\phi) \geq 0.$$

Since

$$\begin{aligned} \mathcal{E}^{\mu_n^\epsilon}(\phi) - \int_E \phi^2 \frac{d\nu_m}{R^{\mu_n^\epsilon}v_m} &\xrightarrow{m \rightarrow \infty} \mathcal{E}^{\mu_n^\epsilon}(\phi) - \int_E \phi^2 \frac{d\nu}{R^{\mu_n^\epsilon}v} \\ &\xrightarrow{\epsilon \rightarrow 1} \mathcal{E}^{\mu_n^1}(\phi) - \int_E \phi^2 \frac{d\nu}{R^{\mu_n^1}v} \\ &\xrightarrow{n \rightarrow \infty} \mathcal{E}^\mu(\phi) - \int_E \phi^2 \frac{d\nu}{R^\mu v}, \end{aligned}$$

we have this lemma. □

Lemma 4.3 leads us to an extension of the inequality (17).

Corollary 4.4. *It holds that*

$$\int_E \phi^2 d\nu \leq \|R^\mu v\|_\infty \mathcal{E}^\mu(\phi), \quad \phi \in \mathcal{D}(\mathcal{E}) \cap C_0(E).$$

Lemma 4.5. *Let $u \in \mathcal{D}_e(\mathcal{E}) \cap L^\infty(E; m)$ is bounded below by a positive constant on each compact set. Then φ/u belongs to $\mathcal{D}(\mathcal{E})$ for any $\varphi \in \mathcal{D}(\mathcal{E}) \cap C_0(E)$.*

Proof. Let $\varphi \in \mathcal{D}(\mathcal{E}) \cap C_0(E)$ and suppose that $u \geq c > 0$ on $\text{supp}[\varphi]$. Let $\{u_n\} \subset \mathcal{D}(\mathcal{E}) \cap C_0(E)$ be an approximating sequence of u . We may suppose $\sup_n \|u_n\|_\infty \leq \|u\|_\infty$. Then since by [9, Theorem 1.4.2 (ii)]

$$\mathcal{E}(u_n \varphi)^{1/2} \leq \|u_n\|_\infty \mathcal{E}(\varphi)^{1/2} + \|\varphi\|_\infty \mathcal{E}(u_n)^{1/2},$$

we have $\sup_n \mathcal{E}(u_n \varphi) < \infty$. On account of [18, 1.6.1'], $u\varphi$ is in $\mathcal{D}_e(\mathcal{E})$ and so in $\mathcal{D}(\mathcal{E})$ because $\mathcal{D}_e(\mathcal{E}) \cap L^2(E; m) = \mathcal{D}(\mathcal{E})$.

Since for $(x, y) \in \text{supp}[\varphi] \times \text{supp}[\varphi]$

$$\begin{aligned} \left| \frac{\varphi(x)}{u(x)} \right| &\leq c^{-1} |\varphi(x)| \\ \left| \frac{\varphi(x)}{u(x)} - \frac{\varphi(y)}{u(y)} \right| &\leq 2c^{-1} |\varphi(x) - \varphi(y)| + c^{-2} |u(x)\varphi(x) - u(y)\varphi(y)|, \end{aligned}$$

we have this lemma by the same argument as in the proof of [9, Theorem 6.3.2]. □

[8, Theorem 10.2] yields the next lemma.

Lemma 4.6. *Let $\mu = \mu^+ - \mu^- \in \mathcal{K}_{loc}(X) - \mathcal{K}(X)$ and $u \in \mathcal{D}_e(\mathcal{E}) \cap L^\infty(E; m)$ be a function bounded below by a positive constant on each compact. If u satisfies $\mathcal{E}^\mu(u, \varphi) = 0$ for any $\varphi \in \mathcal{D}(\mathcal{E}) \cap C_0(E)$, then $(\mathcal{E}^\mu, \mathcal{D}(\mathcal{E}^\mu))$ is positive semi-definite.*

Proof. The function u is a *generalized eigenfunction* corresponding to the *generalized eigenvalue* 0 in [8, Definition 9.1]. Note that by Lemma 4.5, φ/u is a bounded function in $\mathcal{D}(\mathcal{E}^\mu)$ with compact support. Then, applying [8, Theorem 10.2], we have

$$\mathcal{E}^\mu(\varphi) = \mathcal{E}^\mu(u(\varphi/u)) = \int_{E \times E} u(x)u(y)d\Gamma(\varphi/u) \geq 0,$$

where $\Gamma(\varphi/u)$ is the positive measure on $E \times E$ defined in [8, Subsection 3.2]. □

Lemma 4.7. *Let $\nu \in \mathcal{K}_{loc}^\mu(X)$ and $\nu_m = \nu(\cdot \cap K_m)$. Then $R^\mu \nu_m$ belongs to $\mathcal{D}_e(\mathcal{E}^\mu)$ for any m .*

Proof. Since for $\phi \in \mathcal{D}(\mathcal{E}) \cap C_0(E)$

$$\int_E |\phi| d\nu_m \leq \nu(K_m)^{1/2} \left(\int_E \phi^2 d\nu_m \right)^{1/2} \leq \mu(K_m)^{1/2} \|R^\mu \nu_m\|_\infty^{1/2} \mathcal{E}^\mu(\phi)^{1/2}$$

by Corollary 4.4 and $\|R^\mu \nu_m\|_\infty < \infty$ by $\nu \in \mathcal{K}_{loc}^\mu(X)$, we have this lemma. □

Lemma 4.8. *If $\nu \in \mathcal{K}_{loc}^\mu(X)$ is of finite energy with respect to $R^\mu(x, y)$,*

$$(21) \quad \iint_{E \times E} R^\mu(x, y)\nu(dx)\nu(dy) < \infty,$$

then $R^\mu \nu$ belongs to $\mathcal{D}_e(\mathcal{E}^\mu)$.

Proof. Since $R^\mu \nu_m \in \mathcal{D}_e(\mathcal{E}^\mu) \uparrow R^\mu \nu(x)$ for any $x \in E$ as $m \rightarrow \infty$ and

$$\begin{aligned} \sup_m \mathcal{E}^\mu(R^\mu \nu_m) &= \sup_m \int_E R^\mu \nu_m d\nu_m = \sup_m \iint_{K_m \times K_m} R^\mu(x, y)\nu(dx)\nu(dy) \\ &\leq \iint_{E \times E} R^\mu(x, y)\nu(dx)\nu(dy) < \infty. \end{aligned}$$

By Banach-Saks Theorem (cf.[4, Theorem A.4.1]) there exists a subsequence $\{K_{m_l}\} \subset \{K_m\}$ such that

$$\frac{R^\mu \nu_{m_1} + R^\mu \nu_{m_2} + \cdots + R^\mu \nu_{m_l}}{l} = R^\mu \left(\frac{(1_{K_{m_1}} + 1_{K_{m_2}} \cdots + 1_{K_{m_l}})}{l} \nu \right) \longrightarrow R^\mu \nu$$

with \mathcal{E}^μ -strongly, and thus Lemma 4.7 implies this lemma. □

For $\mu \in \mathcal{K}_{loc}(X) - \mathcal{K}_{loc}(X)$ and $\nu \in \mathcal{K}_{loc}^\mu(X)$, define

$$(22) \quad \nu^\mu = \frac{\nu}{R^\mu \nu}, \quad \mu^\nu = \mu - \nu^\mu.$$

Then μ^ν is in $\mathcal{K}_{loc}(X) - \mathcal{K}_{loc}(X)$ by Corollary 4.2 and $(\mathcal{E}^{\mu^\nu}, \mathcal{D}(\mathcal{E}) \cap C_0(E))$ is positive semi-definite by Lemma 4.3. Hence by [20, Theorem 4.2] we can define the Schrödinger form with potential μ^ν , the closure $(\mathcal{E}^{\mu^\nu}, \mathcal{D}(\mathcal{E}^{\mu^\nu}))$ of $(\mathcal{E}^{\mu^\nu}, \mathcal{D}(\mathcal{E}) \cap C_0(E))$ and its extended Schrödinger space $\mathcal{D}_e(\mathcal{E}^{\mu^\nu})$.

Lemma 4.9. *If $u \in \mathcal{D}_e(\mathcal{E}^{\mu^+})$, then*

$$\mathcal{E}^{\mu^y}(u) = \mathcal{E}^{\mu}(u) - \int_E u^2 d\nu^{\mu}.$$

Proof. Noting $u \in \mathcal{D}_e(\mathcal{E})$, there exists an \mathcal{E}^{μ^+} -Cauchy sequence $\{u_n\} \subset \mathcal{D}(\mathcal{E}) \cap C_0(E)$ such that $u_n \rightarrow u$ q.e. Since $\mathcal{E}^{\mu^y}(u) \leq \mathcal{E}^{\mu}(u) \leq \mathcal{E}^{\mu^+}(u)$, $u \in \mathcal{D}(\mathcal{E}) \cap C_0(E)$, $\{u_n\}$ is also an approximating sequence of u in $\mathcal{D}_e(\mathcal{E}^{\mu})$ and $\mathcal{D}_e(\mathcal{E}^{\mu^y})$. In particular, u is in $\mathcal{D}_e(\mathcal{E}^{\mu}) \subset \mathcal{D}_e(\mathcal{E}^{\mu^y})$, and thus $u \in L^2(E; \nu^{\mu})$ by Lemma 4.3. Hence we have

$$\mathcal{E}^{\mu^y}(u) = \lim_{n \rightarrow \infty} \mathcal{E}^{\mu^y}(u_n) = \lim_{n \rightarrow \infty} \left(\mathcal{E}^{\mu}(u_n) - \int_E u_n^2 d\nu^{\mu} \right) = \mathcal{E}^{\mu}(u) - \int_E u^2 d\nu^{\mu}. \quad \square$$

Lemma 4.10. *It holds that*

$$\mathcal{E}^{\mu^y}(R^{\mu} \nu_m) = \mathcal{E}^{\mu}(R^{\mu} \nu_m) - \int_E (R^{\mu} \nu_m)^2 d\nu^{\mu}.$$

Proof. Let $\{\epsilon_n\}$ be a positive sequence such that $\epsilon_n \uparrow 1$ as $n \rightarrow \infty$ and denote by μ'_n the measure $\mu_n^{\epsilon_n}$ defined in Lemma 4.3. Put $u_n = R^{\mu'_n} \nu_m$. Then u_n is in $\mathcal{D}_e(\mathcal{E}^{\mu^+})$ as shown in the proof of Lemma 4.3. Since

$$\mathcal{E}^{\mu}(u_n) \leq \mathcal{E}^{\mu'_n}(u_n) = \int_E u_n d\nu_m \leq \int_E R^{\mu} \nu_m d\nu_m < \infty,$$

There exists a subsequence $\{u_{n_k}\}$ of $\{u_n\}$ such that

$$v_k := \frac{u_{n_1} + u_{n_2} + \cdots + u_{n_k}}{k} \in \mathcal{D}_e(\mathcal{E}^{\mu^+})$$

is an approximating sequence of $R^{\mu} \nu_m$ in $\mathcal{D}_e(\mathcal{E}^{\mu})$ and $v_k(x) \uparrow R^{\mu} \nu_m(x)$ for any $x \in E$.

Noting that $\{v_k\}$ is also an approximating sequence of $R^{\mu} \nu_m$ in $\mathcal{D}_e(\mathcal{E}^{\mu^y})$, we have by Lemma 4.9

$$\mathcal{E}^{\mu^y}(R^{\mu} \nu_m) = \lim_{k \rightarrow \infty} \mathcal{E}^{\mu^y}(v_k) = \lim_{k \rightarrow \infty} \left(\mathcal{E}^{\mu}(v_k) - \int_E v_k^2 d\nu^{\mu} \right) = \mathcal{E}^{\mu}(R^{\mu} \nu_m) - \int_E (R^{\mu} \nu_m)^2 d\nu^{\mu}. \quad \square$$

Let \mathcal{K}_C^{μ} be the set of measures in $\mathcal{K}_{loc}^{\mu}(X)$ satisfying (7). For $\nu \in \mathcal{K}_C^{\mu}$ there exists a sequence $\{K_m\}_{m=1}^{\infty} \subset C$ such that $K_m \uparrow E$ and

$$(23) \quad \sup_m \iint_{E \times E} R^{\mu}(x, y) \nu_m(dx) \nu_m^c(dy) < \infty,$$

where $\nu_m^c(A) = \nu(K_m^c \cap A)$. If a measures $\nu \in \mathcal{K}_{loc}^{\mu}(X)$ of finite energy with respect to R^{μ} , then it satisfies (23).

Lemma 4.11. *If $\nu \in \mathcal{K}_C^{\mu}$, then $R^{\mu} \nu$ is in $\mathcal{D}_e(\mathcal{E}^{\mu^y})$.*

Proof. For $\nu \in \mathcal{K}_C^{\mu}$

$$\int_E R^{\mu} \nu_m d\nu = \int_E R^{\mu} \nu_m d\nu_m + \int_E R^{\mu} \nu_m d\nu_m^c < \infty$$

because

$$\int_E R^\mu v_m d\nu_m = \mathcal{E}^\mu(R^\mu v_m) < \infty$$

by Lemma 4.7.

By Lemma 4.10 we have

$$\begin{aligned} \mathcal{E}^{\mu^\nu}(R^\mu v_m) &= \mathcal{E}^\mu(R^\mu v_m) - \int_E (R^\mu v_m)^2 d\nu^\mu \\ &= \int_E R^\mu v_m d\nu_m - \int_E (R^\mu v_m)^2 d\nu^\mu \\ &= \int_E R^\mu v_m d\nu - \int_E R^\mu v_m d\nu_m^c - \int_E \frac{(R^\mu v_m)^2}{R^\mu v_m + R^\mu v_m^c} d\nu. \end{aligned}$$

The right hand side equals

$$\begin{aligned} (24) \quad & \int_E \left(\frac{R^\mu v_m (R^\mu v_m + R^\mu v_m^c) - (R^\mu v_m)^2}{R^\mu v_m + R^\mu v_m^c} \right) d\nu - \int_E R^\mu v_m d\nu_m^c \\ &= \int_E \frac{R^\mu v_m R^\mu v_m^c}{R^\mu v_m + R^\mu v_m^c} d\nu - \int_E R^\mu v_m d\nu_m^c \\ &= \int_E \frac{R^\mu v_m R^\mu v_m^c}{R^\mu v_m + R^\mu v_m^c} d\nu_m + \int_E \left(\frac{R^\mu v_m R^\mu v_m^c}{R^\mu v_m + R^\mu v_m^c} - R^\mu v_m \right) d\nu_m^c. \end{aligned}$$

Since

$$\frac{R^\mu v_m R^\mu v_m^c}{R^\mu v_m + R^\mu v_m^c} \leq R^\mu v_m^c, \quad \frac{R^\mu v_m R^\mu v_m^c}{R^\mu v_m + R^\mu v_m^c} \leq R^\mu v_m,$$

the right hand side of (24) is less than or equal to $\int_E R^\mu v_m^c d\nu_m$. Therefore, we see from (23) that

$$\sup_m \mathcal{E}^{\mu^\nu}(R^\mu v_m) \leq \sup_m \int_E R^\mu v_m^c d\nu_m < \infty.$$

Since $R^\mu v_m \rightarrow R^\mu v$, this lemma follows from Lemma 4.7. □

The next lemma is obtained in the same argument as in [20, Lemma 5.3].

Lemma 4.12. For $v \in \mathcal{K}_C^\mu$

$$\mathcal{E}^{\mu^\nu}(R^\mu v, \varphi) = 0, \quad \varphi \in \mathcal{D}(\mathcal{E}) \cap C_0(E).$$

Proof. Since $\sup_m \mathcal{E}^{\mu^\nu}(R^\mu v_m) < \infty$, there exists a subsequence $\{K_{m_l}\} \subset \{K_m\}$ such that

$$R^\mu \left(\frac{(1_{K_{m_1}} + 1_{K_{m_2}} \cdots + 1_{K_{m_l}})}{l} v \right) \rightarrow R^\mu v$$

\mathcal{E}^{μ^ν} -strongly.

Let $\phi_l := (1_{K_{m_1}} + 1_{K_{m_2}} \cdots + 1_{K_{m_l}})/l$. For a fixed $\varphi \in \mathcal{D}(\mathcal{E}) \cap C_0(E)$ we can assume $\text{supp}[\varphi] \subset K_{m_1}$. By the same argument as in Lemma 4.10, we have

$$\mathcal{E}^{\mu^\nu}(R^\mu(\phi_l v) + \varphi) = \mathcal{E}^\mu(R^\mu(\phi_l v) + \varphi) - \int_E (R^\mu(\phi_l v) + \varphi)^2 d\nu^\mu,$$

and thus

$$\mathcal{E}^{\mu^\nu}(R^\mu(\phi_l\nu), \varphi) = \mathcal{E}^\mu(R^\mu(\phi_l\nu), \varphi) - \int_E R^\mu(\phi_l\nu)\varphi d\nu^\mu.$$

Hence

$$\begin{aligned} \mathcal{E}^{\mu^\nu}(R^\mu\nu, \varphi) &= \lim_{l \rightarrow \infty} \mathcal{E}^{\mu^\nu}(R^\mu(\phi_l\nu), \varphi) \\ &= \lim_{l \rightarrow \infty} \left(\mathcal{E}^\mu(R^\mu(\phi_l\nu), \varphi) - \int_E R^\mu(\phi_l\nu)\varphi d\nu^\mu \right). \end{aligned}$$

Note that $R^\mu(\phi_l\nu) \in D_e(\mathcal{E}^\mu)$ by Lemma 4.7. Then since

$$\lim_{l \rightarrow \infty} \mathcal{E}^\mu(R^\mu(\phi_l\nu), \varphi) = \lim_{l \rightarrow \infty} \int_E \varphi \phi_l d\nu = \int_E \varphi d\nu$$

and by the monotone convergence theorem

$$\lim_{l \rightarrow \infty} \int_E R^\mu(\phi_l\nu)\varphi d\nu^\mu = \int_E R^\mu\nu \cdot \varphi \frac{d\nu}{R^\mu\nu} = \int_E \varphi d\nu,$$

we have this lemma. \square

The next theorem is an extension of [20, Theorem 5.4].

Theorem 4.13. *If $\nu \in \mathcal{K}_C^\mu$, then $R^\mu\nu$ is a ground state of $(\mathcal{E}^{\mu^\nu}, \mathcal{D}(\mathcal{E}^{\mu^\nu}))$, consequently, $(\mathcal{E}^{\mu^\nu}, \mathcal{D}(\mathcal{E}^{\mu^\nu}))$ is critical.*

Proof. Since $R^\mu\nu$ belongs to $D_e(\mathcal{E}^{\mu^\nu})$, there exists a sequence $\{\varphi_n\} \subset \mathcal{D}(\mathcal{E}) \cap C_0(E)$ such that φ_n converges \mathcal{E}^{μ^ν} -strongly to $R^\mu\nu$. Hence

$$\mathcal{E}^{\mu^\nu}(R^\mu\nu) = \lim_{n \rightarrow \infty} \mathcal{E}^{\mu^\nu}(R^\mu\nu, \varphi_n) = 0$$

by Lemma 4.12. \square

Corollary 4.14. *There exists no non-trivial positive function ψ such that*

$$(25) \quad \int_E u^2 d(\nu^\mu + \psi m) \leq \mathcal{E}^\mu(u, u), \quad u \in \mathcal{D}(\mathcal{E}) \cap C_0(E).$$

Proof. If (25) holds, then

$$\int_E u^2 \psi dm \leq \mathcal{E}^{\mu^\nu}(u) = 0, \quad u \in \mathcal{D}(\mathcal{E}) \cap C_0(E).$$

Since $R^\mu\nu$ is in $D_e(\mathcal{E}^{\mu^\nu})$, there exists an approximating sequence $\{u_n\} \subset \mathcal{D}(\mathcal{E}) \cap C_0(E)$. We then have

$$\int_E (R^\mu\nu)^2 \psi dm \leq \liminf_{n \rightarrow \infty} \int_E u_n^2 \psi dm \leq \liminf_{n \rightarrow \infty} \mathcal{E}^{\mu^\nu}(u_n) = \mathcal{E}^{\mu^\nu}(R^\mu\nu) = 0,$$

and so $\psi = 0$ m -a.e. because $R^\mu\nu > 0$ by the irreducibility of X . \square

Corollary 4.14 tells us that ν^μ is a *critical Hardy weight* for $(\mathcal{E}^\mu, \mathcal{D}(\mathcal{E}^\mu))$ ([6], [10]).

A Hardy weight ν^μ is called *optimal at infinity* if for any $K \in \mathcal{C}$

$$\lambda \int_E u^2 d\nu^\mu \leq \mathcal{E}^\mu(u), \quad u \in \mathcal{D}(\mathcal{E}) \cap C_0(K^c),$$

then $\lambda \leq 1$.

Lemma 4.15. *If $\nu \in \mathcal{K}_C^\mu$ satisfies that*

$$(26) \quad \iint_{K^c \times E} R^\mu(x, y) \nu(dx) \nu(dy) = \infty \text{ for any } K \in \mathcal{C},$$

then ν^μ is optimal at infinity.

Proof. Denote $h = R^\mu \nu$. Since h is a ground state of $(\mathcal{E}^{\mu^\nu}, \mathcal{D}(\mathcal{E}^{\mu^\nu}))$ by Theorem 4.13, h is $p_t^{\mu^\nu}$ -invariant, $p_t^{\mu^\nu} h = h$, where $p_t^{\mu^\nu}$ is the semigroup associated with $(\mathcal{E}^{\mu^\nu}, \mathcal{D}(\mathcal{E}^{\mu^\nu}))$. Denote by $(\mathcal{E}^h, \mathcal{D}(\mathcal{E}^h))$ the Dirichlet form generated by h -transform of $(\mathcal{E}^{\mu^\nu}, \mathcal{D}(\mathcal{E}^{\mu^\nu}))$:

$$\mathcal{E}^h(u) = \mathcal{E}^{\mu^\nu}(uh), \quad u \in \mathcal{D}(\mathcal{E}^h) = \{u \mid uh \in \mathcal{D}(\mathcal{E}^{\mu^\nu})\}.$$

Since h is in $\mathcal{D}_e(\mathcal{E}^{\mu^\nu})$, there exists a sequence $\{h_n\} \subset \mathcal{D}(\mathcal{E}) \cap C_0(E)$ such that $0 \leq h_n \uparrow h$ and $\mathcal{E}^{\mu^\nu}(h - h_n) \rightarrow 0$ as $n \rightarrow \infty$. Then $\{g_n := h_n/h\}$ is an approximating sequence of $1 \in \mathcal{D}_e(\mathcal{E}^h)$.

Suppose that there exist $F \in \mathcal{C}$ and $\epsilon > 0$ such that for any $u \in \mathcal{D}(\mathcal{E}) \cap C_0(F^c)$

$$(27) \quad \mathcal{E}^\mu(u) \geq (1 + \epsilon) \int_{F^c} u^2 d\nu^\mu, \quad u \in \mathcal{D}(\mathcal{E}) \cap C_0(F^c).$$

Let G_1, G_2 be relatively compact open set such that $F \subset G_1 \subset \overline{G_1} \subset G_2 \subset \overline{G_2} \subset E$. Let φ be a function in $\mathcal{D}(\mathcal{E}) \cap C_0(E)$ such that $0 \leq \varphi \leq 1$, $\varphi(x) = 1$ on $x \in \overline{G_1}$ and $\text{supp}[\varphi] \subset G_2$. Put $\psi = (1 - \varphi)$. Then $h_n \psi \in \mathcal{D}(\mathcal{E}) \cap C_0(F^c)$, and so by (27)

$$(28) \quad \epsilon \int_E (h_n \psi)^2 \frac{d\nu}{h} \leq \mathcal{E}^{\mu^\nu}(h_n \psi).$$

Then we have by [9, Theorem 1.4.2 (ii)]

$$\mathcal{E}^{\mu^\nu}(h_n \psi) = \mathcal{E}^h\left(\frac{h_n}{h} \psi\right) \leq 2 \left(\mathcal{E}^h(h_n/h) + \mathcal{E}^h(\psi) \right),$$

and so

$$\sup_n \int_E (h_n \psi)^2 \frac{d\nu}{h} \leq \frac{2}{\epsilon} \left(\sup_n \mathcal{E}^h(h_n/h) + \mathcal{E}^h(\psi) \right) < \infty$$

on account of (28). Hence

$$\int_{\overline{G_2}^c} h d\nu = \int_{\overline{G_2}^c} \lim_{n \rightarrow \infty} (h_n \psi)^2 \frac{d\nu}{h} \leq \lim_{n \rightarrow \infty} \int_E (h_n \psi)^2 \frac{d\nu}{h} < \infty,$$

and thus

$$\iint_{\overline{G_2}^c \times E} R^\mu(x, y) d\nu(x) d\nu(y) = \int_{\overline{G_2}^c} h d\nu < \infty,$$

which is contradictory to (26). □

If $\nu \in \mathcal{K}_C^\mu$ satisfies the inequality (26), then the ground state $R^\mu \nu$ of $(\mathcal{E}^{\mu^\nu}, \mathcal{D}(\mathcal{E}^{\mu^\nu}))$ does not belong to $L^2(E; \mu^\nu)$ and so ν^μ is a null-critical Hardy weight for $(\mathcal{E}^\mu, \mathcal{D}(\mathcal{E}^\mu))$. Therefore, we have

Theorem 4.16. *If $\nu \in \mathcal{K}_C^\mu$ satisfies*

$$\iint_{K^c \times E} R^\mu(x, y) \nu(dx) \nu(dy) = \infty \text{ for any } K \in \mathcal{C},$$

then the measure ν^μ defined in (22) is a optimal Hardy weight for $(\mathcal{E}^\mu, \mathcal{D}(\mathcal{E}^\mu))$.

REMARK 4.17. The measure $\nu(dx) := |x|^{-(d+\alpha)/2} dx$ satisfies (26) with respect to the Green kernel $|x - y|^{\alpha-d}$, $\alpha < d$, the 0-resolvent of the symmetric α -stable process because $(|y|^{\alpha-d} * |y|^{-(d+\alpha)/2})(x) = C|x|^{(\alpha-d)/2}$ and $|x|^{(\alpha-d)/2} \cdot |x|^{-(d+\alpha)/2} = |x|^{-d}$; however ν satisfies (23) ([20, Example 5.6]). Hence ν is an optimal Hardy weight for the Dirichlet form of symmetric α -stable process.

ACKNOWLEDGEMENTS. The author would like to thank the referee for a useful comment on Lemma 4.15.

References

- [1] S. Albeverio and Z.-M. Ma: *Perturbation of Dirichlet forms-lower semiboundedness, closability, and form cores*, J. Funct. Anal. **99** (1991), 332–356.
- [2] J. Cao, A. Grigor'yan and L. Liu: *Hardy's inequality and Green function on metric measure spaces*, J. Funct. Anal. **281** (2021), Paper No. 109020, 78 pp.
- [3] Z.-Q. Chen: *Gaugeability and conditional gaugeability*, Trans. Amer. Math. Soc. **354** (2002), 4639–4679.
- [4] Z.-Q. Chen and M. Fukushima: *Symmetric Markov Processes, Time Change, and Boundary Theory*, London Math. Soc. Monographs Series **35**, Princeton University Press, Princeton, 2012.
- [5] E.B. Davies: *Heat Kernels and Spectral Theory*, Cambridge Univ. Press, Cambridge, 1989.
- [6] B. Devyver, M. Fraas and Y. Pinchover: *Optimal Hardy weight for second-order elliptic operator: an answer to a problem of Agmon*, J. Funct. Anal. **266** (2014), 4422–4489.
- [7] P.J. Fitzsimmons: *Hardy's inequality for Dirichlet forms*, J. Math. Anal. Appl. **250** (2000), 548–560.
- [8] R.L. Frank, D. Lenz and D. Wingert: *Intrinsic metrics for non-local symmetric Dirichlet forms and applications to spectral theory*, J. Funct. Anal. **266** (2014), 4765–4808.
- [9] M. Fukushima, Y. Oshima and M. Takeda: *Dirichlet Forms and Symmetric Markov Processes*, 2nd ed., Walter de Gruyter, Berlin, 2011.
- [10] M. Keller, Y. Pinchover and F. Pogorzelski: *Criticality theory for Schrödinger operators on graphs*, J. Spectr. Theory **10** (2020), 73–114.
- [11] D. Kim and K. Kuwae: *Analytic characterizations of gaugeability for generalized Feynman-Kac functionals*, Trans. Amer. Math. Soc. **369** (2017), 4545–4596.
- [12] H. Kovarik and Y. Pinchover: *On minimal decay at infinity of Hardy-weights*, Commun. Contemp. Math. **22** (2020), Paper No. 1950046, 18 pp.
- [13] M. Kurniawaty, K. Kuwae and K. Tsuchida: *On the doubly Feller property of resolvent*, Kyoto J. Math. **57** (2017), 637–654.
- [14] Y. Miura: *Optimal Hardy inequalities for Schrödinger operators based on symmetric stable processes*, J. Theoret. Probab. **36** (2023), 134–166.
- [15] M. Schmidt: *(Weak) Hardy and Poincaré inequalities and criticality theory*, Dirichlet forms and related topics, Springer Proc. Math. Stat. **394** Springer, Singapore, 2022, 421–460.
- [16] B. Schmuland: *Extended Dirichlet spaces*, C. R. Math. Acad. Sci. Soc. R. Can. **21** (1999), 146–152.
- [17] B. Schmuland: *Positivity preserving forms have the Fatou property*, Potential Anal. **10** (1999), 373–378.
- [18] M. Silverstein: *Symmetric Markov processes*, Lecture Notes in Mathematics **426**, Springer-Verlag, Berlin-New York, 1974.
- [19] P. Stollmann and J. Voigt: *Perturbation of Dirichlet forms by measures*, Potential Anal. **5** (1996), 109–138.

- [20] M. Takeda and T. Uemura: *Criticality of Schrödinger forms and recurrence of Dirichlet forms*, Trans. Amer. Math. Soc. **376** (2023), 4145–4171.

Department of Mathematics
Kansai University
Yamatecho, Suita, 564–8680
Japan
e-mail: mtakeda@kansai-u.ac.jp