

Title	Identification of a novel arthritis-associated osteoclast precursor macrophage regulated by FoxM1			
Author(s)	Hasegawa, Tetsuo; Kikuta, Junichi; Sudo, Takao et al.			
Citation	Nature Immunology. 2019, 20(12), p. 1631–1643			
Version Type	АМ			
URL	https://hdl.handle.net/11094/93182			
rights	© 2019, The Author(s), under exclusive licence to Springer Nature America, Inc.			
Note				

The University of Osaka Institutional Knowledge Archive : OUKA

https://ir.library.osaka-u.ac.jp/

The University of Osaka

1 Supplementary information



- 3 Supplementary Figure 1. Protocol for isolating pannus (hypertrophied
- 4 synovium) in arthritic mice

1	a, Schematic diagram showing that hypertrophied synovium exists behind the
2	patellar ligament and on the "bare area" of the femur, where bone is exposed to
3	the synovium without a cartilage covering. Pa; patella, Fe; femur, Ti; tibia. The
4	red lesion indicates the hypertrophied synovium on the bare area.
5	b , After removal of the biceps femoris muscle, the quadriceps femoris muscles
6	including the vastus intermedius muscle were pinched and lifted with tweezers.
7	The quadriceps femoris muscles and patellar ligament, including the patella,
8	were isolated from the knee joint under a stereoscopic microscope. The
9	hypertrophied synovium is visible on bare areas of the femur (arrowheads) and
10	isolated without damaging the bone.
11	c, Schematic diagram showing that hypertrophied synovium exists behind the
12	Achilles tendon. Ti; Tibia, Ta; Talus, Cal; Calcaneus. The red lesion indicates the
13	hypertrophied synovium.
14	d, After removal of the ankle joint tendons, including the Achilles tendon, the
15	hypertrophied synovium can be detected around the talus (arrowheads) and
16	isolated without damaging the bone.



harmonic generation (SHG) and cell counting by Imaris software. Bar, 70 µm.

1	b , Immunohistochemistry of CX ₃ CR1-EGFP transgenic mice healthy knee joints
2	stained with Ab against CD11b (BV421). Bars, 70 μm.
3	c , Schematic diagram showing the experimental design for production of bone
4	marrow chimeric mice with wild-type hematopoietic cells.
5	d, Representative confocal images of knee joints showing recipient-derived
6	EGFP $^+$ macrophages in the synovium (S) attached to the meniscus (M) with PI
7	and SHG at multiple time points post-transplantation. Bars, 70 $\mu m.$
8	e, Quantification of recipient-derived EGFP ⁺ cells in the synovium as
9	percentages of total nucleated cells at the indicated time points following BM
10	transplantation. Symbols represent individual mice.
11	f, Schematic diagram showing the experimental design for the production of
12	bone marrow chimeric mice with CX ₃ CR1-EGFP transgenic hematopoietic cells.
13	g, Representative confocal images of knee joints showing donor-derived EGFP ⁺
14	macrophages in the synovium (S) attached to the meniscus (M). Bars, 70 μm.

h, Quantification of donor-derived EGFP⁺ cells in the synovium as percentages 1 of total nucleated cells at the indicated time points following bone marrow 2 transplantation. 3 confocal of CX₃CR1-EGFP⁺ Representative images cells and 4 i, TRAP-tdTomato⁺ osteoclasts in arthritic knee joints of double transgenic mice 5 6 (Control) and bone marrow chimeric mice (BMT). Bo: bone, L: patellar ligament. Arrowheads indicate bone erosion. Bars, 200 and 50 µm. 7 8 One-way ANOVA with Bonferroni's post hoc test (e, h). Mean ± S.E.M. for each 9 group. Symbols represent individual mice.



c, Histograms represent percentages of CX₃CR1-EGFP and TRAP-tdTomato⁺ 1 cells in the synovium, blood, and bone marrow of wild-type parabionts. 2 Quantification of CX₃CR1-EGFP⁺ cells in blood was conducted by FACS. Mean 3 ± S.E.M. for each group. Symbols represent individual mice. 4 d, Representative confocal images of CX₃CR1-EGFP⁺ cells 5 and 6 TRAP-tdTomato⁺ osteoclasts in arthritic knee joints from indicated parabionts.

7 $\,$ Bars, 200 and 50 $\mu m.$



1

2 Supplementary Figure 4. Comparison of CD45⁺F4/80⁺ cells in synovium

³ and other organs.

1	a, Phenotypic marker comparison of conventional OP-containing population in
2	bone marrow (BM-OP; $CX_3CR1^{lo}Ly6C^{hi}$ cells) and R3 cells in the inflamed
3	synovium.
4	b , Representative plots of CD45 ⁺ F4/80 ⁺ cells from various of organs of
5	CX ₃ CR1-EGFP transgenic mice.
6	c , Representative TRAP staining images of RANKL-induced osteoclastogenesis
7	and May-Giemsa staining of CX ₃ CR1 ^{lo} Ly6C ^{hi} cells and CX ₃ CR1 ^{hi} Ly6C ^{int} cells
8	from BM in CIA and non-CIA mice. Bars, 200 and <mark>20 μm</mark> .

d, Quantification of nuclei in multinucleated cells within the visual field in **c**.



Supplementary Figure 5. Definition of R2' and R3' fractions based on the 2 expression level of F4/80. 3 Both R2 and R3 fractions may contain transitional states. R2 (CX₃CR1^{lo}Ly6C^{hi}) 4 and R3 (CX₃CR1^{hi}Ly6C^{int}) cells in the inflamed synovium were further gated on 5 F4/80 to discriminate transitional status and define R2' 6 to (CX₃CR1^{lo}Ly6C^{hi}<u>F4/80^{int}</u>) and R3' (CX₃CR1^{hi}Ly6C^{int}<u>F4/80^{hi}</u>), respectively. R3' is 7

- 8 defined as "fully differentiated R3", and R2' is defined as "basal state R2".
- 9

1

10







1	Supplementary Figure 7. Transcriptional profiling of R1, R2', and R3' cells					
2	by RNA-Seq (related to Fig. 4).					
3	a , Downstream effects analysis of R2'/R1 cells and b , R3'/R2' cells by RNA-Seq.					
4	c, Enrichment analysis of mitochondrial translation and oxidative					
5	phosphorylation in R3' cells compared with R2' cells.					
6	d , Upstream analysis of FoxM1 in R3'/R2' cells by RNA-Seq.					



2 Supplementary Figure 8. Effects of thiostrepton *in vitro* and *in vivo* (related

1 to Fig. 4).

2	a , Histogram plots of annexin-V positive bone marrow macrophages treated with
3	10 ng/ml M-CSF and DMSO or thiostrepton for 48 hours.
4	b , Histogram plots displaying proliferation of bone marrow macrophages treated
5	with 10 ng/ml M-CSF and DMSO or thiostrepton for 48 hours, determined by
6	CellTrace Violet signal.
7	c, RT-PCR analysis of IL-1, IL-6, RANKL, and TNF expression in synovial
8	tissues from CIA mice treated with vehicle or 50 mg/kg thiostrepton. Vehicle or
9	thiostrepton were injected intraperitoneally every other day for 2 weeks before
10	sacrifice.
11	d, Representative TRAP staining images of RANKL-induced osteoclastogenesis
12	of synovial CX ₃ CR1 ^{hi} Ly6C ^{int} cells (R3 cells) and CX ₃ CR1 ^{lo} Ly6C ^{hi} cells from BM
13	(BM-OP) treated with thiostrepton.
14	e, Quantification of nuclei in multinucleated cells within the visual field depicted
15	in d . Bars, 200 μm.
16	f , Body weight of WT mice treated with vehicle or 50 mg/kg thiostrepton from 5

1	weeks to 10 weeks of age. Vehicle or thiostrepton were injected intraperitoneally
2	twice a week for 5 weeks.
3	g, Axial view of femur metaphyseal region in 10-week-old mice treated with
4	vehicle or thiostrepton as described in f . Bar, 1 mm.
5	h, Quantification results of micro CT analysis; bone volume/total volume (BV/TV),
6	cortical bone mineral density, trabecular bone mineral density, trabecular
7	number (Tb.N), trabecular thickness (Tb.Th), and trabecular space (Tb.Sp) were
8	measured.





³ erosion on micro-CT (related to Figs. 4, 7).

a, 3D reconstructions of hind paws were scored at six anatomical sites: talus,
navicular bone, medial cuneiform bone, and the bases of the first, second, and
third metatarsals.
b, Erosions were scored on a scale of 0–3 (0 = normal, 1 = pitting, 2 =
full-thickness holes in small–medium areas, 3 = full-thickness holes in medium–

9 large areas) with a maximum score of 18.

- **c-d**, Intra-observer and inter-observer reproducibility were r = 0.9831 and
- 2 0.9883, respectively. Scatter dots represent individual mice.



- 1 d. Representative TRAP staining images of bone marrow macrophages
- 2 electroporated with CMV-T7-FoxM1 plasmid or mock plasmid. Cells were cultured
- ³ with 100 ng/ml RANKL and M-CSF at the indicated concentrations. Bars, 200 μm.
- 4 **e**, Quantification of nuclei within multinucleated cells within the visual field in **d**.
- 5 Unpaired two-tailed t test (a, e). Mean ± S.E.M. for each group. Symbols
- 6 represent individual mice.



2 Supplementary Figure 11. Isolation of mature osteoclasts differentiated

3 from R3 cells.

1

a, Representative TRAP staining images of RANKL-induced osteoclastogenesis
 of synovial CX₃CR1^{hi}Ly6C^{int} cells (R3 cells) and nuclei quantification of
 mononuclear cells and multinucleated mature osteoclasts using Imaris software.
 Bars, 200 μm.

b, Schematic diagram depicting the procedure for isolation of mononuclear cells 1 and multinucleated mature osteoclasts using temperature-responsive cell 2 cultureware, RepCell. 3 c, Representative plots of isolated cells from RepCell. FSC¹⁰SSC¹⁰ population 4 (P1) and FSC^{hi}SSC^{hi} population (P2) were analyzed for TRAP-tdTomato and 5 Hoechst 33342. FSChiSSChitdTomato+ population with multiple nuclei were 6 gated as mature osteoclasts. Shaded region indicates control values from 7 8 wild-type mice. d, RT-PCR analysis of FoxM1 expression in R3 cells and resultant osteoclasts. 9 Unpaired two-tailed *t* test (d). Mean ± S.E.M. for each group. Symbols represent 10 individual mice in **a** and a single experiment in **d**. 11







2 Supplementary Figure 13. Histological examination of inflamed knee joints

³ in FoxM1^{fl/fl} Rosa26^{CreERT2} mice (related to Fig. 7).

- a, Histological examination of knee joints from CAIA mice treated with oil control,
- 5 tamoxifen, tamoxifen plus adoptive transfer of FoxM1^{+/+}CX₃CR1⁺ monocytes or
- 6 FoxM1^{-/-}CX₃CR1⁺monocytes. BM: bone marrow; S: synovium. Bars, 300 and
- 7 **100 µm**.
- 8 **b**, Number of osteoclasts (N. Oc) per visual field at the sites of bone erosions.
- 9 Symbols represent individual mice and values represent the average count of
- 10 three different sections.
- 11 One-way ANOVA with Bonferroni's post hoc test. Mean ± S.E.M. for each group.
- 12



2 Supplementary Figure 14. Physiological bone remodelling was not

- 3 affected by global FoxM1 deletion.
- 4 a, Body weight of FoxM1^{fl/fl}Rosa26^{CreERT2} mice treated with oil control or

5 tamoxifen from 6 weeks of age; 2 mg tamoxifen was injected intraperitoneally 3

⁶ days in a row from 6 weeks of age, and was repeated from 10 weeks of age.

7 b, Axial view of the femur metaphyseal region of 14-week-old

8 FoxM1^{fl/fl}Rosa26^{CreERT2} mice treated with oil control or tamoxifen. Bar, 1 mm.

9 **c,** Quantification of the micro CT analysis: bone volume/total volume (BV/TV),

10 cortical bone mineral density, trabecular bone mineral density, trabecular

number (Tb.N), trabecular thickness (Tb.Th), and trabecular space (Tb.Sp) were

- 1 measured in the femur metaphyseal region of 14-week-old
- 2 FoxM1^{fl/fl}Rosa26^{CreERT2} mice treated with oil control or tamoxifen.
- 3





a, RT-PCR analysis of FoxM1 expression in CX₃CR1⁺Ly6C^{hi} bone marrow cells

5 (BM-OP) from FoxM1^{fl/fl} and LysM-Cre;FoxM1^{fl/fl} mice.

6 **b**, Body weight of FoxM1^{fl/fl} and LysM-Cre:FoxM1^{fl/fl} mice.

7 c, Axial view of the femur metaphyseal region of 8-week-old FoxM1^{fl/fl} and



9 **d**, Quantification results of micro CT analysis; bone volume/total volume (BV/TV),

- 10 cortical bone mineral density, trabecular bone mineral density, trabecular
- number (Tb.N), trabecular thickness (Tb.Th), and trabecular space (Tb.Sp) were

1 measured in the femur metaphyseal region of 8-week-old FoxM1^{fl/fl} and

2 LysM-Cre:FoxM1^{fl/fl} mice.

3



Supplementary Figure 16. FoxM1 is dispensable for R2 to R3 cell
 differentiation.

a, Schematic diagram showing that synovial R2 cells from CAIA mice treated

⁵ with corn oil or tamoxifen were sorted into Nunclon Sphere plates and incubated

```
6 for 48 hours with 10 ng/ml M-CSF.
```

7 b, Flow cytometry plots of R2 cells incubated with 10 ng/ml M-CSF and

8 percentage of resultant CX₃CR1^{hi}Ly6C^{int} cells (R3).

9 Unpaired two-tailed *t* test (b). Mean ± S.E.M. for each group. Symbols represent

10 individual mice.



Supplementary Figure 17. Schematic diagram of the differentiation
 trajectory of inflammatory OPs in arthritis.

4 CX₃CR1^{lo}Ly6C^{hi} cells in the blood (R1) trans-migrate into the synovium as 5 CX₃CR1^{lo}Ly6C^{hi}F4/80^{int} cells (R2'), which express chemokines (*Ccl2, Ccl3, Ccl4,* 6 *Cxcl1, Cxcl2*), inflammatory cytokines (*Tnf, Il-1, Il-6*), and *Vegfa*. A high level of 7 M-CSF in the inflamed synovium up-regulates FoxM1 in the R2' subset and 8 induces differentiation into R3' cells (AtoMs), the osteoclast precursors (OPs) in 9 arthritis. R3' cells differentiate into osteoclasts upon RANKL-stimulation in 10 pannus to cause bone erosions.



- 2 Supplementary Figure 18. Gel source data



1 Supplementary Figure 19. Gating strategy for different tissues.

2	Viable cells were enriched by scatter plots and doublet cells were excluded
3	based on FSC-H/FSC-W. After gating on CD45 ⁺ cells, R2 and R3 cells in the
4	CIA synovium were defined according to CX ₃ CR1 and Ly6C. These populations
5	were further gated by F4/80 to sort R2' and R3' cells.

Gene set				
ACP5	FOS	PDGFB		
ATP6V0D2 GLO1		PPARG		
CALCR	GPC3	PPARGC1B		
CA2	ITGB3 SEMA4D			
CD109	JUN	SH3PXD2A		
CKB	JUNB	SPHK1		
CSF1	MAPK1	SPNS2		
CSF1R	MAPK11	SRC		
CTHRC1	MAPK12	TFRC		
CTNNB1 MAPK14		TM7SF4		
CTSK MITF		TNFRSF11A		
C200RF123	MMP9	TNFRSF11B		
EFNA2 NFATC1		TRAF2		
EFNB2 NFKB1 TRAF		TRAF6		
E2F1	NFKB2 TREM2			
FAM20C	OSCAR			
FARP2	OSTM1			

2 Supplementary Table 1. Gene set related to osteoclast differentiation

3 modified from the Broad Institute Molecular Signatures Database

Case	Diagnosis	Sex	Age	Onset of disease	Treatment	Seropositivity
1	RA	F	65	1987	MTX	RF
2	RA	F	56	2012	MTX	ACPA
3	RA	F	76	1994	MTX	RF, ACPA
4	RA	F	86	2003	PSL, Tac	RF, ACPA
5	RA	F	69	2010	MTX	RF
6	RA	F	77	2001	PSL, Tac, ADA	-
7	RA	F	89	2003	PSL, SASP, BUC, GOL	RF, ACPA
8	RA	F	67	2010	SASP	RF, ACPA
9	RA	F	76	1994	MTX, Tac	RF, ACPA
10	RA	M	87	2014	PSL, MTX, Tac	RF, ACPA
11	RA	F	58	2010	MTX	-
12	RA	F	42	2013	ADA	RF, ACPA
13	RA	F	69	1993	PSL, MTX, BOL	RF, ACPA
14	RA	F	69	2007	none	-
15	RA	M	67	1989	ETN	RF, ACPA
16	RA	F	71	1998	MTX, PSL, IGU	RF
17	RA	M	69	2003	MTX, ABT	RF, ACPA
18	RA	F	55	2018	MTX	RF, ACPA
19	RA	F	66	1992	SASP, PSL	RF, ACPA
20	RA	F	74	2018	SASP	-
21	RA	F	79	1994	Tac, TCZ	-
22	RA	F	74	2002	Tac, BUC	-
23	RA	M	54	2017	MTX, PSL	RF

Supplementary Table 2. RA patient clinical information.

MTX: methotrexate; PSL: prednisolone; Tac: tacrolimus; SASP:
salazosulfapyridine; BUC: bucillamine; IGU: iguratimod; ETN: etanercept; ABT:
abatacept; ADA: adalimumab; GOL: golimumab; TCZ: tocilizumab; RF:
rheumatoid factor; ACPA: anti-citrullinated protein antibody.

Supplementary Video 1. Ex vivo incubation of inflamed synovium from 1 double transgenic mice (CX₃CR1-EGFP/TRAP-tdTomato) (related to Fig. 1). 2 3 Harvested inflamed synovium from double transgenic mice (CX₃CR1-EGFP/TRAP-tdTomato) was incubated with 100 ng/ml RANKL and 10 4 ng/ml M-CSF stimulation. Sequential images of the same visual field were 5 acquired by BioStation IM-Q (Nikon). CX₃CR1-EGFP and TRAP-tdTomato are 6 shown as green and red, respectively, on transmission images. Playback speed is 7 8 6000X.