



Title	On a generalization of QF-3' rings
Author(s)	Kurata, Yoshiki; Katayama, Hisao
Citation	Osaka Journal of Mathematics. 1976, 13(2), p. 407-418
Version Type	VoR
URL	<a href="https://doi.org/10.18910/9322">https://doi.org/10.18910/9322</a>
rights	
Note	

*The University of Osaka Institutional Knowledge Archive : OUKA*

<https://ir.library.osaka-u.ac.jp/>

The University of Osaka

## ON A GENERALIZATION OF QF-3' RINGS\*

Dedicated to Professor Kiiti Morita for the celebration of his sixtieth birthday.

YOSHIKI KURATA AND HISAO KATAYAMA

(Received February 20, 1975)

A ring  $R$  with identity is called left QF-3' if the injective hull  $E(R)$  of the left  $R$ -module  $R$  is torsionless. This class of rings and the other related generalizations of quasi-Frobenius rings have been studied by a number of authors.

Recently, Jans [7] has given a torsion theoretic characterization of left QF-3' rings (cf. also Kato [8] and Tsukerman [14]). The purpose of this paper is, generalizing this idea, to consider a module theoretic generalization of left QF-3' rings. We shall say that a left  $R$ -module  $Q$  is QF-3' if its injective hull  $E(Q)$  is torsionless with respect to  $Q$ , i.e.,  $E(Q)$  can be embedded in a direct product of copies of  $Q$ .

The main theorem of §1 will give some equivalent conditions for  $Q$  to be QF-3'.

In §2, we shall discuss basic properties of QF-3'  $R$ -modules and study a relation between QF-3'  $R$ -modules and cogenerators for  $R$ -mod.

We shall treat, in §3, QF-3'  $R$ -modules with zero singular submodule. We shall give some results relating the notions of  $Q$ -torsionless  $R$ -modules and non-singular  $R$ -modules. In particular we shall show that, if  $Q$  is faithful, these notions coincide if and only if  $Q$  is QF-3' and has zero singular submodule. We shall also give another characterization of a QF-3'  $R$ -module with zero singular submodule making use of its injective submodules.

After completed this paper, we found that the similar results were obtained by Bican [2] and wrought a slight change in the paper.

Throughout this paper,  $R$  will denote an associative ring with identity and  $R$ -mod the category of unital left  $R$ -modules and  $R$ -homomorphisms. We shall deal only with left  $R$ -modules and so  $R$ -modules will mean unital left  $R$ -modules.  $E(M)$  will always denote the injective hull of a left  $R$ -module  $M$  and  $r_M(*)$  the right annihilator of  $*$  in  $M$ .

---

\*<sup>o</sup>) The authors are indebted to the referee for helpful suggestions.

### 1. Preliminaries

A subfunctor  $r$  of the identity functor of  $R\text{-mod}$  is called a preradical of  $R\text{-mod}$ . It is called idempotent if  $r(r(M))=r(M)$  and a radical if  $r(M/r(M))=0$  for all  $R$ -modules  $M$ . To each preradical  $r$  we associate two classes of  $R$ -modules, namely

$$\mathbf{T}(r) = \{M \mid r(M)=M\} \text{ and } \mathbf{F}(r) = \{M \mid r(M)=0\} .$$

In case a preradical  $r$  is idempotent and is a radical, the pair  $(\mathbf{T}(r), \mathbf{F}(r))$  forms a torsion theory for  $R\text{-mod}$  in the sense of [5].

In the class of all preradicals of  $R\text{-mod}$ , there is a partial ordering in which  $r_1 \leq r_2$  means that  $r_1(M) \subset r_2(M)$  for all  $R$ -modules  $M$ . For each preradical  $r$  there exists a largest idempotent preradical  $\hat{r}$  smaller than or equal to  $r$  and a smallest radical  $\bar{r}$  larger than or equal to  $r$ . It is easy to see that  $\mathbf{T}(\hat{r})=\mathbf{T}(r)$  and  $\mathbf{F}(\bar{r})=\mathbf{F}(r)$ . Moreover, if  $r$  is idempotent, then so is  $\bar{r}$ , and  $\hat{r}$  is a radical if  $r$  is.

Let  $Q$  be an  $R$ -module and let us define

$$k_Q(M) = \bigcap_{f \in \text{Hom}_R(M, Q)} \text{Ker}(f)$$

for each  $R$ -module  $M$ . Then  $k_Q$  is a radical of  $R\text{-mod}$  such that  $k_Q(Q)=0$ . Moreover it is a unique maximal one of those preradicals  $r$  of  $R\text{-mod}$  for which  $r(Q)=0$ , and  $k_{E(Q)}$  is a unique maximal one of those left exact radicals  $r$  of  $R\text{-mod}$  for which  $r(Q)=0$ . As is well-known, every left exact radical of  $R\text{-mod}$  is of the form  $k_E$  for some injective  $R$ -module  $E$ . For example, we can take  $E$  as the direct product of injective hulls of all cyclic torsion-free  $R$ -modules (e.g., see [11]).

Since  $k_Q \leq k_{Q'}$  for each submodule  $Q'$  of  $Q$  and  $k_{E(Q)}$  is idempotent, we have  $k_{E(Q)} \leq \hat{k}_Q \leq k_Q$ .

The class  $\mathbf{T}(k_Q)$  coincides with the class  $\{M \mid \text{Hom}_R(M, Q)=0\}$  and is closed under taking homomorphic images, direct sums and extensions. So this is a torsion class in  $R\text{-mod}$  and the corresponding torsion-free class coincides with  $\mathbf{F}(\hat{k}_Q)$ . On the other hand, the class of  $R$ -modules  $\mathbf{F}(k_Q)$  is not a torsion-free class in general. This is closed under taking submodules and direct products, but not extensions in general (see e.g. [16, Example B]). As is well-known,  $Q$  is a cogenerator for  $\mathbf{F}(k_Q)$ . Moreover, an  $R$ -module  $M$  is in  $\mathbf{F}(k_Q)$  if (and only if) it can be embedded in a direct product of copies of  $Q$ . However, for simple  $R$ -modules we have

**Proposition 1.1.** *A simple  $R$ -module  $S$  is in  $\mathbf{F}(k_Q)$  if and only if there exists an  $R$ -monomorphism of  $S$  into  $Q$ .*

The proof is easy and so we will omit it.

As was mentioned above,  $\mathbf{F}(k_Q)$  is not closed under taking extensions.

The following proposition shows when it is closed under taking extensions. Evidently this is the case if  $Q$  is injective.

**Proposition 1.2.** *The following conditions on an  $R$ -module  $Q$  are equivalent :*

- (1)  $\mathbf{F}(k_Q)$  is closed under taking extensions, i.e., it becomes a torsion-free class.
- (2)  $k_Q = \hat{k}_Q$ , i.e.,  $k_Q$  is idempotent.
- (3)  $\mathbf{F}(k_Q) = \mathbf{F}(\hat{k}_Q)$ .

Bican [2] has obtained the same result independently, and so we will omit the proof.

The class  $\mathbf{T}(k_Q)$  is a torsion class, but it is not, in general, closed under taking submodules (e.g., see [16]). Concerning this, we have

**Proposition 1.3.** *For an  $R$ -module  $Q$ , the following conditions are equivalent :*

- (1)  $\mathbf{T}(k_Q)$  is closed under taking submodules.
- (2)  $\hat{k}_Q = k_{E(Q)}$ , i.e.,  $\hat{k}_Q$  is left exact.
- (3)  $\mathbf{T}(k_Q) = \mathbf{T}(k_{E(Q)})$ .

Proof. (1) $\Rightarrow$ (2). Suppose that  $\mathbf{T}(k_Q)$  is closed under taking submodules. Then, since  $\hat{k}_Q \leq k_Q$ ,  $\hat{k}_Q(Q) = 0$  and, since the corresponding torsion-free class  $\mathbf{F}(\hat{k}_Q)$  of  $\mathbf{T}(k_Q)$  is closed under taking injective hulls, we have  $\hat{k}_Q(E(Q)) = 0$ . So  $\hat{k}_Q \leq k_{E(Q)}$  and hence  $\hat{k}_Q = k_{E(Q)}$ .

(2) $\Rightarrow$ (3) is clear and since  $k_{E(Q)}$  is left exact, (3) implies (1).

This proposition was also proved in Bican [2] by a different method.

Combining this with Proposition 1.2, we have

**Theorem 1.4.** *The following conditions on an  $R$ -module  $Q$  are equivalent :*

- (1)  $(\mathbf{T}(k_Q), \mathbf{F}(k_Q))$  forms a hereditary torsion theory for  $R$ -mod.
- (2)  $\mathbf{T}(k_Q)$  is closed under taking submodules and  $\mathbf{F}(k_Q)$  is closed under taking extensions.
- (3)  $k_Q = k_{E(Q)}$ .
- (4)  $k_Q$  is left exact.
- (5)  $\mathbf{F}(k_Q)$  is closed under taking injective hulls.
- (6)  $\mathbf{F}(k_Q)$  is closed under taking essential extensions.
- (7)  $\mathbf{F}(k_Q)$  contains an injective  $R$ -module  $M$  with  $k_M(Q) = 0$ .
- (8)  $E(Q) \in \mathbf{F}(k_Q)$ .
- (9)  $\mathbf{T}(k_Q) = \mathbf{T}(k_{E(Q)})$  and  $\mathbf{F}(k_Q) = \mathbf{F}(k_{E(Q)})$ .

Proof. Here we show only that (7) implies (8). The proof of the other is easy. Since  $k_M(Q) = 0$ ,  $Q \subset \prod M$ , a direct product of copies of  $M$ , and hence  $E(Q) \subset \prod M$ .  $\mathbf{F}(k_Q)$  is closed under taking direct products and submodules and so we have  $E(Q) \in \mathbf{F}(k_Q)$ .

The equivalence of (3), (4) and (8) was also proved in Bican [2]. In case  $Q=R$ , the equivalence of these conditions, except for (1), (4), (5) and (9), was shown by Colby and Rutter [4], Jans [7], and Kato [8].

## 2. QF-3' R-modules

Recall that a ring  $R$  is left QF-3' if the injective hull of the  $R$ -module  $R$  is torsionless, i.e.,  $k_R(E(R))=0$ . Recently, Jans [7] has shown that  $R$  is left QF-3' if and only if  $\mathbf{F}(k_R)$  is closed under taking extensions and  $\mathbf{T}(k_R)$  is closed under taking submodules (cf. also Kato [8] and Tsukerman [14]). From this point of view we now make the following definition.

**DEFINITION.** An  $R$ -module  $Q$  is called QF-3' if  $Q$  satisfies each one of the conditions of Theorem 1.4.

It follows from this definition that every injective  $R$ -module is QF-3'. The following example pointed out by Tsukerman without proof shows that there exist non-injective QF-3'  $R$ -modules.

**EXAMPLE 2.1.** Every direct sum of injective  $R$ -modules is QF-3'.

To see this, let  $Q=\sum_{\lambda\in\Lambda}\oplus Q_\lambda$  be a direct sum of injective  $R$ -modules. Then  $\mathbf{F}(k_{Q_\lambda})\subset\mathbf{F}(k_Q)$  for all  $\lambda$  and hence  $\prod_{\lambda\in\Lambda}Q_\lambda\in\mathbf{F}(k_Q)$ . Since  $Q\subset E(Q)\subset\prod_{\lambda\in\Lambda}Q_\lambda$ ,  $E(Q)\in\mathbf{F}(k_Q)$  and thus  $Q$  is QF-3'.

**Proposition 2.2.** (1) *Every direct product of QF-3'  $R$ -modules is QF-3'.*  
 (2) *Every direct sum of QF-3'  $R$ -modules is QF-3'.*

**Proof.** (1) Let  $Q=\prod_{\lambda\in\Lambda}Q_\lambda$  be a direct product of QF-3'  $R$ -modules. Then  $\mathbf{F}(k_{Q_\lambda})\subset\mathbf{F}(k_Q)$  for all  $\lambda$  and hence  $\prod_{\lambda\in\Lambda}E(Q_\lambda)\in\mathbf{F}(k_Q)$ . Since  $Q\subset E(Q)\subset\prod_{\lambda\in\Lambda}E(Q_\lambda)$ ,  $E(Q)\in\mathbf{F}(k_Q)$  and thus  $Q$  is QF-3'. The proof of (2) is similar to that of (1) and so it will be omitted.

It should be noted that, as we shall show later, direct summands of a QF-3'  $R$ -module need not be QF-3' in general.

**Proposition 2.3.** *Every essential extension of a QF-3'  $R$ -module is QF-3'.*

**Proof.** Suppose that  $Q$  is QF-3' and  $Q'$  is an essential extension of  $Q$ . Then we can assume that  $Q\subset Q'\subset E(Q)$  and hence we have  $k_{E(Q)}\leq k_{Q'}\leq k_Q$ . By Theorem 1.4,  $k_Q=k_{Q'}$  and thus  $Q'$  is QF-3' again by Theorem 1.4.

It follows from this that every rational extension of a QF-3'  $R$ -module is also QF-3'. This appeared in [13] for left QF-3' rings.

Let  $Q$  be an  $R$ -module. As is easily seen,  $Q$  is faithful if and only if  $k_Q(R)=0$  and this is so if and only if  $k_Q\leq k_R$ . On the other hand,  $Q$  is torsionless if and only if  $k_R(Q)=0$ , or equivalently,  $k_R\leq k_Q$ . Therefore if  $Q$  is both

faithful and torsionless, then we have  $k_Q = k_R$ . Applying Theorem 1.4, we have

**Theorem 2.4.** *For a ring  $R$ , the following conditions are equivalent :*

- (1)  $R$  is a left QF-3' ring.
- (2) The  $R$ -module  $R$  is QF-3'.
- (3) There exists a QF-3'  $R$ -module  $Q$  which is both faithful and torsionless.

**Proposition 2.5.** *Let  $Q$  be an  $R$ -module.*

(1) *If  $Q$  is a QF-3'  $R$ -module with non-zero socle, then the injective hull of every simple submodule of  $Q$  is isomorphic to a submodule of  $Q$ .*

(2) *If the injective hull of every cyclic submodule of  $Q$  is isomorphic to a submodule of  $Q$ , then  $Q$  is QF-3'.*

Proof. (1) Let  $S$  be a simple submodule of  $Q$ . Take  $x(\neq 0)$  in  $E(S)$ . Then there exists  $ax(\neq 0)$  in  $Rx \cap S$ .  $S$  is in  $F(k_Q)$  and so  $E(S)$  is in  $F(k_Q)$  by Theorem 1.4. We can find an  $R$ -homomorphism  $f: E(S) \rightarrow Q$  such that  $f(ax) \neq 0$ . Hence we have  $f(S) \neq 0$  and  $f$  must be a monomorphism.

(2) Take  $x(\neq 0)$  in  $E(Q)$ . There exists  $ax(\neq 0)$  in  $Rx \cap Q$ . By assumption,  $E(Rax) \subset Q$ , and the inclusion mapping  $Rax \rightarrow E(Rax)$  can be extended to an  $R$ -homomorphism  $f: E(Q) \rightarrow Q$  such that  $f(x) \neq 0$ , which shows that  $Q$  is QF-3'.

Clearly, for a direct sum  $Q$  of injective  $R$ -modules, the injective hull of every cyclic submodule is isomorphic to a submodule of  $Q$  and so (2) above gives another proof of Example 2.1.

As an immediate consequence of this proposition, we have at once

**Corollary 2.6.** *For an  $R$ -module  $Q$  with non-zero socle, the following conditions are equivalent :*

- (1)  $Q$  is indecomposable and QF-3'.
- (2)  $Q = E(S)$  for every simple submodule  $S$  of  $Q$ .
- (3)  $Q = E(Q')$  for every non-zero submodule  $Q'$  of  $Q$ .

**Proposition 2.7.** *Let  $Q$  be an  $R$ -module. If every cyclic submodule of  $Q$  is QF-3', then  $Q$  is itself QF-3'.*

Proof. Take  $x(\neq 0)$  in  $E(Q)$  and claim that there exists an  $R$ -homomorphism  $f^*: E(Q) \rightarrow Q$  with  $f^*(x) \neq 0$ . Choose an element  $a$  in  $R$  such that  $ax(\neq 0)$  is in  $Q$ . Then we have  $Rax \subset E(Rax) \subset E(Q)$  and  $E(Q) = E(Rax) \oplus Q_1$  for some submodule  $Q_1$  of  $E(Q)$ . By assumption,  $Rax$  is QF-3' and so there exists an  $R$ -homomorphism  $f: E(Rax) \rightarrow Rax$  such that  $f(ax) \neq 0$ . Then it is easy to see that the composition  $f^*: E(Q) \rightarrow Q$  of  $f$  and the projection mapping  $E(Q) \rightarrow E(Rax)$  has the desired property.

As a direct consequence of this, we see that if every cyclic  $R$ -module is QF-3', then every  $R$ -module is also QF-3'. This was proved by Tsukerman

[14] under the assumption that  $R$  is left hereditary.

Recall that an  $R$ -module  $Q$  is a cogenerator for  $R\text{-mod}$  if  $F(k_Q) = R\text{-mod}$ . Therefore, a cogenerator for  $R\text{-mod}$  is necessarily QF-3'. We now consider the question of when a QF-3'  $R$ -module becomes a cogenerator for  $R\text{-mod}$ . To do this we shall prove

**Proposition 2.8.** *For an  $R$ -module  $Q (\neq 0)$ , the following conditions are equivalent :*

- (1)  $Q$  contains a copy of every simple  $R$ -module.
- (2) Every simple  $R$ -module belongs to  $F(k_Q)$ .
- (3) For every simple  $R$ -module  $S$ ,  $\text{Hom}_R(S, Q) \neq 0$ .
- (4) For every maximal left ideal  $m$  of  $R$ ,  $r_Q(m) \neq 0$ .
- (5) For every proper left ideal  $m$  of  $R$ ,  $r_Q(m) \neq 0$ .
- (6) For every non-zero finitely generated  $R$ -module  $M$ ,  $\text{Hom}_R(M, Q) \neq 0$ .
- (7) For every non-zero cyclic  $R$ -module  $M$ ,  $\text{Hom}_R(M, Q) \neq 0$ .
- (8)  $E(Q)$  is a cogenerator for  $R\text{-mod}$ .
- (9) Every non-zero injective  $R$ -module  $M$  with  $k_M(Q) = 0$  is a cogenerator for  $R\text{-mod}$ .

Proof. We shall show only (7) $\Rightarrow$ (8) $\Rightarrow$ (9) $\Rightarrow$ (1).

(7) $\Rightarrow$ (8). Let  $M$  be an  $R$ -module. Take  $x (\neq 0)$  in  $M$ . Then by assumption there exists a non-zero  $R$ -homomorphism  $f: Rx \rightarrow Q$ . Since  $E(Q)$  is injective, it can be extended to an  $R$ -homomorphism  $f': M \rightarrow E(Q)$  and  $f'(x) = f(x) \neq 0$ . This shows that  $E(Q)$  is a cogenerator for  $R\text{-mod}$ .

(8) $\Rightarrow$ (9). Since  $k_M(Q) = 0$ ,  $Q \subset \prod M$ , a direct product of copies of  $M$ , and hence  $E(Q) \subset \prod M$ . Then we have  $R\text{-mod} = F(k_{E(Q)}) \subset F(k_{\prod M})$ . It follows that  $\prod M$  is a cogenerator for  $R\text{-mod}$  and so is  $M$  by [12, Lemma 1].

(9) $\Rightarrow$ (1). Since  $E(Q)$  is a cogenerator for  $R\text{-mod}$ , for every simple  $R$ -module  $S$ , there exists a non-zero  $R$ -homomorphism  $f: S \rightarrow E(Q)$ .  $S$  is simple, so  $f$  must be a monomorphism. Since  $f(S) \cap Q \neq 0$ ,  $f(S) \cap Q = f(S)$  and hence  $f(S)$  is contained in  $Q$ .

An  $R$ -module satisfying (1) and (8) was called lower distinguished by Azumaya [1] and a quasi-cogenerator by Morita [10] respectively.

Generalizing results due to Kato [8], Jans [6] and Sugano [12], we have

**Theorem 2.9.** *The following conditions on an  $R$ -module  $Q$  are equivalent :*

- (1)  $Q$  is a cogenerator for  $R\text{-mod}$ .
- (2)  $Q$  is QF-3' and contains a copy of every simple  $R$ -module.
- (3)  $\sum_{\lambda \in \Lambda} \oplus E(S_\lambda) \in F(k_Q)$ , where  $\{S_\lambda\}_{\lambda \in \Lambda}$  be a complete set of representatives for the isomorphism classes of simple  $R$ -modules.
- (4) There exists a cogenerator for  $R\text{-mod}$  contained in  $F(k_Q)$ .
- (5) Every  $R$ -module  $M$  with  $k_M(Q) = 0$  is a cogenerator for  $R\text{-mod}$ .

(6)  $Q$  is faithful QF-3' and  $\mathbf{F}(k_Q)$  is closed under taking homomorphic images.

Proof. (1) $\Rightarrow$ (2) $\Rightarrow$ (3) follow from Proposition 1.1 and Theorem 1.4 and (3) $\Rightarrow$ (4) and (5) $\Rightarrow$ (6) are easy.

(4) $\Rightarrow$ (5). Let  $N$  be a cogenerator for  $R$ -mod contained in  $\mathbf{F}(k_Q)$  and let  $M$  be an  $R$ -module with  $k_M(Q)=0$ . Then we have  $k_M \leq k_Q \leq k_N$  and  $\mathbf{F}(k_N)=R$ -mod. Hence  $\mathbf{F}(k_M)=R$ -mod as desired.

(6) $\Rightarrow$ (1). By assumption, there exists a class  $\mathbf{T}$  of  $R$ -modules such that  $(\mathbf{T}(k_Q), \mathbf{F}(k_Q), \mathbf{T})$  forms a 3-fold torsion theory for  $R$ -mod in the sense of [9]. It follows from Lemma 2.1 of [9] that  $k_Q(M)=k_Q(R) \cdot M$  for each  $R$ -module  $M$ . Hence it results that  $\mathbf{F}(k_Q)=R$ -mod since  $Q$  is faithful.

### 3. Non-singular QF-3' R-modules

In case the singular submodule  $Z(Q)=0$ , we can give a simple criterion for  $Q$  being QF-3'.

**Theorem 3.1.** *Let  $Q$  be an  $R$ -module with  $Z(Q)=0$ . Then  $Q$  is QF-3' if and only if  $\mathbf{T}(k_Q)$  is closed under taking submodules.*

This was also obtained by the same method in Bican [2] and we will omit the proof.

As is well-known, the functor  $Z$  of  $R$ -mod which assigns to each  $R$ -module  $M$  its singular submodule  $Z(M)$  is a left exact preradical of  $R$ -mod. It is to be noted that, for this preradical,  $\mathbf{F}(Z)$  is nothing but the torsion-free class of the so-called Goldie torsion theory. We shall now give other characterizations of non-singular QF-3'  $R$ -modules by means of the functor  $Z$ . To do this, we first prove the following which appeared in Colby and Rutter [4] for the case  $Q=R$ .

**Proposition 3.2.** *The following conditions on an  $R$ -module  $Q$  are equivalent:*

- (1)  $Z(Q)=0$ .
- (2)  $\mathbf{F}(k_Q) \subset \mathbf{F}(Z)$ .
- (3)  $\mathbf{T}(Z) \subset \mathbf{T}(k_Q)$ .

Proof. (1) $\Rightarrow$ (2). Since  $Z \leq k_Q$ , we have  $\mathbf{F}(k_Q) \subset \mathbf{F}(Z)$ .

(2) $\Rightarrow$ (3). Let  $M$  be in  $\mathbf{T}(Z)$ . Take  $f$  in  $\text{Hom}_R(M, Q)$  and  $x$  in  $M$ . Then, since  $\text{Ann}_R(x)$  is essential in  $R$ , so is  $\text{Ann}_R(f(x))$  and hence  $f(x)$  is in  $Z(Q)$ . But by assumption (2)  $Z(Q)=0$  and this implies that  $M$  is contained in  $\mathbf{T}(k_Q)$ .

(3) $\Rightarrow$ (1). Since  $Z$  is an idempotent preradical,  $Z(Q)$  is in  $\mathbf{T}(Z)$  and hence is in  $\mathbf{T}(k_Q)$ . This shows that  $\text{Hom}_R(Z(Q), Q)=0$  and  $Z(Q)=0$ .

**Lemma 3.3.** *Let  $Q$  be a faithful  $R$ -module. Then we have*

- (1)  $\mathbf{T}(k_{E(Q)}) \subset \mathbf{T}(Z)$ , and
- (2)  $\mathbf{F}(Z) \subset \mathbf{F}(k_{E(Q)})$ .

*Proof.* (1) For every  $R$ -module  $M$  in  $\mathbf{T}(k_{E(Q)})$  and every element  $x$  in  $M$ , we shall claim that  $\text{Ann}_R(x)$  is essential in  $R$ . Suppose that  $\mathfrak{m}$  is a non-zero left ideal in  $R$  such that  $\text{Ann}_R(x) \cap \mathfrak{m} = 0$ . Define  $f: \mathfrak{m}x \rightarrow R$  such that  $f(ax) = a$  for  $a \in \mathfrak{m}$ . Clearly this is a well defined  $R$ -homomorphism. Let  $a$  be a non-zero element of  $\mathfrak{m}$ . Then there exists an  $R$ -homomorphism  $g: R \rightarrow E(Q)$  such that  $g(a) \neq 0$  since  $E(Q)$  is faithful. The composition map  $g \circ f: \mathfrak{m}x \rightarrow E(Q)$  can be extended to an  $R$ -homomorphism  $h: M \rightarrow E(Q)$  and  $h(ax) = g(f(ax)) = g(a) \neq 0$ . Thus we have  $\text{Hom}_R(M, E(Q)) \neq 0$ , but this is a contradiction. Similarly we can show that (2) holds.

It follows from Lemma 3.3 that, if  $Q$  is faithful and non-singular, then  $E(Q)$  is a cogenerator for  $\mathbf{F}(Z)$ . However, we can show that this is also true for more general QF-3'  $R$ -modules.

**Theorem 3.4.** *For a faithful  $R$ -module  $Q$ , the following conditions are equivalent:*

- (1)  $Q$  is QF-3' and  $Z(Q) = 0$ .
- (2)  $\mathbf{T}(k_Q) = \mathbf{T}(Z)$ .
- (3)  $\mathbf{F}(k_Q) = \mathbf{F}(Z)$ .
- (4)  $k_Q = Z$ .
- (5)  $Q$  is a cogenerator for  $\mathbf{F}(Z)$ .

*Proof.* (1)  $\Rightarrow$  (2) and (1)  $\Rightarrow$  (3) follow from Proposition 3.2 and Lemma 3.3.

(2)  $\Rightarrow$  (1). By Proposition 3.2,  $Z(Q) = 0$ . Since  $\mathbf{T}(Z)$  is closed under taking submodules, so is  $\mathbf{T}(k_Q)$ . Therefore,  $Q$  is QF-3' by Theorem 3.1.

(3)  $\Rightarrow$  (1). By Proposition 3.2,  $Z(Q) = 0$ . Since  $\mathbf{F}(Z)$  is closed under taking injective hulls, so is  $\mathbf{F}(k_Q)$ . Therefore,  $Q$  is QF-3' by Theorem 1.4.

(4)  $\Rightarrow$  (1) follows from Theorem 1.4 since  $Z$  is left exact. So we assume (2) and also (3). By Proposition 3.2,  $Z(Q) = 0$  and we have  $Z \leq k_Q$ .  $\mathbf{F}(k_Q) = \mathbf{F}(Z)$  is closed under taking extensions and so by Proposition 1.2  $k_Q$  is idempotent. For each  $R$ -module  $M$ ,  $k_Q(M) \in \mathbf{T}(k_Q) = \mathbf{T}(Z)$  and  $k_Q(M) = Z(k_Q(M)) \subset Z(M)$ . Therefore we have  $k_Q \leq Z$ .

(3)  $\Leftarrow$  (5). The fact that  $Q$  is a cogenerator for  $\mathbf{F}(Z)$  means that  $Z(Q) = 0$  and  $\mathbf{F}(Z) \subset \mathbf{F}(k_Q)$ , or equivalently  $\mathbf{F}(Z) = \mathbf{F}(k_Q)$  by Proposition 3.2. This completes the proof of the theorem.

In [4], it was given a similar characterization, except for (4) and (5), of non-singular left QF-3 rings in case these are semi-primary, and (4) may be viewed as a generalization of a result of [15].

In Proposition 2.3 we have shown that every essential extension of a QF-3'

$R$ -module is QF-3'. However, in case it is non-singular, we have

**Corollary 3.5.** *Let  $Q$  be a faithful QF-3'  $R$ -module and let  $Q'$  be a non-singular  $R$ -module such that  $Q \subset Q'$ . Then  $Q'$  is also QF-3'.*

Proof. By Proposition 3.2 and Theorem 3.4,  $F(Z) = F(k_Q) \subset F(k_{Q'}) \subset F(Z)$ . Hence we have  $F(k_{Q'}) = F(Z)$  and  $Q'$  is QF-3'.

As another corollary to this theorem, we have

**Corollary 3.6.** *For a ring  $R$  with  $Z({}_R R) = 0$  and its maximal ring of left quotients  $Q$ , the following conditions are equivalent :*

- (1) *Every non-singular  $R$ -module is torsionless, i.e.,  $R$  is a cogenerator for  $F(Z)$ .*
- (2)  *$R$  is a left QF-3' ring.*
- (3)  *${}_R Q$  is torsionless.*

Recently, Cateforis [3] has given a necessary and sufficient condition for a non-singular  $R$ -module to be a cogenerator for  $F(Z)$ . The following theorem is motivated by his Theorem 1.1, and provides alternative characterizations of non-singular QF-3'  $R$ -modules to that given in Theorem 3.4.

**Theorem 3.7.** *For a non-singular  $R$ -module  $Q$ , the following conditions are equivalent :*

- (1)  *$Q$  is faithful and QF-3'.*
- (2)  *$Q$  contains non-zero injective submodules and the sum  $Q^*$  of all such injective submodules is faithful.*
- (3) *There exists a faithful submodule  $Q_0$  of  $Q$  such that  $Q_0$  contains the injective hull of every one of its finitely generated submodules.*

Before proving the theorem, we shall quote Lemma 0.2 of [3] and give its proof for the sake of completeness.

**Lemma 3.8.** *If  $A$  is an injective  $R$ -module and  $B$  is a non-singular  $R$ -module, then, for every  $R$ -homomorphism  $f: A \rightarrow B$ , both  $\text{Ker}(f)$  and  $\text{Im}(f)$  are injective.*

Proof. Since  $A$  is injective, we can assume that  $\text{Ker}(f) \subset E(\text{Ker}(f)) \subset A$ . Take  $x (\neq 0)$  in  $E(\text{Ker}(f))$  and  $a (\neq 0)$  in  $R$ . If  $ax = 0$ , then  $a$  is in  $Ra \cap \text{Ann}_R(f(x))$ . If  $ax \neq 0$ , then we can find  $ba x (\neq 0)$  in  $Rax \cap \text{Ker}(f)$  for some  $b$  in  $R$ . Since  $f(bax) = 0$  and  $ba \neq 0$ ,  $Ra \cap \text{Ann}_R(f(x)) \neq 0$ . At any rate, we have  $Ra \cap \text{Ann}_R(f(x)) \neq 0$  and hence  $\text{Ann}_R(f(x))$  is essential in  $R$ .  $f(x)$  is then in  $Z(B) = 0$ . Therefore,  $x$  is in  $\text{Ker}(f)$  which shows that  $\text{Ker}(f) = E(\text{Ker}(f))$ .

Proof of Theorem 3.7. (1) $\Rightarrow$ (2). By assumption,  $\text{Hom}_R(E(Q), Q) \neq 0$

and so by Lemma 3.8  $Q$  contains certainly non-zero injective submodules. Moreover  $k_{Q^*}(E(Q))=k_Q(E(Q))$  again by Lemma 3.8. Hence we have  $k_{Q^*}(Q)=0$  which implies that  $k_{Q^*}\leq k_Q$  and  $Q^*$  is faithful. (Moreover in this case  $k_Q=k_{Q^*}$  holds.)

(2) $\Rightarrow$ (3). For every finite family  $\{M_1, M_2, \dots, M_n\}$  of non-zero injective submodules of  $Q$ ,  $\sum_{i=1}^n M_i$  is a homomorphic image of an injective  $R$ -module  $\sum_{i=1}^n \oplus M_i$  and so by Lemma 3.8 it is also injective. It follows from this that  $Q^*$  contains the injective hull of every one of its finitely generated submodules.

(3) $\Rightarrow$ (1). By Proposition 2.5  $Q_0$  is QF-3'.  $Q_0$  is faithful and  $Q$  is non-singular, so by Corollary 3.5  $Q$  is also QF-3'. (Here we shall point out that  $k_Q=k_{Q_0}$  holds. To see this it is sufficient to show that  $k_{Q_0}(Q)=0$ . Take  $x(\neq 0)$  in  $E(Q)$ . Then  $\text{Ann}_R(x)$  is not essential in  $R$  so we can find  $a(\neq 0)$  in  $R$  such that  $Ra \cap \text{Ann}_R(x)=0$ . Since  $ax$  is a non-zero element of  $E(Q)$ , there exists some  $bax(\neq 0)$  in  $Rax \cap Q$ .  $ba$  is a non-zero element in  $R$  and  $Q_0$  is faithful and so for some  $x_0$  in  $Q_0$  we have  $bax_0 \neq 0$ . Then the mapping  $f: Rbax \rightarrow Rbax_0$  given by  $f(rbax)=rbax_0$ , for  $r$  in  $R$ , is a well-defined  $R$ -homomorphism. By assumption,  $E(Rbax_0) \subset Q_0$  and so  $f$  has an extension  $f^*: E(Q) \rightarrow Q_0$  and  $f^*(x) \neq 0$ . Thus  $k_{Q_0}(E(Q))=0$  and  $k_{Q_0}(Q)=0$ .)

To illustrate the theorem, we shall give some examples.

EXAMPLE 3.9. Let  $R$  be the ring of  $2 \times 2$  upper triangular matrices over a field  $K$ . Then it is a faithful non-singular left module over itself. It has only one non-zero injective left ideal, namely

$$\begin{pmatrix} 0 & K \\ 0 & K \end{pmatrix},$$

and this is also a faithful  $R$ -module. Hence  $R$  is a QF-3'  $R$ -module with

$$R^* = \begin{pmatrix} 0 & K \\ 0 & K \end{pmatrix}.$$

There is no faithful left ideal of  $R$  properly contained in  $R^*$ , so we have  $R_0=R^*$ . Moreover  $R=R^* \oplus R'$ , where

$$R' = \begin{pmatrix} K & 0 \\ 0 & 0 \end{pmatrix}$$

and is not QF-3'.

EXAMPLE 3.10. Let  $R$  be as above and  $Q$  the ring of all  $2 \times 2$  matrices over  $K$ . Then  $Q$  is also a faithful non-singular  $R$ -module and is QF-3' since  $Q=E({}_R R)$ . In this case,  $Q=Q^*$  and we may take for  $Q_0$ , for example, as

$$\begin{pmatrix} K & 0 \\ K & 0 \end{pmatrix}, \begin{pmatrix} 0 & K \\ 0 & K \end{pmatrix}, \text{ or } Q = \begin{pmatrix} K & K \\ K & K \end{pmatrix}.$$

Hence the submodule  $Q_0$  in the theorem is not uniquely determined within isomorphisms.

REMARK. Let  $Q$  be a faithful, non-singular QF-3'  $R$ -module. Then there exist faithful submodules  $Q^*$  and  $Q_0$  of  $Q$  with properties mentioned in Theorem 3.7. As was pointed out in the proof of the theorem,  $k_{Q^*} = k_{Q_0} = k_Q$  hold and hence by Theorem 1.4 both  $Q^*$  and  $Q_0$  are also QF-3'. These, as well as  $Q$  and  $E(Q)$ , are faithful, non-singular QF-3'  $R$ -modules. Clearly  $Q^*$  includes  $Q_0$  and moreover it is a unique maximal one of those submodules of  $Q$  which contain the injective hull of every one of its finitely generated submodules. Since each injective submodule of  $Q$  is that of  $Q^*$ , we can conclude that  $Q^*$  coincides with the sum of all non-zero injective submodules of  $Q^*$ , i.e.,  $(Q^*)^* = Q^*$ .

Let us suppose furthermore that every direct sum of non-singular injective  $R$ -modules is injective. For example, we may take a finite dimensional ring  $R$  in the sense that it contains no infinite direct sum of submodules. Then  $Q^*$  is itself injective and hence  $Q$  can be decomposed into a direct sum of submodules  $Q^*$  and  $Q'$ :  $Q = Q^* \oplus Q'$ . Since  $Q^*$  is a unique maximal non-zero injective submodule of  $Q$ , if  $Q' \neq 0$ , then  $Q'$  does not contain any non-zero injective submodule of  $Q$ . Therefore by Lemma 3.8  $\text{Hom}_R(E(Q'), Q') = 0$ . This shows that  $Q'$  can not be QF-3'.

YAMAGUCHI UNIVERSITY

---

### References

- [1] G. Azumaya: *Completely faithful modules and self-injective rings*, Nagoya Math. J. **27** (1966), 697-708.
- [2] L. Bican: *QF-3' modules and rings*, Comment. Math. Univ. Carolinae **14** (1973), 295-303.
- [3] V.C. Cateforis: *Minimal injective cogenerators for the class of modules of zero singular submodule*, Pacific J. Math. **40** (1972), 527-539.
- [4] R.R. Colby and E.A. Rutter, Jr.: *Semi-primary QF-3 rings*, Nagoya Math. J. **32** (1968), 253-258.
- [5] S.E. Dickson: *A torsion theory for abelian categories*, Trans. Amer. Math. Soc. **121** (1966), 223-235.
- [6] J.P. Jans: *Some aspects of torsion*, Pacific J. Math. **15** (1965), 1249-1259.
- [7] ———: *Torsion associated with duality*, Tôhoku Math. J. **24** (1972), 449-452.
- [8] T. Kato: *Torsionless modules*, Tôhoku Math. J. **20** (1968), 233-242.
- [9] Y. Kurata: *On an  $n$ -fold torsion theory in the category  ${}_R\mathbf{M}$* , J. Algebra **22** (1972), 559-572.
- [10] K. Morita: *Localizations in categories of modules. I*, Math. Z. **114** (1970), 121-144.

- [11] B. Stenström: *Rings and Modules of Quotients*, Lecture Notes in Math. **237**, Springer-Verlag, Berlin-Heidelberg-New York, 1971.
- [12] K. Sugano: *A note on Azumaya's theorem*, Osaka J. Math. **4** (1967), 157–160.
- [13] H. Tachikawa: *On left QF-3 rings*, Pacific J. Math. **32** (1970), 255–268.
- [14] G.M. Tsukerman: *Pseudo-injective modules and self-pseudo-injective rings*, Math. Notes. **7** (1970), 220–226.
- [15] D.Y. Wei: *On the concept of torsion and divisibility for general rings*, Illinois J. Math. **13** (1969), 414–431.
- [16] L.E.T. Wu, H.Y. Mochizuki and J.P. Jans: *A characterization of QF-3 rings*, Nagoya Math. J. **27** (1966), 7–13.