



Title	Complex cases with suspected dementia in the community need psychiatric support: Results from a nationwide survey in Japan
Author(s)	Satake, Yuto; Taomoto, Daiki; Suzuki, Maki et al.
Citation	Asian Journal of Psychiatry. 2024, 91, p. 103840
Version Type	AM
URL	https://hdl.handle.net/11094/94005
rights	© 2023. This manuscript version is made available under the CC-BY-NC-ND 4.0 license https://creativecommons.org/licenses/by-nc-nd/4.0/
Note	

The University of Osaka Institutional Knowledge Archive : OUKA

<https://ir.library.osaka-u.ac.jp/>

The University of Osaka

1 The Initial-phase Intensive Support Team for Dementia (IPIST) is a multidisciplinary
2 outreach team that provides intensive initial assessment and support for dementia in Japan,
3 introduced in reference to the UK Memory Service.^{1,2} The Japanese government launched a
4 model project of IPIST in 2013 and subsequently mandated that all municipalities, including
5 cities, towns or villages, establish at least one team, which was completed in September
6 2019. One of the most distinctive features of IPIST is officially acknowledged outreach
7 function. Each IPIST consists of at least one physician and two non-physician professional
8 staff, such as public health nurses and care workers. When IPISTs receive a request, they
9 conduct a home visit, followed by a team meeting, making a support plan, consultation with a
10 medical institution, practical support for the individual's life at home, family support,
11 introduction of public support, and information sharing at community care meetings.³ This
12 process is usually completed within six months and community services and other social
13 resources take over the support. In principle, eligible persons are residents at home aged 40 or
14 older who are suspected of having dementia or who have dementia and who fall into any of
15 the following categories: 1) Those who are not receiving medical or long-term care services,
16 or those who have suspended the services they were receiving, and 2) Those who are
17 receiving medical and long-term care services but are difficult-to-treat because of severe
18 behavioral and psychological symptoms of dementia (BPSD).

19

20 While memory services in the UK do not normally handle cases with complex behavioral or
21 psychological problems,⁴ approximately 40% of IPIST cases in Japan are “complex cases”
22 that are extremely difficult to approach or connect to medical or long-term care.⁵ In order to
23 effectively support the older residents in communities, it is essential to understand the causes
24 of complex cases and optimize IPIST activities accordingly. Therefore, we conducted a
25 nationwide survey on this point.

26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50

From January 14 to March 8, 2021, our original questionnaire was distributed electronically to each IPIST through all 1741 municipalities in Japan and responses were collected. The questionnaire assessed the characteristics of each IPIST, including specialty of team doctors and availability of staff with psychiatric expertise, and which out of 12 categories of causes each complex case had from April to September 2020, allowing multiple choice. The categories formulated by ourselves are listed in Table 1. This study was performed in accordance with the Declaration of Helsinki. This human study was approved by The Ethical Review Board Osaka University Hospital (Suita, Japan) - approval: 20419-3.

We collected responses from 1291 IPISTs. 43.3% of IPISTs had a psychiatrist. 59.4% of the teams had medical staff members with work experience in psychiatric services, including psychiatrists. In a total of 7340 cases, all categories of causes for difficulties were reported (Table 1). Additional information is shown in Supplementary Table 1 and 2.

This survey revealed complex cases of IPIST had many factors requiring psychological and psychiatric intervention. The three refusal issues of “refusal of public support”, “refusal to see a doctor” and “refusal of IPISTs' visiting” together accounted for 42.8% of all answers; “BPSD”, “complications of mental illness” and “mental illness in the persons living with them” together accounted for 26.1%. The reason for not wanting to receive public support is suspected to be related to psychological concerns about not wanting to undermine the sense of independence in one's life,⁶ and some caregivers may refuse to receive support by various individual factors.⁷ Greater stigma leads to refusal of dementia diagnostic assessment.⁸ Tailored approaches to take their needs and explain the requirements of assessment and support through establishing a trusting relationship by skillful professionals may alleviate

51 these refusals. BPSD and mental illness found in the clients and their families must need
52 psychiatric intervention. Our results indicating the high prevalence of those problems were
53 consistent with the previous reports about the outreach psychiatric service in Tokyo.⁹
54 Fortunately, psychiatrists were the most common team doctors, with as many as 59.4% of all
55 teams having medical experts with clinical experience in psychiatry, including psychiatrists.
56 IPIST's human resources may have been naturally optimized to some extent to reflect the
57 needs of the frontline for complex cases.

58

59 The current study revealed that the Japanese outreach team for dementia suspected
60 community dwelling older people faced a lot of complex cases with issues that would be
61 related to psychological and psychiatric problems. With disease modifying therapies
62 becoming a treatment option, the need for early detection of dementia patients including
63 prodromal stage with psychiatric symptoms will increase. Besides, assistive technologies
64 which may be an option to support patients with neurocognitive disorders in near future
65 probably also need supportive visiting staffs¹⁰. Mental health professionals with interpersonal
66 relationship-building skills and knowledge of mental illness will play more crucial role in
67 dementia related outreach fields.

68

69 **References:**

- 70 1 Banerjee S, Willis R, Matthews D, Contell F, Chan J, Murray J. Improving the quality
71 of care for mild to moderate dementia: an evaluation of the Croydon Memory Service Model.
72 *Int J Geriatr Psychiatry* 2007; **22**:782–8.
- 73 2 Department of Health. 291591a/Living well with dementia: A National Dementia
74 Strategy 2009.
75 https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment_data
76 [/file/168220/dh_094051.pdf](https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment_data/file/168220/dh_094051.pdf) (accessed Jun 27, 2023).
- 77 3 Takeda A. Outline of Initial-Phase Intensive Support Team. *Neurological*
78 *Therapeutics* 2017; **34**: 294–7.
- 79 4 Rubinsztein JS, van Rensburg MJ, Al-Salihy Z, Girling D, Lafortune L,
80 Radhakrishnan M, et al. A memory clinic v. traditional community mental health team
81 service: comparison of costs and quality. *BJPsych Bull* 2015; **39**: 6–11.
- 82 5 Awata S. Initial-phase intensive support team and community general support center:
83 From the perspective of coordination and networking of social support. *Nippon Ronen*
84 *Igakkai Zasshi* 2020; **57**: 22–7.
- 85 6 Abdi S, Spann A, Borilovic J, De Witte L, Hawley M. Understanding the care and
86 support needs of older people: a scoping review and categorisation using the WHO
87 international classification of functioning, disability and health framework (ICF). *BMC*
88 *Geriatr* 2019; **19**: 195.
- 89 7 Zwingmann I, Dreier-Wolfgramm A, Esser A, Wucherer D, Thyrian JR, Eichler T, et
90 al. Why do family dementia caregivers reject caregiver support services? Analyzing types of
91 rejection and associated health-impairments in a cluster-randomized controlled intervention
92 trial. *BMC Health Serv Res* 2020; **20**: 121.

93 8 Fowler NR, Frame A, Perkins AJ, Gao S, Watson DP, Monahan P, et al. Traits of
94 patients who screen positive for dementia and refuse diagnostic assessment. *Alzheimers*
95 *Dement Diagn Assess Dis Monit* 2015; **1**: 236–41.

96 9 Ito K, Okamura T, Tsuda S, Ogisawa F, Awata S. Characteristics of complex cases of
97 community-dwelling older people with cognitive impairment: A classification and its
98 relationships to clinical stages of dementia. *Geriatr Gerontol Int* 2022; **22**: 997–1004.

99 10 Lee-Cheong S, Amanullah S, Jardine M. New assistive technologies in dementia and
100 mild cognitive impairment care: A PubMed review. *Asian J Psychiatry* 2022; **73**: 103135.

101

102 **Table 1. Causes for difficulties in IPISTs' complex cases**

Causes	n	%
Refusal of public support	1433	19.5
BPSD	1176	16.0
Refusal to see a doctor	1166	15.9
Refusal of IPISTs' visiting	543	7.4
Complications of mental illness	534	7.3
Complaints from neighbors	518	7.1
Financial issues	484	6.6
No caregivers	453	6.2
Others	363	4.9
Trash-house	314	4.3
Mental illness in the persons living with them	205	2.8
Abuse	151	2.1
Total	7340	100.0

103 Abbreviation: BPSD, behavioral and psychological symptoms of dementia; IPIST, Initial-
 104 phase Intensive Support Team for Dementia. The percentages above represent values for the
 105 total number of causes for difficulties collected.