



Title	Markov processes associated with certain integro-differential operators
Author(s)	Komatsu, Takashi
Citation	Osaka Journal of Mathematics. 1973, 10(2), p. 271-303
Version Type	VoR
URL	https://doi.org/10.18910/9509
rights	
Note	

The University of Osaka Institutional Knowledge Archive : OUKA

<https://ir.library.osaka-u.ac.jp/>

The University of Osaka

MARKOV PROCESSES ASSOCIATED WITH CERTAIN INTEGRO-DIFFERENTIAL OPERATORS

TAKASHI KOMATSU

(Received June 27, 1972)

Introduction. Let us consider a linear operator L :

$$(1) \quad L v(s, x) = \frac{1}{2} \sum_{ij} a_{ij}(s, x) D_i D_j v(s, x) + \sum_i b_i(s, x) D_i v(s, x) \\ + \int [v(s, x+u) - v(s, x) - I_{(|u| \leq 1)}(u, \nabla v(s, x))] S(s, x, du),$$

where $D_i = \partial / \partial x_i$, $\nabla = (D_i)$, a is a non-negative definite $d \times d$ -matrix, b is a d -vector and S is a *Lévy kernel*, that is, a kernel satisfying $S(s, x, \{0\}) = 0$ and

$$\int_{\mathbb{R}^d} |u|^2 \wedge 1 S(s, x, du) < \infty$$

In case $S \equiv 0$, the Markov process on \mathbb{R}^d having L as its generator can be constructed by solving Ito's stochastic equation

$$(2) \quad dX_t = a(t, X_t)^{1/2} dB_t + b(t, X_t) dt,$$

where B_t is the d -dimensional Brownian motion. It is well known that if a and b are continuous in (s, x) , then there exists at least a solution of (2). Roughly speaking, the problem whether the equation (2) has a unique solution in the sense of probability law corresponds to the analytical problem whether the equation

$$\left(\lambda - \frac{\partial}{\partial s} - L \right) v = f$$

has solutions v for each λ (\geq some λ_0) and for sufficiently wide class of functions f . We know that if a is continuous in (s, x) and is positive definite, then the above equation has a solution v (in the sense of distribution) for each $\lambda > 0$ and test function f . Using this fact, Strook-Varadhan [8] proved that the equation (2) has a unique solution in the sense of probability law. Their method is basically analytic but it also needs some probabilistic arguments. They introduced and made use of certain martingale equations equivalent to the equation (2).

In this paper, we shall consider the case where S is not always 0. In §2, three equivalent martingale formulations of the stochastic equation associated with the operator L are defined, and we call each of them '*the (a, b, S) -stochastic equation*'. The main aim of this paper is to prove the uniqueness and existence of the solution of the (a, b, S) -stochastic equation. Our results are as follows. To explain them, let us introduce some conditions:

I. for each bounded domain $D \subset R_+ \times R^d$, there exist positive constants a_1 and a_2 such that $a_1 |\theta|^2 \leq (\theta, a(s, x)\theta) \leq a_2 |\theta|^2$ for each $\theta \in R^d$ and $(s, x) \in D$,
 II. $\limsup_{x' \rightarrow x} \sum_{i,j} |a_{ij}(s, x') - a_{ij}(s, x)| = 0$ for each T and $x \in R^d$; $b(s, x)$ is locally bounded; and, for each bounded domain $D \subset R_+ \times R^d$, there exists a measure $\bar{S}(du)$ such that

$$\int |u|^2 \wedge 1 \bar{S}(du) < \infty$$

and $\bar{S}(du) - S(s, x, du)$ is a non-negative measure for $(s, x) \in D$,

III. for each bounded interval $[0, T]$, there exists a constant K such that

$S(s, x, \{|u| > 1\}) \leq K$ and

$$|(x, b(s, x))| + \text{trace } a(s, x) + \int_{|u| \leq 1} |u|^2 S(s, x, du) \leq K(1 + |x|^2 \log^+ |x|)$$

for all $(s, x) \in [0, T] \times R^d$. (We shall give a more general condition in §5.)

If conditions I and II are satisfied, then the (a, b, S) -stochastic equation has at most a solution; if condition I, II and III are satisfied, then there exists uniquely a solution of the equation; and if condition II and III are satisfied and a, b, S are continuous in x , then the equation has at least a solution.

The proof of the uniqueness is progressed in the same way as in Strook-Varadhan [8]. But, in our case, there arise some difficulties, for the condition II is very weak and $a(s, x)$ is not always continuous in s . Our proof of the existence is based on Hille-Yosida's semi-group theory. The merit of the way is in the fact that one can weaken continuity condition for a, b and S (especially with respect to the time variable s).

Finally we should mention that Tsuchiya [10] disposed of a similar problem in a different context. In his case, $a = a(x)$, $b = b(x)$ and $S = K(x, u) |u|^{-d-\alpha} du$ where K is a positive and bounded function and $1 < \alpha < 2$, and he considered two cases: i) $a(x)$ is positive definite, ii) $a(x)$ is identically 0. In Case i), his results are included in ours. But Case ii) is quite different; Tsuchiya solved the problem by making use of a perturbation method based on the α -stable process.

I wish to give my thanks to Professor T. Watanabe who kindly gave me many suggestions in the course of my research.

1. Notation and preliminaries

We shall consider the space W_t^s of trajectories consisting of right continuous functions admitting limits from the left, defined on $[s, t] \subset R_+$, with values in R^d . The value of a function w at time t is denoted by $x_t(w)$. Let W_t^s be the σ -field generated by $(x_\tau; s \leq \tau \leq t)$. The function space W_t^s becomes a complete separable metric space by the Billingsley metric and the σ -field W_t^s coincides with the σ -field of Borel sets of the metric space. Let W^s be the space of functions w on $[s, \infty)$ satisfying $w(\tau \wedge t) \in W_t^s$ for each $t \geq s$. The σ -field W_t^s is identified with the σ -field in W^s generated by $(x_\tau; s \leq \tau \leq t)$.

Lemma 1.1. (Skorokhod [7]) *Let Q be a family of probability measures on (W_t^s, W_t^s) . If the family Q satisfies the condition:*

$$\lim_{l \rightarrow \infty} \sup_{Q \in Q} Q(\sup_{\tau} |x_\tau| > l) = 0 \text{ and}$$

$$\lim_{\delta \downarrow 0} \sup_{Q \in Q} \sup_{|\tau - \tau'| \leq \delta} Q(|x_\tau - x_{\tau'}| > \varepsilon) = 0 \text{ for any } \varepsilon > 0,$$

then it is possible to pick up a sequence (Q^n) from Q and is possible to construct a sequence (X_τ^n) of processes and a process X_τ which are defined on a certain probability space (Ω', F', P') such that a) for each n , the processes (x, Q^n) and (X_τ^n, P') are equivalent, b) the process X_τ is stochastically continuous and the sequence (X_τ^n) of random variables converges to the random variable X_τ in probability for any τ .

We call $T(w)$ an s -stopping time if $s \leq T$ and the set $\{T \leq t\}$ is W_t^s -measurable for all $t \geq s$. The σ -field W_T^s , defined as the collection of sets $\{A \in W^s; A \cap (T \leq t) \in W_t^s \text{ for all } t \geq s\}$, coincides with the σ -field generated by $(x_{t \wedge T}; t \geq s)$. The σ -field W_T^s admits 'regular conditional probabilities'. (see Parthasarathy [6])

Lemma 1.2. (Strook-Varadhan [8]) *Let T be a finite s -stopping time. Let Q' be a probability measure on (W^s, W^s) , and for each $w \in W^s$, let Q_w'' be a probability measure on $(W^s, W^{T(w)})$ where $W^{T(w)}$ is the σ -field generated by $(x_t(w'); t \geq T(w))$. Suppose that*

- 1) $Q_w''(w'; x_{T(w)}(w) = x_{T(w)}(w')) = 1$ for each $w \in W$,
- 2) for each $t \geq s$ and $A \in W_t^s$, $w \mapsto Q_w''(A) I_{\{T(w) \leq t\}}$ is W_t^s -measurable.

Then there exists a unique probability measure Q on (W^s, W^s) such that $Q = Q'$ on W_T^s , and for almost all w (w.r.t. Q') the regular conditional probability of Q given the σ -field W_T^s coincides with Q_w'' on $W^{T(w)}$.

Let (Ω, F, P) be a complete probability space with a non-decreasing and right continuous family $(F_t)_{t \geq 0}$ of sub- σ -fields of F . Without loss of generality, we can suppose that the σ -field F_0 contains all the negligible sets of F . Moreover, we assume that the family (F_t) has no time of discontinuity. In the rest of this

section, we suppose that each real valued process is adapted to the family (F_t) and its paths are right continuous, have limits from the left and equals 0 at time 0. We shall pick up some notation and remarks from Meyer [5]- I.

Notation and remarks. We say that a process X_t is '*natural*' if $\Delta X_T \equiv X_T - X_{T-} = 0$ for any totally inaccessible stopping time T . Each martingale is *quasi-left-continuous*, for the family (F_t) has no time of discontinuity. Therefore each martingale has no common jump with any natural process. We say that a process X_t is '*locally integrable*' if there exists a sequence (T_n) of stopping times such the $T_n \uparrow \infty$ as $n \rightarrow \infty$ and each stopped process $X_{t \wedge T_n}$ is uniformly integrable.

1) We denote by A_{loc} the space of all processes whose total variations on any bounded intervals are finite. Each process $X_t \in A_{loc}$ is decomposed to the sum of a continuous process and a purely discontinuous process as follows:

$$X_t = [X_t - \sum_{\tau \leq t} \Delta X_\tau] + \sum_{\tau \leq t} \Delta X_\tau \quad \text{where } \Delta X_\tau = X_\tau - X_{\tau-}$$

We denote the subspace of A_{loc} consisting of all continuous (resp. purely discontinuous) processes by A_{loc}^c (resp. A_{loc}^d). The decomposition $A_{loc} = A_{loc}^c + A_{loc}^d$ is always direct, and the decomposition $A_{loc}^n = A_{loc}^c + A_{loc}^{nd}$ is also direct, where the spaces with superfix '*n*' express the subspaces consisting of natural processes. In the sequel, we shall consider the subspace $A_{loc}^q = \{A_t \in A_{loc}; A_t \text{ is quasi-left continuous}\}$ instead of the space A_{loc} . If $A_t \in A_{loc}^q$ is locally integrable, then we can uniquely choose $B_t \in A_{loc}^c$ such that $A_t - B_t$ is a locally integrable martingale. We denote the process B_t (resp. $A_t - B_t$) by $\langle A_t \rangle$ (resp. ${}^c A_t$).

2) We denote the space of all locally integrable martingales by M_{loc} . Let X_t be in M_{loc} and let a sequence (T_n) of stopping times carry all the jumps of X_t , that is to say, $P[T_n = T_m < \infty] = 0$ for any $n \neq m$ and $\{(w, t); X_t(w) \neq X_{t-}(w)\} \subset \bigcup_{n=1}^{\infty} (\text{graph } T_n)$. We call the process $\sum_{n=1}^{\infty} {}^c(\Delta X_{T_n} I_{(T_n \leq t)})$, which is formally denoted by ${}^c(\sum_{\tau \leq t} \Delta X_\tau)$, the '*purely discontinuous local martingale*' part of X_t . Let M_{loc}^c (resp. M_{loc}^d) be the subspace consisting of all continuous (resp. purely discontinuous) local martingales, then we have $M_{loc} = M_{loc}^c + M_{loc}^d$ (direct sum). If M_t and N_t in M_{loc} are locally square integrable, then there exists a unique $A_t \in A_{loc}^c$ such that $M_t N_t - A_t \in M_{loc}$. The process A_t is denoted by $\langle M_t, N_t \rangle$.

Every process $X_t = M_t + A_t$ with $M_t \in M_{loc}$ and $A_t \in A_{loc}$ is called a '*weak local semi-martingale*'.

Let $C^n(D)$ (resp. $C^{n,b}(D)$) be the space of functions on a domain D whose r -th ($0 \leq r \leq n$) derivatives are continuous (resp. bounded and continuous). The formula of change of variables on semi-martingale (see Kunita-Watanabe [4] and Meyer [5]) gives a base of the discussions in this paper.

Theorem 1.3. (Kunita-Watanabe's formula) *Let $X_t=(X_t^i)$ be a d -dimensional weak local semi-martingale such that*

$$X_t^i - X_0^i = M_t^i + N_t^i + A_t^i + B_t^i \text{ with } M_t^i \in \mathbf{M}_{loc}^c, N_t^i \in \mathbf{M}_{loc}^d, A_t^i \in \mathbf{A}_{loc}^c \text{ and } B_t^i \in \mathbf{A}_{loc}^d,$$

and that $\Delta N_t^i \Delta B_t^j = 0$ for all i, j and t . If $F(x) \in C^2(R^d) \cap C^{1,b}(R^d)$, then the process $F(X_t) - F(X_0)$ has the unique decomposition:

$F(X_t) - F(X_0) = M_t' + N_t' + A_t'$ with $M_t' \in \mathbf{M}_{loc}^c$, $N_t' \in \mathbf{M}_{loc}^d$ and $A_t' \in \mathbf{A}_{loc}^c$, where

$$\begin{aligned} M_t' &= \sum_{i=1}^d \int_0^t D_i F(X_\tau) dM_\tau^i, \\ N_t' &= {}^c \left(\sum_{\substack{\tau \leq t \\ \Delta N_\tau \neq 0}} [F(X_\tau) - F(X_{\tau-})] \right) + {}^c \left(\sum_{\substack{\tau \leq t \\ \Delta B_\tau \neq 0}} [F(X_\tau) - F(X_{\tau-})] \right), \\ A_t' &= \int_0^t \left(\frac{1}{2} \sum_{i,j=1}^d D_i D_j F(X_\tau) d\langle M_\tau^i, M_\tau^j \rangle + \sum_{i=1}^d D_i F(X_\tau) dA_\tau^i \right) \\ &\quad + \left\langle \sum_{\substack{\tau \leq t \\ \Delta N_\tau \neq 0}} [F(X_\tau) - F(X_{\tau-})] - \sum_{i=1}^d D_i F(X_{\tau-}) \Delta N_\tau^i \right\rangle + \left\langle \sum_{\substack{\tau \leq t \\ \Delta B_\tau \neq 0}} [F(X_\tau) - F(X_{\tau-})] \right\rangle. \end{aligned}$$

The following lemma is useful.

Lemma 1.4. *If a process X_t is uniformly integrable and $E[X_T] = 0$ for any bounded stopping time T , then the process is a martingale.*

Proof. Let $s < t$ and $A \in \mathbf{F}_s$. Considering the stopping time $T = sI_A + tI_{A^c}$, we have $E[X_T] = E[X_s I_A] + E[X_t I_{A^c}] = 0$. On the other hand $E[X_t] = E[X_t I_A] + E[X_t I_{A^c}] = 0$. Thus, $E[X_t I_A] = E[X_s I_A]$.

Notation. The spaces \mathbf{A}_{loc} , \mathbf{M}_{loc} etc. are also defined for d -dimensional real valued processes (resp. complex valued processes) in such a way that each coordinate (resp. real and imaginary parts of each coordinate) belongs to the spaces \mathbf{A}_{loc} , \mathbf{M}_{loc} etc. of the previous sense. We often denote the spaces \mathbf{A}_{loc} , \mathbf{M}_{loc} etc. and the expectation E with respect to $(\Omega, \mathbf{F}, \mathbf{F}_t, P)$ by $\mathbf{A}_{loc}(\mathbf{F}_t, P)$, $\mathbf{M}_{loc}(\mathbf{F}_t, P)$ etc. and by E_P respectively, lest we should get confused.

Let S and S' be kernels. It is simply denoted by $S \leq S'$ the fact that $S' - S$ is non-negative. And $|S - S'|$ stands for the total variation of the signed measure $S' - S$.

2. Martingale problems of stochastic equations

Let (Ω, \mathbf{F}, P) and $(\mathbf{F}_t)_{t \geq 0}$ be the object as we stated in §1. Let us choose

a d -dimensional Brownian motion B_t , a *Poisson random measure* $p(dt, dz)$ on $R_+ \times R^d$ with parameter $dt \times dz/|z|^{d+1}$ and the *Poisson martingale measure* $q(dt, dz)$. These are defined as follows: for each Borel set A in $R^d - \{0\}$, the process $p([0, t] \times A)$ is a Poisson process adapted to (F_t) which is independent of B_t and satisfies that

$$E[p([0, t] \times A)] = \int_0^t \int_A \frac{d\tau dz}{|z|^{d+1}}.$$

And $q(dt, dz) = p(dt, dz) - dt dz/|z|^{d+1}$.

For each $s \in R_+$, $x, z \in R^d$, let $\sigma(s, x)$ denote a $d \times d$ -matrix, $b(s, x)$ and $c(s, x, z)$ d -vectors. We suppose that σ , b and c are Borel measurable. Let T be an s -stopping time and let $(X_t; t \in [s, T])$ be an R^d -valued right continuous process admitting limits from the left and adapted to (F_t) , where $[s, T] = \{t < \infty; s \leq t \leq T\}$. From now on, we shall assume these conditions without any assignment.

DEFINITION. We call a process X_t a solution of the *original* (σ, b, c) -stochastic equation starting from (s, x) , constructed over $(\Omega, F, F_t, P, B_t, p, q)$ if

$$\begin{aligned} \text{S0)} \quad X_t = x &+ \int_s^t \sigma(\tau, X_\tau) dB_\tau + \int_s^t b(\tau, X_\tau) d\tau + \int_s^t \int c^1(\tau, X_\tau, z) q(d\tau, dz) \\ &+ \int_s^t \int c_1(\tau, X_\tau, z) p(d\tau, dz) \quad \text{for all } t \in [s, T], \end{aligned}$$

where $c^1 = cI_{(|c| \leq 1)}$ and $c_1 = cI_{(|c| > 1)}$.

We assume that

$$\int_s^t \left\{ \text{trace}(\sigma\sigma^*)(\tau, X_\tau) + |b(\tau, X_\tau)| + \int |c(\tau, X_\tau, z)|^2 \wedge 1 \frac{dz}{|z|^{d+1}} \right\} d\tau < \infty$$

for all $t \in [s, T]$, where the domain of integration by dz is the set $R^d - \{0\}$.

Let $a(s, x)$ be a measurable and non-negative definite $d \times d$ -matrix and $S(s, x, du)$ be a *Lévy kernel*. Let us introduce an integro-differential operator L defined by, for each $v \in C^{2,b}(R_+ \times R^d)$,

$$\begin{aligned} Lv(s, x) &= \frac{1}{2} \sum_{ij} a_{ij} D_i D_j v(s, x) + \sum_i b_i D_i v(s, x) \\ &+ \int [v(s, x+u) - v(s, x) - I_{(|u| \leq 1)}(u, \nabla v(s, x))] S(s, x, du). \end{aligned}$$

Set $\Phi_\theta(s, x) = e^{-i(\theta, x)} L e^{i(\theta, \cdot)}(s, x)$. Then we have

$$\Phi_\theta(s, x) = -\frac{1}{2}(\theta, a(s, x)\theta) + i(\theta, b(s, x)) + \int [e^{i(\theta, u)} - 1 - I_{(|u| \leq 1)} i(\theta, u)] S(s, x, du).$$

Now suppose that $a = \sigma\sigma^*$ (σ^* stands for the transposed matrix of σ),

$$S(s, x, D) = \int_{c(s, x, z) \in D} \frac{dz}{|z|^{d+1}} \text{ for each open set } D \subset R^d - \{0\},$$

and that $(X_t, P; t \in [s, T])$ is a solution of the original (σ, b, c) -stochastic equation starting from (s, x) . Then we have

$$v(t, X_t) - v(s, x) - \int_s^t \left(\frac{\partial}{\partial \tau} + L \right) v(\tau, X_\tau) d\tau \in M_{loc}(F_t, P)$$

for all $v \in C^{2,b}(R_+ \times R^d)$. This can be verified by the Kunita-Watanabe formula (Th. 1.3.).

Notation and remarks. Let $J(dt, du)$ be the random measure defined by

$$J(dt, du) = \sum_{s \in dt} I_{\{dX_s \in du - \{0\}\}}.$$

Then it holds that

$$E \left[\int_s^{T'} \int h(t, X_{t-}, u) J(dt, du) \right] = E \left[\int_s^{T'} \int h(t, X_t, u) S(t, X, du) dt \right].$$

for any non-negative and measurable function $h(s, x, u)$ and s -stopping time T' ($T' \leq T$); where if any one of the members is integrable, so is the other. Set, for $p=1, 2$,

$$H^p = \left\{ h(s, x, u); \int_s^t |h(\tau, X_\tau, u)|^p S(\tau, X_\tau, du) d\tau < \infty \text{ a. e. for all } t \in [s, T] \right\}.$$

If h is a function of H^1 , then the purely discontinuous process

$$\int_s^t h(\tau, X_\tau, u) J(d\tau, du)$$

is locally integrable. By Lemma 1.4, we have

$$\left\langle \int_s^t h(\tau, X_\tau, u) J(d\tau, du) \right\rangle = \int_s^t \int h(\tau, X_\tau, u) S(\tau, X_\tau, du) d\tau.$$

Set ${}^c J(dt, du) = J(dt, du) - S(t, X_t, du)dt$. Then stochastic integrals by ${}^c J$ are local martingales, therefore we shall say that ${}^c J$ is a 'martingale measure'. If h is a function of H^2 , then the locally square integrable martingale:

$$\int_s^t h(\tau, X_\tau, u) {}^c J(d\tau, du) = {}^c \left(\sum_{\substack{s < \tau \leq t \\ \Delta X_\tau \neq 0}} h(\tau, X_{\tau-}, \Delta X_\tau) \right)$$

is defined. And if h_1 and h_2 are elements of H^2 , then we have

$$\left\langle \int_s^t h_1 {}^c J, \int_s^t h_2 {}^c J \right\rangle = \int_s^t h_1 h_2 S d\tau.$$

In order to give the martingale formulation of stochastic equation, we need the following theorem. (a, b, S are given and L, Φ_θ are defined as before.)

Theorem 2.1. *Let $(X_t, P; t \in [s, T])$ be a process such that $\lim_{l \rightarrow \infty} P[\sup_{\tau \leq t} |X_\tau| > l] = 0$ and*

$$\int_s^t \{ \text{trace } a(\tau, X_\tau) + |b(\tau, X_\tau)| + \int |u|^2 \wedge 1 S(\tau, X_\tau, du) \} d\tau < \infty$$

for all $t \in [s, T]$. Then the following three conditions are equivalent.

S1) *For any $v(s, x) \in C^{2,b}(R_+ \times R^d)$,*

$$v(t, X_t) - v(s, x) - \int_s^t \left(\frac{\partial}{\partial \tau} + L \right) v(\tau, X_\tau) d\tau \in \mathbf{M}_{loc}(\mathbf{F}_t, P).$$

S2) *For all $\theta \in R^d$,*

$$\exp[i(\theta, X_t - x) - \int_s^t \Phi_\theta(\tau, X_\tau) d\tau] - 1 \in \mathbf{M}_{loc}(\mathbf{F}_t, P).$$

S3) *X_t is a weak local semi-martingale which has the following property:*

a) *for any positive measurable function h and for any s -stopping time T' ($T' \leq T$),*

$$E \left[\int_s^{T'} h(\tau, X_\tau, u) J(d\tau, du) \right] = E \left[\int_s^{T'} h(\tau, X_\tau, u) S(\tau, X_\tau, du) d\tau \right]$$

if a measure J of jumps of the process X_t is defined as before. That is to say, $J(dt, du) - S(t, X_t, du) dt$ is a martingale measure (which we shall denote by ${}^c J(dt, du)$).

b) *there exists a d -dimensional process $M_t = (M_t^i)$ such that*

$$M_t^i \in \mathbf{M}_{loc}^c(\mathbf{F}_t, P), \quad \langle M_t^i, M_t^j \rangle = \int_s^t a_{ij}(\tau, X_\tau) d\tau,$$

and the process X_t can be decomposed as follows:

$$X_t = x + M_t + \int_s^t b(\tau, X_\tau) d\tau + \int_s^t \int_{|u| \leq 1} u {}^c J(d\tau, du) + \int_s^t \int_{|u| > 1} u J(d\tau, du).$$

Proof. 1° Condition S1) implies the next condition:

S2') *for all $\theta \in R^d$,*

$$\exp[i(\theta, X_t - x)] - 1 - \int_s^t \exp[i(\theta, X_\tau - x)] \Phi_\theta(\tau, X_\tau) d\tau \in \mathbf{M}_{loc}(\mathbf{F}_t, P).$$

By the Kunita-Watanabe formula, it is easily proved that Condition S2) and S2') are equivalent.

2° Next, we shall show that Condition S2') implies Condition S3). Set

$$T_n = \inf \{t \in [s, T]; |X_t| > n\}.$$

Each stopped process $X_{t \wedge T_n}$ can be decomposed as follows:

$$X_{t \wedge T_n} = Y_t^n + \Delta X_{T_n} I_{(t \geq T_n)}, \sup_t |Y_t^n| \leq n$$

If $|\theta| < 2\pi/n$, then $i(\theta, Y_t^n) = \log \exp [i(\theta, Y_t^n)]$. Since the process $\exp [i(\theta, Y_t^n)]$ is a weak local semi-martingale, the process $i(\theta, Y_t^n)$ is a weak local semi-martingale. From the fact that $T_n \uparrow T$, the process X_t is a weak local semi-martingale. It is easy to show that the process X_t has no natural jump, in other words, the process X_t is quasi-left-continuous. Thus the process X_t is decomposed as follows:

$$X_t = x + M_t + A_t + N_t + B_t, \quad M_t \in \mathbf{M}_{loc}^c, \quad A_t \in \mathbf{A}_{loc}^c, \quad N_t \in \mathbf{M}_{loc}^d, \quad B_t \in \mathbf{A}_{loc}^d$$

We suppose that $|\Delta N_t| \leq 1$ (resp. $|\Delta B_t| > 1$) if $\Delta N_t \neq 0$ (resp. $\Delta B_t \neq 0$).

By the Kunita-Watanabe formula, we have

$$e^{i(\theta, X_t - x)} - 1 = [\text{a local martingale taking value 0 at time } s]$$

$$\begin{aligned} & + \int_s^t e^{i(\theta, X_\tau - x)} \left[-\frac{1}{2} d\langle (\theta, M_\tau), (\theta, M_\tau) \rangle + i(\theta, dA_\tau) \right] \\ & + \left\langle \sum_{\substack{s < \tau \leq t \\ \Delta X_\tau \neq 0}} e^{i(\theta, X_\tau - x)} [e^{i(\theta, \Delta X_\tau)} - 1 - i(\theta, \Delta N_\tau)] \right\rangle. \end{aligned}$$

Condition S2') and the uniqueness of Meyer's decomposition imply that, for each $\theta \in R^d$, the process

$$\begin{aligned} Z_t^\theta = & \left[-\frac{1}{2} \langle (\theta, M_t), (\theta, M_t) \rangle + i(\theta, A_t) + \left\langle \sum_{\substack{s < \tau \leq t \\ \Delta X_\tau \neq 0}} (e^{i(\theta, \Delta X_\tau)} - 1 - i(\theta, \Delta N_\tau)) \right\rangle \right] \\ & - \int_s^t \Phi_\theta(\tau, X_\tau) d\tau \end{aligned}$$

is identically zero. An elementary computation shows that

$$\begin{aligned} & \left\langle \sum_{\substack{s < \tau \leq t \\ \Delta X_\tau \neq 0}} \sin^4(\beta, \Delta X_\tau) \cdot e^{i(\alpha, \Delta X_\tau)} \right\rangle - \int_s^t \sin^4(\beta, u) \cdot e^{i(\alpha, u)} S(\tau, X_\tau, du) d\tau \\ & = (Z_t^{4\beta+\alpha} - 4Z_t^{2\beta+\alpha} + 6Z_t^\alpha - 4Z_t^{-2\beta+\alpha} + Z_t^{-4\beta+\alpha})/2^4 = 0. \end{aligned}$$

Thus, for each s -stopping time T' ($T' \leq T$),

$$\mathbb{E} \left[\int_s^{T'} \int \sin^4(\beta, u) \cdot e^{i(\alpha, u)} J(dt, du) \right] = \mathbb{E} \left[\int_s^{T'} \int \sin^4(\beta, u) \cdot e^{i(\alpha, u)} S(t, X_t, du) dt \right].$$

(If any one of the members is integrable, so is the other.) Set

$$S_n = \inf \{t \in [s, T]; \int_s^t |u|^2 \wedge 1 S(\tau, X_\tau, du) d\tau > n\}, \text{ and}$$

$H = \{\text{bounded measurable function } h(u) \text{ on } R^d;$

$$E \left[\int_s^{S_n} \sin^4(\beta, u) h(u) J(dt, du) \right] = E \left[\int_s^{S_n} \sin^4(\beta, u) h(u) S(t, X_t, du) dt \right].$$

We note that the class $C = \{\exp [i(\alpha, u)]; \alpha \in R^d\}$ is included in H . It is easy to show that H is closed under the formation of limits of uniform or bounded monotone sequences. Since C is closed under the multiplication and contains 1, H contains all the $\sigma(C)$ -measurable bounded functions, where $\sigma(C)$ is the σ -field generated by the functions of C . Thus H contains all the bounded measurable functions. As β and n are arbitrary, we have

$$E \left[\int_s^{T'} h(u) J(dt, du) \right] = E \left[\int_s^{T'} h(u) S(t, X_t, du) dt \right]$$

for each non-negative measurable function h on R^d . Property S3)-a) follows immediately from this fact. On the other hand it holds that

$$\begin{aligned} 0 = \operatorname{Re} Z_t^g &= -\frac{1}{2} \langle (\theta, M_t), (\theta, M_t) \rangle + \frac{1}{2} \int_s^t (\theta, a(\tau, X_\tau) \theta) d\tau \\ &+ \langle \int_s^t (\cos(\theta, u) - 1) J(d\tau, du) \rangle - \int_s^t \int (\cos(\theta, u) - 1) S(\tau, X_\tau, du) d\tau, \end{aligned}$$

that is to say,

$$\langle (\theta, M_t), (\theta, M_t) \rangle = \int_s^t (\theta, a(\tau, X_\tau) \theta) d\tau.$$

It is immediate to show that

$$A_t = \int_s^t b(\tau, X_\tau) d\tau, N_t = \int_s^t \int_{|u| \leq 1} u^c J(d\tau, du), B_t = \int_s^t \int_{|u| > 1} u J(d\tau, du).$$

3° By the Kunita-Watanabe formula, it is easily proved that Condition S3) implies Condition S1). Q.E.D.

DEFINITION. We say that a process $(X_t, P; t \in [s, T])$ is a solution of *the martingale problem of the (a, b, S) -stochastic equation* starting from (s, x) if X_t satisfies either one of the conditions of Theorem 2. 1.

A solution of the original stochastic equation is also a solution of the associated martingale problem. We can prove the converse of this fact under some restriction for a and S .

Theorem 2.2 *Let (Ω, F_t, P, X_t) be a martingale solution of the (a, b, S) -*

stochastic equation starting from (s, x) . If $a(s, x)$ is strictly positive, and the kernel $S(s, x, du)$ has no point mass and its support is $R^d - \{0\}$, then there exist σ, c , and $(B_t, p(dt, dz), q(dt, dz))$ such that

a) σ and c are Borel measurable, $\sigma\sigma^* = a$ and

$$\int_{c(s, x, z) \in D} \frac{dz}{|z|^{d+1}} = S(s, x, D) \text{ for each open set } D \subset R^d - \{0\}.$$

b) B_t (resp. p, q) is a Brownian motion (resp. a Poisson random measure, a Poisson martingale measure) on the probability space (Ω, F_t, P) . B_t, p and q satisfy the same condition as we stated at the beginning of this section.

c) X_t is a solution of the original (σ, b, c) -stochastic equation starting from (s, x) constructed over $(\Omega, F_t, P, B_t, p, q)$.

We omit the proof, since we never make use of the original stochastic equation.

3. Operator L and transition probabilities

I. (Some inequalities for solutions of a parabolic equation)

Let $a(s)$ be a measurable $d \times d$ symmetric matrix such that

$$a_1 |\theta|^2 \leq (\theta, a(s)\theta) \leq a_2 |\theta|^2, \quad 0 < a_1 \leq a_2 < \infty,$$

for all $\theta \in R^d$ and $s \in R_+$. We shall consider the parabolic equation:

$$\left(\lambda - \frac{\partial}{\partial s} - \frac{1}{2} \sum_{ij} a_{ij}(s) D_i D_j \right) g(s, x) = f(s, x), \quad \lambda > 0.$$

Define an operator G_λ acting on a suitable class of functions on $R_+ \times R^d$ by

$$G_\lambda f(s, x) = \int_0^\infty \int_{R^d} \frac{\det U(s, t)}{(2\pi t)^{d/2}} e^{-|U(s, t)y|^2/2t - \lambda t} f(s+t, x-y) dt dy$$

where $U(s, t) = \left(\frac{1}{t} \int_s^{s+t} a(\tau) d\tau \right)^{-1/2}$.

We say that a function f on $R_+ \times R^d$ belongs to the class C_K^∞ (or is a test function) if and only if f is a $C^\infty(R_+ \times R^d)$ -function with compact support. We can verify that if $f \in C_K^\infty$, then $g(s, x) = G_\lambda f(s, x)$ is a solution of the parabolic equation. Set

$$J_\lambda(s, t, x) = \frac{\det U(s, t)}{(2\pi t)^{d/2}} e^{-|U(s, t)x|^2/2t - \lambda t}.$$

Then we have, for each $f \in C_K^\infty$,

$$G_\lambda f(s, x) = \int_0^\infty \int J_\lambda(s, t, y) f(s+t, x-y) dt dy,$$

$$D_i G_\lambda f(s, x) = \int_0^\infty \int D_i J_\lambda(s, t, y) f(s+t, x-y) dt dy.$$

Moreover, for the set $B_\varepsilon = \{(t, y) \in R_+ \times R^d; t > \varepsilon \text{ or } |U(s, t)y|^2 > \varepsilon\}$, the integral:

$$\iint_{B_\varepsilon} D_i D_j J_\lambda(s, t, y) f(s+t, x-y) dt dy$$

exists for each $\varepsilon > 0$ and it converges to a certain function as $\varepsilon \downarrow 0$. Let us denote the limit by I_λ^{ij} . We can verify that

$$D_i D_j G_\lambda f(s, x) = I_\lambda^{ij} f(s, x) + c^{ij}(s) f(s, x)$$

where $c^{ij}(s)$ is a function on R_+ such that there exists a finite upper bound of $|c^{ij}|$ independent of λ and f . (see Bers-John-Schechter [1] p-226)

REMARK. We can verify that if $a(s)$ is continuous, then the functions $G_\lambda f$, $D_i G_\lambda f$, $D_i D_j G_\lambda f$ and $\frac{\partial}{\partial s} G_\lambda f$ are continuous in (s, x) for $f \in C_K^\infty$, and that these functions converge to 0 as $s + |x|^2 \rightarrow \infty$.

Let us introduce some norms for functions on $R_+ \times R^d$.

$$\|f\| = \sup_{s, x} |f(s, x)|$$

$$H^\alpha(f) = \sup_{s, s', x'} \frac{|f(s, x) - f(s, x')|}{|x - x'|^\alpha} \quad (0 < \alpha < 1)$$

$$\|f\|_{L^p} = \left(\int_0^\infty \int |f(s, x)|^p ds dx \right)^{1/p} \quad (1 < p < \infty)$$

In the sequel, these α and p are fixed. The following theorem can be proved by a similar way to the proof by Jones [2] and in Bers *et. al.* [1].

Theorem 3.1 *If f is a function of C_K^∞ and $\lambda r^2 \geq 1$, then*

- 1) $H^\alpha(I_\lambda^{ij} f) \leq c$, $H^\alpha(f)$ and $\|I_\lambda^{ij} f\| \leq c$, $r^\alpha H^\alpha(f)$,
- 2) $\|I_\lambda^{ij} f\|_{L^p} \leq c$, $\|f\|_{L^p}$,

where the c 's stand for certain constants independent of λ , f and r .

REMARK. Let $\|f\|_{C^\alpha}$ be the norm $\|f\| + r^\alpha H^\alpha(f)$. Then there exist constants c_α and c_p depending only on d, α, p, a_1 and a_2 such that

$$\sum_{ij} \|D_i D_j G_\lambda f\|_{C^\alpha} \leq c_\alpha \|f\|_{C^\alpha} \text{ and } \sum_{ij} \|D_i D_j G_\lambda f\|_{L^p} \leq c_p \|f\|_{L^p}$$

for $f \in C_K^\infty$ and $\lambda r^2 \geq 1$.

In the following two lemmas, we suppose that $\lambda r^2 \geq 1$ and $f \in C_K^\infty$, and

the c 's stand for certain constants independent of λ , f , and r . We will omit the proofs of these lemmas, because these are not so difficult.

Lemma 3.2

- 1) $\|G_\lambda f\| \leq r^2 \|f\|$ and $\|D_i G_\lambda f\| \leq c \cdot r \|f\|$,
- 2) $H^\alpha(G_\lambda f) \leq r^2 H^\alpha(f)$ and $H^\alpha(D_i G_\lambda f) \leq c \cdot r H^\alpha(f)$,
- 3) $|G_\lambda f|_{L^p} \leq r^2 |f|_{L^p}$ and $|D_i G_\lambda f|_{L^p} \leq c \cdot r |f|_{L^p}$.

Lemma 3.3

- 1) If $2-d/p > 0$, then $\|G_\lambda f\| \leq c \cdot r^{2-d/p} |f|_{L^p}$,
and if $1-d/p > 0$, then $\|D_i G_\lambda f\| \leq c \cdot r^{2-d/p} |f|_{L^p}$.
- 2) If $1+\alpha-d/p > 0$, then $H^{1-\alpha}(G_\lambda f) \leq c \cdot r^{1+\alpha-d/p} |f|_{L^p}$,
and if $\alpha-d/p > 0$, then $H^{1-\alpha}(D_i G_\lambda f) \leq c \cdot r^{\alpha-d/p} |f|_{L^p}$.
- 3) $|D_i G_\lambda f(\cdot, \cdot + u) - D_i G_\lambda f(\cdot, \cdot)|_{L^p} \leq c \cdot |u| |f|_{L^p}$.

II. (*A priori estimate for the operator L*)

Let $a(s, x)$, $b(s, x)$, $S(s, x, du)$ be as follows:

- a) there exist positive constants a_1 and a_2 such that

$$a_1 |\theta|^2 \leq (\theta, a(s, x)\theta) \leq a_2 |\theta|^2$$

for all $\theta \in R^d$ and $(s, x) \in R_+ \times R^d$,

- b) $\sup_{x, s} |b(s, x)| < \infty$,

- c) there exists a measure $\bar{S}(du)$ such that

$$\int |u|^2 \wedge 1 \bar{S}(du) < \infty \text{ and } S(s, x, du) \leq \bar{S}(du) \text{ for all } (s, x) \in R_+ \times R^d.$$

Let L be an operator defined by

$$\begin{aligned} Lv(s, x) = & \frac{1}{2} \sum_{ij} a_{ij}(s, x) D_i D_j v(s, x) + \sum_i b_i(s, x) D_i v(s, x) \\ & + \int [v(s, x+u) - v(s, x) - I_{(|u| \leq 1)}(u, \nabla v(s, x))] S(s, x, du). \end{aligned}$$

Let x_0 be an arbitrary but fixed point of R^d and set

$$L_0 = \frac{1}{2} \sum_{ij} a_{ij}(s, x_0) D_i D_j \text{ and } T_\lambda = (L - L_0) G_\lambda,$$

where G_λ is an operator defined similarly to that in Subsection I using $a(s, x_0)$ in place of $a(s)$. And denote by K_λ a formal expression $G_\lambda[I - T_\lambda]^{-1}$.

Assumption. Let c_a and c_p be the constants defined in Subsection I.

$$(A_a) \quad c_a \max_{ij} \sup_{s, x} |a_{ij}(s, x) - a_{ij}(s, x_0)| \leq 1.$$

$$(A_p) \quad c_p \max_{ij} \sup_{s, x} |a_{ij}(s, x) - a_{ij}(s, x_0)| \leq 1.$$

Lemma 3.4 *If Assumption (A_p) is satisfied, then there exists a function $\sigma_p(r)$ on R_+ such that $\lim_{r \downarrow 0} \sigma_p(r) = 0$ and*

$$\begin{aligned} & \frac{1}{2} \max_{ij} \sup_{x,s} |a_{ij}(s, x) - a_{ij}(s, x_0)| \sum_{ij} |D_i D_j G_\lambda f|_{L^p} + \max_i \|b_i\| \sum_i |D_i G_\lambda f|_{L^p} \\ & + \left| \int |G_\lambda f(s, x+u) - G_\lambda f(s, x) - I_{(|u| \leq 1)}(u, \nabla G_\lambda f(s, x))| \bar{S}(du) \right|_{L^p} \\ & \leq \left(\frac{1}{2} + \sigma_p(r) \right) |f|_{L^p}. \end{aligned}$$

In particular, $|T_\lambda f|_{L^p} \leq \left(\frac{1}{2} + \sigma_p(r) \right) |f|_{L^p}$ for $f \in C_K^\infty$ and $\lambda r^2 \geq 1$.

Proof. In this proof, the c 's stand for certain constants independent of λ , f and r . For simplicity, we set $a'(s, x) = a(s, x) - a(s, x_0)$ and $g(s, x) = G_\lambda f(s, x)$.

By Theorem 3.1 and Lemma 3.2, we have

$$\begin{aligned} & \frac{1}{2} \max_{ij} \|a'_{ij}\| \sum_{ij} |D_i D_j g|_{L^p} + \max_i \|b_i\| \sum_i |D_i g|_{L^p} \\ & \leq \frac{1}{2} \max_{ij} \|a'_{ij}\| c_p |f|_{L^p} + \max_i \|b_i\| c_r |f|_{L^p} \leq \left(\frac{1}{2} + c_r \right) |f|_{L^p}. \end{aligned}$$

On the other hand,

$$\begin{aligned} & \left| \int_{|u| > 1} |g(s, x+u) - g(s, x)| \bar{S}(du) \right|_{L^p}^p \\ & \leq \int_0^\infty \int ds dx \left(\int_{|u| > 1} \bar{S}(du) \right)^{p/q} \int_{|u| > 1} |g(s, x+u) - g(s, x)|^p \bar{S}(du) \quad \left(\frac{1}{p} + \frac{1}{q} = 1 \right) \\ & \leq c_2 \int_{|u| > 1} (2|g|_{L^p})^p \bar{S}(du) \leq c_3 |g|_{L^p}^p \leq c(r^2 |f|_{L^p})^p \quad (\text{by Lemma 3.2}). \end{aligned}$$

Moreover,

$$\begin{aligned} & \left| \int_{|u| \leq r} |g(s, x+u) - g(s, x) - (u, \nabla g(s, x))| \bar{S}(du) \right|_{L^p}^p \\ & \leq \int_0^\infty \int ds dx \int_0^1 d\theta \left(\int_{|u| \leq r} |\nabla g(s, x+\theta u) - \nabla g(s, x)| \bar{S}(du) \right)^p \\ & \leq \int_0^\infty \int ds dx \int_0^1 d\theta \left(\int_{|u| \leq r} |u|^2 \bar{S}(du) \right)^{p/q} \int_{|u| \leq r} \left(\frac{|\nabla g(s, x+\theta u) - \nabla g(s, x)|}{|u|} \right)^p \bar{S}(du) \\ & \leq \left(\int_{|u| \leq r} |u|^2 \bar{S}(du) \right)^{p/q} \int_0^1 d\theta \int_{|u| \leq r} \left(\frac{|\nabla g(s, x+\theta u) - \nabla g(s, x)|_{L^p}}{|u|} \right)^p \bar{S}(du) \\ & \leq c \left(\int_{|u| \leq r} |u|^2 \bar{S}(du) \right)^p |f|_{L^p}^p \quad (\text{by Lemma 3.3-3}). \end{aligned}$$

Similarly we have

$$\begin{aligned} & \left| \int_{r < |u| \leq 1} |g(s, x+u) - g(s, x) - (u, \nabla g(s, x))| \bar{S}(du) \right|_{L^p}^p \\ & \leq \int_0^\infty \int dx ds \int_0^1 d\theta \left(\int_{r < |u| \leq 1} |u| |\nabla g(s, x+\theta u) - \nabla g(s, x)| \bar{S}(du) \right)^p \end{aligned}$$

$$\begin{aligned} &\leq \left(\int_{r < |u| \leq 1} |u| \bar{S}(du) \right)^{p/q} \int_0^1 d\theta \left(\int_{r < |u| \leq 1} 2|u| |\nabla g|_{L^p} \bar{S}(du) \right) \\ &\leq c. \left(\int_{r < |u| \leq 1} |u| \bar{S}(du) \right)^p (r|f|_{L^p})^p \quad (\text{by Lemma 3.2}). \end{aligned}$$

Combining these inequalities, we see that

$$\begin{aligned} & \left| \int_{|u| \leq 1} g(s, x+u) - g(s, x) - (u, \nabla g(s, x)) \bar{S}(du) \right|_{L^p} \\ & \leq c. \int_{|u| \leq 1} |u|^2 \left(\frac{r}{|u|} \wedge 1 \right) \bar{S}(du) |f|_{L^p}. \end{aligned}$$

The right-hand side of this inequality tends to 0 as $r \downarrow 0$. Thus we can choose $\sigma_p(r)$ of the form

$$\sigma_p(r) = c. \left(r + \int_{|u| \leq 1} |u|^2 \left(\frac{r}{|u|} \wedge 1 \right) \bar{S}(du) \right). \quad \text{Q.E.D.}$$

The kernel S is said to be continuous if

$$\lim_{s' \rightarrow s, x' \rightarrow x} \int |u|^2 \wedge 1 |S(s', x', du) - S(s, x, du)| = 0 \quad \text{for all } (s, x) \in R_+ \times R^d.$$

Lemma 3.5. Suppose that Assumption (A_a) is satisfied. If $a(s, x)$, $b(s, x)$ and $S(s, x, du)$ are continuous in (s, x) , and if there exists a constant h_a such that

$$\begin{aligned} & \max_{ij} |a_{ij}(s, x') - a_{ij}(s, x)| + \max_i |b_i(s, x') - b_i(s, x)| \\ & + \int |u|^2 \wedge 1 |S(s, x', du) - S(s, x, du)| \leq h_a |x' - x|^a \quad \text{for any } s, x', x, \end{aligned}$$

then there exists a function $\sigma_a(r)$ on R_+ such that $\lim_{r \downarrow 0} \sigma_a(r) = 0$ and

$$|T_\lambda f|_{C^p} \leq \left(\frac{1}{2} + \sigma_a(r) \right) |f|_{C^p}$$

for $f \in C_K^\infty$ and $\lambda r^2 \geq 1$.

We can prove this lemma by means of Theorem 3.1 and Lemma 3.2. Although the computation is rather long, it is a routine work; so we omit the proof. We only mention that we can choose $\sigma_a(r)$ of the form

$$\sigma_a(r) = c. \left(r + r^a h_a + \int_{|u| \leq 1} |u|^2 \left(\frac{r}{|u|} \wedge 1 \right) \bar{S}(du) \right).$$

Let us introduce two norms for functions on $R_+ \times R^d$.

$$\begin{aligned} |f|_{L^{2,p}} &= |f|_{L^p} + \sum_i |D_i f|_{L^p} + \sum_{ij} |D_i D_j f|_{L^p}, \\ |f|_{C^{2+\alpha}} &= \|f\| + \sum_i \|D_i f\| + \sum_{ij} \|D_i D_j f\| + \sum_{ij} H^\alpha(D_i D_j f) \end{aligned}$$

Completing the space C_K^∞ by the norms $|\cdot|_{L^p}$, $|\cdot|_{L^{2,p}}$, $|\cdot|_{C_1^\alpha}$ and $|\cdot|_{C^{2+\alpha}}$, we get Banach spaces L^p , $L^{2,p}$, C_*^α and $C_*^{2+\alpha}$, respectively.

REMARKS. 1) Let Assumption (A_p) be satisfied and let r_p be a positive constant such that $\sigma_p(r_p) < \frac{1}{2}$. Then $|T_\lambda|_{L^p} < 1$ for $\lambda \geq r_p^{-2}$. Therefore the operators

$$[I - T_\lambda]^{-1}: L^p \rightarrow L^p \quad \text{and} \quad K_\lambda: L^p \rightarrow L^{2,p}$$

are well defined,

2) Suppose that the conditions in Lemma 3.5 are satisfied, and let r_α be a positive constant such that $\sigma_\alpha(r_\alpha) < \frac{1}{2}$. Then $|T_\lambda|_{C_*^\alpha} < 1$ for $\lambda \geq r_\alpha^{-2}$. Therefore, considering the first remark in Subsection I, the operators

$$[I - T_\lambda]^{-1}: C_*^\alpha \rightarrow C_*^\alpha \quad \text{and} \quad K_\lambda: C_*^\alpha \rightarrow C_*^{2+\alpha}$$

are well defined.

Theorem 3.6. 1) Suppose that Assumption (A_p) is satisfied. If $f \in L^p$ and $\lambda \geq r_p^{-2}$, then $v = K_\lambda f \in L^{2,p}$ and this is a solution (in the distribution sense) of the equation

$$\left(\lambda - \frac{\partial}{\partial s} - L \right) v = f.$$

Moreover if $p > d$ and $\frac{d}{p} < \alpha < 1$, then there exists a constant c (independent of f) such that

$$\|v\| + \sum_i \|D_i v\| + \sum_i H^{1-\alpha}(D_i v) \leq c \|f\|_{L^p}.$$

2) Suppose that conditions in Lemma 3.5 are satisfied. If f is a C_*^α -function such that the support of the function $\sup_x |f(s, x)|$ is a compact set in R_+ , then $v = K_\lambda f$ is a $C_*^{2+\alpha}$ -function satisfying the equation

$$\left(\lambda - \frac{\partial}{\partial s} - L \right) v = f$$

for any $\lambda > 0$, and $\|v\| \leq 1/\lambda \|f\|$.

Proof. 1) Set $f' = [I - T_\lambda]^{-1} f$. Then $f' \in L^p$ and $K_\lambda f = G_\lambda f'$. Therefore

$$\left(\lambda - \frac{\partial}{\partial s} - L \right) v = \left[\left(\lambda - \frac{\partial}{\partial s} - L_0 \right) - (L - L_0) \right] G_\lambda f' = [I - T_\lambda] f' = f.$$

The rest of the assertion follows from Lemma 3.3.

2) Set $\lambda' = r_\alpha^{-2}$. If f is a function concerned with, then the $C_*^{2+\alpha}$ -function $K_\lambda f$:

$$K_\lambda f(s, x) = e^{-(\lambda' - \lambda)s} K_{\lambda'} f_1(s, x) \quad \text{with} \quad f_1(s, x) = e^{(\lambda' - \lambda)s} f(s, x)$$

is well defined for each $\lambda > 0$. It is easy to show that $v = K_\lambda f$ satisfies the equation

$$\left(\lambda - \frac{\partial}{\partial s} - L\right)v = f.$$

By the maximum principle of parabolic-type equation, we see easily the last assertion. Q.E.D.

III. (Construction of transition probabilities)

Let $a, b, S, a_1, a_2, \bar{S}, c_a, c_p, r_p$ and other notations be the same as in Subsection II. In this subsection, we suppose that Assumption (A_a) and (A_p) are satisfied. Let us denote the Radon-Nikodym derivative of $S(s, x, du)$ given $\bar{S}(du)$ by $K(s, x, u)$.

It is easy to construct a sequence (a^n, b^n, S^n) satisfying a)–e).

- a) $a_1|\theta|^2 \leq (\theta, a^n(s, x)\theta) \leq a_2|\theta|^2$ for all θ and $(s, x) \in R_+ \times R^d$, and a^n satisfies Assumption (A_a) and (A_p) .
- b) $\|a_{ij}^n\| \leq \|a_{ij}\|$, $\|b_i^n\| \leq \|b_i\|$ and there exists K^n such that $0 \leq K^n(s, x, u) \leq 1$ and $S^n(s, x, du) = K^n(s, x, u)\bar{S}(du)$.
- c) There exists a bounded set $D^n \subset R_+ \times R^d$ such that $a^n(s, x) - a^n(s, x_0) = b^n(s, x) - b^n(s, x_0) = 0$ for $(s, x) \notin D^n$.
- d) a^n, b^n and S^n are continuous in (s, x) , and there exists a constant h^n such that

$$\begin{aligned} & \sum_{ij} |a_{ij}^n(s, x') - a_{ij}^n(s, x)| + \sum_i |b_i^n(s, x') - b_i^n(s, x)| \\ & + \int |u|^2 \wedge 1 |K^n(s, x', u) - K^n(s, x, u)| \bar{S}(du) \leq h^n |x' - x|^\alpha \end{aligned}$$

for all s, x', x .

- e) $a_{ij}^n(s, x) \rightarrow a_{ij}(s, x)$, $b_i^n(s, x) \rightarrow b_i(s, x)$ a.e. with respect to $dsdx$, and $K^n(s, x, u) \rightarrow K(s, x, u)$ a.e. with respect to $dsdx\bar{S}(du)$.

We define operators L^n and K_λ^n similarly to L and K_λ respectively by using (a^n, b^n, S^n) in place of (a, b, S) .

Lemma 3.7. *Let $f \in L^p \cap L^\infty$ and let (f^m) be a sequence of C_*^α -functions with compact supports such that $f^m \rightarrow f$ in L^p and $\|f^m\| \leq \|f\|$. Then $K_\lambda^n f^m$ is a $C_*^{2+\alpha} \cap L^{2,p}$ -function. And if $\lambda \geq r_p^{-2}$, then*

$$|K_\lambda^n f^m - K_\lambda^n f|_{L^{2,p}} \rightarrow 0 \text{ as } m \rightarrow \infty, \quad |K_\lambda^n f - K_\lambda f|_{L^{2,p}} \rightarrow 0 \text{ as } n \rightarrow \infty.$$

Moreover if $2p > d$ and $\lambda \geq r_p^{-2}$, then

$$\|K_\lambda^n f^m - K_\lambda^n f\| \rightarrow 0 \text{ as } m \rightarrow \infty, \quad \|K_\lambda^n f - K_\lambda f\| \rightarrow 0 \text{ as } n \rightarrow \infty.$$

Proof. By Theorem 3.6 we see that $K_\lambda^n f^m \in C_*^{2+\alpha} \cap L^{2,p}$. It is immediate to show that $\lim_{m \rightarrow \infty} |K_\lambda^n f^m - K_\lambda^n f|_{L^{2,p}} = 0$ for $\lambda \geq r_p^{-2}$. By Lemma 3.3, this implies

that $\lim_{n \rightarrow \infty} \|K_\lambda^n f^n - K_\lambda^n f\| = 0$ for $\lambda \geq r_p^{-2}$ and $2p > d$. On the other hand, for each $L^{2,p}$ -function v , $\lim_{n \rightarrow \infty} \|(L^n - L)v\|_{L^p} = 0$. In fact, for example,

$$\begin{aligned} I^n &= \iint ds dx \left[\int |v(s, x+u) - v(s, x) - I_{(|u| \leq 1)}(u, \nabla v(s, x))| |S^n(s, x, dv) - S(s, x, du)| \right]^p \\ &= \iint ds dx \left[\int |u|^2 \wedge 1 h(s, x, u) |K^n(s, x, u) - K(s, x, u)| \bar{S}(du) \right]^p \\ &\leq c \cdot \int |u|^2 \wedge 1 \left[\iint h(s, x, u)^p |K^n(s, x, u) - K(s, x, u)|^p ds dx \right] \bar{S}(du), \end{aligned}$$

where the c stands for a constant independent of n and

$$h(s, x, u) = |v(s, x+u) - v(s, x) - I_{(|u| \leq 1)}(u, \nabla v(s, x))| / (|u|^2 \wedge 1).$$

It is easy to show that $\sup_u |h(s, x, u)|_{L^p} < \infty$. As $|K^n - K| \leq 2$ and $K^n \rightarrow K$ a.e., we conclude that $\lim_{n \rightarrow \infty} I^n = 0$. In particular, $\lim_{n \rightarrow \infty} \|(L^n - L)K_\lambda\|_{L^p} = 0$ for $\lambda \geq r_p^{-2}$. Since $K_\lambda^n f = K_\lambda [I - (L^n - L)K_\lambda]^{-1} f$, we have $\lim_{n \rightarrow \infty} \|K_\lambda^n f - K_\lambda f\|_{L^{2,p}} = 0$ for $\lambda \geq r_p^{-2}$. By Lemma 3.3, we have also $\lim_{n \rightarrow \infty} \|K_\lambda^n f - K_\lambda f\| = 0$ for $\lambda \geq r_p^{-2}$ and $2p > d$.

Q.E.D.

Lemma 3.8. *Let v be $C_*^{2+\alpha}$ -function with compact support and let*

$$f = \left(\lambda - \frac{\partial}{\partial s} - L \right) v \quad (\lambda > 0).$$

Then f is an $L^p \cap L^\infty$ -function such that the support of $\sup_x |f(s, x)|$ is compact, and $v = K_\lambda f$.

Proof. Set

$$g^n = \left(\lambda - \frac{\partial}{\partial s} - L^n \right) v = f - (L^n - L)v.$$

Then g^n is an $L^p \cap C_*^\alpha$ -function such that the support of $\sup_x |g^n(s, x)|$ is compact. Thus

$$v = K_\lambda^n g^n = K_\lambda^n (f - (L^n - L)v).$$

From $\lim_{n \rightarrow \infty} \|(L^n - L)v\|_{L^p} = 0$, which have proved in the proof of Lemma 3.7, it follows that $\lim_{n \rightarrow \infty} \|K_\lambda^n (L^n - L)v\|_{L^{2,p}} = 0$. In fact, in the case $\lambda \geq r_p^{-2}$, this follows from Theorem 3.6; and in the case $\lambda < r_p^{-2}$, it follows from Theorem 3.6 and the next remark that if f' is an L^p -function such that the support of $\sup_x |f'(s, x)|$ is compact, then we have

$$K_\lambda^n f'(s, x) = e^{-(\lambda' - \lambda)s} K_{\lambda'}^n f'_1(s, x) \quad \text{with} \quad f'_1(s, x) = e^{(\lambda' - \lambda)s} f'(s, x),$$

where $\lambda' = r_p^{-2}$. On the other hand $\lim_{n \rightarrow \infty} \|K_\lambda^n f - K_\lambda f\|_{L^2, p} = 0$ by Lemma 3.7 (and the above remark). Thus $v = K_\lambda f$. Q.E.D.

Let C_0 be the class of all continuous function $f(s, x)$ on $R_+ \times R^d$ such that $\lim_{s+|x|^2 \rightarrow \infty} f(s, x) = 0$. In the rest of this section, we assume that $2p > d$.

By Lemma 3.7 and Theorem 3.6, we see that if $\lambda \geq r_p^{-2}$ and $f \in L^p \cap L^\infty$, then $K_\lambda f$ is an $L^{2,p} \cap C_0$ -function and

$$\|K_\lambda f\| = \lim_{n, m} \|K_\lambda^n f^m\| \leq \lim_m \frac{1}{\lambda} \|f^m\| \leq \frac{1}{\lambda} \|f\|,$$

where (f^m) is a sequence approximating f , having the same properties as in Lemma 3.7. Let K_λ' be a unique extension of K_λ onto C_0 . Then $\|K_\lambda'\| \leq 1/\lambda$. It is easy to show that

$$K_\lambda^n f^m - K_\mu^n f^m = -(\lambda - \mu) K_\lambda^n K_\mu^n f^m \quad (\lambda, \mu \geq 0), \text{ and } K_\lambda^n f^m \geq 0 \text{ if } f^m \geq 0.$$

This implies that, for each $\lambda, \mu \geq r_p^{-2}$,

$$K_\lambda' - K_\mu' = -(\lambda - \mu) K_\lambda' K_\mu' \text{ and } K_\lambda' \geq 0.$$

Set $K = K_\lambda (L^p \cap C_0)$ (which is independent of λ), and let $A = \frac{\partial}{\partial s} + L$. The family K is dense in C_0 . In fact, let v and f be the functions considered in Lemma 3.8 and let (f^m) be a sequence approximating f which has been considered in Lemma 3.7. Then we have $\lim_{m \rightarrow \infty} \|K_\lambda f^m - v\| = \lim_{m \rightarrow \infty} \|K_\lambda f^m - K_\lambda f\| = 0$ for $\lambda \geq r_p^{-2}$ (by Lemma 3.7 and Lemma 3.8). This implies that K is dense in C_0 . It is easy to show that

$$(\lambda - A)K_\lambda = I \text{ on } L^p \cap C_0, \text{ and } K_\lambda(\lambda - A) = I \text{ on } K.$$

Therefore

$$(\lambda - A)K_\lambda' = K_\lambda'(\lambda - A) = I \text{ on } K.$$

By Hille-Yosida's semi-group theory, there exists a closed extension A' of the operator A and a positive contraction semi-group (P_t) on C_0 whose infinitesimal generator is A' and whose resolvents are (K_λ') . That is,

$$K_\lambda' f = (\lambda - A')^{-1} f = \int_0^\infty e^{-\lambda t} P_t f dt, \quad f \in C_0 \text{ and } \lambda \geq r_p^{-2}.$$

Let $\rho(\xi)$ be a $C^\infty(R^1)$ -function such that $0 \leq \rho(\xi) \leq 1$, $\rho(\xi) = 1$ for $|\xi| \leq \frac{1}{2}$ and $\rho(\xi) = 0$ for $|\xi| \geq 1$.

Lemma 3.9. Suppose that the domain of the operator P_t is extended to $C^{0,b}(R_+ \times R^d)$ by such a way that $P_t f = \lim_{m \rightarrow \infty} P_t f^m$ for each $f \in C^{0,b}(R_+ \times R^d)$ where

$$f^m(s, x) = f(s, x) \rho\left(\frac{s}{m^2}\right) \rho\left(\frac{|x|}{m}\right).$$

Then 1) $P_t 1 = 1$, and 2) if f is a bounded continuous function on R_+ with compact support, then $P_t f(s, x) = f(s+t)$ ($P_t f$ is independent of x).

Proof. 1) Let $1^m = \rho(s/m^2) \rho(|x|/m)$. Then we have

$$\left(\lambda - \frac{\partial}{\partial s} - L\right) 1^m = \lambda 1^m + g^m, \quad g^m \in L^p \cap L^\infty \text{ and } \lim_{m \rightarrow \infty} \|g^m\| = 0.$$

By Lemma 3.8, $1^m = \lambda K_\lambda 1^m + K_\lambda g^m$. Since $\|K_\lambda g^m\| \leq \frac{1}{\lambda} \|g^m\|$, it holds that $\lim_{m \rightarrow \infty} K_\lambda 1^m = \frac{1}{\lambda}$. And so we have

$$\int_0^\infty e^{-\lambda t} P_t 1 dt = \int_0^\infty e^{-\lambda t} \lim_{m \rightarrow \infty} P_t 1^m dt = \lim_{m \rightarrow \infty} \int_0^\infty e^{-\lambda t} P_t 1^m dt = \lim_{m \rightarrow \infty} K_\lambda' 1^m = \frac{1}{\lambda},$$

for $\lambda \geq r_p^{-2}$. Since $P_t 1 \leq 1$, it must hold that $P_t 1 = 1$.

2) Let $h(s) = \int_s^\infty e^{-\lambda(t-s)} f(t) dt = \int_0^\infty e^{-\lambda t} f(s+t) dt$ and $h^m(s, x) = h(s) \rho(|x|/m)$. Then we have

$$\left(\lambda - \frac{\partial}{\partial s} - L\right) h^m = f(s) \rho\left(\frac{|x|}{m}\right) + g^m, \quad g^m \in L^p \cap L^\infty \text{ and } \lim_{m \rightarrow \infty} \|g^m\| = 0.$$

By a similar way to the item 1), we have

$$\int_0^\infty e^{-\lambda t} P_t f dt = \lim_{m \rightarrow \infty} K_\lambda' f^m = h(s) = \int_0^\infty e^{-\lambda t} f(s+t) dt,$$

for all $\lambda \geq r_p^{-2}$. This implies that $P_t f(s, x) = f(s+t)$.

Q.E.D.

There exists a kernel $p_t'(s, x; d\tau, dy)$ such that

$$P_t f(s, x) = \int p_t'(s, x; d\tau, dy) f(\tau, y)$$

for each bounded continuous function f on $R_+ \times R^d$. By Lemma 3.9, we see that

$$p_t'(s, x; R_+, dy) = p_t'(s, x; \{t+s\}, dy), \quad p_t'(s, x; R_+, R^d) = 1.$$

Set $p_t'(s, x; R_+, dy) = p(s, x; t+s, dy)$. Then $p(s, x; t, dy)$ is a kernel such that

- a) $p(s, x; t, R^d) = 1$,
- b) $p(s, x; s, dy) = \delta_x(dy)$ (δ -measure at x),
- c) $p(s, x; t, dy) = \int p(s, x; r, dz) p(r, z; t, dy)$ for $s < r < t$, and
- d) $P_{t-s} f(s, x) = \int p(s, x; t, dy) f(y)$ for each $f \in C^{0,b}(R^d)$.

4. Uniqueness of martingale solution

In this section, we suppose that the following condition is satisfied.

Condition (U)

- 1) For each bounded domain D , there exist constants $0 < a_1 \leq a_2$ such that

$$a_1 |\theta|^2 \leq (\theta, a(s, x)\theta) \leq a_2 |\theta|^2$$

for all $\theta \in R^d$ and $(s, x) \in D$, Moreover $\lim_{x' \rightarrow x} \sup_{s \leq T} \sum_{ij} |a_{ij}(s, x') - a_{ij}(s, x)| = 0$ for any T .

- 2) $b(s, x)$ is locally bounded.

- 3) For each bounded domain D , there exists a measure $S(du)$ such that

$$\int |u|^2 \wedge 1 \bar{S}(du) < \infty, \text{ and } S(s, x, du) \leq \bar{S}(du) \text{ for } (s, x) \in D.$$

Let T be a positive constant, and $\rho(\xi)$ the function defined in §3—III. Let y be an arbitrary but fixed point of R^d , and set

$${}^R a(s, x) = a(s \wedge T, y) + [a(s \wedge T, x) - a(s \wedge T, y)] \rho\left(\frac{|x-y|}{R}\right),$$

$${}^R b(s, x) = b(s \wedge T, x) \rho\left(\frac{|x-y|}{R}\right),$$

$${}^R S(s, x, du) = S(s \wedge T, x, du) \rho\left(\frac{|x-y|}{R}\right).$$

In §3—I, we have learnt that if $a_1 |\theta|^2 \leq (\theta, a(s)\theta) \leq a_2 |\theta|^2$ for all θ and s , then there exist constants $c_p(a_1, a_2)$ and $c_\alpha(a_1, a_2)$ such that

$$\sum_{ij} |D_i D_j G_\lambda|_{L^p} \leq c_p(a_1, a_2), \quad \sum_{ij} |D_i D_j G_\lambda|_{C_r^\alpha} \leq c_\alpha(a_1, a_2) \quad (\lambda r^2 \geq 1),$$

where G_λ is the Green operator associated with the parabolic operator

$$\lambda - \frac{\partial}{\partial s} - \frac{1}{2} \sum_{ij} a_{ij}(s) D_i D_j.$$

Let \underline{a} and \bar{a} be positive constants such that

$$\underline{a} |\theta|^2 \leq (\theta, {}^R a(s, x)\theta) \leq \bar{a} |\theta|^2,$$

for any $\theta \in R^d$, $R \leq 1$, $(s, x) \in R_+ \times R^d$. There exists a positive constant $R(y) \leq 1$ such that

$$c_\alpha(\underline{a}, \bar{a}) \vee c_p(\underline{a}, \bar{a}) \max_{ij} \sup_{s, s', x} |{}^{R(y)} a_{ij}(s, x') - {}^{R(y)} a_{ij}(s, x)| \leq 1.$$

It is possible to assume, without any loss of generality, that $R(y)$ is measurable and $1/R(y)$ is locally bounded.

As we have proved in §3—III, there exists a system of transition probabilities $p(s, x; t, dy)$ associated with the operator L with coefficients $({}^{R(\mathcal{Y})}a, {}^{R(\mathcal{Y})}b, {}^{R(\mathcal{Y})}S)$. Therefore it is possible to construct a strong Markov process $(x_t, Q_{s,x}; t \in [s, \infty))$ on the measurable space $(W^s, \mathcal{W}^s, \mathcal{W}_t^s)$ such that

$$E_{s,x}[f(x_t) | \mathcal{W}_r^s] = \int p(r, x_r; t, dy) f(y) \quad (s \leq r \leq t)$$

for each bounded Borel function f on R^d .

In the next lemma, $({}^{R(\mathcal{Y})}a, {}^{R(\mathcal{Y})}b, {}^{R(\mathcal{Y})}S)$ are simply denoted by (a, b, S) . And L and K_λ are the operators associated with these coefficients.

Lemma 4.1. *The strong Markov process $(x_t, Q_{s,x})$ is a martingale solution of the (a, b, S) -stochastic equation starting from (s, x) .*

Proof. Suppose that $2p > d$, $\lambda \geq r_p^{-2}$ and $f \in L^p \cap C_0$ (see §3.) Then we have

$$K_\lambda f(s, x) = \int_s^\infty e^{-\lambda(t-s)} E_{s,x}[f(t, x_t)] dt.$$

Set $\rho_N(x) = \rho(|x|/N)$ and

$$\mathbf{H} = \left\{ f \in L^\infty; K_\lambda(f\rho_N)(s, x) = E_{s,x} \left[\int_s^\infty e^{-\lambda(t-s)} (f\rho_N)(t, x_t) dt \right] \right\}.$$

Then \mathbf{H} contains 1 and all test functions. If $f_n \in \mathbf{H}$ and $f_n \rightarrow f$ in sup-norm, then $f_n \rho_N \rightarrow f \rho_N$ in L^p and in sup-norm, and hence $f \in \mathbf{H}$. Similarly, if $f_n \in \mathbf{H}$, $f_n \geq 0$ and $f_n \uparrow f (f \in L^\infty)$, then $f \in \mathbf{H}$. Therefore \mathbf{H} must contain all the bounded measurable functions. From this fact, it is easily verified that if $\lambda \geq r_p^{-2}$ and $f \in L^p \cap L^\infty$, then

$$K_\lambda f(s, x) = E_{s,x} \left[\int_s^\infty e^{-\lambda(t-s)} f(t, x_t) dt \right].$$

This fact and the Markov property of $(x_t, Q_{s,x})$ imply that the process

$$e^{-\lambda(t-s)} K_\lambda f(t, x_t) - K_\lambda f(s, x) + \int_s^t e^{-\lambda(\tau-s)} f(\tau, x_\tau) d\tau$$

is a square integrable martingale for each $\lambda \geq r_p^{-2}$ and $f \in L^p \cap L^\infty$. Let v be a C^2 -function with compact support and let

$$\left(\lambda - \frac{\partial}{\partial s} - L \right) v = f.$$

Then $f \in L^p \cap L^\infty$ and $v = K_\lambda f$ (by Lemma 3.8). Thus the process

$$e^{-\lambda(t-s)} v(t, x_t) - v(s, x) + \int_s^t e^{-\lambda(\tau-s)} f(\tau, x_\tau) d\tau$$

is a square integrable martingale. It is easy to show that this property holds for any $C^{2,b}(R_+ \times R^d)$ -function v . Moreover the property $\lim_{t \rightarrow \infty} Q_{s,x}[\sup_t |x_t| > l] = 0$

follows easily from the above facts. Consequently, the process $(x_t, Q_{s,x})$ is a martingale solution of the (a, b, S) -stochastic equation starting from (s, x) .

Q.E.D.

REMARK. In the above proof, we used the constant r_p which was not well identified. Here we shall give a discussion upon this. In this remark, let us denote $(^{R(y)}a, ^{R(y)}b, ^{R(y)}S)$ by (a, b, S) . And let G_λ^z be the Green operator associated with the parabolic operator

$$\lambda - \frac{\partial}{\partial s} - \frac{1}{2} \sum_{ij} a_{ij}(s, z) D_i D_j = \lambda - \frac{\partial}{\partial s} - L^z.$$

There exists a measure $\bar{S}(du)$ (depending on $R(y)$) such that

$$\int |u|^2 \wedge 1 \bar{S}(du) < \infty, \quad S(s, x, du) \leq \bar{S}(du) \text{ for each } (s, x) \in R_+ \times R^d.$$

The definition of $R(y)$ shows the existence of positive constants γ and r_p such that $0 < \gamma < 1$ and

$$\begin{aligned} & \frac{1}{2} \max_{ij} \|a_{ij}(s, x) - a_{ij}(s, z)\| \left\| \sum_{ij} D_i D_j G_\lambda^z f \right\|_{L^p} + \max_i \|b_i\| \left\| \sum_i D_i G_\lambda^z f \right\|_{L^p} \\ & + \left\| \int G_\lambda^z f(s, x+u) - G_\lambda^z f(s, x) - I_{(|u| \leq 1)}(u, \nabla G_\lambda^z f(s, x)) \bar{S}(du) \right\|_{L^p} \leq \gamma \|f\|_{L^p} \end{aligned}$$

for each $\lambda \geq r_p^{-2}$, $f \in L^p$ and $z \in R^d$.

In the following theorem, (a, b, S) means the coefficient $(^{R(y)}a, ^{R(y)}b, ^{R(y)}S)$, and $G_\lambda^z, \bar{S}(du)$, γ and r_p are the objects defined in the above remark.

Theorem 4.2. *The martingale solution of the (a, b, S) -stochastic equation starting from (s, x) exists uniquely, and this is a strong Markov process.*

Proof. 1° Let $(x_t, Q_{s,x})$ be any martingale solution of the (a, b, S) -stochastic equation starting from (s, x) . Let us define an operator V_λ acting on bounded measurable functions f on $R_+ \times R^d$ by the formula:

$$V_\lambda f(s, x) = E_{s,x} \left[\int_s^\infty e^{-\lambda(t-s)} f(t, x_t) dt \right].$$

We shall prove that if $p > d$, $\lambda \geq r_p^{-2}$ and $f \in L^p \cap L^\infty$, then $K_\lambda f(s, x) = V_\lambda f(s, x)$, where K_λ is the operator associated with the coefficient (a, b, S) . From this fact, the assertion of the theorem follows immediately.

2° There exists a Brownian motion B_t such that

$$\begin{aligned} x_t = x & + \int_s^t a(\tau, x_\tau)^{1/2} dB_\tau + \int_s^t b(\tau, x_\tau) d\tau \\ & + \int_s^t \int_{|u| \leq 1} u^c J(d\tau, du) + \int_s^t \int_{|u| > 1} u J(d\tau, du) \end{aligned}$$

where J is the measure of jumps of x_t and cJ is the martingale measure associated with J and S (see §2). Let π_n be a function satisfying $\pi_n(t) = s + \frac{\nu}{n}$ when $s + \frac{\nu}{n} < t \leq s + \frac{\nu+1}{n}$, and define a new process x_t^n by the formula:

$$\begin{aligned} x_t^n = x + \int_s^t a(\tau, x_{\pi_n(\tau)})^{1/2} dB_\tau + \int_s^t b(\tau, x_\tau) d\tau \\ + \int_s^t \int_{1/n < |u| \leq 1} u {}^cJ(d\tau, du) + \int_s^t \int_{|u| > 1} u J(d\tau, du). \end{aligned}$$

Let V_λ^n be an operator defined by

$$V_\lambda^n f(s, x) = E_{s,x} \left[\int_s^\infty e^{-\lambda(t-s)} f(t, x_t^n) dt \right].$$

It is not so difficult to prove that, for each $\varepsilon > 0$ and $T' < \infty$,

$$\lim_{\delta \downarrow 0} \sup_{\substack{0 \leq t, t' \leq T' \\ |t-t'| < \delta}} Q_{s,x} [|x_t - x_{t'}| > \varepsilon] = 0.$$

This property and the martingale inequality imply that

$$\lim_{n \rightarrow \infty} Q_{s,x} \left[\sup_{0 \leq t \leq T'} |x_t^n - x_t| > \varepsilon \right] = 0.$$

Thus we have $\lim_n V_\lambda^n f(s, x) = V_\lambda f(s, x)$ for each bounded continuous function f .

3° We shall prove that there exists a constant N_λ^n such that $\|V_\lambda^n f\| \leq N_\lambda^n \|f\|_{L^p}$ for each $f \in C_*^\alpha \cap L^p$, $p > d$ and $\lambda \geq r_p^{-2}$.

Set $v(z; s, x) = G_\lambda^z f(s, x)$ for $f \in C_*^\alpha \cap L^p$, then $v(z; \cdot, \cdot) \in C_*^{2+\alpha} \cap L^{2,p}$ for each $z \in R^d$. Let us denote $s + \frac{\nu}{n}$ by t_ν and $v(x_{t_\nu}; s, x)$ by $v_\nu(s, x)$. Then the process

$$\begin{aligned} e^{-\lambda(t-s)} v_\nu(t, x_t^n) - e^{-\lambda(t_\nu-s)} v_\nu(t_\nu, x_{t_\nu}^n) + \int_{t_\nu}^t e^{-\lambda(\tau-s)} f(\tau, x_\tau^n) d\tau \\ + \int_{t_\nu}^t e^{-\lambda(\tau-s)} \left\{ \sum_i b_i(\tau, x_\tau) D_i v_\nu(\tau, x_\tau^n) + \int_{|u| > 1/n} [v_\nu(\tau, x_\tau^n + u) - v_\nu(\tau, x_\tau^n) \right. \\ \left. - I_{(|u| \leq 1)}(u, \nabla v_\nu(\tau, x_\tau^n))] S(\tau, x_\tau, du) \right\} d\tau \end{aligned}$$

is a square integrable martingale on the time interval $(t_\nu, t_{\nu+1}]$. Therefore,

$$\begin{aligned} \|V_\lambda^n f\| \leq \sum_{\nu=0}^\infty (e^{-\lambda(t_{\nu+1}-s)} - e^{-\lambda(t_\nu-s)}) \sup_z \|v(z; \cdot, \cdot)\| \\ + \int_s^\infty e^{-\lambda(t-s)} \left\{ \sum_i \|b_i\| \sup_z \|D_i v(z; \cdot, \cdot)\| + 2 \sup_z \|v(z; \cdot, \cdot)\| \int_{|u| > 1} \bar{S}(du) \right. \\ \left. + 2 \sum_i \sup_z \|D_i v(z; \cdot, \cdot)\| n \int_{|u| \leq 1} |u|^2 \bar{S}(du) \right\} dt. \end{aligned}$$

Since, by Lemma 3.3, there exists a constant c independent of f and z such that

$$\sum_i \|D_i v(z; \cdot, \cdot)\| + \|v(z; \cdot, \cdot)\| \leq c \|f\|_{L^p},$$

there exists a constant N_λ^n such that $\|V_\lambda^n f\| \leq N_\lambda^n \|f\|_{L^p}$ for each $f \in C_*^\alpha \cap L^p$.

4° We shall prove that there exists a constant N_λ such that $\|V_\lambda f\| \leq N_\lambda \|f\|_{L^p}$ for each $f \in C_*^\alpha \cap L^p$ provided that $\lambda \geq r_p^{-2}$ ($p > d$).

Let $f \in C_*^\alpha \cap L^p$ and $v(s, x) = G_\lambda^z f(s, x)$ where z is an arbitrary but fixed point of R^d .

Set

$$\begin{aligned} h(s, x) = & \max_{ij} \|a_{ij}(s, x) - a_{ij}(s, z)\| \sum_{ij} |D_i D_j v(s, x)| + \max_i \|b_i\| \sum_i |D_i v(s, x)| \\ & + \int \|v(s, x+u) - v(s, x) - I_{(|u| \leq 1)}(u, \nabla v(s, x))\| \bar{S}(du). \end{aligned}$$

Since the process

$$\begin{aligned} & e^{-\lambda(t-s)} v(t, x_t^n) - v(s, x) + \int_s^t e^{-\lambda(\tau-s)} f(\tau, x_\tau^n) d\tau \\ & - \int_s^t e^{-\lambda(\tau-s)} \left\{ \frac{1}{2} \sum_{ij} (a_{ij}(\tau, x_{\pi_n(\tau)}) - a_{ij}(\tau, z)) D_i D_j v(\tau, x_\tau^n) + \sum_i b_i(\tau, x_\tau) D_i v(\tau, x_\tau^n) \right. \\ & \left. + \int_{1/n < |u|} [v(\tau, x_\tau^n + u) - v(\tau, x_\tau^n) - I_{(|u| \leq 1)}(u, \nabla v(\tau, x_\tau^n))] S(\tau, x_\tau, du) \right\} d\tau \end{aligned}$$

is a square integrable martingale, we have $\|V_\lambda^n f\| \leq \|v\| + \|V_\lambda^n h\|$. Let N_λ^n be the smallest constant such that $\|V_\lambda^n f\| \leq N_\lambda^n \|f\|_{L^p}$ for any $C_*^\alpha \cap L^p$ -function f , and let N_λ be a constant such that $\|G_\lambda^z f\| \leq (1-\gamma)N_\lambda \|f\|_{L^p}$ for any $L^p \cap L^\infty$ -function f . Since $\|h\|_{L^p} \leq \gamma \|f\|_{L^p}$, we have

$$\|V_\lambda^n f\| \leq (1-\gamma)N_\lambda \|f\|_{L^p} + N_\lambda^n \|h\|_{L^p} \leq (1-\gamma)N_\lambda \|f\|_{L^p} + \gamma N_\lambda^n \|f\|_{L^p}$$

This implies that $N_\lambda^n \leq (1-\gamma)N_\lambda + \gamma N_\lambda^n$, and so $N_\lambda^n \leq N_\lambda$. Therefore, for any $f \in C_*^\alpha \cap L^p$,

$$\|V_\lambda f\| = \lim_n \|V_\lambda^n f\| \leq N_\lambda \|f\|_{L^p}$$

5° The inequality $\|V_\lambda f\| \leq N_\lambda \|f\|_{L^p}$ holds good for any $L^p \cap L^\infty$ -function f . This may be proved by making use of general results in measure theory. Therefore we omit the proof.

6° Let $p > d$, $\lambda \geq r_p^{-2}$, $f \in C_*^\alpha \cap L^p$ and set $v = G_\lambda^z$ where z is an arbitrary fixed point. Then the process

$$e^{-\lambda(t-s)} v(t, x_t) - v(s, x) + \int_s^t e^{-\lambda(\tau-s)} \left(\lambda - \frac{\partial}{\partial \tau} - L \right) v(\tau, x_\tau) d\tau$$

is a square integrable martingale. Therefore

$$v(s, x) = E_{s,x} \left[\int_s^\infty e^{-\lambda(t-s)} \left(\lambda - \frac{\partial}{\partial t} - L \right) v(t, x_t) dt \right] = V_\lambda (I - T_\lambda^s) f(s, x),$$

where $T_\lambda^s = (L - L^s) G_\lambda^s$. On the other hand $G_\lambda^s = K_\lambda (I - T_\lambda^s)$. Hence $K_\lambda g = V_\lambda g$ for each function g of the form: $g = (I - T_\lambda^s) f, f \in C_*^\alpha \cap L^p$. The set of such functions g is dense in $L^p \cap L^\infty$ with L^p -norm, and so $K_\lambda f = V_\lambda f$ for each $f \in L^p \cap L^\infty$ for $p > d$ and $\lambda \geq r_p^{-2}$. Q.E.D

REMARK. Let ζ be a bounded s -stopping time and $p > d$, then

$$K_\lambda f(\zeta, x_\zeta) = E_{s,x} \left[\int_\zeta^\infty e^{-\lambda(t-\zeta)} f(t, x_t) dt \mid \mathcal{W}_\zeta^s \right] \text{ a.e. } (Q_{s,x})$$

for each $f \in L^p \cap L^\infty$ and $\lambda \geq r_p^{-2}$. (This can be proved similarly to the above theorem.)

In the following lemma, we do not assume that (a, b, S) satisfy Condition (U). (And (a, b, S) does not mean the coefficient $(^{R(y)}a, ^{R(y)}b, ^{R(y)}S)$.)

Lemma 4.3 *We assume that (a, b, S) satisfy the condition*

$$\max \|a_{i,j}\| + \max \|b_i\| + \left\| \int |u|^2 \wedge 1 S(s, x, du) \right\| < \infty.$$

Let T be a bounded s -stopping time. Let Q' be a probability measure on (W^s, \mathcal{W}^s) such that $(x_t, Q'; t \in [s, T])$ is a martingale solution of the (a, b, S) -stochastic equation starting from (s, x) . Suppose that Q_w'' ($w \in W^s$) is a probability measure on $(W^s, \mathcal{W}^{T(w)})$ a. e. w (Q') such that $(x_t, Q_w''; t \in [T(w), \infty))$ is a martingale solution of the (a, b, S) -stochastic equation starting from $(T(w), x_{T(w)})$, and $w \rightsquigarrow Q_w''(A)I_{(T(w) \leq t)}$ is \mathcal{W}_t^s -measurable for each $A \in \mathcal{W}^t$ ($t \geq s$). Then there exists a unique probability measure Q on (W^s, \mathcal{W}^s) such that $Q = Q'$ on \mathcal{W}_T^s and the regular conditional disribution of Q given \mathcal{W}_T^s equals Q_w'' on $\mathcal{W}^{T(w)}$. If Q is this probability, then the process $(x_t, Q; t \in [s, \infty))$ is a martingale solution of the (a, b, S) -stochastic equation starting from (s, x) .

Proof. The first assertion is the conclusion of Lemma 1.2. Extend Q_w'' onto \mathcal{W}^s so that $Q[A \mid \mathcal{W}_T^s] = Q_w''[A]$ for all $A \in \mathcal{W}^s$. Then, for each $A \in \mathcal{W}^s$ and $B \in \mathcal{W}_T^s$, we have

$$Q[A \cap B] = \int_B Q_w''[A] Q'(dw).$$

Let $\Phi_\theta(s, x)$ be Ito's differential associated with (a, b, S) (see the paragraph under the first definition in §2), and set

$$M_r^t = \exp[i(\theta, x_t - x_r) - \int_r^t \Phi_\theta(\tau, x_\tau) d\tau], \quad s \leq r \leq t.$$

From the assumption, $M_{T \wedge t}^s$ (resp. $M_t^{T \wedge t}$) is a square integrable martingale with respect to (W_t^s, Q') (resp. $(W_{T \vee t}^s, Q'_w)$). Let $A \in \mathcal{W}_r^s$ be of the form: $A = A_v^1 \cap A_v^2$, where

$$A_v^1 = \{x_{s_1} \in F_1, \dots, x_{s_v} \in F_v\}, A_v^2 = \{x_{s_{v+1}} \in F_{v+1}, \dots, x_{s_n} \in F_n\}, s \leq s_1 < \dots < s_n \leq r.$$

Then we have,

$$\begin{aligned} \int_{(r > T)} Q'_w[I_A M_t^s] Q' &= \sum_v \int_{(s_v \leq T < s_{v+1})} Q'_w[I_A M_t^s] Q' \\ &= \sum_v \int_{(s_v \leq T < s_{v+1}) \cap A_v^1} M_T^s Q'_w[M_t^T I_{A_v^2}] Q' \\ &= \sum_v \int_{(s_v \leq T < s_{v+1}) \cap A_v^1} M_T^s Q'_w[M_r^T I_{A_v^2}] Q' \\ &= \sum_v \int_{(s_v \leq T < s_{v+1})} Q'_w[I_A M_r^s] Q' = \int_{(T < r)} Q'_w[I_A M_r^s] Q'. \end{aligned}$$

on the other hand,

$$\begin{aligned} \int_{(r \leq T)} Q'_w[I_A M_t^s] Q' &= \int_{(r \leq T) \cap A} M_{T \wedge t}^s Q'_w[M_t^T I_A^s] Q' \\ &= \int_{(r \leq T) \cap A} M_{T \wedge t}^s Q' = \int_{(r \leq T) \cap A} M_t^s Q'. \end{aligned}$$

Combining these equalities, we have $E_Q[I_A M_t^s] = E_Q[I_A M_r^s]$. This equality holds good for each $A \in \mathcal{W}_r^s$, provided that $s \leq r \leq t$. This implies that M_t^s is a square integrable martingale (for each $\theta \in R^d$) with respect to (W_t^s, Q) . Thus $(x_t, Q; t \in [s, \infty))$ is a martingale solution of the (a, b, S) -stochastic equation. Q.E.D.

Theorem 4.4. *If (a, b, S) satisfy Condition (U), then a martingale solution $(x_t, Q_{s,x}; t \in [s, T])$ of the (a, b, S) -stochastic equation starting from (s, x) is uniquely determined for any $(s, x) \in R_+ \times R^d$. It is all the same when T is an s -stopping time with respect to the family (W_t^s) .*

Proof. For the simplicity let us suppose that T is a constant.

1° Set $T_1 = \inf \{t \in [s, T]; |x_t - x| > 1/2 R(x)\}$. Let $(x_t, Q'; t \in [s, T])$ be a martingale solution of the (a, b, S) -stochastic equation starting from (s, x) . Then $(x_t, Q'; t \in [s, T_1])$ is a martingale solution of the $(^{R(x)}a, ^{R(x)}b, ^{R(x)}S)$ -stochastic equation starting from (s, x) . By Lemma 4.1, there exists a martingale solution $(x_t, Q''_{s',x'}; t \in [s', \infty))$ of the $(^{R(x)}a, ^{R(x)}b, ^{R(x)}S)$ -stochastic equation starting from (s', x') . Since the probability measure $Q''_{s',x'}$ is Borel measurable in (s', x') , $Q''_w = Q''_{T_1(w), x_{T_1(w)}}$ satisfies the conditions of Lemma 4.3. And so, there exists a martingale solution $(x_t, Q; t \in [s, \infty))$ of the $(^{R(x)}a, ^{R(x)}b, ^{R(x)}S)$ -stochastic equation starting from (s, x) such that $Q = Q'$ on $\mathcal{W}_{T_1}^s$ and $Q[\cdot | \mathcal{W}_{T_1}^s] = Q''_w[\cdot]$ on \mathcal{W}^{T_1} . By Theorem 4.2, a martingale solution (x_t, Q) is uniquely determined. Thus the

restricted measure $Q' | \mathcal{W}_{T_1}^s$ is uniquely determined.

2° Let us define a non-decreasing sequence of s -stopping times by

$$T_{n+1} = \inf\{t \in (T_n, T]; |x_t - x_{T_n}| > 1/2 R(x_{T_n})\}.$$

Then $\lim_{n \rightarrow \infty} Q'[T_n < T] = 0$ holds, for the process $(x_t, Q', t \in [s, T])$ is right continuous, stochastically bounded and $1/R(y)$ is locally bounded. By applying the same method used in 1° for the $(R(x_{T_1})a, R(x_{T_1})b, R(x_{T_1})S)$ -stochastic equation, it can be shown that the restricted measure $Q' | \mathcal{W}_{T_2}^s$ is uniquely determined. Inductively, we conclude that $Q' | \mathcal{W}_{T_n}^s$ is uniquely determined for each n . Q.E.D.

5. Existence of martingale solutions

Let T be a positive constant. We shall introduce a new condition.

Condition (B)

- 1) There exists a constant K such that $S(s, x, \{|u| > 1\}) \leq K$ for all $(s, x) \in [0, T] \times R^d$
 - 2) There exists a $C^1(R_+)$ -function $k(\xi)$ satisfying the following conditions.
 - a) $k(0) > 0$, $k'(\xi) \geq 0$ and $k(\xi)$ is a concave function.
 - b) $\int_1^\infty \frac{d\xi}{\xi k(\xi)} = \infty$.
 - c) $|x, b(s, x)| + \text{trace } a(s, x) + \int_{|u| \leq 1} |u|^2 S(s, x, du) \leq k(|x|)(1 + |x|^2)$ for all $(s, x) \in [0, T] \times R^d$.
 - 3) For each bounded domain $D \subset R^d$, there exists a measure $\bar{S}(du)$ such that $\int |u|^2 \wedge 1 \bar{S}(du) < \infty$ and $S(s, x, du) \leq \bar{S}(du)$ for $(s, x) \in [0, T] \times D$.
- (The function $\log(e + \xi)$ is an example satisfying a) and b) of Condition (B)—2.)

Let $\rho(\xi)$ be the function defined in §3—III. Set

$${}^N a(s, x) = a(s \wedge T, 0) + (a(s \wedge T, x) - a(s \wedge T, 0))\rho(|x|/N),$$

$${}^N b(s, x) = b(s \wedge T, x)\rho(|x|/N) \text{ and } {}^N S(s, x, du) = S(s \wedge T, x, du)\rho(|x|/N).$$

Lemma 5.1. *Let (a, b, S) satisfy Condition (B), and let $(x_t, {}^N Q; t \in [s, T])$ be a martingale solution of the $({}^N a, {}^N b, {}^N S)$ -stochastic equation starting from (s, x) . Then*

- 1) $\lim_{t \rightarrow \infty} \sup_N {}^N Q[\sup_t |x_t| > l] = 0$, and
- 2) $\lim_{\delta \rightarrow 0} \sup_N \sup_{|t-t'| \leq \delta} {}^N Q[|x_t - x_{t'}| > \varepsilon] = 0$ for each $\varepsilon > 0$.

Proof. 1° Set $T_0 = s$, $T_{n+1} = \inf\{t \in (T_n, T]; |\Delta x_t| > 1\}$. Since

$${}^N Q[T_n < T] = {}^N Q\left[\int_s^T \int_{|u|>1} J(dt, du) > n\right] \leq \frac{(T-s)K}{n}$$

where J is the measure of jumps of x_t (see §2), we have $\lim_{n \rightarrow \infty} \sup_N {}^N Q[T_n < T] = 0$.

If the property $\lim_{l \rightarrow \infty} \sup_N {}^N Q[\sup_t |x_t \wedge T_n| > l] = 0$ implies the property $\lim_{l \rightarrow \infty} \sup_N {}^N Q[\sup_t |x_t \wedge T_{n+1}| > l] = 0$, then assertion 1) of the lemma holds.

2° There exists a non-negative function $h(\xi) \in C^2(R_+)$ such that $h'(+0) = h''(+0) = 0$ and

$$h(\xi) = \int_0^\xi \frac{\eta d\eta}{(1+\eta^2)k(\eta)} \quad \text{on } [1, \infty).$$

Let ${}^N L$ be an operator defined by

$$\begin{aligned} {}^N L v(s, x) &= \frac{1}{2} \sum_{ij} {}^N a_{ij}(s, x) D_i D_j v(s, x) + \sum_i {}^N b_i(s, x) D_i v(s, x) \\ &+ \int_{|u| \leq 1} \{v(s, x+u) - v(s, x) - (u, \nabla v(s, x))\} {}^N S(s, x, du). \end{aligned}$$

An elementary computation shows that there exist constants H and H_1 such that

- i) $|{}^N L h(|x|)| \leq H$ and
- ii) $h'(|x|)^2 \frac{(x, {}^N a(s, x)x)}{|x|^2} + \int_{|u| \leq 1} (h(|x+u|) - h(|x|))^2 {}^N S(s, x, du) \leq H_1$

for each $(s, x) \in [0, T] \times R^d$ and N .

3° Let us denote $(t \vee T_n) \wedge T_{n+1}$ by t_n . Let us introduce a new process y_t :

$$y_t = y_{T_n} - \int_{T_n}^{t_n} \int_{|u|>1} J(d\tau, du).$$

Then the process y_t has the Meyer decomposition (with respect to the measure ${}^N Q$)

$$y_t = x_{T_n} + \int_{T_n}^{t_n} dM_\tau^N + \int_{T_n}^{t_n} {}^N b(\tau, x) d\tau + \int_{T_n}^{t_n} \int_{|u| \leq 1} u^c J^N(d\tau, du),$$

where

$$\begin{aligned} M_t^{N,i} &\in \mathbf{M}_{loc}^c(W_t^s, {}^N Q), \quad M_t^{N,i} M_t^{N,j} - \int_s^t {}^N a_{ij}(\tau, x_\tau) d\tau \in \mathbf{M}_{loc}^c(W_t^s, {}^N Q) \text{ and} \\ {}^c J^N(dt, du) &= J(dt, du) - {}^N S(t, x_t, du) dt. \end{aligned}$$

From inequality i) and the Kunita-Watanabe formula, we have

$$h(|y_t|) \leq h(|x_{T_n}|) + \int_{T_n}^{t_n} h'(|x_\tau|) \frac{(x_\tau, dM_\tau^N)}{|x_\tau|} \\ + \int_{T_n}^{t_n} \int_{|u| \leq 1} (h(|x_t + u|) - h(|x_\tau|)) {}^c J^N(d\tau, du) + H(t_n - T_n).$$

Let us denote the right-hand side of the inequality by z_t , then the process z_t is a positive submartingale with respect to the measure ${}^N Q$. By the martingale inequality,

$${}^N Q[\sup_t |y_t| > l | \mathbf{W}_{T_n}^s] = {}^N Q[\sup_t h(|y_t|) \geq h(l) | \mathbf{W}_{T_n}^s] \\ \leq \frac{4}{h(l)^2} (2h(|x_{T_n}|)^2 + 3H^2 T^2 + 2H_1 T).$$

Since $h(\infty) = \infty$, the right-hand side of the above inequality tends to 0 as $l \rightarrow \infty$.

On the other hand, $\sup_t |x_{t_n}| \leq \sup_t |y_t| + |dx_{T_{n+1}}|$, and

$${}^N Q[|dx_{T_{n+1}}| > l | \mathbf{W}_{T_n}^s] = {}^N E\left[\int_{T_n}^{T_{n+1}} \int_{|u| > l} {}^N S(t, y_t, du) dt | \mathbf{W}_{T_n}^s\right] \\ \leq T \sup_{0 \leq s \leq T, |x| \leq \sup_t |y_t|} \int_{|u| > l} {}^N S(s, x, du).$$

By Condition (B)–3), it holds that

$$\lim_{l \rightarrow \infty} \sup_N \sup_{(s, x) \in [0, T] \times D} \int_{|u| > l} {}^N S(s, x, du) = 0 \text{ for each bounded domain } D \subset R^d.$$

Thus, if $\lim_{l \rightarrow \infty} \sup_N {}^N Q[|x_{T_n}| > l] = 0$, then $\lim_{l \rightarrow \infty} \sup_N {}^N Q[\sup_t |y_t| > l] = 0$; and if $\lim_{l \rightarrow \infty} \sup_N {}^N Q[\sup_t |y_t| > l] = 0$, then $\lim_{l \rightarrow \infty} \sup_N {}^N Q[\sup_t |x_{t_n}| > l] = 0$. Thus assertion 1) is verified.

4° Set $U_0 = s$ and $U_{n+1} = \inf \{t \in (U_n, T]; |x_t| > n\}$. Then $\lim_{n \rightarrow \infty} \sup_N {}^N Q[U_n < T] = 0$.

The process $(x_t, {}^N Q)$ has the Meyer decomposition:

$$x_t - x = \int_s^t dM_\tau^N + \int_s^t {}^N b(\tau, x_\tau) d\tau + \int_s^t \int_{|u| \leq 1} u {}^c J^N(d\tau, du) + \int_s^t \int_{|u| > 1} u J(d\tau, du).$$

Therefore, for each $\varepsilon > 0$ and $s \leq t < t' \leq T$,

$${}^N Q[|x_{t'} - x_t| > \varepsilon] \leq {}^N Q\left[\left|\int_{t \wedge U_n}^{t' \wedge U_n} \{dM_\tau^N + {}^N b(\tau, x_\tau) d\tau + \int_{|u| \leq 1} u {}^c J^N(d\tau, du)\}\right| > \varepsilon\right] \\ + {}^N Q\left[\int_t^{t'} \int_{|u| > 1} J(d\tau, du) \neq 0\right] + {}^N Q[U_n < T].$$

The second term of the right-hand side of this inequality tends to 0 uniformly in N as $|t-t'| \downarrow 0$ because of the following inequality

$${}^N Q \left[\int_t^{t'} \int_{|u|>1} u J(d\tau, du) \neq 0 \right] \leq {}^N E \left[\int_t^{t'} \int_{|u|>1} J(d\tau, du) \right] \leq K(t'-t).$$

By Condition (B)—2), the first term also tends to 0 uniformly in N as $|t-t'| \downarrow 0$. Consequently, the second assertion of this lemma holds good. Q.E.D.

Theorem 5.2. *If (a, b, S) satisfies Condition (U) and (B), then there exists (uniquely) a martingale solution $(x_t, Q_{s,x}; t \in [s, T])$ of the (a, b, S) -stochastic equation starting from (s, x) . And the solution is a strong Markov process.*

Proof. By Theorem 4.2 and Lemma 4.3 (see also the proof of Theorem 4.4), it is possible to construct (uniquely) a martingale solution $(x_t, {}^N Q_{s,x}; t \in [s, T])$ of the $({}^N a, {}^N b, {}^N S)$ -stochastic equation starting from (s, x) . Let $T_0 = s$, $T_{N+1} = \inf\{t \in (T_N, T]; |x_t| > N/2\}$. By the uniqueness of the martingale solution of the $({}^N a, {}^N b, {}^N S)$ -stochastic equation, we have ${}^N Q_{s,x} = {}^{N+1} Q_{s,x}$ on the σ -field $\mathcal{W}_{T_N}^s$. Thus there exists a probability measure $Q_{s,x}$ on the σ -field $\mathcal{W}_{\bigvee T_N}^s$ such that $Q_{s,x} = {}^N Q_{s,x}$ on the σ -field $\mathcal{W}_{T_N}^s$. The process $(x_t, Q_{s,x}; t \in [s, \bigvee T_N])$ is a martingale solution of the (a, b, S) -stochastic equation starting from (s, x) . Since, by Lemma 5.1,

$$\lim_{N \rightarrow \infty} Q_{s,x}[T_N < T] = \lim_{N \rightarrow \infty} \sup_{M \geq N} {}^M Q_{s,x}[T_N < T] \leq \lim_{N \rightarrow \infty} \sup_M {}^M Q_{s,x}[\sup_t |x_t| > N/2] = 0,$$

the process $(x_t, Q_{s,x})$ is a martingale solution of the (a, b, S) -stochastic equation on the time interval $[s, T]$. It is easy to show that this process is a strong Markov process. Q.E.D.

Condition (C)

- 1) $\lim_{x' \rightarrow x} \sup_{0 \leq s \leq T} \sum_{ij} |a_{ij}(s, x') - a_{ij}(s, x)| = 0$ for each $x \in R^d$.
- 2) $b(s, x)$ is locally bounded, and

$$\lim_{x' \rightarrow x} \left\{ \sum_i |b_i(s, x') - b_i(s, x)| + \int |u|^2 \wedge 1 |S(s, x', du) - S(s, x, du)| \right\} = 0$$

for all $(s, x) \in [0, T] \times R^d$.

Lemma 5.3. *If Condition (B) and (C) are satisfied, then there exists a martingale solution $(x_t, Q_{s,x}; t \in [s, T])$ of the $({}^N a, {}^N b, {}^N S)$ -stochastic equation starting from (s, x) .*

Proof. We shall omit the super-prefix N of ${}^N a$, ${}^N b$, and ${}^N S$ in this proof.

1° If $a^{(m)} = a + 1/m \cdot I$, then there exists a martingale solution $(x_t, Q^{(m)}_{s,x})$ of the $(a^{(m)}, b, S)$ -stochastic equation starting from (s, x) , by Theorem 5.2. Similarly to the proof of Lemma 5.1, we can prove that

- 1) $\lim_{l \rightarrow \infty} \sup_m Q^{(m)} [\sup_t |x_t| > l] = 0$, and
 2) $\lim_{\delta \downarrow 0} \sup_m \sup_{|t'-t| \leq \delta} Q^{(m)} [|x_{t'} - x_t| > \varepsilon] = 0$ for any $\varepsilon > 0$.

By Lemma 1.1, it is possible to extract a subsequence $Q^n = Q^{(m_n)}$ from the sequence $Q^{(m)}$ and it is possible to construct a sequence $(X_t, X_t^n; n=1, 2, \dots)$ of processes on a certain probability space $(\Omega, \mathbf{F}_t, P)$ such that the processes (x_t, Q^n) and (X_t^n, P) are equivalent for each n , and the random sequence X_t^n converges in probability to X_t for each $t \in [s, T]$. Let $(x_t, Q_{s,x})$ be the process on the base space $(W^s, \mathbf{W}^s, \mathbf{W}_t^s)$ equivalent to the process (X_t, P) .

2° Let $\Phi_\theta(s, x)$ be Ito's differential associated with (a, b, S) (defined in the paragraph under the first definition of §2). Then

$$E_P[\{\exp[i(\theta, X_t^n - X_r^n)] - \int_r^t \Phi_\theta(\tau, X_\tau^n) d\tau - \frac{1}{m_n} |\theta|^2 (t-r) - 1\} f_1(X_{s_1}^n) \cdots f_k(X_{s_k}^n)] = 0.$$

for any $s \leq s_1 \leq \dots \leq s_k \leq r < t$, and for any $f_1, \dots, f_k \in C^{0,b}(R^d)$. Therefore

$$\begin{aligned} & E_P [\{\exp[i(\theta, X_t - X_r)] - \int_r^t \Phi_\theta(\tau, X_\tau) d\tau - 1\} f_1(X_{s_1}) \cdots f_k(X_{s_k})] \\ & \leq c \lim_{n \rightarrow \infty} E_P \left[\int_r^t |\Phi_\theta(\tau, X_\tau^n) - \Phi_\theta(\tau, X_\tau)| d\tau \right] \end{aligned}$$

where c is a constant depending only on $\|f_j\|$ ($j=1, \dots, k$), $t-r$ and $\|\Phi_\theta\|$.

By Condition (C), $\lim_{n \rightarrow \infty} E_P[|\Phi_\theta(\tau, X_\tau^n) - \Phi_\theta(\tau, X_\tau)|] = 0$ for each $\tau \in [r, t]$. Thus,

$$\lim_{n \rightarrow \infty} \int_r^t E_P[|\Phi_\theta(\tau, X_\tau^n) - \Phi_\theta(\tau, X_\tau)|] d\tau = 0.$$

Consequently, for any $s \leq r < t$ and $\theta \in R^d$,

$$E_P [\exp(i(\theta, X_t - X_r) - \int_r^t \Phi_\theta(\tau, X_\tau) d\tau) | \mathbf{F}_r] = 1.$$

This implies that the process (X_t, P) or $(x_t, Q_{s,x})$ is a martingale solution of the (a, b, S) -stochastic equation starting from (s, x) . Q.E.D.

Theorem 5.4 *If Condition (B) and (C) are satisfied, then there exists a martingale solution $(x_t, Q_{s,x}; t \in [s, T])$ of the (a, b, S) -stochastic equation starting from (s, x) .*

Proof. By Lemma 5.3, we can construct a martingale solution $(x_t, {}^N Q_{s',x'}^s; t \in [s', T])$ of the $({}^N a, {}^N b, {}^N S)$ -stochastic equation starting from (s', x') for each $(s', x') \in [s, T] \times R^d$. Set $T_0 = s$ and $T_{N+1} = \inf \{t \in (T_N, T]; |x_t| > N/2\}$. By Lemma 4.3, there exists a probability measure ${}^2 Q'_{s,x}$ on the σ -field $\mathbf{W}_{T_2}^s$ such that ${}^2 Q'_{s,x} = {}^1 Q_{s,x}$ on the σ -field $\mathbf{W}_{T_1}^s$ and ${}^2 Q'_{s,x}[\cdot | \mathbf{W}_{T_1}^s] = {}^2 Q_{T_1, x_{T_1}}[\cdot]$ on the

σ -field $\mathcal{W}_{T_2}^{T_1}$. Inductively, we can construct a sequence ${}^N Q'_{s,x}$ of probability measures on the σ -fields $\mathcal{W}_{T_N}^s$ such that ${}^{N+1} Q'_{s,x} = {}^N Q'_{s,x}$ on the σ -field $\mathcal{W}_{T_N}^s$ and ${}^{N+1} Q'_{s,x}[\cdot | \mathcal{W}_{T_N}^s] = {}^{N+1} Q_{T_N, x T_N}[\cdot]$ on the σ -field $\mathcal{W}_{T_{N+1}}^{T_N}$. Let $Q_{s,x}$ be the probability measure on the σ -field $\mathcal{W}_{T_N}^s$ such that $Q_{s,x} = {}^N Q'_{s,x}$ on the σ -field $\mathcal{W}_{T_N}^s$. The method used in the proof of Theorem 5.2. yields us the fact that the process $(x_t, Q)_{s,x}$ is a martingale solution of the (a, b, S) -stochastic equation starting from (s, x) Q.E.D.

OSAKA CITY UNIVERSITY

References

- [1] L. Bers, F. Jones and M. Schechter: Partial Differential Equations, Lectures in Appl. Math. III, Wiley, 1964.
- [2] F. Jones: *A class of singular integrals*, Amer. J. Math. **86** (1964), 441–462.
- [3] V. Krylov: *On Ito's stochastic integral equations*, Theor. Probability Appl. **14** (1969), 330–336.
- [4] H. Kunita and S. Watanabe: *On square integrable martingales*, Nagoya Math. J. **30** (1967), 209–245.
- [5] P. A. Meyer: *Intégrales stochastiques I, II*, Séminaire de probabilités, Univ. Strasbourg, (1966–'67).
- [6] K. R. Parthasarathy: Probability Measures on Metric Spaces, Academic Press, 1967.
- [7] V. Skorokhod: Studies in the Theory of Random Processes (translation), Addison-Wesley, 1965.
- [8] W. Strook and R. Varadhan: *Diffusion processes with continuous coefficients I, II*, Comm. Pure Appl. Math. **22** (1969), 345–400, 479–530.
- [9] H. Tanaka: *Existence of diffusions with continuous coefficients*, Mem. Fac. Sci. Kyushu Univ. Ser. A, **18** (1964), 89–103.
- [10] M. Tsuchiya: *Stochastic integral equation of jump type*, (in Japanese), R.I.M.S. Kokyuroku, 138, 1973.

