



Title	Sequential C-F Bond Transformation of the Difluoromethylene Unit in Perfluoroalkyl Groups: A Combination of Fine-Tuned Phenothiazine Photoredox Catalyst and Lewis Acid
Author(s)	Sugihara, Naoki; Nishimoto, Yoshihiro; Osakada, Yasuko et al.
Citation	Angewandte Chemie – International Edition. 2024, 63(14), p. e202401117
Version Type	VoR
URL	https://hdl.handle.net/11094/95279
rights	This article is licensed under a Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International License.
Note	

The University of Osaka Institutional Knowledge Archive : OUKA

<https://ir.library.osaka-u.ac.jp/>

The University of Osaka

Photoredox Catalysis

Sequential C–F Bond Transformation of the Difluoromethylene Unit in Perfluoroalkyl Groups: A Combination of Fine-Tuned Phenothiazine Photoredox Catalyst and Lewis Acid

Naoki Sugihara, Yoshihiro Nishimoto,* Yasuko Osakada, Mamoru Fujitsuka, Manabu Abe, and Makoto Yasuda*

Abstract: A sequential process via photoredox catalysis and Lewis acid mediation for C–F bond transformation of the CF₂ unit in perfluoroalkyl groups has been achieved to transform perfluoroalkylarenes into complex fluoroalkylated compounds. A phenothiazine-based photocatalyst promotes the defluoroaminoxylation of perfluoroalkylarenes with (2,2,6,6-tetramethylpiperidin-1-yl)oxyl (TEMPO) under visible light irradiation, affording the corresponding aminoxylated products. These products undergo a further defluorinative transformation with various organosilicon reagents mediated by AlCl₃ to provide highly functionalized perfluoroalkyl alcohols. Our novel phenothiazine catalyst works efficiently in the defluoroaminoxylation. Transient absorption spectroscopy revealed that the catalyst regeneration step is crucial for the photocatalytic aminoxylation.

The carbon–fluorine (C–F) bond transformation in perfluoroalkyl compounds not only is an important synthetic method in organic chemistry,^[1] but also is an urgent issue to solve PFAS (polyfluoroalkyl substances) environmental problems.^[2] Numerous C–F bond activation protocols have been reported for single C–F bond transformations of perfluoroalkyl group.^[3,4] However, a sequential C–F bond transformation of a difluoromethylene unit (–CF₂–) into two different functional groups remains underdeveloped (Figure 1A) despite being an important clue to the solution of PFAS problems. This is because the harsh reaction con-

ditions needed to cleave robust C–F bonds cause the undesired installation of the same functional group. In fact, dialkoxylation,^[5] dimethylation^[6] and dichlorination^[6] of a CF₂ moiety have been reported. To avoid installing the same groups, amino alcohols were used in the aminoalkoxylation of α -perfluoroalkyl ketones in a three-component tandem reaction (Figure 1B, a).^[7] Recently, two distinguished reactions were reported: a sequential defluorinative alkylation of trifluoroacetyl compounds by a radical mechanism (Figure 1B, b)^[8] and a coupling reaction of 1,1-difluoroalkyl compounds (RCF₂R') with Grignard reagents and chlorosilanes or alkyl tosylates by CrCl₂ catalysis via chromium carbenoid species (Figure 1B, c).^[9] Neither method is applicable to the transformation of longer perfluoroalkyl compounds (RCF₂(CF₂)_nR'). Herein we propose a reaction design based on a sequential process via radical and ionic paths (Figure 1C). The primary substitution of F with RO groups involves C–F bond activation by photocatalysis^[4g] and capture of the perfluoroalkyl radical by an oxyl radical.^[10] Then the second transformation employs a Lewis acid and nucleophiles. Because the reaction mechanism includes an oxonium intermediate, diverse nucleophiles can be introduced. Based on our proposed design using a dual activation system, in this study we achieved a sequential C–F bond transformation of perfluoroalkylarenes via aminoxylation with a fine-tuned phenothiazine photocatalyst and aminoxyl radical reagent followed by AlCl₃-mediated nucleophilic substitution with organosilicon reagents (Figure 1D).

[*] N. Sugihara, Dr. Y. Nishimoto, Prof. Dr. M. Yasuda
Department of Applied Chemistry, Graduate School of Engineering
Osaka University
2-1 Yamadaoka, Suita, Osaka 565-0871, Japan
E-mail: nishimoto@chem.eng.osaka-u.ac.jp
yasuda@chem.eng.osaka-u.ac.jp


Dr. Y. Nishimoto, Dr. Y. Osakada, Prof. Dr. M. Fujitsuka,
Prof. Dr. M. Yasuda
Innovative Catalysis Science Division, Institute for Open and
Transdisciplinary Research Initiatives (ICS-OTRI)
Osaka University
2-1 Yamadaoka, Suita, Osaka 565-0871, Japan
Dr. Y. Osakada, Prof. Dr. M. Fujitsuka
SANKEN (The Institute of Scientific and Industrial Research)
Osaka University
Mihogaoka 8-1, Ibaraki, Osaka 567-0047, Japan

Dr. Y. Osakada
Institute for Advanced Co-Creation Studies
Osaka University
Yamadagaoka 1-1, Suita, Osaka 565-0871, Japan




Prof. Dr. M. Abe
Department of Chemistry, Graduate School of Advanced Science
and Engineering
Hiroshima University
Higashi-Hiroshima, Hiroshima 739-8526, Japan

© 2024 The Authors. Angewandte Chemie International Edition published by Wiley-VCH GmbH. This is an open access article under the terms of the Creative Commons Attribution Non-Commercial NoDerivs License, which permits use and distribution in any medium, provided the original work is properly cited, the use is non-commercial and no modifications or adaptations are made.


A. Sequential C–F bond transformation of perfluoroalkanes to different functional groups



B. Reported works

a. Aminoalkoxylation with aminoalcohols (Loh and Shen)⁷b. Sequential defluoroalkylation by a radical mechanism (Houk and Wang)⁸c. Three-component cross coupling (Chen and Zeng)⁹

C. Reaction design



D. This work




Figure 1. Sequential C–F bond transformation of perfluoroalkyl compounds.

Firstly, we investigated reaction conditions for the photo-catalyzed aminoxylation using 4-phenyl(perfluorobutyl)benzene **1a** ($E_{\text{red}} = -2.06$ V vs SCE) and (2,2,6,6-tetramethylpiperidin-1-yl)oxyl (TEMPO) (**2**) under visible light irradiation (370 nm) (Table 1). The use of Ir(ppy)₃ resulted in no reaction due to the low reducing ability of excited Ir(ppy)₃ (Entry 1).^[4e] We focused on phenothiazines exhibiting higher reducing abilities than Ir(ppy)₃. Their photocatalytic activities can be tuned by a substituent effect.^[11] *N*-Phenylphenothiazine **PC1** exhibited a catalytic activity to mediate the aminoxylation, giving product **3a** (Entry 2). To access more negative redox potentials, phenothiazine **PC2** was used, leading to an improved yield (Entry 3). It should be noted that newly developed phenothiazine **PC3** bearing diisopropylamino and two methyl groups showed the best catalytic activity (Entry 4), suggesting the effect of the two Me groups is crucial to the aminoxylation. *N*-Dimethylphenylphenothiazine **PC4** was less effective, which indicates the increase of reducing ability by an amino group is significant (Entry 5). Next, to utilize much lower reduction


Table 1: Optimization of photo-catalyzed aminoxylation of perfluoroalkylarene **1a** with TEMPO **2**.^[4]

Entry	Catalyst	E_{ox}^* (V vs SCE)	Yield
1	Ir(ppy) ₃	-1.73	0%
2	PC1	-2.45	35%
3	PC2	-2.68	42%
4	PC3	-2.57	60%
5	PC4	-2.47	38%
6 ^[b]	4-MeOC ₆ H ₄ SH	-3.31	0%
7 ^[c]	Mes-Acr-BF ₄	-3.36	trace
8 ^[d]	PC3	-2.57	0%
9	none	–	0%
10 ^[e]	PC3	-2.57	82% (71%) ^[f]

[a] **1a** (0.4 mmol), **2** (0.8 mmol), catalyst (0.004 mmol), MeCN (2 mL), irradiation with 370 nm LEDs at 35 °C for 4 h. Yields were determined by ¹H NMR spectroscopy using an internal standard. [b] 4-MeOC₆H₄SH (0.08 mmol), HCO₂Cs (0.8 mmol), DMSO (2 mL) irradiation with 427 nm LEDs at 35 °C for 24 h. [c] Mes-Acr-BF₄ (0.04 mmol), NⁱPr₂Et (1.2 mmol), MeCN (1.3 mL), irradiation with 390 nm LEDs at 35 °C for 24 h. [d] No irradiation. [e] **PC3** (0.02 mmol), 24 h. [f] Isolated yield.

potential of photocatalysts, we applied thiolate catalysis^[12] and consecutive photo-induced electron transfer (conPET) system^[13] to this reaction. Thiolate catalysis resulted in no reaction, and **1a** was hardly converted (Entry 6). In the conPET system, **1a** was completely consumed, but only a trace amount of **3a** was obtained according to complicated products (Entry 7). In this case, the high reducing ability of the active catalytic species generated by conPET would cause the undesired overreduction or side-reactions. Both photocatalyst and photo-irradiation were essential for the reaction progress (Entries 8 and 9). Finally, under the optimized conditions using the 5 mol % amount of **PC3**, **3a** was obtained in 82 % yield (Entry 10). Further optimization for addition amount of TEMPO **2** and solvent screening is described in the Supporting Information.^[14]

Scheme 1 depicts a plausible mechanism for the aminoxylation of **1** with TEMPO **2** catalyzed by phenothiazine **PC**. The photoexcited **PC*** reduces **1** via single electron transfer (SET), affording radical anion **A** and radical cation **PC**^{•+}. Mesolysis of a C–F bond affords benzyl radical **B**.^[4g] Then, **B** associates with **2** to give product **3**. **PC** ($E(\text{PC}^{3+}/\text{PC}3) =$



Scheme 1. Proposed mechanism for photo-catalyzed aminoxylation of perfluoroalkylarenes.

0.61 V vs SCE) is regenerated by single electron reduction with **2** ($E(2^+/2) = 0.62$ V vs SCE).^[15] A by-product, *N*-oxoammonium cation **C** captures F^- , suppressing the retro-reaction from **B** to **A**.^[16] HRMS and ^{19}F NMR confirmed *N*-oxoammonium fluoride **D** was generated.^[17] The appropriate reduction potential of **PC3*** achieves selective reduction of starting material **1** and not product **3**, realizing a single C–F bond transformation without overreduction and side-reactions.^[18]

We focused on the fact that our developed **PC3** exhibited a more efficient catalytic activity than **PC2** despite the lower reducing ability of **PC3*** than **PC2*** (Table 1).^[19] We conducted mechanistic studies to understand the origin of the high activity of **PC3**. First, the fluorescence quenching experiments of **PC*** with 4-trifluoromethyl(perfluoroethyl)benzene **1e** were performed. Stern–Volmer plots determined that the rate constants of the dynamic quenching of excited singlet species **PC2*** and **PC3*** were $5.3 \times 10^{10} \text{ M}^{-1} \text{ s}^{-1}$ and $3.1 \times 10^{10} \text{ M}^{-1} \text{ s}^{-1}$, respectively (Figures S2 and S4).^[20,21] The 1st SET between **1e** and **PC*** is a diffusion-controlled process,^[22] which indicates that the catalytic turnover is independent of the reducing ability of **PC***.^[23] We then considered the 2nd SET between **PC**** and TEMPO for the catalyst regeneration. The sub-microsecond transient absorption spectroscopy using laser flash photolysis method at 355 nm (4 mJ/pulse, 4 ns pulse-width) was conducted for a mixture of PC, 4-phenyl(perfluoroethyl)benzene **1n**, and TEMPO to monitor the generation and quenching of **PC****.^[24] The quenching rate constant of **PC3**** ($3.2 \times 10^4 \text{ M}^{-1} \text{ s}^{-1}$) was found to be much larger than that of **PC2**** ($0.93 \times 10^2 \text{ M}^{-1} \text{ s}^{-1}$) (Figures S19 and S20).^[25] Thus, the fast regeneration of **PC3** dominates the catalytic turnover (Figure 2A). Next, Gibbs free energy changes (ΔG_r) and reorganization energies (λ) in the 2nd SET were obtained by DFT calculation to estimate activation energies (ΔG^\ddagger) according to Marcus-Hush theory (Figure 2B).^[26,27] While two ΔG_r values are almost identical (3.8 and 3.7 kcal/mol), λ value for **PC3** (38.8 kcal/mol) is lower than that for **PC2** (42.3 kcal/mol). Finally, ΔG^\ddagger for





Figure 2. (A) Quenching rate constants of **PC*** and **PC****. (B) Activation energies (ΔG^\ddagger), Gibbs free energy changes (ΔG_r), reorganization energies (λ), and bent angles (θ) in the 2nd SET by DFT calculation studies ((U)ωB97XD/6-31 + G(d,p)/SMD(acetonitrile)).

PC3 (11.6 kcal/mol) is lower than that for **PC2** (12.6 kcal/mol). This trend in ΔG^\ddagger is consistent with that in the catalyst regeneration rate. Thus, we focused on the geometry change of PC because a smaller λ value leads to the decrease of ΔG^\ddagger for SET. The planar structure of phenothiazine backbone in **PC**** changes to the bent one upon reduction (Figure 2B). This bending is the dominant geometry change and would be deeply related to λ values. We adopted the bent angle (θ) in Figure 2B right to represent the extent of a bent structure of PC. The smaller value of θ in **PC3** ($\theta = 14.5^\circ$) shows the smaller geometry change in the reduction of **PC3**** compared to **PC2** ($\theta = 18.0^\circ$). The steric repulsion of Me groups of **PC3** suppresses bending of a phenothiazine backbone, eventually decreasing ΔG^\ddagger . A catalyst design including both fast catalyst regeneration and effective photo-excited reduction potential is achieved by the rigidity of molecular structure and the introduction of an electron-donating group.


Using the determined optimal reaction conditions, we explored the substrate scope of this aminoxylation (Scheme 2). Electron withdrawing groups such as CN, CO_2Me , and CONMe_2 were available for the transformation (**3b**, **3c**, **3d**). The CF_3 -substituted perfluoroalkylarene selectively underwent the aminoxylation in the perfluoroalkyl group (**3e**).^[28] Silyl and boryl substituents were also tolerated (**3f** and **3g**). It is noted that perfluoroalkylarenes with electron-donating groups smoothly underwent aminoxylation (**3h**, **3i**, and **3j**). The development of **PC3** with high reducing ability overcame the limitation of the substrate scope in our previous report for defluoroallylation.^[4g] Substrates including pyridine, benzofuran, or naphthalene



Scheme 2. Substrate scope of perfluoroalkylarenes in the aminoxylation with TEMPO.^[a] [a] **1a** (0.4 mmol), **2** (0.8 mmol), **PC3** (0.02 mmol), MeCN (2 mL), irradiation with 370 nm LEDs at 35 °C for 24 h. Isolated yields are shown. [b] **2a** (1.2 mmol) and **PC3** (0.04 mmol).

moieties efficiently gave the corresponding products (**3k**, **3l**, and **3m**). The reaction of perfluoroethylarene also gave **3n** in a moderate yield. Perfluoroalkyl-substituted pyridines were feasible substrates, and various functional groups such as OMe, OH, NH₂, and acetal groups were compatible with the present reaction (**3o**, **3p**, **3q**, and **3r**). A quinoline-based substrate afforded desired product **3s** in 68% yield. The substrate with two C₄F₉ groups underwent single aminoxylation to give product **3t**. Reactions of perfluoroalkylphenanthrene and -pyrene gave no products (**3u** and **3v**).^[29]

Next, Lewis acid mediators and nucleophilic coupling partners were surveyed for the selective C–F bond transformation of aminoxylated compounds **3** via an ionic path^[30] (Tables S3 and S4). The combination of AlCl₃ and organosilicon reagents was found to be appropriate in the transformation (Scheme 3). After isolation of **3a**, which was provided by aminoxylation between **1a** and **2** (Table 1, Entry 10), **3a** was treated with allyltrimethylsilane (**4a**) in the presence of AlCl₃. The reaction gave allylated alcohol




Scheme 3. Second defluorinative transformation mediated by a Lewis acid.^[a] [a] **3** (0.2 mmol), **4** (1.0 mmol), and AlCl₃ (0.4 mmol) in CHCl₃ (2 mL) at room temperature for 6 h. Isolated yields are shown. [b] **4h** (1 mL).

5a in 74% yield, in which the amino group was removed on the O atom (see below). The CN and CO₂Me groups were tolerated in this allylation (**5b** and **5c**). Various organosilicon nucleophiles were applicable to this C–F bond transformation. Methallylsilane **4b**, silyl enol ethers **4c** and **4d**, silyl ketene acetal **4e**, and alkynylsilane **4f** provided functionalized perfluoroalkyl alcohols **5d**, **5e**, **5f**, **5g**, and **5h**, respectively. An organotin reagent, methallyltributyltin (**4g**) also acted well as a nucleophile. Toluene (**4h**) was a suitable nucleophile for the Friedel–Crafts reaction to give product **5i**. The reduction with HSiEt₃ smoothly proceeded to yield defluorinated product **5j**. On the other hand, vinylsilane **4j** and silyl ketene imine **4k** were not applicable.


Using the radical and ionic methods to realize two types of C–F bond activation, we demonstrated a one-pot transformation of a CF₂ unit via aminoxylation and allylation reactions (Scheme 4A). After aminoxylation of perfluoroalkylarene **1a** with **2** using **PC3** and 370 nm LED light, the crude product was treated with **4a** and AlCl₃ to give product **5a** in high yield. The one-pot aminoxylation/allylation was also successful using silyl ketene acetal **4e** (Scheme 4B).

Scheme 5 illustrates a proposed mechanism for AlCl₃-mediated C–F bond allylation of **3** with allylsilane **4a**. AlCl₃


A. Aminoxylation/allylation



B. Aminoxylation/alkylation



Scheme 4. One-pot sequential C–F bond transformation of CF₂ unit.^[a]
[a] **1a** (0.4 mmol), **2** (0.8 mmol), **PC3** (0.02 mmol), MeCN (2 mL). Then, **4** (2.0 mmol), AlCl₃ (1.2 mmol), CHCl₃ (4 mL). Yields were determined by ¹H NMR spectroscopy using an internal standard.



Scheme 5. Proposed mechanism for C–F bond transformation mediated by AlCl₃.

abstracts F[−] to give oxonium intermediate **E**. Then *N*-chloroamine and AlFCl₂ are eliminated to give ketone **F**. AlCl₃ activates **F**, and **4a** adds to a carbonyl group, affording **G**.^[31] Finally, hydrolysis of **G** yields product **5**. The generation of **F** was confirmed when **3** was treated with AlCl₃ in the absence of nucleophiles (Scheme S4). Other typical Lewis acids^[30] were not effective (Table S3). AlCl₃ can mediate abstraction of fluoride ion of **3** and activation of a less basic carbonyl group of **F** due to high Lewis acidity. In terms of the intermediacy of ketone **F**, our procedure has an impact on the synthesis of perfluoroalkyl ketones from PFAS via defluorination. Traditional methods such as defluorination of fully-perfluorinated alkylbenzenes,^[32] perfluoroalkyl-substituted anilines^[33] and -enamines,^[34] and α-hydroperfluoroalkanoic acid esters^[5,35] have problems such as narrow substrate scopes. Especially for the synthesis of aryl ketones **F**, available substrates were extremely limited. In this report, compounds **3** can be synthesized and used as synthetic equivalents for **F** with the wide substrate scope and the high compatibility of functional groups. Our process is an efficient synthetic method of functionalized perfluoroalkyl alcohols like **5** from PFAS.

In summary, a combination of photoredox catalysis and Lewis acid activation realizes sequential C–F bond transformation of a CF₂ unit in perfluoroalkylarenes. Functionalized perfluoroalkyl alcohols were synthesized by phenothiazine-catalyzed photo-induced defluoroaminoxylation with TEMPO and subsequent AlCl₃ mediated substitution of a F atom with various carbon nucleophiles. Mechanistic studies revealed that the rigidity of molecular structure and the introduction of an electron-donating group is important in catalyst design to achieve fast catalyst regeneration and effective photo-excited reduction potential.

Acknowledgements

This work was financially supported by JSPS KAKENHI (JP23K17845 (M.Y.), JP23H04906 (M.F.)). This work was supported by the Grant-in-Aid for Transformative Research Areas (A) JP21H05212 (M.Y.) Digitalization-driven Transformative Organic Synthesis (Digi-TOS) from the Ministry of Education, Culture, Sports, Science & Technology, Japan and JST CREST Grant No. JPMJCR20R3 (M.Y.) and JPMJCR18R4 (M.A.), Japan. We thank the members of the Radiation Laboratory of SANKEN, Osaka University, for running the linear accelerator. N.S. acknowledges support from the JSPS Fellowship for Young Scientists (JP22J10775). Y.N. acknowledges support from Research Encouragement Grants of The Asahi Glass Foundation.

Conflict of Interest

The authors declare no conflict of interest.

Data Availability Statement

The data that support the findings of this study are available in the supplementary material of this article.

Keywords: Fluorine • Phenothiazine • Photochemistry • Radical reactions • Radical ions

- [1] a) R. E. Banks, B. E. Smart, J. C. Tatlow, *Organofluorine Chemistry: Principles and Commercial Applications*; Plenum, **1994**; b) P. Jeschke, *ChemBioChem* **2004**, *5*, 570–589; c) K. Uneyama, *Organofluorine Chemistry*; Blackwell Publishing, **2006**; d) K. Müller, C. Faeh, F. Diederich, *Science* **2007**, *317*, 1881–1886; e) S. M. Ametamey, M. Honer, P. A. Schubiger, *Chem. Rev.* **2008**, *108*, 1501–1516; f) S. Purser, P. R. Moore, S. Swallow, V. Gouverneur, *Chem. Soc. Rev.* **2008**, *37*, 320–330; g) P. Kirsch, *Modern Fluoroorganic Chemistry: Synthesis, Reactivity, Applications*, 2nd ed.; Wiley-VCH, **2013**; h) E. P. Gillis, K. J. Eastman, M. D. Hill, D. J. Donnelly, N. A. Meanwell, *J. Med. Chem.* **2015**, *58*, 8315–8359; i) Y. Zhu, J. Han, J. Wang, N. Shibata, M. Sodeoka, V. A. Soloshonok, J. A. S. Coelho, F. D. Toste, *Chem. Rev.* **2018**, *118*, 3887–3964.
- [2] a) C. E. Schaefer, C. Andaya, A. Urtiaga, E. R. McKenzie, C. P. Higgins, *J. Hazard. Mater.* **2015**, *295*, 170–175; b) A. M.

- Trautmann, H. Schell, K. R. Schmidt, K. M. Mangold, A. Tiehm, *Water Sci. Technol.* **2015**, *71*, 1569–1575; c) X. Liang, J. Cheng, C. Yang, S. Yang, *Chem. Eng. J.* **2016**, *298*, 291–299; d) R. K. Singh, S. Fernando, S. F. Baygi, N. Multari, S. M. Thagard, T. M. Holsen, *Environ. Sci. Technol.* **2019**, *53*, 2731–2738; e) M. J. Bentel, Y. Yu, L. Xu, Z. Li, B. Wong, Y. Men, J. Liu, *Environ. Sci. Technol.* **2019**, *53*, 3718–3728; f) B. N. Nzeribe, M. Crimi, S. M. Thagard, T. M. Holsen, *Rev. Environ. Sci. Bio/Technol.* **2019**, *49*, 866–915; g) M. J. Bentel, Z. Liu, Y. Yu, J. Gao, Y. Men, J. Liu, *Environ. Sci. Technol. Lett.* **2020**, *7*, 351–357; h) A. Leeson, T. Thompson, H. F. Stroo, R. H. Anderson, J. Speicher, M. A. Mills, J. Willey, C. Coyle, R. Ghosh, C. Lebrón, C. Patton, *Environ. Toxicol. Chem.* **2021**, *40*, 24–36; i) M. J. Krause, E. Thoma, E. Sahle-Damesessie, B. Crone, A. Whitehill, E. Shields, B. Gullett, *Environ. Eng.* **2022**, *148*, 05021006; j) B. Trang, Y. Li, X.-S. Xue, M. Ateia, K. N. Houk, W. R. Dichtel, *Science* **2022**, *377*, 839–845.
- [3] Recent reviews about C–F bond transformation of perfluoroalkyl compounds: a) W. Xu, Q. Zhang, Q. Shao, C. Xia, M. Wu, *Asian J. Org. Chem.* **2021**, *10*, 2454–2472; b) L. Zhou, *Molecules* **2021**, *26*, 7051; c) S. Li, W. Shu, *Chem. Commun.* **2022**, *58*, 1066–1077; d) Z. Wang, Y. Sun, L.-Y. Shen, W.-C. Yang, F. Mengb, P. Li, *Org. Chem. Front.* **2022**, *9*, 853–873; e) Y. Nishimoto, N. Sugihara, M. Yasuda, *Synthesis* **2022**, *54*, 2765–2777; f) S. Yoshida, *Chem. Rec.* **2023**, *23*, e202200308; g) L. V. Hooker, J. S. Bander, *Angew. Chem. Int. Ed.* **2023**, *62*, e202308880.
- [4] Selected papers about photoredox catalyzed C–F bond transformation: a) K. Chen, N. Berg, R. Gschwind, B. König, *J. Am. Chem. Soc.* **2017**, *139*, 18444–18447; b) H. Wang, N. T. Jui, *J. Am. Chem. Soc.* **2018**, *140*, 163–166; c) C. Luo, J. S. Bandar, *J. Am. Chem. Soc.* **2019**, *141*, 14120–14125; d) D. B. Vogt, C. P. Seath, H. Wang, N. T. Jui, *J. Am. Chem. Soc.* **2019**, *141*, 13203–13211; e) J. B. I. Sap, N. J. W. Straathof, T. Knauber, C. F. Meyer, M. Medebielle, L. Buglioni, C. Genicot, A. A. Trabanco, T. Noel, C. W. Am Ende, V. Gouverneur, *J. Am. Chem. Soc.* **2020**, *142*, 9181–9187; f) Y. C. Luo, F. F. Tong, Y. Zhang, C. Y. He, X. Zhang, *J. Am. Chem. Soc.* **2021**, *143*, 13971–13979; g) N. Sugihara, K. Suzuki, Y. Nishimoto, M. Yasuda, *J. Am. Chem. Soc.* **2021**, *143*, 9308–9313; h) S.-S. Yan, S.-H. Liu, L. Chen, Z.-Y. Bo, K. Jing, T.-Y. Gao, B. Yu, Y. Lan, S.-P. Luo, D.-G. Yu, *Chem* **2021**, *7*, 3099–3113; i) S. Ghosh, Z.-W. Qu, S. Pradhan, A. Ghosh, S. Grimme, I. Chatterjee, *Angew. Chem. Int. Ed.* **2022**, *61*, e202115272; j) C. Liu, N. Shen, R. Shang, *Nat. Commun.* **2022**, *13*, 354–361; k) S. E. Wright, J. S. Bandar, *J. Am. Chem. Soc.* **2022**, *144*, 13032–13038; l) J. H. Ye, P. Bellotti, C. Heusel, F. Glorius, *Angew. Chem. Int. Ed.* **2022**, *61*, e202115456; m) J. Wang, Y. Wang, Y. Liang, L. Zhou, L. Liu, Z. Zhang, *Angew. Chem. Int. Ed.* **2023**, *62*, e202215062; n) W.-J. Yue, R. Martin, *Angew. Chem. Int. Ed.* **2023**, *62*, e202310304.
- [5] S. Thiebaut, C. Gerardin, J. Amos, C. Selve, *J. Fluorine Chem.* **1995**, *73*, 179–184.
- [6] J. Terao, M. Nakamura, N. Kambe, *Chem. Commun.* **2009**, 6011–6013.
- [7] X.-Q. Chu, D. Ge, M.-L. Wang, W. Rao, T.-P. Loh, Z.-L. Shen, *Adv. Synth. Catal.* **2019**, *361*, 4082–4090.
- [8] Y.-J. Yu, F.-L. Zhang, T.-Y. Peng, C.-L. Wang, J. Cheng, C. Chen, K. N. Houk, Y.-F. Wang, *Science* **2021**, *371*, 1232–1240.
- [9] S. Wang, L. Long, X. Zhang, L. Ling, H. Chen, X. Zeng, *Angew. Chem. Int. Ed.* **2023**, *62*, e202312856.
- [10] D. Leifert, A. Studer, *Chem. Rev.* **2023**, *123*, 10302–10380.
- [11] a) E. H. Discekici, N. J. Treat, S. O. Poelma, K. M. Mattson, Z. M. Hudson, Y. Luo, C. J. Hawker, J. R. de Alaniz, *Chem. Commun.* **2015**, *51*, 11705–11708; b) F. Speck, D. Rombach, H.-A. Wagenknecht, *J. Org. Chem.* **2019**, *15*, 52–59; c) F. Weick, N. Hagmeyer, M. Giraud, B. Dietzek-Ivanšić, H.-A. Wagenknecht, *Chem. Eur. J.* **2023**, e202302347.
- [12] C. Liu, K. Li, R. Shang, *ACS Catal.* **2022**, *12*, 4103–4109.
- [13] I. A. MacKenzie, L. Wang, N. P. R. Onuska, O. F. Williams, K. Begam, A. M. Moran, B. D. Dunietz, D. A. Nicewicz, *Nature* **2020**, *580*, 76–81.
- [14] See Supporting Information Tables S1 and S2.
- [15] Z.-H. Wang, P.-S. Gao, X. Wang, J.-Q. Gao, Z.-T. Xu, Z. He, C. Ma, T.-S. Mei, *J. Am. Chem. Soc.* **2021**, *143*, 15599–15605.
- [16] Y. J. Chen, W. H. Deng, J. D. Guo, R. N. Ci, C. Zhou, B. Chen, X. B. Li, X. N. Guo, R. Z. Liao, C. H. Tung, L. Z. Wu, *J. Am. Chem. Soc.* **2022**, *144*, 17261–17268.
- [17] See Supporting Information Figures S35 and S36.
- [18] Excited **PC3*** selectively reduces starting material **1** in prior to product **3** in terms of reduction potential. For example, $E(\mathbf{1a}/\mathbf{1a}^{\bullet-}) = -2.06$ V vs SCE, $E(\mathbf{3a}/\mathbf{3a}^{\bullet-}) = -2.31$ V vs SCE.
- [19] In contrast to **PC2** and **PC3**, **PC1** and **PC4** did not work in the aminoxylation of **1j**, which exhibited lower reduction potential than **1a**, due to the lower reducing abilities of **PC1*** and **PC4*** (Tables S6 and S7). This is because that the 1st SET is slower due to lower reducing abilities of **PC1*** and **PC4*** caused by the lack of N^iPr_2 group. Thus, the catalytic activities of **PC1** and **PC4** are lower than those of **PC2** and **PC3**. The Marcus analysis of **PC1** and **PC4** as well as **PC2** and **PC3** is shown in the Supporting Information.
- [20] Figures S2 and S4 show the Stern–Volmer fluorescence quenching studies of **PC2** and **PC3** with **1e** (see Supporting Information).
- [21] Sub-nano second laser flash photolysis was conducted to determine the lifetimes of singlet states of **PC2** and **PC3**. The singlet lifetime of **PC2** was $^1\tau_0 = 2.4$ ns ($1/^1\tau = k_d = 4.1 \times 10^8$ s $^{-1}$) and that of **PC3** was $^1\tau_0 = 2.5$ ns ($1/^1\tau = k_0 = 4.0 \times 10^8$ s $^{-1}$) (see Supporting Information Figures S22 and S24).
- [22] The diffusion-controlled rate constant of acetonitrile is $k_{diff} = 1.9 \times 10^{10}$ L mol $^{-1}$ s $^{-1}$. M. Montalti, A. Credi, L. Prodi, M. T. Gandolfi, *Handbook of Photochemistry*, 3rd ed.; CRC Press, **2006**.
- [23] The quenching rate constants of excited singlet species of PC (1k_q) with **1e** is much larger than the inter-system-crossing rates of PC (Scheme S1). Thus, the triplet species has a negligible contribution to this reaction. In fact, open air conditions gave the target product in high yield (Scheme S3).
- [24] The absorption of radical anion **A** was not observed due to being out of the monitoring range.
- [25] Figures S19 and S20 show the Stern–Volmer plots for $1/\tau$ of $PC^{\bullet+}$ versus [TEMPO **2**].
- [26] Recently, in several papers, activation energies for single electron transfer processes, in organic reactions were estimated by Marcus-Hush theory. See selected papers: a) O. López-Estrada, H. G. Laguna, C. Barrueta-Flores, C. Amador-Bedolla, *ACS Omega* **2018**, *3*, 2130–2140; b) M.-C. Fu, R. Shang, B. Zhao, B. Wang, Y. Fu, *Science* **2019**, *363*, 1429–1434; c) Tiffany O. Paulisch, F. Strieth-Kalthoff, C. Henkel, Lena Pitzer, D. M. Guldi, F. Glorius, *Chem. Sci.* **2020**, *11*, 731–736; d) K. Matsumoto, M. Nakajima, T. Nemoto, *J. Org. Chem.* **2020**, *85*, 11802–11811; e) S. Sil, A. S. Bhaskaran, S. Chakraborty, B. Singh, R. Kuniyil, S. K. Mandal, *J. Am. Chem. Soc.* **2022**, *144*, 22611–22621; f) C. Ma, J. Shen, C. Qu, T. Shao, S. Cao, Y. Yin, X. Zhao, Z. Jiang, *J. Am. Chem. Soc.* **2023**, *145*, 20141–20148; g) M. Wang, R. Rowshanpour, L. Guan, J. Ruskin, P. M. Nguyen, Y. Wang, Q. A. Zhang, R. Liu, B. Ling, R. Woltornist, A. M. Stephens, A. Prasad, T. Dudding, T. Lectka, C. R. Pitts, *J. Am. Chem. Soc.* **2023**, *145*, 22442–22455.
- [27] See Supporting Information for the derivation of ΔG_r , λ , and ΔG^{\ddagger} .
- [28] DFT calculation study revealed that the alkyl radical intermediate generated via C–F bond mesolysis at perfluoroalkyl group was more stable than one generated via the mesolysis at

- CF₃ group and that the activation energy for perfluoroalkyl group is lower than that for CF₃ group (Scheme S6).
- [29] The elimination of fluoride ion from the corresponding radical anion intermediate would be very slow because a negative charge in the radical anion is effectively delocalized in a large π -conjugation system.
- [30] Selected papers about fluoride abstraction by Lewis acids: a) T. Ooi, D. Uruguchi, N. Kagashima, K. Maruoka, *Tetrahedron Lett.* **1997**, 38, 5679–5682; b) T. Stahl, H. F. T. Klare, M. Oestreich, *ACS Catal.* **2013**, 3, 1578–1587; c) K. Fuchibe, H. Hatta, K. Oh, R. Oki, J. Ichikawa, *Angew. Chem. Int. Ed.* **2017**, 56, 5890–5893; d) G. Meißner, K. Kretschmar, T. Braun, E. Kemnitz, *Angew. Chem. Int. Ed.* **2017**, 56, 16338–16341; e) K. Fuchibe, R. Oki, H. Hatta, J. Ichikawa, *Chem. Eur. J.* **2018**, 24, 17932–17935; f) K. Fuchibe, T. Fushihara, J. Ichikawa, *Org. Lett.* **2020**, 22, 2201–2205; g) F. Wang, Y. Nishimoto, M. Yasuda, *J. Am. Chem. Soc.* **2021**, 143, 20616–20621.
- [31] Selected reviews for Mukaiyama Aldol reaction and Hosomi-Sakurai allylation; P. V. Ramachandran, D. R. Nicponski and P. D. Gagare in *Comprehensive Organic Synthesis (Second Edition)*, Vol. 2 (Ed.: P. Knochel), Elsevier, Amsterdam, **2014**, pp. 72–147; S. Kobayashi, Y. Yamashita, W. J. Yoo, T. Kitanosono, J. F. Soulé in *Comprehensive Organic Synthesis (Second Edition)*, Vol. 2 (Ed.: P. Knochel), Elsevier, Amsterdam, **2014**, pp. 396–450.
- [32] Y. V. Zonov, V. M. Karpov, V. E. Platonov, *J. Fluorine Chem.* **2007**, 128, 1058–1064.
- [33] Q. L. Zhou, Y. Z. Huang, *J. Fluorine Chem.* **1988**, 39, 87–98.
- [34] a) Y. Z. Huang, Q. L. Zhou, *J. Org. Chem.* **1987**, 52, 3552–3558; b) W. Han, Y. L. Chen, X. Tang, J. Zhou, M. Ma, Z. L. Shen, X. Q. Chu, *Green Chem.* **2023**, 25, 9672–9679.
- [35] a) C. Portella, M. Iznaden, *Synthesis* **1991**, 1013–1014; b) M. A. Kurykin, I. M. Vol'pin, L. S. German, *J. Fluorine Chem.* **1996**, 80, 9–12.

Manuscript received: January 16, 2024

Accepted manuscript online: February 21, 2024

Version of record online: ■■, ■■

